

FUV Scattering by Dust in Orion

P. Shalima¹ *, N. V. Sujatha¹ †, Jayant Murthy¹ ‡, Richard Conn Henry² §
and
David J. Sahnou² ¶

¹Indian Institute of Astrophysics, Bangalore 560034, India

²Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, MD 21218

Accepted. Received; in original form

ABSTRACT

We have modelled diffuse far-ultraviolet spectrum observed by *FUSE* near M42 as scattering of starlight from the Trapezium stars by dust in front of the nebula. The dust grains are known to be anomalous in Orion with $R_V = 5.5$ and these are the first measurements of the FUV optical properties of the grains outside of “normal” Milky Way dust. We find an albedo varying from 0.3 ± 0.1 at 912 Å to 0.5 ± 0.2 at 1020 Å which is consistent with theoretical predictions.

Key words: dust, extinction – ultraviolet: ISM.

1 INTRODUCTION

The Orion Nebula (M42) was first observed as one of the brightest diffuse sources in the ultraviolet (UV) sky by Carruthers & Opal (1977). They identified this light to be due to the radiation from the bright Orion stars scattered by dust in the Orion Molecular Cloud. Further observations allowed Wen & O’Dell (1995) to suggest a model for the M42 region in which M42, itself, is a thin blister of ionized gas in front of the Orion Molecular Cloud. The scattering arises in a neutral sheet in front of the Nebula known as the Veil (O’Dell et al. 1992) where there is a deficiency of small grains compared to the diffuse ISM (Baade & Minkowski 1937; Costero & Peimbert 1970; Cardelli & Clayton 1988).

The first observations of far ultraviolet (FUV: 905 – 1187 Å) emission in this region were made by Murthy et al. (2005) who used serendipitous pointings of the *Far Ultraviolet Spectroscopic Explorer (FUSE)* to find intensities as high as 3×10^5 ph cm⁻² s⁻¹ sr⁻¹ Å⁻¹. In this work, we have modelled these observations in order to determine the optical properties of the dust grains in the FUV. We find that the albedo of these grains varies from 0.3 ± 0.1 at 912 Å to 0.5 ± 0.2 at 1020 Å which is consistent with the theoretical predictions of Draine (2003).

Table 1. *FUSE* observations

Data set	l (deg)	b (deg)	Time (s)	Intensity(1100Å) ph cm ⁻² s ⁻¹ sr ⁻¹ Å ⁻¹
S4054601	208.8	-19.3	10565	$2.93 \pm 0.03 (\times 10^5)$
S4054602	208.8	-19.3	5696	$2.95 \pm 0.04 (\times 10^5)$

2 OBSERVATIONS AND MODEL

As part of the *FUSE* S405/505 program, regions of nominally blank sky were observed to allow the instrument to thermalize before realignment of the spectrograph mirrors. Murthy & Sahnou (2004) found diffuse radiation in many of these locations with the brightest being the two in the vicinity of M42 (Murthy et al. 2005). The locations and brightnesses of these two observations are listed in Table 1 and the Digital Sky Survey (DSS) map of M42 and the surrounding region is shown in Fig. 1 with the location of the *FUSE* observation superimposed (square).


The spectrum of the diffuse light is shown in Fig. 2. Although only 1.5' from HD 36981, Murthy et al. (2005) have shown that the observed emission could not be scattered radiation from that star because the broad photospheric Ly β absorption line in the stellar spectrum is not reflected in the diffuse spectrum and suggested that the radiation was instead due to scattering of the light from the Trapezium cluster of stars. Indeed, 65% of the radiation at the scattering location is provided by θ^1 Ori C alone and 99% by the four Trapezium stars in Table 2.

The amount of light scattered by the dust depends on the scattering properties of the interstellar dust grains

* E-mail: shalima@iiaap.res.in
 † E-mail: sujaskm@yahoo.com
 ‡ E-mail: jmurthy@yahoo.com
 § E-mail: henry@jhu.edu
 ¶ E-mail: sahnou@pha.jhu.edu


Table 2. Properties of stars near target

HD No.	Name	l (deg)	b (deg)	angular distance (')	Sp. Type ^a	d^b (pc)	Flux at location (1100 Å) ($\times 10^5$ ph cm ⁻² s ⁻¹ Å ⁻¹)
37022	θ^1 Ori C	209.01	-19.38	12.7	O6pe	450	1500
37041	θ^2 Ori A	209.05	-19.37	14.8	O9.5V	450	370
37020	θ^1 Ori A	209.01	-19.39	12.9	B0.5V	450	150
37023	θ^1 Ori D	209.01	-19.38	12.7	B0.5V _p	450	250

^a obtained from the SIMBAD astronomical database^b Hipparcos catalogue**Figure 1.** DSS map of the region with the brightest stars (asterisks) and the *FUSE* location (filled square) overplotted.

(albedo, cross-section and scattering phase function), their number density and distribution, and the relative geometry of the stars and dust. We have used the Henyey-Greenstein function (Henyey & Greenstein 1941) for the scattering phase function. Although this is a purely empirical function which may not represent the true scattering, particularly for strongly forward scattering grains (Draine 2003), it is the most prevalent in the literature.

We have used the overall morphology of M42 and the surrounding medium described in O’Dell (2001) to derive the scattering geometry (Fig. 3) for our location. Light from the Trapezium stars passes behind the foreground H I sheet (Orion’s Veil) and is scattered by dust in the edges of the H I sheet at point A, about 3 ± 1 pc from θ^1 Ori C. Assuming

**Figure 2.** Observed diffuse spectrum from Murthy et al. (2005). The assumed dust scattering continuum is overplotted as a dark line.

that the Veil is always parallel to the molecular cloud, this distance corresponds to the distance of the Veil (Abel et al. 2004) at point A.

The strong interstellar absorption line at Ly β indicates a total column density of $N(\text{H I}) = (6.3 \pm 0.1) \times 10^{20}$ cm⁻² (using the “H2OOLS” package of McCandliss (2003)) but 21 cm observations (Condon et al. 1998) show a much smaller column density of 4.5×10^{19} cm⁻² ($\tau_{912-1020 \text{ \AA}} \sim 0.029-0.033$) at Point A¹. Most of the absorption therefore arises in the medium between the Trapezium cluster of stars and the scattering location. Interestingly, the column density along the direct line of sight to the Trapezium stars ($N(\text{H}) = 3.9 \times 10^{21}$ cm⁻²; Shuping & Snow (1997)) is much higher than that seen in our scattered light observation indicating that we are actually observing light from θ^1 Ori C reflected around the foreground clouds.

In order to convert the H I column densities into ef-

¹ The 21 cm intensities have been converted to HI column densities using the ratio of van der Werf & Goss (1989) for material in the envelope of the molecular cloud.




Figure 3. Schematic representation of the distribution of dust at the location showing the path (arrow) taken by the observed photons from the Trapezium stars towards the observer. The figure is not to scale.

fective dust densities, we have used the dust cross-sections per hydrogen atom tabulated by Draine (2003) for an $R_V (= A_V/E(B-V))$ of 5.5, characteristic of the interstellar dust in Orion (Cardelli et al. 1989; Fitzpatrick 1999). We note that Draine’s cross-sections assume the standard gas to dust ratio of Bohlin et al. (1978); however, this is 2.06 times higher than the ratio observed in Orion (Shuping & Snow 1997). We have therefore reduced the dust cross-sections per hydrogen atom by this factor.

In summary, we have assumed that the observed emission is due to scattering of the light from the Trapezium stars by a scattering layer of thickness $4.5 \times 10^{19} \text{ cm}^{-2}$ at a distance of approximately 3 pc from θ^1 Ori C. The radiation from the Trapezium stars has been attenuated by a column density of $6.3 \times 10^{20} \text{ cm}^{-2}$ using an extinction curve corresponding to $R_V = 5.5$.

The observed spectrum (Fig. 2) includes many lines, both absorption and emission, of molecular hydrogen (France & McCandliss 2005) as well as the H I Lyman series of absorption lines. In order to deduce the level of the dust scattered emission, we masked out these features and applied a 50 point median filter to the data (dark line in Fig. 2). We then calculated the dust scattered radiation as a function of the optical constants a and g using single scattering (since $\tau < 1$) and compared the intensities with the observations to constrain the values of the parameters. Beyond 1020 \AA the observed radiation is contaminated by fluorescent emission of H_2 because of which we were unable to derive the optical constants.

3 RESULTS AND DISCUSSION

We have plotted our derived albedos as circles in Fig. 4 corresponding to $g = 0.55$. Because we only have observations at a single location, where the scattering angle is about 48° ,




Figure 4. Allowed values of a (circles) corresponding to $g = 0.55$, as a function of wavelength for dust cross-sections reduced by a factor of 2 compared to the $R_V = 5.5$ model. The minimum and maximum values corresponding to $g = 0.4$ and $g = 0.8$ respectively are marked as “x”. The solid line represents the theoretical values of Draine (2003).

it was not possible for us to constrain g and we assumed the value, $g = 0.55 \pm 0.25$ derived by Sujatha et al. (2005) for locations in Ophiuchus. The error bars on the albedos for each value of g , include the observational errors as well as the uncertainties in the exact location of the scattering cloud. Within the assumed range of g , the albedo varies from 0.3 ± 0.1 at 912 \AA to 0.5 ± 0.2 at 1020 \AA . The maximum allowed albedo corresponds to $g=0.8$ while the minimum corresponds to $g=0.4$.

As mentioned above, the dust grains are anomalous in Orion with a depleted population of small grains with respect to the general interstellar medium perhaps due to the destruction and selective acceleration of the small grains by the stellar radiation combined with their coagulation into large grains (Cardelli & Clayton 1988). However, the measured albedo is really a weighted average over all the different dust sizes and hence is only mildly dependent on R_V in the FUV. In fact, we do find that our derived albedo is comparable to other determinations in the same wavelength region (Table 3).

Calzetti et al. (1995) has derived the albedo and g for IC 435 in Orion using data from the International Ultraviolet Explorer (*IUE*). They obtained a high albedo of 0.8 corresponding to $g = 0.75$ at 1200 \AA for dust with an R_V of 5.3. However, as suggested by Burgh et al. (2002), if this is due to the low optical depths used by them, using the correct optical depths would result in low albedos similar to those obtained for other regions despite the difference in R_V .

Table 3. Previous determinations

Location	Wavelength (Å)	Albedo	g
NGC 7023	1000	0.42 ± 0.04	0.75^1
NGC 2023	1100	0.35 ± 0.05	0.85^2
Ophiuchus	1100	0.40 ± 0.10	0.55 ± 0.25^3

¹ Witt et al. (1993)² Burgh et al. (2002)³ Sujatha et al. (2005)

4 CONCLUSIONS

We have modelled the intense diffuse light observed near M42 by Murthy et al. (2005) as starlight from the Trapezium stars scattered by interstellar dust in Orion's Veil, a sheet of neutral hydrogen in front of the Orion Nebula. Most of the absorption seen in the spectrum is due to material between the Trapezium and the scattering location and is much less than the absorption along the direct line of sight to the Trapezium.

If we fix g at 0.55, the albedo of the interstellar grains increases from a value of 0.3 ± 0.02 at 912 \AA to 0.5 ± 0.03 at 1020 \AA close to previously observed values but with a different R_V . On the other hand, if we assume a g of 0.85, as observed by Burgh et al. (2002), the albedo increases from 0.40 ± 0.02 to 0.68 ± 0.03 . We have restricted our analysis up to 1020 \AA since molecular hydrogen fluorescence contaminates the spectrum longward of 1020 \AA . These observations are the first of dust grains with an R_V significantly different from the Galactic norm in the FUV; however we do not find a significant difference in the optical properties.

ACKNOWLEDGMENTS

We thank an anonymous referee for useful comments and suggestions which has helped in improving the quality and presentation of the paper. We thank Kevin France for a careful reading of the manuscript and for useful suggestions. We also thank Dr. Sunetra Giridhar and Dr. P. Manoj for helpful discussions. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France.

REFERENCES

- Abel N. P., Brogan C. L., Ferland G. J., O'Dell C. R., Shaw G., Troland T. H., 2004, *ApJ*, 609, 247
- Baade W., Minkowski R. L., 1937, *ApJ*, 86, 119
- Bohlin R. C., Savage B. D., Drake J. F., 1978, *ApJ*, 224, 132
- Burgh E. B., McCandliss S. R., Feldman P. D., 2002, *ApJ*, 575, 240
- Calzetti D., Bohlin R. C., Gordon K. D., Witt A. N., Bianchi L., 1995, *ApJ*, 446, L97
- Cardelli J. A., Clayton G. C., 1988, *AJ*, 95, 516
- Cardelli J. A., Clayton G. C., Mathis J. S., 1989, *ApJ*, 345, 245
- Carruthers G. R., Opal C. B., 1977, *ApJ*, 217, 95
- Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J., 1998, *AJ*, 115, 1693
- Costero R., Peimbert M., 1970, *BOTT*, 5, 229
- Draine B. T., 2003, *ApJ*, 598, 1017
- Fitzpatrick E. L., 1999, *PASP*, 111, 63
- France K., McCandliss S. R., 2005, *ApJL*, 629, L97
- Henry L. C., Greenstein J. L., 1941, *ApJ*, 93, 70
- McCandliss S. R., 2003, *PASP*, 115, 651
- Murthy J., Sahnou D. J., 2004, *ApJ*, 615, 315
- Murthy J., Sahnou D. J., Henry R. C., 2005, *ApJ*, 618, 99
- O'Dell C. R., 2001, *ARA&A*, 39, 99
- O'Dell C. R., Walter D. K., Dufour R. J., 1992, *ApJ*, 399, L67
- Shuping R. Y., Snow T. P., 1997, *ApJ*, 480, 272
- Sujatha N. V., Shalima P., Murthy J., Henry R. C., 2005, *ApJ*, 633, 257
- van der Werf P. P., Goss W. M., 1989, *A&A*, 224, 209
- Wen Z., O'Dell C. R., 1995, *ApJ*, 438, 784
- Witt A. N., Petersohn J. K., Holberg J. B., Murthy J., Dring A., Henry R. C., 1993, *ApJ*, 410, 714