

# INTER STELLAR SHOCKED REGION CHEMISTRY CERTAIN REACTIONS BETWEEN INTERSTELLAR MOLECULES AND O AND N ATOMS

K.K.Ghosh, S.S.Saleem and K.N. Kutty  
Kavalur Observatory, Alangayam 635 701, India

**ABSTRACT.** The reactions between O and N atoms and certain interstellar molecules ( $\text{H}_2\text{S}$ ,  $\text{NH}_3$ ,  $\text{H}_2\text{CO}$ ,  $\text{CH}_3\text{OH}$ ,  $\text{C}_2\text{H}_5\text{OH}$ ) are studied using the dynamical flowing after glow method. It is found that NH and OH radicals are formed which can be used as a diagnostic probe to differentiate between interstellar shocked and heated regions.

## 1. INTRODUCTION



The propagating shock waves, generated by interstellar winds, expanding HII regions, and supernovae remnants will lead to the rapid collision-induced dissociation of heavy homonuclear molecules like  $\text{N}_2$  and  $\text{O}_2$  in shocked regions whereas the dissociation occurs slowly in heated region (1). These active dissociated atoms will react with ambient molecules and the resulting chemical species may be used as diagnostic probes to differentiate between interstellar shocked and heated regions.

## 2. EXPERIMENTAL

We have carried out certain experiments between  $\text{O}(^1\text{D})$  and  $\text{N}(^2\text{P})$  atoms and certain interstellar molecules ( $\text{H}_2\text{S}$ ,  $\text{NH}_3$ ,  $\text{H}_2\text{CO}$ ,  $\text{CH}_3\text{OH}$ ,  $\text{C}_2\text{H}_5\text{OH}$ ). The experimental set up and procedures in detail is described elsewhere (2,3)

## 3. RESULTS AND DISCUSSION

The wavelengths of observed spectra which are mainly due to NH and OH radicals are presented in Table I. The number densities of excited species were obtained using the method of De Vos (4) and Larrabee (5). The chemical reaction between  $\text{O}(^1\text{D})$ ,  $\text{N}(^2\text{P})$  and the reactants may be written as:



where  $\text{R} = \text{H}_2\text{S}, \text{NH}_3, \text{H}_2\text{CO}, \text{CH}_3\text{OH}, \text{C}_2\text{H}_5\text{OH}$

The rate coefficients for the above reactions (1), (2) are of the order of  $2 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$  and  $8 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$  respectively. From energy considerations it can be shown that  $\text{O}(^1\text{D})$  and  $\text{N}(^2\text{P})$  can produce  $\text{OH}(A^2\Sigma^+)$  and  $\text{NH}(d^1\Sigma)$  radicals whereas the ground state of  $\text{O}(^3\text{P})$  and  $\text{N}(^4\text{S})$  can produce  $\text{OH}(X^2\Pi)$  and  $\text{NH}(^3\Sigma^-)$  radicals only.

From the experiment it is found that NH and OH will be formed other than CN and CO radicals when dissociated N and O atoms react with the reactants like  $\text{H}_2\text{S}$ ,  $\text{NH}_3$ ,  $\text{H}_2\text{CO}$ ,  $\text{CH}_3\text{OH}$ ,  $\text{C}_2\text{H}_5\text{OH}$ . Thus the abundances of NH and OH will be enhanced in the shocked region.

The enhancement of the abundances of NH and OH radicals may be used as a diagnostic probe to differentiate between the interstellar shocked and heated regions.

TABLE I. NH and OH band systems of glow

NH band system					
Observed Wavelength	Observed Intensity	Standard Wavelength	Standard Intensity	Transition ( $d^1\Sigma \rightarrow c^1\Pi$ )	
2516.0	Ms	2516.3	5	( $v'=1$ )	( $v''=1$ )
2557.0	S	2557.3	10	( $v'=0$ )	( $v''=0$ )
2683.0	Ms	2683.4	5	( $v'=0$ )	( $v''=1$ )
OH band system					
3064.8	Vs	3063.6	10	( $v'=0$ )	( $v''=0$ )
3122.4	W	3121.7	-	( $v'=1$ )	( $v''=1$ )
3185.7	W	3184.8	-	( $v'=2$ )	( $v''=2$ )
3428.8	W	3428.1	-	( $v'=0$ )	( $v''=1$ )

Vs, Very strong; S, Strong; Ms Medium Strong; W, Weak.

#### References

1. Dalgarno, A., and Roberge, W.G. 1979, Ap.J.(Letters), 233, L25.
2. Ghosh, S.N., Jain, S.K., and Gupta, S.K. 1962 Proc.Phys.Soc. (London) 79, 207.
3. Ghosh, K.K. 1984, Physica 124C, 395.
4. De Vos, J.C. 1954, Physica 20, 690.
5. Larrabee, R.D. 1959, J.Opt.Soc.Am. 49, 619.