

## INTERPRETATION OF THE "THIRD HARMONIC" OF THE SOLAR MAGNETIC CYCLE

M.H.GOKHALE and J.JAVARAIAH  
 Indian Institute of Astrophysics  
 Bangalore 560 034  
 India

**ABSTRACT.** The peak at  $\sim 3/22 \text{ y}^{-1}$  seen for all odd-degree axisymmetric modes in the SHF analysis of 'nominal magnetic field' seems to be an artifact of some non-linearity between the field magnitude and the sunspot occurrence probability.

### 1. Introduction

We have an on-going project of analysing in various ways the distribution of the sunspot occurrence probability (and a 'nominal' magnetic field defined therefrom), on the sun's surface and in time, using Greenwich data for sunspot groups. We have shown earlier that the spherical harmonic Fourier (SHF) analysis of these quantities yields approximately constant amplitudes and constant phases for their odd-degree axisymmetric SHF components during 1874-1976 (over which the data is available on a magnetic tape provided by H.Balthasar). Detailed results of the SHF analysis are discussed in earlier papers (Gokhale, Javaraiah and Hiremath, 1989, Gokhale and Javaraiah 1990).

In this paper we investigate the behaviour and significance of a third harmonic peak at  $\sim 3/22 \text{ y}^{-1}$  which is quite significantly present in the spectra of all the odd degree axisymmetric modes with respect to the temporal frequency. These peaks are unlikely to be caused by the asymmetry in the 11 year cycle of the data size, since the magnitude of the nominal field is normalised with respect to the data size in each interval of analysis. Similar periodicity is seen in the SHF analysis of magnetogram data (Stenflo and Vogel 1986) and the hydrogen-alpha spectroheliogram data 1917-1985 (Gokhale, et al 1990). It is also seen in the Fourier analysis of solar rotation (Singh and Prabhu, 1985). It is therefore important to see if these 'third harmonic' peaks represent independent periodicity in the nominal magnetic field. For this purpose we have determined the correlation between the amplitudes and phases of the third harmonic components with those of the corresponding first harmonic components. The presence of high correlation between the amplitudes (and a stronger correlation between the phases) of the first and the third harmonics show that the third harmonic peaks do not represent any independent periodicity. It seems they result from some residual nonlinearity between the nominal field and the data size.

## 2. Data and Method of Analysis

The data used consists of the heliographic latitudes  $b_i$ , longitudes  $\phi_i$  and times of observation  $t_i$  of all the sunspot groups observed during 1874-1976 which are included in the ledgers I and II of the Greenwich Photoheliographic Results. The  $t_i$  are converted to values  $\tau_i$  of the time measured from the zero hour of 1 January 1874. The sunspot occurrence probability  $p(\theta, \phi, t)$  during any sufficiently long time interval  $(T_1, T_2)$  is defined as

$$p(\theta, \phi, t) = \frac{1}{N(T_1, T_2)} \delta(\mu - \mu_i, \phi - \phi_i, \tau - \tau_i)$$

in the neighbourhood of the points  $(\mu_i, \phi_i, \tau_i)$  in the  $(\mu, \phi, t)$  space, and

$$p(\theta, \phi, t) = 0 \quad \text{elsewhere}$$

where  $\mu = \cos\theta$ ,  $\tau = (t - T_1)/(T_2 - T_1)$ ,  $\delta$  represents a delta function,  $N(T_1, T_2)$  is the number of data points given by sunspot groups during in the time interval  $(T_1, T_2)$ . It may be noted that this way the data of each sunspot group gets weighted by the number of days it is observed. The 'nominal magnetic field' 'B'  $(\theta, \phi, t)$  is defined as

$$'B'(\theta, \phi, t) = \pm p(\theta, \phi, t)$$


where the signs  $\pm$  are chosen according to Hale's laws of magnetic polarities, taking care to separate the data from the old and the new cycles during the overlaps of successive cycles.

## 3. Presence of the "Third Harmonics" with constant amplitudes and phases


In an earlier paper (Gokhale and Javaraiah, 1990) we have plotted the SHF amplitudes of 'B'  $(\theta, \phi, t)$  during the interval 1902-1954, as functions of temporal frequency  $\nu$ , for axisymmetric ( $m=0$ ) modes of odd degrees of  $\ell=23$ . All these plots show a very high 'primary' peak at  $\nu = 1/22y^{-1}$  as expected. But all of them also show a smaller "third harmonic" peak at  $\sim 3/22 y^{-1}$ , which, in the case of modes  $\ell > 9$ , is comparable to the primary peak. For  $\ell = 13, 15$  it is as high as the primary peak. Similar concentration of SHF power at the third harmonic is also seen in the results obtained by Stenflo and Vogel (1986) from the magnetogram data and by Gokhale et al (1990) from the hydrogen-alpha spectroheliogram data 1915-1985. The '7 year' periodicity is present even in the analysis of solar rotation using CaII K spectroheliograms (Singh and Prabhu 1985). We find that for  $\ell \geq 13$  the amplitudes and the phases of the third harmonic mode are approximately constant during all the 82 intervals of 22 y length, from 1874-1895 to 1955-1976, each of which is shifted with respect to the previous intervals by one year (Fig.1).

## 4. Interpretation

It may be thought that the 'third harmonic' peaks result from the asymmetry of the 11 year half cycles of the data size. However this is not



**Figure 1** (a) Amplitude ratio  $A_3/A_1$  and (b) the phase difference  $\delta_3 - \delta_1$  as functions of time represented by the interval number. Symbols:  $\ell=1$ : filled circle;  $\ell=3$ : open circle,.....  $\ell=21$ : letter 'phi'.



**Figure. 2** (a) Amplitude  $A_3$  of the third harmonic versus amplitude  $A_1$  of the first harmonic, and (b) phase  $\delta_3$  of the third harmonic versus phase  $\delta_1$  of the first harmonic, for the mode  $\ell=1, m=0$ .

true, since the nominal field 'B' is normalised to the size of the data as shown by the factor  $1/N$  ( $T_1, T_2$ ) in equation (1). In order to see if the third harmonic is an independent SHF mode in the time dependence of 'B' we plotted for each odd degree  $\ell$ , the amplitudes  $A_3$  (and the 'initial' phases  $\delta_3$ ) of the third harmonic mode against the amplitude  $A_1$  (initial phase  $\delta_1$ ) of the first harmonic mode. We find, that for each odd degree upto  $\ell = 13$  (upto which we have done our computations), there is a strong correlation between the amplitudes (and the initial phases) of the third and the first harmonics. In figures 2(a) and 2(b) we illustrate the correlations  $A_3$  versus  $A_1$  and  $\delta_3$  versus  $\delta_1$  for the mode ( $\ell=1, m=0$ ).

For all the odd modes upto  $\ell = 13$ , the correlation coefficients are  $\sim 0.49$  between  $A_1$  and  $A_3$  and  $\sim 0.85$  to  $0.95$  between  $\delta_1$  and  $\delta_3$ .

This shows that the third harmonic peaks in the SHF power are unlikely to represent an independent set of modes. They result most probably from some non-linearity between the magnitude of the analyzed function 'B' ( $\theta, \phi, t$ ) and the true sunspot occurrence probability.

### References

- Gokhale M.H., Javaraiah, J. and Hiremath, K.M., 1990, "Study of sun's 'hydro-magnetic' oscillations using sunspot data" in Stenflo, J.O. (ed) "Solar Photosphere: Structure, Convection and Magnetic Fields", IAU Symp.138, (in press).
- Gokhale M.H. and Javaraiah, J., 1990 "Sunspot activity as originating in interference of sun's global MHD oscillations", Mon.Not.Roy.Astron.Soc. (in press).
- Gokhale, M.H., Makarov, V.I., Sivaraman, K.R., Javaraiah, J., Tavastsherna, K.S., Vasileva, V.V., 1990 "Global modes of solar magnetic fields derived from H-alpha synoptic charts", (in preparation).
- Singh, J. and Prabhu, T.P., 1985 "Variations of the solar rotation rate derived from Ca+K plage areas", Solar Phys. **97**, 302.
- Stenflo, J.O. and Vogel, M 1986 "Global resonances in the evolution of solar magnetic fields" Nature, **319**, 285.

### DISCUSSION

DALSGAARD: The presence of the third harmonic may simply reflect the non-sinusoidal shape of the sunspot index as function of time. This is supported by the constancy of the amplitude ratios and phase difference. The situation is equivalent to Fourier analysis of Cepheid light curves.

**GOKHALE:** Certainly one does expect a third harmonic due to the non-sinusoidal nature of the sunspot cycle, or due to a non-linear dependence of the 'data' variable (P) on a basic variable such as the magnetic field (B). That is why we have tested the correlation of the amplitudes and phases of the third harmonic with respect to those of the first harmonic. However, we have *normalized* the sunspot probability (p) to the *total* number of data points in each 22-year sample, thereby filtering out the time-dependence of the data size (if p depends linearly on B). Since we still see the third harmonic, with different amplitudes for different odd  $\ell$  ( $\leq 9$ ), but the amplitude and phase for each  $\ell$  correlates with those of the first harmonic for the same  $\ell$ , we conclude that there must be some non-linearity between p and B. As for the independence of the third and first harmonics for  $\ell > 9$ , further analysis will be needed.