

**DIURNAL VARIATION OF  $f_0F_2$  AT EQUATORIAL LATITUDES AND COUNTER ELECTROJET**

ONE well documented feature of the diurnal variation of  $f_0F_2$  at equatorial latitudes is the occurrence of two maxima, one in the morning and the other in the evening with a prominent trough around midday, termed as noon 'bite-out'<sup>1-5</sup>. The asymmetry of the bite-out, *i.e.*, the relative amplitudes of the morning and evening maxima exhibit a significant dependence on the phase of the sunspot cycle. During the periods of low sunspot activity, the amplitude of the evening maximum is prominent while during periods of high sunspot activity the morning maximum gains prominence<sup>6,7</sup>. The physical mechanisms responsible for this long term behaviour of the asymmetry of the bite-out are yet to be established.

It is well known that the intense band of *eastward* ionospheric currents at E-region altitudes, over and in the vicinity of the dip equator, referred to as electrojet, gives rise to the pronounced enhancement in the diurnal range of the H-component of the earth's geomagnetic field monitored on ground. The electric field (east-west) at E-region levels associated with the electrojet is transferred along the highly conducting magnetic lines of force to higher altitudes where it interacts with the horizontal geomagnetic field to set up an upward directed (during daytime)  $E \times B$  drift of plasma. Thus the electrojet current strength, on which the  $E \times B$  drift depends, governs to a large extent the behaviour of the equatorial F-region<sup>8-11</sup>. It is well established now that on occasions, designated as 'Counterelectrojet' (CEJ) events<sup>12</sup>, the electrojet current gets reversed for a while during daytime, *i.e.*, current is westward, resulting in a depression of the H-component below the mean midnight level<sup>13</sup>. These CEJ events are closely associated with several phenomena in the lower and upper equatorial ionosphere, the most prominent among them being the disappearance of  $E_{sq}$  traces on bottomside ionograms and reversal of the E-region horizontal drift velocities, clearly indicating a reversal in the electrojet direction<sup>14-16</sup>. The probable effect of the CEJ events is to reverse or reduce the  $E \times B$  vertical drift (due to a reversal of the E-region electric field) and hence to increase the F-region electron densities at and close to the dip equator<sup>11</sup>. The occurrence of CEJ events is noticed to be more frequent during afternoon hours under conditions of low geomagnetic and solar activity<sup>17</sup>. It is thus quite possible that the pronounced evening maximum in the diurnal variation of  $f_0F_2$  at equatorial latitudes, observable during low sunspot activity, might be due to the frequent occurrence of afternoon CEJ events. In fact, such an explanation has already been put forward<sup>18</sup> although systematic studies have not been made. It is therefore felt worthwhile to examine this possibility.

The present study is based on the ionogram and magnetogram data at Kodaikanal (77.5° E, 10.2° N, dip 3.5°), situated well within the electrojet region, for the period January 1963 to December 1964 corresponding to low sunspot activity (mean sunspot number = 19). The analysis is mainly confined to geomagnetically quiet days (AP index < 6) as the occurrence of CEJ is more frequent during quiet periods. The objective is to study in comparison the mean diurnal variation of  $f_0F_2$  on normal quiet days and CEJ days. If the frequent occurrence of afternoon CEJ events is indeed responsible for the prominent evening maximum in  $f_0F_2$ , then one would expect a significant increase in  $f_0F_2$  values during CEJ days compared to normal quiet days. Careful examination of the quarter-hourly ionogram and magnetogram data revealed that out of the 310 quiet days that occurred during the 24 month period considered, CEJ manifested on 73 days. Following current practice<sup>19</sup>, the presence of CEJ is inferred if, at any particular time, there is a negative fluctuation in the value of  $\Delta H$  at Kodaikanal (within the electrojet) minus  $\Delta H$  at Alibag (outside the electrojet), the reference value being the mean midnight level, coincident with disappearance of  $E_{sp}$  on ionograms. Occasions wherein there is only an obvious reduction in the value of  $H$  at Kodaikanal but not large enough to give a negative value of  $\Delta H$  at Kodaikanal minus  $\Delta H$  at Alibag, but associated with the disappearance of  $E_{sp}$ , referred to as 'partial counter-electrojet' events<sup>20</sup>, have also been taken into consideration.

In Fig. 1 is shown the mean diurnal variation of  $f_0F_2$  corresponding to the 237 normal quiet days (solid curve). The presence of noon 'bite-out' with a prominent evening peak is quite evident. In Fig. 1




FIG. 1. Mean diurnal variation of  $f_0F_2$  at Kodaikanal corresponding to normal quiet days (solid curve) and CEJ days (solid circles) during the period January 1963 to December 1964. The vertical bars represent the standard deviations of the mean values of  $f_0F_2$  for the CEJ days.

is also shown the mean diurnal variation of  $f_0F_2$  corresponding to the 73 CEJ days. A perusal of Fig. 1 shows that there is an apparent increase in  $f_0F_2$  values in the evening period during CEJ days compared to normal quiet days. The difference in  $f_0F_2$  values is however not statistically significant as may be seen from the standard deviations of the mean values. It has been reported earlier that increases in  $f_0F_2$  occur during period of CEJ<sup>11,21</sup>. Examination of  $f_0F_2$  behaviour during selected days of strong CEJ by the authors also showed an increase in  $f_0F_2$  values during the evening period but only on some of the days. It is thus quite evident that the prominent evening peak in the diurnal variation of  $f_0F_2$  at equatorial latitudes during low sunspot activity conditions is not entirely due to the frequent occurrence of afternoon CEJ events. Further work is therefore required to assess the other physical mechanisms involved.

Indian Institute of Astrophysics,  
Kodaikanal 624 103, May 17, 1978.

J. HANUMATH SASTRI.  
K. SASIDHARAN.

- Berkner, L. V. and Wells, H. W., *Terr. Magn. Atmos. Elect.*, 1934, **39**, 215.
- Appleton, E. V., *Nature*, 1946, **157**, 691.
- Bailey, D. K., *Terr. Magn. Atmos. Elect.*, 1948, **53**, 35.
- Maeda, H., *Rept. Ionospheric Res. Japan*, 1955, **9**, 60.
- Rastogi, R. G., *J. Geophys. Res.*, 1959, **64**, 727.
- Rao, B. C. N., *Ibid.*, 1963, **68**, 2541.
- Olatunji, E. O., *Ann. Geophys.*, 1966, **22**, 393.
- Dunford, E., *J. atmos. terr. Phys.*, 1970, **32**, 421.
- Mac Dougall, J. N., *Radio Sci.*, 1969, **4**, 805.
- Rastogi, R. G. and Rajaram, G., *Indian J. Pure appl. Phys.*, 1971, **9**, 531.
- Rajaram, G. and Rastogi, R. G., *Ind. J. Radio-space Phys.*, 1974, **3**, 323.
- Gouin, P. and Mayaud, P. N., *Ann. Geophys.*, 1967, **23**, 41.
- Onwumechilli, A. and Akasof, S. I., *J. Geomag. Geoelect.*, 1972, **24**, 161.
- Hutton, R. and Oyinloye, J. D., *Ann. Geophys.*, 1970, **26**, 921.
- Rastogi, R. G., Chandra, H. and Chakravarty, S. C., *Proc. Ind. Acad. Sci.*, 1971, **74**, 62.
- *Ibid.*, 1975, **81**, 80.
- Chandra, H., *J. Inst. Telecom. Engrs., New Delhi*, 1973, **19**, 539.
- , Rajaram, G. and Rastogi, R. G., *Ind. J. Radio-space Phys.*, 1973, **2**, 243.
- Fambitakoye, O., Rastogi, R. G., Tabbagh, J. and Vila, P., *J. atmos. terr. Phys.*, 1973, **35**, 1119.
- Rastogi, R. G., *J. geophys. Res.*, 1974, **79**, 1503.
- Bhargava, B. N. and Subrahmanyam, R. V., *Proc. Ind. Acad. Sci.*, 1964, **60**, 271.