

EVIDENCE OF V-BAND POLARIMETRIC SEPARATION OF CARBON STARS AT HIGH GALACTIC LATITUDE

ARUNA GOSWAMI¹, SREEJA S. KARTHA¹, AND ASOKE K. SEN²

¹ Indian Institute of Astrophysics, Bangalore 560034, India; aruna@iiap.res.in

² Department of Physics, Assam University, Silchar, Assam, India

Received 2010 July 19; accepted 2010 September 2; published 2010 September 22

ABSTRACT

Polarization is an important indicator of stellar evolution, especially for stars evolving from red giant stage to planetary nebulae. However, not much is known about the polarimetric properties of the carbon-enhanced metal-poor (CEMP) stars, although they have been well studied in terms of photometric as well as low- and high-resolution spectroscopy. We report here first-ever estimates of *V*-band polarimetry of a group of CEMP stars. *V*-band polarimetry was planned, as the *V* band is known to show maximum polarization among *BVRI* polarimetry for any scattering of light caused due to dust. Based on these estimates the program stars show a distinct classification into two: one with $p\% < 0.4$ and the other with $p\% > 1$. Stars with circumstellar material exhibit a certain amount of polarization that may be caused by scattering of starlight due to circumstellar dust distribution into non-spherically symmetric envelopes. The degree of polarization increases with asymmetries present in the geometry of the circumstellar dust distribution. Our results reflect upon these properties. While the sample size is relatively small, the polarimetric separation of the two groups ($p\% < 0.4$ and $p\% > 1$) is very distinct; this finding, therefore, opens up an avenue of exploration with regard to CEMP stars.

Key words: stars: carbon – stars: chemically peculiar – stars: late-type – stars: low-mass

1. INTRODUCTION

A characteristic property of stars evolving from red giant stage to planetary nebulae (PNe) is polarization; observed polarization reaches a maximum during the proto-planetary stage and gradually decreases as the star evolves into a PN (Johnson & Jones 1991). Polarimetric studies of post-asymptotic giant branch (AGB) stars by Trammell et al. (1994) and Parthasarathy & Jain (1993) have helped to identify aspherical morphology shells that originate during mass-loss phase as common features of these objects. Quasi-periodic variations in linear polarization are known to be observed in many late-type variables. However, not much work has been done on polarization of carbon stars; we report here *V*-band polarimetric measurements of a sample of carbon stars at high Galactic latitude. Polarization of carbon stars are known to show a flatter wavelength dependence than the oxygen-rich stars. Stars with circumstellar materials exhibit a certain amount of linear polarization. The degree of polarization increases with asymmetries present in the geometry of the shells. By modeling the observed polarization, it is possible to study the geometry and structure of the shells. Polarization models of circumstellar grain scattering shows that graphites do not adequately explain the mean polarimetric behavior of carbon stars (Raveendran 1991). Rowan-Robinson & Harris (1983) have shown that graphites also do not adequately explain the IR spectra of circumstellar dust envelopes around carbon-rich objects. It was suggested by Czyzak et al. (1982) that graphite formation in circumstellar envelopes is not a very likely process and its likely that carbon grains exist most likely as amorphous carbon.

While Parthasarathy et al. (2005) find no significant variation in polarization for HD 100746 at *U*, *B*, *I* bands, etc., Raveendran (1991) finds mild wavelength dependence of polarization in the case of a few carbon stars of Mira variables. In particular, the polarization was found to increase systematically toward the red in RT Pup, UU Aur shows a dip in the polarization curve around the *V* band, and X Vel shows a larger polarization in the *R* band than in the *V* band. Multiband

polarimetric observations by Serkowski et al. (1975) showed that the interstellar polarization has a peak occurring at a median wavelength $\lambda = 0.545 \mu\text{m}$. The maximum effect of interstellar polarization can be felt, therefore, in the yellow spectral region. It is important therefore to undertake follow-up observations on several occasions to study the variations in polarization particularly for those that show significant polarization in the *V* band. Our present sample includes faint high latitude carbon stars from Christlieb et al. (2001). We have studied a large fraction of this sample based on low- as well as high-resolution spectroscopy (Goswami 2005; Goswami et al. 2006, 2007, 2010b; Goswami & Aoki 2010a). A primary characteristics of these stars is that they provide evidence of nucleosynthesis of the low- and intermediate-mass stars as most of them are polluted by now-extinct AGB stars. The basic parameters of the program stars are listed in Table 1.

2. OBSERVATIONS AND DATA REDUCTION

Observations were made using the 2 m telescope of IUCAA, Pune using the polarimeter (IMPOL) attached to it. The polarimeter available at the cassegrain focus of the 2 m IUCAA telescope has a rotating half-wave plate (HWP) and Wollaston prism through which light passes before forming a pair of images of an object on the CCD (Ramprakash et al. 1998). The HWP can rotate in several discrete steps, such that its fast axis makes angles (α) with some reference direction (generally celestial north–south). The light which is transmitted out of the Wollaston prism forms two images of any celestial source on the CCD, with the ordinary and extraordinary set of rays. For each object observations were taken at four different positions of the HWP: 0° , 22.5° , 45° , and 67.5° .

Among *BVRI* polarimetry, the *V* band is known to show the maximum polarization (Serkowski et al. 1975) and is ideal for the detection of polarization caused due to scattering by dust. Moreover, because of a low photon flux in the *B* band, the polarization measurements in the *B* band often have large errors. For these reasons, we have conducted only *V*-band polarimetry

Table 1
Basic Parameters of the Program Stars

Stars	R.A.(2000)	Decl.(2000)	l	b	V	J	H	K_s	Date of Obs.
HE 1027–2501	10 29 29.5	–25 17 16.2	266.6829	+27.4161	12.70	10.627	9.896	9.722	2009 Apr 30
HD 94851*	10 56 44.2	–20 39 52.6	269.8196	+34.7281	9.27	8.869	8.816	8.771	2009 Apr 30
HD 100764	11 35 42.7	–14 35 36.6	276.8586	+44.4107	8.73	7.048	6.60	6.513	2009 Apr 30
HE 1152–0355	11 55 06.1	–04 12 24.0	277.3246	+55.8385	11.43	9.339	8.665	8.429	2009 Apr 30
HE 1305+0007	13 08 03.8	–00 08 47.4	311.9430	+62.4340	12.22	10.247	9.753	9.600	2009 Apr 30
HE 1429–0551	14 32 31.3	–06 05 00.2	343.0186	+48.7605	12.60	10.734	10.272	10.066	2009 Apr 30
HE 1523–1155	15 26 41.0	–12 05 42.6	351.8699	+35.6301	13.22	11.372	10.846	10.748	2009 Apr 30
HD 147084*	16 20 38.1	–24 10 09.5	352.3279	+18.0503	4.55	2.222	1.899	1.681	2009 Apr 30
LP 625–44	16 43 14.0	–01 55 30.2	015.1463	+27.1749	11.85	10.432	10.058	10.825	2009 Apr 30
HD 160529*	17 41 59.0	–33 30 13.7	355.7021	–01.7322	6.77	3.547	3.056	2.790	2009 Apr 30
HD 168986	18 23 10.1	–15 36 36.5	015.8506	–00.9879	9.21	6.901	6.379	6.184	2009 Apr 30
HD 196944	20 40 46.0	–06 47 50.6	039.5635	–27.3703	8.41	7.017	6.626	7.504	2009 Apr 30
HD 198269	20 48 36.7	+17 50 23.7	063.2273	–15.9516	8.12	6.078	5.505	5.385	2009 Apr 30
HD 206983	21 46 09.3	–15 14 39.7	038.6462	–45.4725	9.45	6.934	6.247	6.035	2009 Apr 28
HD 209621	22 04 25.1	+21 03 08.9	078.6453	–27.1175	8.86	6.661	6.045	5.913	2009 Apr 28

Note. The stars marked with * are polarization standard stars.

Table 2
V-band Polarimetry of Polarization Standard Stars

Star Names (1)	HJD (2)	$p\%$ (3) ^a	θ° (4) ^a	$p\%$ (5)	θ° (6)	Remarks (7)
HD 94851	2454952.13280	0.11 ± 0.12		0.057 ± 0.018 0.134 ≤ 0.1		Turnshek et al. (1990) Rautela et al. (2004) Leonard et al. (2005)
HD 147084	2454952.25726	4.12 ± 0.02	32.2	4.16 ± 0.01 4.17 ± 0.008 4.19 ± 0.03 4.23 4.21	32.1 ± 1.9 32.9 32.4 ± 0.9 33.5 31.3	McDavid (1999) Turnshek et al. (1990) Chavero et al. (2006) Clarke et al. (1998) Weitenbeck (2004)
HD 160529	2454952.46513	7.45 ± 0.02	20.9	7.35 ± 0.55 7.3 ± 0.54 7.52	20 20.1	Reiz & Franco (1998) Serkowski (1974) Clarke et al. (1998)

Notes. Estimates listed in Columns 5 and 6 are from references listed in Column 7.

^a Our estimates of $p\%$ and offset corrected position angles (magnitudes).

for the program stars. In addition to the program stars we have observed three polarization standard stars and one unpolarized standard star for polarimetric calibration.

Data reduction was carried out using various tasks in IRAF. The task PHOT in APPHOT was used to measure the stellar flux. Zero polarization standard stars were observed to check for any possible instrumental error, which proved to be smaller than 0.1%. For stars which show low degree of polarization, measurement accuracy is an important issue. The measured errors are smaller than 0.1% for majority of the observed stars.

3. RESULTS AND DISCUSSIONS

V-band polarimetric measurements for the standard polarized and unpolarized stars are presented in Table 2. Estimates from literature are also listed for a comparison. A close agreement between the estimated and literature values lends support to the reliability of our results. The estimated error in polarization is less than or near about 0.1% for all stars (Table 3) except for the star HD 209621 for which the error is ± 0.23 . Since polarimetric error in our case is photon noise dominated, in the case of very bright stars this error becomes close to zero. Our program stars include objects brighter as well as fainter than HD 209621.

Thus, the error in HD 209621 can be considered to be a typical error value.

The polarization standard stars HD 147084 and HD 160529, both observed on the same night, show an offset of about $\sim -10^\circ$; the near equality of the offsets is as expected for consistency, and a similar offset was also reported earlier by Sen et al. (2000). The offset is essentially a result of constraints in instrumental mounting, such as constraint in mounting the polarimeter with the axis of fixed HWP making an angle 0° with celestial north–south axis (this will give 0 offset). We have applied an offset of -10° to the estimated position angles of all the stars observed on the same night as that for HD 147084 and HD 160529. The expected final values of position angles (θ°), obtained with the necessary offset corrections are listed in Table 3.

The linear polarization ($p\%$), with the errors in the measurements of the program stars, is presented (Table 3) without any corrections applied to the observed values for interstellar polarization. For stars with zero polarization ($\leq 0.1\%$) the position angles are not defined, hence position angles are listed only for stars with non-zero polarization. The interstellar polarization is presumed to be negligible because of their high Galactic latitudes. The interstellar reddening, listed in Table 3, for the program stars is small. For the program stars for which color excess

Table 3
V-band Polarimetry of Carbon Stars

Star Names	$E(B - V)$	T_{eff}	$\log g$	[Fe/H]	Reference	HJD	$p\%$	θ°
HD 209621	0.09	4500	2.0	-1.93	1	2454950.47784	0.11 ± 0.23	
HD 206983	...	4200	1.4	-1.43	2	2454950.48532	0.06 ± 0.13	
HD 100764	0.02	4850	2.2	-0.6	3	2454952.14664	0.33 ± 0.06	4.0
HE 1152-0355	0.026	4000	1.0	-1.3	4	2454952.15514	0.24 ± 0.09	30.5
HE 1027-2501	0.05	2454952.18021	1.13 ± 0.11	12.4
HE 1305+0007	0.038	4750	2.0	-2.01	4	2454952.22191	1.26 ± 0.04	41.9
HE 1429-0551	0.05	4700	1.5	-2.47	5	2454952.27449	1.31 ± 0.06	28.9
LP 625-44	0.08	5500	2.5	-2.70	6	2454952.31728	1.06 ± 0.03	8.0
HE 1523-1155	0.05	4800	1.6	-2.15	5	2454952.35635	0.85 ± 0.08	30.3
HD 168986	2454952.39775	0.36 ± 0.03	32.5
HD 196944	0.0	5250	1.8	-2.25	7	2454952.47013	0.09 ± 0.02	
HD 198269	<0.07	4460	...	-1.56	8	2454952.47835	0.27 ± 0.02	39.5

References. (1) Goswami & Aoki 2010a; (2) Drake & Pereira 2008; (3) Dominy 1984; (4) Goswami et al. 2006; (5) Aoki et al. 2007; (6) Aoki et al. 2002; (7) Van Eck et al. 2001; (8) Lee 1974.




Figure 1. Upper panel: observed V-band percentage polarization for the program stars vs. interstellar $E(B - V)$. The solid line represents the mean interstellar polarization ($\sim 3E(B - V)$). Middle panel: observed V-band percentage polarization of the program stars w.r.t. their metallicities [Fe/H]. Polarization does not seem to have any co-relation w.r.t. metallicities of the stars. Bottom panel: the observed V-band percentage polarizations of the program stars are plotted w.r.t. their Galactic latitudes. Estimated polarizations are larger for objects with positive Galactic latitudes; however, there are two outliers (shown with triangles) that show weaker polarization. These two objects (HD 100764 and HE 1152-0355 with [Fe/H] ~ -0.6 and -1.3 , respectively) are not so metal-poor compared to the other stars that show larger polarization.

have been known, it is instructive to compare the spatial distribution of the color excess with polarization. In Figure 1 (upper panel), we have plotted the V-band polarimetric estimates of the stars ($p\%$) against interstellar reddening $E(B - V)$ along with the mean interstellar polarization $\sim 3E(B - V)$; Parthasarathy et al. 2005). Estimated V-band polarization is found to be larger than the mean interstellar contribution to the polarization in most of the cases indicating that they are intrinsically polarized.

Polarization does not seem to have any co-relation w.r.t. metallicity ([Fe/H]) of the stars (Figure 1, middle panel). The observed V-band percentage polarizations are larger for high Galactic latitude objects (Figure 1, lower panel); however

there are two outliers (shown with triangles in Figure 1, lower panel) that are high Galactic latitude objects but show weaker polarization. These two objects (HD 100764 and HE 1152-0355 with [Fe/H] ~ -0.6 and -1.3 , respectively) are not so metal-poor compared to the other stars that show larger polarization. It has been observed that the objects clearly fall into two distinct groups; five objects showing polarization below 0.4% and the other five showing polarization above 1.0%. HD 206983 and HD 196944 in our sample do not show significant evidence of linear polarization in the V band. The properties of the individual stars are discussed below.

3.1. HD 206983, HD 196944

HD 206983 is listed in the CH star catalog of Bartkevicius (1996). Recent high-resolution analysis of HD 206983 by Drake & Pereira (2008) shows this object to be a Barium star with an effective temperature of 4200 K and a metallicity of [Fe/H] = -1.43 . Estimated $p\%$ is 0.06 ± 0.13 . The s -process element-rich star HD 196944 is one of the first stars recorded to have shown very high abundance of lead (Pb). This star, also known as a lead star, is hotter (~ 5200 K) and more metal-poor ([Fe/H] = -2.25) than HD 206983 (Aoki et al. 2002; Van Eck et al. 2001). Both C and N abundances are enhanced ([C/Fe] = 1.2, [N/Fe] = 1.3). This is a high-velocity object with estimated radial velocity -174.76 ± 0.36 km s $^{-1}$, the abundance distribution observed in this star is interpreted as resulting from now-extinct companion AGB star. HD 196944 is mentioned in SIMBAD as a star with envelop of CH; the envelop might have resulted from as the companion star evolved through AGB and PN phases suffering mass loss. The estimated polarization ($p\% = 0.09 \pm 0.02$) is not significant; it is also likely that the non-detection of polarization would perhaps mean that the CH envelop is symmetrically structured. No previous polarization estimates are available for these two objects.

3.2. HD 100764, HD 168986, HD 198269, HD 209621, and HE 1152-0355

Estimated $p\%$ for these objects are, respectively, 0.330 ± 0.06 , 0.358 ± 0.03 , 0.269 ± 0.02 , 0.114 ± 0.23 , and 0.245 ± 0.09 .

Among these objects only HD 100764, a non-pulsating R-type carbon star, has previous polarimetric measurements in literature. Percentage polarization estimates in the V band on two

occasions by Parthasarathy et al. (2005) are, respectively, 0.20 ± 0.05 (JD 2,447,971) and 0.42 ± 0.14 (JD 2,448,398). Multiband photopolarimetry of HD 100764 shows no significant variations in $p\%$ in B , V , U , and I bands. No information is available on time variability of polarization for this star. Our estimated $p\% = 0.33 \pm 0.06$ is not too different from the previous estimates.

The reddening in the direction of HD 100764 ($E(B - V) = 0.02$; Eggen 1972) is consistent with the high Galactic latitude of the star. The observed V -band polarization is greater than the mean interstellar contribution to the polarization, indicating that the star is intrinsically polarized. Richer (1975) derived the absolute visual magnitude of $M_v = +0.3$ for this object that corresponds to a distance of 500 pc with $E(B - V) = 0.02$. From a detailed spectroscopic analysis Dominy (1984) found $T_{\text{eff}} = 4850$ K, $\log g = 2.2$, and a metallicity $[\text{Fe}/\text{H}] = -0.6$ for this object.

HD 100764 shows significant far-infrared excess fluxes at $12 \mu\text{m}$, $25 \mu\text{m}$, $60 \mu\text{m}$, and $100 \mu\text{m}$ with uncertainties of 4%, 8%, 12%, and 12%, respectively (Beichmann et al. 1985). The silicate dust shell around HD 100764 is believed to absorb the radiation from the star and re-emit the radiation in the infrared. Modeling of its circumstellar dust by Skinner (1994) revealed a massive dusty disk. Parthasarathy (1991) suggested that HD 100764 is a star with a detached cold dust shell and that the optical region fluxes of HD 100764 do not show evidence for significant reddening, and argued that the dust shells are optically thin with relatively large dust grain size and dust is confined in the form of thin disc around the star and hence shows less reddening. The silicate dust shells around this star may be the result of mass loss experienced by the star during core helium flash, which has taken place recently. The presence of dust shell around HD 100764 also suggests that it has experienced significant mass loss in the recent past.

The star HD 168986 is listed in the Barium star catalog of Lü (1991). Barium stars are however not known to have dust shells or discs around them (Dominy et al. 1986). This star mentioned as a peculiar star in SIMBAD shows an estimated $p\% = 0.36 \pm 0.03$, that seems to represent the intrinsic polarization property of the star. No previous polarization estimates are available for this object.

HD 198269 is listed in the CH star catalog of Bartkevicius (1996). In SIMBAD, this star is mentioned as a star with envelop of CH. Lee (1974) gave an upper limit to its distance estimate as ~ 750 pc that corresponds to $M_v = -1.66$, an upper limit to the absolute magnitude. This is a non-variable CH star of type R with a mass of about $0.8 M_{\odot}$ (Wallerstein 1973) and is similar in chemical compositions in comparison to other CH stars. With $E(B - V) \leq 0.07$ mag for this star, the estimated $p\% = 0.27 \pm 0.29$ is marginally higher than the mean interstellar contribution to the polarization. No previous polarization estimates are available for this object.

HD 209621 and HE 1152–0355, both are confirmed CH stars. Surface composition of these two objects show abundance patterns that are consistent with the abundances generally noticed in CH stars, essentially arising from s -processing (Goswami & Aoki 2010a; Goswami et al. 2006). HD 209621 shows high abundances of r -process element Eu as well as of the third-peak s -process element Pb. Eu and Pb are, however, not detected in the spectrum of HE 1152–0355. Both these objects show larger enhancement of the second-peak s -process elements as compared to those of the first-peak s -process elements. Both are high-velocity objects. HD 209621 is a radial

velocity variable with a period of 407.4 days (McClure & Woodsworth (1990); however, radial velocity variability is not yet confirmed for HE 1152–0355. Our V -band polarimetric estimates indicate no significant polarization from HD 209621, and HE 1152–0355 shows polarization that is marginally above the mean interstellar contribution to the polarization.

3.3. HE 1027–2501, HE 1305+0007, HE 1429–0551, LP 625–44, and HE 1523–1155

These objects show significant polarization with $p\% \geq 1$, except for HE 1523–1155 for which we have obtained $p\% = 0.849 \pm 0.08$. Reddening estimates $E(B - V)$ range from 0.03 to 0.08 for these stars; the values of $E(B - V)$ for the HE stars are taken from Beers et al. (2007).

A low carbon isotopic ratio ($^{12}\text{C}/^{13}\text{C} < 10$) for HE 1027–2501 indicates that the star is on the first ascent of the giant branch wherein the material transferred from the now unseen companion has been mixed into the CN-burning region of the CH star or constitute a minor fraction of the envelop mass of the CH star. Such low values are believed to be due to convection which dredges up the products of internal CNO cycle to the stellar atmosphere in the ascending red giant branch (RGB). When the star reaches the AGB stage, fresh ^{12}C may be supplied from the internal He-burning layer to the stellar surface, leading to an increase of $^{12}\text{C}/^{13}\text{C}$ ratio.

HE 1305+0007 shows enhancement of both r -process and s -process elements including lead. The second-peak s -process elements are more enhanced than the first-peak s -process elements. This is a low-metallicity and high-velocity object ($V_r = +217.8 \pm 1.5 \text{ km s}^{-1}$); its atmospheric parameters are consistent with a present location on the RGB (Goswami et al. 2006).

The star LP 625–44 is a carbon- and s -process-element-rich very metal-poor subgiant. Abundance estimate derived using the O I triplet around 7770 \AA shows excess of oxygen by a factor of 10 (Aoki et al. 2002); and Na enhancement by about a factor of 50 in comparison to HD 140283 (a metal-poor subgiant with normal abundance ratio) and a high Mg abundance ($[\text{Mg}/\text{Fe}] = 1.12 \pm 0.24$). High abundance of Na suggests that hydrogen burning in the ^{22}Ne -rich layer in an AGB star must have produced the abundance pattern of this object. The Pb enhancement shown by LP 625–44 is not high enough to be placed in the group of lead stars. The abundance ratio of s -process elements at the second peak (La, Ce, and Nd) to that at the third peak (Pb) in LP 625–44 is significantly higher (by a factor of five) than that in the s -process element-rich lead star HD 196944 (Van Eck et al. 2001). Unlike CH stars, the radial velocity of this object is low ($\sim 30 \text{ km s}^{-1}$) and the variation of the radial velocity is expected to be for about 200 days (Aoki et al. 2000).

Like LP 625–44, HE 1429–0551, and HE 1523–1155 are also low-velocity objects (~ -44.9 and $\sim -46.03 \text{ km s}^{-1}$, respectively). Both show high abundances of C, N, and Mg relative to Fe. Estimated carbon isotopic ratio of $^{12}\text{C}/^{13}\text{C}$ for HE 1429–0551 is high $\sim 30^{+20}_{-10}$ (Aoki et al. 2007). Such high ratios are generally noticed in C–N stars. With a marginal difference in the molecular band depths, the spectra of the star HE 1523–1155 closely resemble the spectrum of HD 5223, a well-known CH giant (Goswami 2005). Both the HE stars show high barium abundance with respect to Fe ($[\text{Ba}/\text{Fe}] = 1.57$ and 1.72 , respectively), and $\log(L/L_{\odot})$ are, respectively, 2.56 and 2.50 (Aoki et al. 2007). The observed V -band polarization estimates are markedly higher than the mean interstellar contribution to the

polarization and seem to represent the stars intrinsic polarization properties.

4. CONCLUSIONS

Our V-band polarimetric observations presented for a sample of 10 carbon-enhanced metal-poor (CEMP) stars show five objects with polarization $p\% > 1$ and the other five with $p\% < 0.4$. While the size of the sample is relatively small, it should be noted that these stars belong to similar group in terms of carbon enhancement and metal deficiency. Thus, a clear separation of these objects into well-separated V-band polarimetric groups is significant. Intrinsic polarization of late-type stars are usually attributed either to the Raleigh scattering of the light emerging from the limb of a star in which case the sources of photometric asymmetry necessary for a net observable polarization are suggested to be non-radial pulsation of the star, variation of temperature over the surface, and the presence of giant convection cells (Harrington 1969; Schwarzschild 1975) or due to scattering by molecules and dust grains in an extended asymmetric circumstellar envelope (Kruszewski et al. 1968; Shawl 1975; Daniel 1978). It seems circumstellar grain scattering is the main mechanism responsible for the observed polarization in the sample of program stars. It would be worthwhile to conduct polarimetric observations for the entire sample of carbon stars from Hamburg ESO survey (Christlieb et al. 2001) for a statistical evaluation of their polarization properties.

Except for HD 100764, our results are the first estimates of polarization for the program stars. While the separation of the stars into two distinct polarization groups is a significant feature, it is not clear at the moment if this is a time-dependent property. While a few of the CEMP stars are confirmed as radial velocity variables and binaries, binarity of others are not yet established. Two important means for establishing variability and binarity of stars are long-term photometric observations as well as long-term spectroscopic radial velocity measurements. Time variation of polarization measurements for these objects would also be an indicator of their variability caused by the morphology of the circumstellar dust clouds that might have originated during AGB mass loss of the companion stars. Thus, the detection of polarization from these objects would lend support to those physical scenarios that consider binarity of the objects to explain their formation mechanism and peculiar abundances of heavy elements.

We thank the staff at IUCAA observatory for time allocation and assistance during the observations. This work made use of the SIMBAD astronomical database, operated at CDS, Strasbourg, France, and the NASA ADS, USA. We thank Prof. A. V. Raveendran for useful discussions. S.S.K. is a JRF in the DST project SR/S2/HEP-09/2007; funding from this project is gratefully acknowledged.

REFERENCES

- Aoki, W., Norris, J. E., Ryan, S. G., Beers, T. C., & Ando, H. 2000, *ApJ*, **536**, L97
- Aoki, W., et al. 2002, *PASJ*, **54**, 427
- Aoki, W., et al. 2007, *ApJ*, **655**, 492
- Bartkevicius, A. 1996, *Balt. Astron.*, **5**, 217
- Beers, T. C., et al. 2007, *ApJS*, **168**, 128
- Beichmann, C. A., Neugebauer, G., Habing, H. J., Clegg, P. E., & Chester, T. J. 1985, *IRAS Point Sources Catalog* (Pasadena, CA: JPL)
- Chavero, C., Gomez, M., Whitney, B. A., & Saffe, C. 2006, *A&A*, **452**, 921
- Christlieb, N., Green, P. J., Wisotzki, L., & Reimers, D. 2001, *A&A*, **375**, 366
- Clarke, D., Smith, R. A., & Yudin, R. V. 1998, *A&A*, **336**, 604
- Czyzak, S. J., Hirth, J. P., & Tabak, R. G. 1982, *Vistas Astron.*, **25**, 337
- Daniel, J. Y. 1978, *A&A*, **67**, 345
- Dominy, J. F. 1984, *ApJS*, **55**, 27
- Dominy, J. F., Lambert, D. L., Gehrz, K. D., & Mozurkewich, D. 1986, *AJ*, **91**, 591
- Drake, N. A., & Pereira, C. B. 2008, *AJ*, **135**, 1070
- Eggen, O. 1972, *MNRAS*, **159**, 403
- Goswami, A. 2005, *MNRAS*, **359**, 531
- Goswami, A., & Aoki, W. 2010a, *MNRAS*, **404**, 253
- Goswami, A., Aoki, W., Beers, T. C., Christlieb, N., Norris, J. E., Ryan, S. G., & Tsangarides, S. 2006, *MNRAS*, **372**, 343
- Goswami, A., Bama, P., Shantikumar, N. S., & Devassy, D. 2007, *Bull. Astron. Soc. India*, **35**, 339
- Goswami, A., Karinkuzhi, D., & Shantikumar, N. S. 2010b, *MNRAS*, **402**, 1111
- Harrington, J. P. 1969, *Astrophys. Lett.*, **3**, 165
- Johnson, J. J., & Jones, T. J. 1991, *AJ*, **101**, 1735
- Kruszewski, A., Gehrels, T., & Serkowski, K. 1968, *AJ*, **73**, 677
- Lee, P. 1974, *ApJ*, **192**, 133
- Leonard, D. C., Li, W., Filippenko, V., Foley, R. J., & Chornock, R. 2005, *ApJ*, **632**, 450
- Lü, P. K. 1991, *AJ*, **101**, 2229
- McClure, R. D., & Woodsworth, A. W. 1990, *ApJ*, **352**, 709
- McDavid, D. 1999, *PASP*, **111**, 494
- Parthasarathy, M. 1991, *A&A*, **247**, 429
- Parthasarathy, M., & Jain, S. K. 1993, in *IAU Symp. 155, Planetary Nebulae*, ed. R. Weinberger & A. Acker (Dordrecht: Kluwer), 353
- Parthasarathy, M., Jain, S. K., & Sarkar, G. 2005, *AJ*, **129**, 2451
- Ramprakash, A. N., Gupta, R., Sen, A. K., & Tandon, S. N. 1998, *A&AS*, **128**, 369
- Rautela, B. S., Joshi, G. C., & Pandey, J. C. 2004, *Bull. Astron. Soc. India*, **32**, 159
- Raveendran, A. V. 1991, *A&A*, **243**, 453
- Reiz, A., & Franco, G. A. P. 1998, *A&AS*, **130**, 133
- Richer, H. B. 1975, *ApJ*, **197**, 611
- Rowan-Robinson, M., & Harris, S. 1983, *MNRAS*, **202**, 797
- Schwarzschild, M. 1975, *ApJ*, **195**, 137
- Sen, A. K., Gupta, R., Ramprakash, A. N., & Tandon, S. N. 2000, *A&AS*, **141**, 175
- Serkowski, K. 1974, in *Planets, Stars, and Nebulae Studied with Photopolarimetry*, ed. T. Gehrels (Tucson, AZ: Univ. Arizona Press), 135
- Serkowski, K., Mathewson, D. S., & Ford, V. L. 1975, *ApJ*, **196**, 261
- Shawl, S. J. 1975, *AJ*, **80**, 595
- Skinner, C. J. 1994, *MNRAS*, **271**, 300
- Trammell, S. R., Dinerstein, H. L., & Goodrich, R. W. 1994, *AJ*, **108**, 984
- Turnshek, D. A., Bohlin, R. C., Williamson, R. L., Lupie, O. L., Koornneef, J., & Morgan, H. D. 1990, *AJ*, **99**, 1243
- Van Eck, S., Goriely, S., Jorissen, A., & Plez, B. 2001, *Nature*, **412**, 793
- Wallerstein, G. 1973, *ARA&A*, **11**, 115
- Weitenbeck, A. J. 2004, *Acta Astron.*, **54**, 87