

RESULT

OF

ASTRONOMICAL OBSERVATIONS

MADE AT

THE HONORABLE

THE EAST INDIA COMPANY'S OBSERVATORY

AT MADRAS

BY

THOMAS GLANVILLE TAYLOR, Esq.

ASTRONOMER TO THE HONORABLE COMPANY.

VOL. I.

FOR THE YEAR 1831.

PRINTED

BY ORDER OF THE

MADRAS GOVERNMENT.

MADRAS:

PRINTED AT THE ORPHAN ASYLUM PRESS.

M.DCCC.XXXII.

P R E F A C E.

PROFESSOR AIRY says* “ I think it may be fairly asserted that the value of unreduced Observations is very small and I do not hesitate to say that if an offer were made of a mass of regular meridional observations, unreduced, I should not think it worth acceptance :” upon such respectable authority I have thought it adviseable to bring before the public in the first instance the result of my observations with the details of the manner in which the reductions have been effected, and for the present to suppress altogether the rough observations depositing for the sake of reference manuscript copies with the Honorable Court of Directors.

~~It is proper I should state that the Instruments with which the Observa-~~
tory is at present supplied (those with which the observations I am about to detail have been made) consist of a five feet Transit Instrument, a Mural Circle of four feet diameter and a five feet Telescope equatorially mounted recently constructed by Dollond on purpose for the Observatory. These Instruments were sent out to this Country in the summer of 1829, but by reason of the absence of the Astronomer they were allowed peacefully to repose in their packing cases till my arrival at Madras on the 15th September 1830 ; from this time to the end of the year I found abundance of employment in superintending the removal of the Instruments hitherto in use (a 20-inch Transit Instrument by Stancliff, and a 12-inch Altitude and Azimuth Instrument by Troughton) in building Piers for the reception of the new Instruments abovementioned and in instructing the native assistants in their duties.

* Preface to Cambridge Observations.

The Observatory as it at present stands was built in the year 1792, having in this interval received alterations of a nature little interfering with its general character as an Observatory, in consequence of which on my taking charge, I had several objections to offer to the construction of the building, some of which were surmountable and were through the liberality of the Madras Government immediately attended to, others which could not so easily be overcome I have been compelled to allow to remain, of these I may mention the smallness of the aperture in the roof (18 inches) which prevents an alteration in the temperature on the outside of the building being communicated readily to the inside, and the unavoidable noise made by the observer at the one instrument disturbing the observer at the other; it has however been my particular care to diminish the effect of these inconveniencies, and I believe they little if at all interfere with the general accuracy of the results. The Observatory consists of ~~but one apartment measuring~~ 65 feet long, 20 feet broad, and 14 feet high; in the centre of, and perfectly insulated from the building, about one foot beneath the floor, is a solid mass of brickwork going down to a depth of 7 feet below the floor, 45 feet long and tapering (Pyramid fashion) from a base of 12 feet, to 6 feet width at top; after a careful examination I was convinced that no foundation I could construct would exceed in stability that of the pile in question which assumed more the nature of a single block of stone than of detached pieces held together by cement; accordingly it is upon the extreme ends of this pyramid that the Transit and Mural Circle Piers are built opposite to two apertures in the roof which had hitherto been used with the old instruments. On the eastern end of the Pyramid are built the Piers for the Transit Instrument; they consist of two pyramidal upright stone Piers resting upon a slab of the same 9 inches thick, $7\frac{1}{4}$ feet long and $2\frac{1}{2}$ feet wide to which they are cemented. See Plate I. These stones were cut under my own direction from blocks of an exceedingly hard quartzous stone brought from the Palaveram Hills known

by the name of blue stone. The Transit Clock is in like manner very steadily supported by a large block of blue stone.

In the centre of the building is a conical stone Pier tapering from a base of 4 feet diameter at the floor to 2 feet at 18 feet high or 4 feet above the roof of the Observatory serving to support a 12-inch Altitude and Azimuth Instrument by Troughton which is enclosed by a circular revolving dome.

The Mural Circle Pier which rests upon the western extremity of the Pyramid consists of $2\frac{1}{2}$ feet of new brick work surmounted by blocks of stone, *a, b, c, d*, which were cut and sent out from England on purpose. Plate II. The five feet Achromatic is exceedingly well and steadily mounted on a mahogany frame armed with brass, and being supplied with ~~two graduated circles and a long axis~~ moving on a graduated arc, it has occasionally been employed as an equatoreal in making rough observations out of the meridian in addition to its other uses in observing Occultations and Eclipses.

In using these Instruments I have adopted the plan of never touching any of the adjusting screws except in cases where the error was inconveniently large, but of computing the corrections due to each particular deviation; in the case of the Transit Instrument the inversion of the axis is rendered unnecessary by two marks one to the North the other to the South of the Observatory, the angular distance between which has been accurately found.

The points to which I have particularly directed my observation are:

1st. The determination of the places of a large Catalogue of Stars from a moderate number of observations, at first 1200 Stars only were

selected as an observing Catalogue, but lately this number has been increased to 2881 being the Catalogue selected by Mr. Bailly, published in the Second Volume of the Astronomical Society's Memoirs.

2ndly. To observations of the Planets at all times when they pass the Meridian between half past 5 o'clock in the morning and half past 12 at night, (the regular hours of observing).

3rdly. To observations of the Moon, Moon Culminating Stars and Occultations.

4thly. To observations of the Eclipses of Jupiter's Satellites.

To the observations of double Stars I have yet not been able to devote any time; although they are a class of Observation of a highly interesting nature, I have preferred for the present not to neglect other duties in their pursuit.

It now remains for me to state that the nature of my appointment, that provides in addition to the Superintendance of the H. C. Observatory at Madras I should be required to assist in the operations of the Great Trigonometrical Survey of India, so that I might calculate with certainty on being absent occasionally several months together from the Observatory; having this in view the plan proper to be followed appeared to be that of procuring assistants whom I could trust to make the observations during my absence; after but short experience with four native assistants I found that this could with great safety be done, accordingly the main mass of the observations here spoken of are made by the native assistants under my superintendance, and lately during my absence for six months in Bengal for the purpose above named the observations were altogether made by the native assistants; by this arrangement when at home I am enabled to attend more particularly to the computation of the observations which could not with any degree of safety be trusted to an assistant, accordingly

I consider myself, and myself only answerable for the accuracy of all the computations which may hereafter follow having in no case permitted computations made by the natives to go unexamined; that the observations and computations may prove accurate and useful, and tend in a degree however slight to advance the cause of Astronomy is my most constant and ardent wish.

T. G. TAYLOR,

H. C. ASTRONOMER.

Madras Observatory,

1st March 1832.

OF THE TRANSIT INSTRUMENT, &c.

THE Transit Instrument as I have already stated was constructed by Dollond, is 61 Inches focal length with a clear aperture of $3\frac{3}{4}$ Inches, this for Observations of the Sun and bright Stars has occasionally been reduced to $2\frac{1}{2}$ Inches, in either case the optical power of the Telescope has given me the greatest possible satisfaction, the focal distance of the eye-piece is Inch 0, 40. giving a power of 150: the eye-end is furnished with one horizontal and five vertical fixed wires, to which is added two wires moveable by a spring micrometer screw with which every day at Sun-rise and Sun-set the deviation of the central wire from the North and South marks has been registered; for the distance between the wires and value of the screw of the micrometer, see "Observations made with the Transit Instrument."

The axis is perforated at one end in the usual way for the admission of light from a lamp at night, which by means of a screw near the eye-piece can be moderated for the observation of faint Stars.

The Y's are each composed of three distinct pieces, each piece being moveable round an axis peculiar to itself; the first piece or Y in which the pivots immediately rest is represented by Fig. 1. Plate II. together with its axis which is a screw with a capstan head working in the Y itself; when the instrument is first brought to bear upon the Y's this piece is allowed to revolve easily and tightened only when the pivots have come to their bearing and the Y's taken up a position convenient to their repose.

The first and second pieces put together are represented Plate II. Fig. 2, where *a a* represent two pins intended to act as a fulcrum to the counterpoises. A.A. Plate II. Fig. 2 and 3, are two pivots on which the whole is supported, by this contrivance the Y's which are filed flat are brought to support the pivots throughout their whole length. Fig. 3, Plate II, represents the western side of the western Y which is adapted exclusively to azimuthal motion and is performed by means of two antagonizing screws B.B. causing the pieces above described to revolve round a centre C (see also Plate I.), placed 10 inches from the face of the Y's themselves.

The eastern *Y* differs from the above only from being exclusively adapted to vertical motion; the pieces B.B. have consequently no place in the eastern *Y* and the frame *a, b, c, d*, which in this case slides in contact with the shoulders on either side is raised or lowered equally by two screws with graduated heads situated at D.D. working in the pieces immediately above, by this contrivance an adjustment for level is effected without altering the position in azimuth. Allusion is made above to counterpoises, it will consequently be expected that some description of them should be given: The counterpoises in question were constructed on a principle which is I believe new, and were to all appearance on a plan likely to ensure success, but after several days of vexation at not meeting with the accordance I could wish among the observations for collimation and level, &c. I was at length tempted to try the effect of removing the counterpoises, the result was that the observations agreed perfectly, and that I have not since used them. The Level for the Transit Instrument is one of great sensibility, in order to find the value of the divisions recourse was had to the Mural Circle on which it was placed and its indications registered corresponding to every 10 seconds through which the Circle was moved, the result was that the reading of the scale corresponded in every part with the reading of the Circle giving each division which measures 0,05 Inch = 0",985. The Transit Clock is the common lead-beat escapement by Shelton with a gridiron Pendulum, it was reported to have gone well previously to my having the care of it, but the rate from day to day which hereafter follows will best bespeak its worth.

OF THE OBSERVATIONS MADE WITH THE TRANSIT INSTRUMENT.

In the first instance it was necessary to know if the pivots of the Transit Instrument were alike in size and if not what the difference of their diameter and the effect it introduced in the Observations for level; for this purpose the following Observations were made and registered in the Level book.

OBJECT GLASS NORTH, CROSS LEVEL EAST, AND ILLUMINATING END OF THE AXIS.

	East.	West.	Diff. or illuminating end largest.
1831	"	"	"
April 3, Mean from 20 reversion.....	0,67 E.	1,25 W.	+ 0,96
April 9, Mean from 20 reversion.....	0,56 W.	0,81 W.	+ 0,125
April 11, Mean from 20 reversion.....	1,11 W.	2,61 W.	+ 0,79
April 14, Mean from 20 reversion.....	1,31 W.	2,20 W.	+ 0,445

Taking the mean we have 0",58 for the apparent excess of the Illuminating or perforated pivot above the other pivot. The angle of the Y's of the transit as well as that of the level is exactly 90°; now if r, r', represent the radii of the illuminating and unilluminating pivots we have

$$\left. \begin{aligned} 2 r \operatorname{cosect.} \left(\frac{90}{2}\right) \\ 2 r' \operatorname{cosect.} \left(\frac{90}{2}\right) \end{aligned} \right\} \text{apparent height of the } \left\{ \begin{array}{l} \text{illuminating pivot} \\ \text{unilluminating pivot,} \end{array} \right.$$

the difference 2,82 (r - r') = 0",58, or the difference of the radii of the pivots = 0",206 = 0,000034 Inches which produces an alteration in the height of the centre of the axis = (r - r') cosect. (90/2) = 0",29, accordingly this quantity must always be applied to the registered error of level as set down in the level book to obtain the true error.

The Intervals between the wires next came under examination, for this purpose the Equatorial Interval was computed from 70 Observations to be

from 1st wire to centre.....	+ 55",036
2d	+ 27",477
4th.....	- 27",478
5th.....	- 54",868

These numbers hold good up to 15th March 1831, when the following note appears in the Transit book.

“ At the commencement of the evening Observations a piece of stick about one inch long and a quarter of an inch wide, was found suspended on the wires, having by its weight broken the central one and bent the others, in consequence of which the observations were given up for the evening, and a new set of cobwebs introduced next morning.”

From March 15th to 22d from 70 Observations the Equatorial Interval was

from 1st wire to centre.....	+ 55",166
2d	+ 27 ,526
4th.....	— 27 ,596
5th.....	— 55 ,080

On March 23d, 1831, the following note appears in the Transit book.

“ On looking at the Sun a piece of wood about $\frac{1}{4}$ inch long appeared suspended on the 4th wire which with others it had broken, this piece of wood as well as that found on the 15th Instant, had evidently been recently picked, the ends shewing the fibres to have been twisted in the act of being broken from the tree by the miscreant hand who introduced it, and the bark on being removed exhibited the sap in a moist state; I am quite at a loss to account for the motives of the individual who can thus act! I introduced a new set of cobwebs to supply the place of those broken this morning.”

From March 23d to April 21st, the Equatorial Interval from 156 Observations was found to be

from 1st wire to centre.....	+ 54",929!
2d	+ 27 ,451
4th.....	— 27 ,501
5th.....	— 55 ,081

April 21st, 1831, the following note appears in the Transit book.

“ On looking through the Telescope for the Observation of the Sun the five vertical and horizontal lines were found broken; I replaced them with cobwebs.”

NOTE. I have great reason to believe that an Invalid Sepoy to whom the keys of the Observatory were intrusted during the absence of the Assistants, is the person who introduced the pieces of stick abovementioned into the Instrument; he was (as I afterwards found out) occasionally intoxicated and was in consequence discharged.

For the Observations of April 21st from 27 Observations the following Equatorial Intervals have been computed :

from 1st wire to centre.....	+ 55 ^s ,100
2d	+ 27 ,747
4th.....	— 27 ,336
5th.....	— 54 ,667

On the 22d April 1831, the following note appears in the Transit book.

“ The central wire appears to have been badly secured yesterday, so that on looking into the Telescope this morning one end had disengaged itself from the varnish ; it was removed and another put in : the 4th wire was also secured with fresh varnish.”

From April 22d to May 2d, 1831, the Equatorial Interval from 148 Observations was found to be

from 1st wire to centre.....	+ 54 ^s ,669
2d	+ 27 ,364
4th.....	— 27 ,597
5th.....	— 55 ,045

On the 2d May 1831, the following note appears in the Transit book.

“ The central wire appeared split into two parts at one end ; I removed all the wires and put in another fresh set.”

From 2d May to 7th May 1831, the Equatorial Interval from 70 Observations was found to be

from 1st wire to centre.....	+ 54 ^s ,635
2d	+ 27 ,642
4th.....	— 27 ,781
5th.....	— 54 ,798

May 7th 1831, the following note appears in the Transit book.

“ Found the fourth wire broken ! In consequence of the ill success attending cobweb lines and the difficulty of procuring them, I have now introduced silk lines from unspun silk ; their appearance is much more uniform than the best cobwebs and they moreover are not acted upon as cobwebs are by the drying properties of the air.”

From May 10th to July 25th, the Equatorial Interval from 100 Observations was found to be

OBSERVATIONS of the time occupied by the POLE STAR to pass from the central wire to

1831	R. D.	R. D.	R. D.	R. D.	R. D.	R. D.	R. D.	R. D.	R. D.
	1 60	1 20	0 80	0 40	0 40	0 80	1 20	1 60	
	East of the Central Wire.				West of the Central Wire.				
	t "		t "			t "			
February	23 2 11,0	1 10,0	1 7,0	
June	10	1 5,5	1 9,5	2 11,5	
	15 2 10,5	1 4,5	1 6,0	2 13,0	
	15 2 12,0	1 6,0	1 4,0	2 10,5	
	19 2 11,8	1 6,0	1 3,2	2 9,7	
	28 2 12,0	1 40,5	1 7,0	0 33,5	0 29,5	1 3,5	1 37,5	2 11,0	
	28 2 10,5	1 38,0	1 6,0	0 34,0	0 31,5	1 4,0	1 37,0	2 9,5	
	29 2 13,5	1 9,5	0 33,0	1 2,5	2 8,5	
	29 2 12,5	1 40,0	1 6,5	0 33,0	0 33,5	1 5,0	1 38,0	2 9,5	
July	1	1 7,0	
	5 2 13,5	1 40,0	1 7,0	0 34,0	0 33,0	1 5,5	1 38,5	2 10,5	
	5 2 19,5	1 36,0	1 4,0	0 31,0	0 33,5	1 7,5	1 40,5	2 14,0	
Mean..	2 11,68	1 38,90	1 6,55	0 33,08	0 32,22	1 5,40	1 38,50	2 10,77	

Now the spaces passed over by the Pole Star in any very small intervals of time from the meridian t, t' , are equal to $15 \sin. P.D. \times t$ and $15 \sin. P.D. \times t'$ respectively, or very nearly so, and for $t - t'$ the time intermediate = $15 \sin. P.D. \times (t - t')$ very nearly; taking the successive difference and using for P.D. $1^\circ 35' 48''$ we get the following values for every successive 40 divisions of the micrometer screw, viz.

R. D.	R. D.	s.	
from 1 60 East to 1 20 East.....		32,78	corresponding to.....13,70
1 20.....0 80.....		32,3513,52
0 80.....0 40.....		33,4713,98
0 40.....Centre Wire.....		33,0813,83
Centre Wire..0 40 West.....		32,2213,46
0 40 West...0 80.....		33,1813,88
0 80.....1 20.....		33,1013,75
1 20.....1 60.....		32,2713,57
R. D.			
∴ 0 40 =	13",711	or 1 Revolution =	34",277

From the few Observations hitherto made of this nature the exact value of the micrometer screw must still be considered as undetermined, but the near agreement of the results seem to justify for the present the assumption that the threads of the screw are very nearly equidistant, and that each revolution does not greatly differ from the mean found above which has consequently been employed in the reductions.

ERROR OF LEVEL.

THE error of Level of the Transit Axis has been determined alone by the Spirit Level from Observations made at intervals of two or three days; the plan I have adopted, is taking the mean of three readings; one with the Y's of the level situated at the eastern extremity of the pivots, the next at the middle, and lastly when removed to the western extremity of the pivots; the mean of these three readings is employed for "Cross Level East," the same is repeated on inverting the level when the Cross Level is West; after this the Level is again inverted or the Cross Level is again placed on the eastern Y and the three readings again repeated; if this third reading does not differ more than $0''.5$ from the first result the mean between it and the first is employed with the second reading to obtain the error of level; but if a difference exceeding this quantity is found (which I need hardly remark does sometimes occur from accidental and unavoidable violence in reversing the level) the readings are repeated again and again until the desired agreement is obtained; the value of the divisions of the level were found previously to its being used by a comparison of the readings of the Mural Circle on which it was placed, with the indications of the level due to intervals of $10''$ through which the circle was moved, from this it appeared that 30,45 divisions of the level are equivalent to $30''$ of space, or each division which measures $\frac{1}{30}$ of an inch = $0''.985^*$: having mislaid the paper in which the details of the observations for this purpose were registered I have thought it advisable to re-examine the level in the manner above mentioned; from the mean of several readings the following results are obtained.

1832, April 12th.			Mural Circle.		Mean.	Resulting Value of 1 division.	
North End. D.	South End. D.	Mean. D.	A	B			
+ 33,6	+ 30,0	+ 31,80	166 42 58,7	1,7	0,20	} — 0,986	
26,6	22,6	24,60	43 6,0	8,6	7,30		
17,5	14,1	15,80	13,4	17,2	15,30	} — 0,960	
+ 4,6	+ 2,0	+ 3,30	25,6	29,0	27,30		
— 7,6	— 10,4	— 9,00	37,6	41,0	39,30	} — 0,976	
— 18,6	— 21,0	— 19,60	48,5	51,7	50,10		
						Mean = 0,967	

* In the reductions of the Observations each division of the level has been assumed equal to one second.

OBSERVATIONS FOR LEVEL.

1831			1831	1831			
	D. H.	Illuminated End.	Error from Level.		D. H.	Illuminated End.	Error from Level.
February	18 1	West	5,18 W.	June	2 1	East	1,23 E.
	19 1	East	3,73 —		4 1	0,51 W.
	20 1	4,38 —		6 1	1,82 E.
	21 1	4,74 —		8 1	0,15 W.
	22 1	5,02 —		11 1	1,33 E.
	23 1	4,68 —		13 1	2,19 —
	24 1	4,37 —		15 1	1,79 —
	25 1	4,22 —		20 1	1,71 —
	28 1	4,13 —		22 1	3,64 —
March	3 1	3,85 —		24 1	3,54 —
	4 1	3,04 —		27 1	4,02 —
	5 1	2,66 —		29 1	3,27 —
	6 1	2,61 —	July	1 1	3,45 —
	7 1	2,76 —		3 1	3,62 —
	12 1	3,09 —		5 1	4,21 —
	15 1	3,71 —		7 1	3,55 —
	16 1	3,66 —		9 1	4,72 —
	18 1	3,65 —		13 1	4,69 —
	20 1	2,22 —		15 1	5,32 —
	23 1	1,81 —		17 1	5,97 —
	25 1	1,80 —		19 1	6,67 —
	28 6 *	0,43 E.		21 1	5,51 —
	29 6 *	0,65 —		23 1	5,96 —
	31 1	0,25 W.		25 1	6,26 —
April	1 1	0,31 —		27 1	5,25 —
	2 1	0,18 —		29 1	5,29 —
	3 6 *	0,13 —		31 1	5,07 —
	4 1	0,43 —	August	2 1	5,75 —
	6 1	0,41 —		4 1	6,27 —
	7 1	0,50 —		6 1	6,16 —
	9 6 *	0,41 —		8 1	6,07 —
	11 6 *	0,15 —		10 1	6,82 —
	14 6 *	2,25 —		12 1	6,12 —
	16 1	1,73 —		14 1	6,84 —
	18 1	0,62 —		16 1	4,55 —
	20 1	0,60 —		18 1	4,05 —
	25 1	0,18 —		20 1	4,53 —
	27 1	0,72 —		22 1	4,66 —
	29 1	0,08 —		24 1	4,75 —
May	1 1	0,30 —		26 1	4,37 —
	3 1	0,05 —		29 1	4,89 —
	5 1	1,65 —		31 1	4,79 —
	7 1	1,21 —	September	2 1	4,79 —
	11 1	0,76 —		4 1	5,83 —
	13 1	0,13 E.		7 1	6,13 —
	15 1	0,29 W.		9 1	6,14 —
	18 1	0,52 E.		11 1	5,18 —
	21 1	0,32 —		13 1	4,94 —
	23 1	0,12 W.		15 1	4,55 —
	25 1	0,12 —		17 1	4,53 —
	27 1	0,17 E.		19 1	4,53 —
	29 1	0,47 —		21 1	4,81 —
	31 1	0,50 —		23 1	4,54 —

* On each of these days the Axis was inverted and brought back again ten times.

			Illuminated End.	Error from Level.				Illuminated End.	Error from Level.
1831				"	1831				"
	D. H.					D. H.	East		
September	24	1	East	4,79 E.	November	11	1	3,36 E.
	26	1	4,01 —		18	1	6,29 —
	28	1	4,36 —		19	1	6,98 —
	30	1	4,64 —		21	1	5,57 —
October	2	1	4,50 —		23	1	5,01 —
	4	1	5,05 —		27	1	6,42 —
	6	1	4,28 —		29	1	6,52 —
	8	1	4,42 —	December	1	1	7,27 —
	10	1	4,71 —		3	1	6,36 —
	12	1	4,41 —		5	1	5,64 —
	14	1	4,33 —		7	1	5,77 —
	16	1	4,10 —		9	1	5,22 —
	18	1	4,76 —		11	1	4,88 —
	20	1	4,42 —		13	1	5,03 —
	22	1	4,67 —		15	1	5,42 —
	24	1	4,85 —		17	1	5,37 —
	26	1	4,03 —		19	1	4,87 —
	28	1	4,75 —		21	1	4,74 —
	30	1	4,57 —		23	1	4,45 —
November	1	1	5,14 —		25	1	4,41 —
	3	1	5,12 —		27	1	5,74 —
	5	1	4,86 —		29	1	4,94 —
	7	1	4,78 —		31	1	5,06 —
	9	1	4,69 —					

N. B.—E. and W. represent that the East or West end of the Axis is too *high* as the case may be.

The above column "Error from Level" is the mean of six or more readings of the Spirit Level taken in the manner already described together with the correction for inequality in the diameter of the pivots found at page 3; thus from the Level book.

1831, February 13.—Illuminating End W. error by Level..... 5",47 W.
 Illuminating Pivot too large..... — 0,29
 True error from Level..... 5,18 W.

These numbers can I think be depended upon to three or four tenth of a second for the day on which they are found; for the intermediate days I have generally employed corresponding intermediate values except in cases where the Axis has been inverted or the Instrument otherwise disturbed.

Now if e represent the true error of level as just found; every observed transit
 above the Pole } requires a correction $+ \left\{ \frac{e \times \sin. \text{Altitude}}{15 \sin. \text{N.P.D.}} \right.$
 below the Pole } $-$
 where e is $+$ for West error of level and $-$ if the error of level is East.

ERROR OF COLLIMATION.



THE position of the line of Collimation was determined at first in the ordinary way by inversion of the axis, measuring with the Micrometer the angular distance of the central wire from a distant mark to the North before and after inversion. In a similar manner it was determined with regard to a pair of cross wires fixed in the focus of a 20 Inch Transit Instrument situated in the southern aperture of the building, the wires being seen distinctly by looking through the object glass. By the original plan it was intended thus to obtain a mark at 180° distance from the North meridian mark whereby the position of the line of Collimation would have at once been obtained but a bad level and insecure Y's rendering the results doubtful the plan was abandoned: the following Observations are registered in the Collimation book 1831, February 18th.

Illuminating End West.	Illuminating End East.	Error of Collimation.
"	"	"
North Meridian Mark...30,30 W.....	29,48 W.....	0,41 W.
South Collimator.....13,88 E.	13,71 E.	0,08 E.

On a repetition.

"	"	"
North Meridian Mark...30,16 W.....	31,36 W.....	0,60 E.
South Collimator.....16,11 E.	13,54 E.	1,29 E.

Hence it appears that the illuminating pivot being East the central wire describes a small circle 0',40 to the West of the Meridian or great circle which the Telescope describes:

The want of agreement in these results added to the circumstance that an inversion of the axis is very seldom performed without an alteration resulting both in azimuth and level, rendered it desirable to do away with the necessity of inversion; for this purpose since the 6th March two meridian marks have been employed one situated about a mile to the North and the other about the same distance to the South of the Observatory. The angular distance between these was determined by frequent inversion of the axis from the following Observations.

	No. of Invers.	Illuminating End E.		Illuminating End W.		Collimation by	
		North mark	South mark	North mark	South mark	North mark	South mark
		"	"	"	"	"	"
1831							
March 28	5	28,59 W.	46,41 E.	21,77 W.	56,85 E.	3,41 W.	5,22 W.
29	5	28,23 —	45,22 —	18,34 —	56,79 —	4,94 —	5,78 —
April 11	3	26,35 —	44,90 —	19,84 —	52,58 —	3,26 —	3,84 —
„ 3	3	26,62 —	44,73 —	18,85 —	52,98 —	3,88 —	4,12 —
11	4	25,66 —	45,65 —	18,10 —	53,61 —	3,78 —	3,98 —
„ 3	3	25,19 —	44,78 —	18,22 —	54,07 —	3,48 —	4,65 —

∴ Now if the Instrument had not been subject to error of Collimation the following results would have been obtained.

	N. obsers.	Illuminating End E or W.		
		North mark	South mark	Difference.
		"	"	"
March 28	5	25,18 W.	51,63 E.	26,45
29	5	23,29 —	51,00 —	27,71
April 11	3	23,09 —	48,74 —	25,65
„ 3	3	22,74 —	48,85 —	26,11
14	4	21,88 —	49,63 —	27,75
„ 3	3	21,71 —	49,43 —	27,72

Giving to each result its proper weight in proportion to the number of times the Instrument was inverted we have for the angular distance between the North and South marks (reckoning to the West about) = $180^\circ + 26''{,}97$ so that if N and S represent the observed deviation of the central wire from the North and South marks respectively, $\frac{N + S - 26''{,}97}{2}$ will represent the error of Collimation.

The observations for determining the error of Collimation which are registered in the Collimation book were made every day at sun-set and sun-rise the marks being at these times seen distinctly and free from all appearance of tremor to which they are subject in the middle of the day. It is proper here to remark that the coincidence of the Micrometer and central wires was determined by bringing the former into contact with the latter on either side and moving the head of the screw until the + and — readings were equal, the accuracy of this adjustment was verified every second or third day, occasion however for readjustment has occurred but seldom, and then only to very small amount. The following results which are the mean between the Observations at Sun-set and Sun-rise have been reduced from the value of the Micrometer screw found at page 8.

1831	North mark.	South- mark.	N. + S.	N S - 26", 97"		Mean.	REMARKS.
				N	S		
June	"	"	"	"	"	"	
4	+ 22,79	- 34,86	- 12,07	+ 7,45			
5	22,79	35,13	12,16	7,41			
6	22,83	35,31	12,48	7,25			
7	23,31	35,31	12,00	7,48			
8	23,35	34,45	11,10	7,93			
9	24,41	34,72	10,31	8,33			
10	24,61	34,21	9,60	8,68			
11	25,09	34,52	9,43	8,77			
12	24,58	34,10	9,52	8,72			
13	24,78	35,55	10,77	8,10			
14	24,34	35,89	11,55	7,71			
15	24,85	35,51	10,66	8,16			
16	25,37	35,65	10,28	8,35			
17	25,02	35,48	10,46	8,25	+ 8,01		
18	23,38	36,13	12,75	7,11			
19	23,38	35,86	12,48	7,25			
20	23,38	36,47	13,09	6,94			
21	23,14	36,20	13,06	6,95			
22	23,89	35,99	12,10	7,43			
23	23,93	36,26	12,33	7,32			
24	23,62	35,72	12,10	7,43			
25	23,45	36,02	12,57	7,20			
26	22,79	36,33	13,54	6,71			
27	22,94	36,23	13,29	6,84			
28	23,21	35,21	12,00	7,48			
29	22,62	35,55	12,93	7,02			
30	23,65	36,02	12,37	7,30			
July	"	"	"	"	"	"	
1	24,06	35,21	11,15	7,91			
2	24,51	35,96	11,45	7,76			
3	22,62	34,52	11,90	7,53	+ 7,26		
4	23,82	33,62	9,80	8,58			
5	23,48	33,25	9,77	8,60			
6	24,44	34,42	9,98	8,50			
7	25,02	33,69	8,67	9,15			
8	23,68	33,28	9,60	8,68			
9	23,99	33,83	9,84	8,56			
10	23,48	33,26	10,48	8,25			
11	23,99	34,21	10,22	8,37			
12	23,72	34,69	10,97	8,00			
13	23,11	33,66	10,55	8,21			
14	24,20	33,59	9,39	8,79			
15	24,99	33,56	8,57	9,20			
16	24,58	33,22	8,64	9,16			
17	23,99	32,72	8,73	9,12			
18	22,86	32,98	10,12	8,42			
19	23,38	33,62	10,24	8,36			
20	23,38						
21	23,79	32,22	8,43	9,27			
22	23,38	32,84	9,46	8,75			
23	22,72	32,29	9,57	8,70			
24	23,31	32,94	9,63	8,67			

1831	North mark.	South mark.	N. + S.	N + S - 26',07"		Mean.	REMARKS.
				N + S	- 2		
July	25	+ 23,41	- 32,53	- 9,12	+ 8,92	+ 8,68	
	26	25,37	32,14	6,77	+ 10,10		
	27	25,85	31,95	6,10	10,43		
	28	25,26	31,95	6,69	10,14		
	29	25,95	31,16	5,21	10,88	+ 10,39	
	30	26,81	30,23	3,42	+ 11,77		
	31	27,94	30,19	2,70	12,13		
August	1	26,81	30,26	3,45	11,76		
	2	27,49	30,26	2,77	12,10		
	3	26,29	31,06	4,77	11,10		
	4	26,81	30,16	3,35	11,81		
	5	26,81	30,16	3,35	11,81		
	6	26,64	30,51	3,87	11,55		
	7	25,98	30,06	4,08	11,45		
	8	26,08	29,07	2,99	11,99		
	9	26,81	29,75	2,94	12,01		
	10	26,26	29,41	3,15	11,91		
	11	26,67					
	12	26,12	30,30	4,18	11,40		
	13	25,78	30,37	4,59	11,19		
	14	25,92	31,06	5,14	10,91		
	15	25,92				+ 11,56	Finding the errors of Azimuth and Collimation inconveniently large, I reduced them to more narrow limits.
	16	28,66	43,02	14,36	+ 6,30		
	17	29,48	42,78	13,30	6,83		
	18	28,21	42,23	14,02	6,48		
	19	28,32	42,53	14,21	6,38		
	20	28,28	42,50	14,22	6,38		
	21	27,93	42,19	14,26	6,38		
	22	27,15	41,82	14,67	6,15		
	23	27,45	41,92	14,47	6,25		
	24	28,14	41,75	13,61	6,68	+ 6,42	The fourth wire found broken.
	25	27,93	41,96	14,03	+ 6,47		
	26	27,97	42,09	14,12	6,42		
	27	27,63	42,38	14,75	6,11		
	28	26,56	42,06	15,50	5,73		
	29	25,74	42,16	16,42	5,28		
	30	26,19	41,92	15,73	5,62		
	31	26,42	42,40	15,98	5,50	+ 5,88	
September	1	26,49	43,67	17,18	+ 4,90		
	2	26,08	43,70	17,62	4,67		
	3	24,71	42,71	18,00	4,48		
	4	25,02	43,33	18,31	4,33		
	6	An extraordinary difference between the readings on the morning of the 6th, is perfectly unaccountable unless a fall of 2½ Inches of rain and a thunder storm during the night preceding had moved the foundation of the Piers, the Collimation screws were secure.

1881	North mark.	South mark.	N. + S.	N + S — 26" 97"	Mean.	REMARKS.
September 7	+ 28,45	I left Madras for Calcutta to assist in the operations of the Great Trigonometrical Survey leaving this class of observation to the care of T. Vytheanathum, the Head Assistant.
8	28,59	— 45,35	— 16,76	+ 5,10		
9	29,92	47,33	17,41	4,78		
10	30,16	46,55	16,39	5,29		
11	30,30	45,39	15,09	5,94	+ 4,94	
12	29,65	42,60	12,95	+ 7,01		
13	29,65	42,30	12,65	7,16		
14	29,58	41,97	12,39	7,29		
15	29,24	40,93	11,69	7,64	+ 7,27	
16	29,96	40,27	10,31	+ 8,33		
17	29,75	39,83	10,08	8,45		
18	29,79	39,63	9,84	8,57		
20	30,03	39,56	9,53	8,72	+ 8,52	
21	30,13	37,95	7,82	+ 9,57		
22	29,85	37,80	7,95	9,51		
23	30,16	38,15	7,99	9,49		
24	30,74	38,08	7,34	9,81		
25	30,23	38,39	8,16	9,41		
26	29,99	38,56	8,57	9,20		
27	30,92	38,73	7,81	9,58		
28	31,12	37,91	6,79	10,09		
29	30,26	38,56	8,30	9,33		
30	28,66	38,15	9,49	8,74		
October 1	30,26	37,36	7,10	9,93		
2	30,26	38,90	8,64	9,16		
3	30,37	36,68	6,31	10,33		
4	31,40	36,23	4,83	11,07		
5	30,16	36,43	6,27	10,35		
6	29,82	36,33	6,51	10,23		
7	29,65					
8	29,45	37,60	8,15	9,41		
9	29,35	38,12	8,77	9,10		
10	29,17	37,36	8,19	9,39	+ 9,67	
11	29,48					
12	29,34	40,82	11,48	+ 7,75		
13	29,48	39,69	10,21	8,38		
14	28,52	38,26	9,74	8,61		
15	28,69	38,08	9,39	8,79		
16	27,42	37,80	10,38	8,30		
17	27,90	39,22	11,32	7,82		
18	27,08	38,42	11,34	7,81		
21	28,14	38,15	10,01	8,48		
22	28,42	37,87	9,45	8,76		
23	28,11					
24	28,11	39,42	10,31	7,83		
25	28,28	39,35	11,07	7,95		

ANGULAR DISTANCE, &c. *continued.*

1831	North mark.	South mark.	N. + S.	$\frac{N + S - 26''.97}{2}$	Mean.	REMARKS.
October 27	+ 27,08	- 38,60	- 11,52	+ 7,72	"	
28	27,22	38,66	11,44	7,76		
29	27,63	38,22	10,59	8,20		
30	27,86	37,19	9,33	8,82		
31	27,05	37,70	10,65	8,16		
November 1	26,98	36,85	9,87	8,55		
2	28,72	36,78	8,06	9,45	+ 8,28	
3	28,21	36,68	8,47	9,25		
4	28,52	38,87	10,35	8,31		
5	26,39	38,49	12,10	7,43		
6	28,01	38,42	10,41	8,28		
7	27,76					
8	26,64	37,84	15,20	7,88		
9	27,59	35,99	8,40	9,28		
10	27,11	36,06	8,95	9,01		
11	26,39	36,23	9,84	8,57		
12	26,71	37,29	10,58	8,20		
13	27,39	36,58	9,19	8,89		
14	26,95	37,91	10,96	8,01		
19	26,91					
20	25,23	36,02	10,79	8,09		
21	25,47	36,54	11,07	7,95		
22	26,05	36,58	10,53	8,22		
23	26,22	37,57	11,35	7,81		
26	24,75	35,65	10,90	8,03		
27	24,34	34,25	9,91	8,53		
29		33,08				
30		34,45				
December 1		33,93	+ 8,34	
2	25,05	38,56	13,51	+ 6,73		
3	24,58	38,63	14,05	6,46		
4	21,62	38,05	16,43	5,27		
5	21,73	37,43	15,70	5,63		
6		38,12				
7	21,46	38,83	17,37	4,80		
8	22,62	38,15	15,53	5,72		
9	22,38	38,80	16,42	5,28		
10	22,18	40,10	17,92	4,53	+ 5,55	
12	23,28	27,97	4,69			
13	24,02	27,97	3,95	11,51		
14	22,97	28,28	5,31	10,83		
15	23,46	28,89	5,43	10,77		
16	24,24	28,52	4,28	11,35		
17	23,48	27,42	3,94	11,51		
18	23,48	29,28	5,80	10,58		
19	23,41	29,24	5,83	10,57		
20	23,79	29,35	5,56	10,70		
21	23,04	29,92	6,88	10,05		
22	22,62	29,14	6,52	10,22	+ 10,81	
23	21,39	30,68	9,49	8,74		
24	21,28	31,36	10,08	8,45		

1831	North mark.	South mark.	N. + S.	$\frac{N + S - 26''.97}{2}$	Mean.	REMARKS.
	"	"	"	"	"	
December 25	+ 21,62	- 30,30	- 8,68	+ 9,15		
26	23,18	29,00	5,82	10,08		
28	23,24					
29	21,52					
30	20,60	31,53	10,93	8,02		
31	19,88	30,85	10,97	8,00	+ 8,74	

In the reduction of the Observations the numbers in column *Mean* have been employed in conjunction with the numbers found at pages 6 and 7 for reducing the mean of the three central or five wires to the central wire ; with these also have been employed the correction for Diurnal Aberration, the effect of which is that to see Stars upon the Meridian the Instrument should move in a small circle parallel to the Meridian and to the eastward of it by a quantity $X \cos. Lat.$ (X being the equatorial constant of diurnal aberration) in this latitude = $0',29$, thus for the first 17 days of the month of June.

Collimation of the central wire.....	= + 8'',01
Correction for 5 wires + $0'',073$	= + 1,10
Diurnal aberration.....	= - 0,29
	<hr/>
Number employed.....	+ 8,82

The column " correction for Collimation" (see the observations) consequently contains the correction..... $+ 8'',82$

$$\frac{15 \cdot \sin. N.P.D.}{}$$

additive to observations above the Pole and subtractive for those below.

ON THE DEVIATION IN AZIMUTH.

The Deviation of the Transit Instrument from the Meridian has been found from the Observations of the two meridian marks already detailed at Pages 7 and 8 for finding the error of Collimation ; the Azimuth of these marks was

determined from observations of the Polar Star at its consecutive passages over the central wire above and below the pole; as this correction is one of great importance my attention was directed most particularly to the making of the observations, it happens however unfortunately that the Polar Star is not visible in the day time once a month in this Latitude, and *when seen*, by reason of the excessive tremor under which Stars appear in the day time the observations cannot for nice purposes be depended upon; it happens moreover very unfortunately that at the time of the year when the Pole Star passes the meridian in the morning and evening when the necessary observations could be made, the prevalence of cloudy weather peculiar to the South West monsoon renders observations very scarce, so that from these united causes in the year 1831, I was able only to obtain five pair of observations of the nature required; these observations are as follows.

Observations of the North Polar Star.

1831	h. m. s.	Corrections for			h. m. s.
		Level.	Collimation.	Clock Error.	
February 18	S.P. 12 59 52,00....	- 2,480....	+ 1,661....	- 33,85....	12 59 17,33
19	Above.. 1 0 5,00....	+ 2,280....	- 1,661....	*19,21....	0 59 46,41
June 15	S.P. 12 59 30,30....	+ 0,850....	- 18,670....	- 7,17....	12 59 5,31
15	Above.. 1 0 27,20....	- 1,090....	+ 18,670....	- 7,59....	1 0 37,19
28	S.P. 12 59 49,59....	+ 1,610....	- 16,720....	- 17,88....	12 59 16,60
28	Above.. 1 0 48,22....	- 2,050....	+ 16,720....	- 18,08....	1 0 44,81
29	S.P. 12 59 51,10....	+ 1,430....	- 16,720....	- 18,35....	12 59 17,46
29	Above.. 1 0 50,17....	- 1,800....	+ 16,720....	- 18,56....	1 0 46,53
July 5	S.P. 13 0 4,22....	+ 1,880....	- 20,270....	- 26,14....	12 59 19,69
5	Above.. 1 1 5,33....	- 2,390....	+ 20,270....	- 26,48....	1 0 56,73

If to the above be applied the error for Azimuth and the corrections for Aberration, Nutation and Precession, &c. we obtain the mean place of the Pole Star for January 1, 1831, as follows.

1831	h. m. s.	Correction for Azimuth.	Aberration, &c.	Mean A.R. Jan. 1, 1831.
February 18	12 59 17,33....	+ 2,356 a	+ 13,09....	—
19	0 59 46,41....	- 2,326 a	+ 13,39....	—
June 15	12 59 5,31....	+ 2,345 a'	- 5,24....	—
15	1 0 37,19....	- 2,315 a'	- 5,61....	—
28	12 59 16,60....	+ 2,345 a''	- 15,16....	—
28	1 0 44,80....	- 2,315 a''	- 15,54....	—
29	12 59 17,46....	+ 2,345 a'''	- 15,94....	—
29	1 0 46,53....	- 2,315 a'''	- 16,33....	—
July 5	12 59 19,69....	+ 2,345 a''''	- 20,41....	—
5	1 0 56,73....	- 2,315 a''''	- 20,79....	—

* The Clock had stoped for about fifteen seconds from some unknown cause but the result here given is in no wise affected thereby.

In the above (where, a , a' &c. are intended to represent the deviations of the Instrument from the Meridian expressed in seconds of *space*) subtracting the first line of each pair from the second we get the following.

	s.	"
February 18 and 19 . . .	29,38 = 4,682 a	or $a = 6,275$
June 15 . . .	91,51 = 4,660 a'	— $a' = 19,637$
28 . . .	87,82 = 4,660 a''	— $a'' = 18,845$
29 . . .	88,68 = 4,660 a'''	— $a''' = 19,030$
July 5 . . .	96,66 = 4,660 a''''	— $a'''' = 20,742$

Applying the above values a , a' , &c. to the observed deviation of the central wire from the North and South marks we obtain their Azimuth from the Meridian as follows.

	Observed deviation of the		Deviation of		Azimuth of	
	central wire.		central wire.		N — S.	
	from	from	from	from	from	from
	North mark.	South mark.	Meridian.	North mark.	South mark.	N — S.
	"	"	"	"	"	"
Feb. 18 and 19 . . .	30,32 E. . . .	57,29 W. . . .	6,275 . .	36,595 W. . .	63,565 E. . .	100,160
June 15 . . .	16,84 —	43,52 —	19,637 . .	36,477 — . . .	63,157 — . . .	99,634
28 . . .	15,95 —	42,47 —	18,845 . .	34,795 — . . .	61,315 — . . .	96,110
29 . . .	15,36 —	42,81 —	19,030 . .	34,390 — . . .	61,840 — . . .	96,230
July 5 . . .	14,81 —	41,92 —	20,742 . .	35,552 — . . .	62,662 — . . .	98,214
			Mean =	35,562 W. . .	62,508 E. . .	98,070

These results at first sight appear very unsatisfactory, but on further inspection it will be found that errors of four or five tenths of a second of space in the values of the error of collimation will altogether reconcile the disagreement: that the error does arrive from this cause, or rather that it is not chargeable on the observations will be made clear on taking the mean of the results above and below the pole, where an agreement is obtained equal to any thing which the nature of the observation will permit.

Mean A.R. January 1, 1831.

<i>h.</i>	<i>m.</i>	<i>s.</i>		<i>s.</i>		<i>h.</i>	<i>m.</i>	<i>s.</i>
0	59	45,11	+ 0,015 a	=	0	59	45,20
		45,82	+ 0,015 a'				46,11
		45,35	+ 0,015 a''				45,63
		45,86	+ 0,015 a'''				46,15
		45,10	+ 0,015 a''''				45,41

The Mean of these $0^h. 59^m. 45,70^s.$ differs nearly two seconds from the Greenwich determination notwithstanding which from the care bestowed upon

the observations and computations (which were made by myself) I have no sort of doubt of its being an accurate determination and in employing it in preference to any other in the computations which now follow, I do it from a consciousness of its being accurate on the one hand, and on the other hand from a determination which I have formed of calling for assistance upon other observations for the sake of reducing my own as seldom as possible, and then only when their use cannot be dispensed with. But to return to our subject let a and a' represent the observed transits of any two Stars corrected for error of Level and Collimation, A , A' , their true apparent Right Ascensions which in the present instance are those given in the Greenwich approximate Catalogue of 720 Stars (with the above exception,) reduced by the numbers given in the Astronomical Society's Catalogue ;

putting c and c' for the terms $\frac{\cos. \text{Altitude}}{\sin. \text{N.P.D.}}$ and x for the error in Azimuth we have $a' - a + (c' - c) x = A' - A$ (supposing the clock keeps sidereal time, otherwise its rate must be allowed for) whence x in seconds of space = $\frac{(A' - A) - (a' - a)}{15 (c' - c)}$ pursuing this course with the Polar Star

(which affords in this latitude a large coefficient) and Stars situated near to the Zenith, we obtain values of x which applied to the corresponding deviations of the central wire from the North and South marks give their Azimuth as follows.

Determination of the Azimuth of the North and South Meridian marks from a comparison of the observed transit of the Polar Star below the Pole with

		NAMES OF STARS.		Deviation of the central wire from		Azimuth of	
				North mark.	South mark.	North mark.	South mark.
1831			"			"	"
February	19	* α Andromedæ.....	$x = 6,10$				
		α Arietis.....	$= 6,35$				
	20	α Cancræ.....	$= 4,39$				
	27	β Virginis.....	$= 6,80$				
		α Aquilæ.....	$= 5,61$				
	28	Procyon.....	$= 4,08$				
March	2	ζ Hydræ.....	$= 7,11$				
		δ Virginis.....	$= 7,15$				
	9	α Hydræ.....	$= 9,94$	25,76	52,74	35,70	62,68
		α Lyræ.....	$= 9,90$	35,66	62,64
	13	π Virginis.....	$= 9,76$	24,84	51,80	34,60	61,56
	17	Regulus.....	$= 10,57$	24,59	51,57	35,16	62,14
		α Lyræ.....	$= 10,41$	35,00	61,98

* Polaris observed above the Pole.

Comparison of the observed transit of *Polaris* with

		NAMES OF STARS.	Deviation of the central wire from		Azimuth of	
			North mark.	South mark.	North mark.	South mark.
1831			North	South	"	"
March	18	γ Leonis..... $x = 10,61$	23,41	50,39	34,02	61,00
		a Lyræ..... $= 10,48$	33,89	60,87
	19	a Lyræ..... $= 11,86$	22,28	49,25	34,14	61,11
	21	ϵ Leonis..... $= 10,47$	23,39	50,35	33,86	60,82
		ζ Aquilæ..... $= 10,24$	33,63	60,59
	22	a Hydræ..... $= 12,91$	22,85	49,82	35,76	62,73
		a Aquilæ..... $= 12,75$	35,60	62,57
	23	χ Leonis..... $= 13,85$	22,73	49,71	36,58	63,56
		a Aquilæ..... $= 13,67$	36,40	63,38
	25	σ Leonis..... $= 13,09$	22,53	49,50	35,62	62,59
		δ Aquilæ..... $= 12,98$	35,51	62,48
	26	γ Aquilæ..... $= 13,46$	23,05	50,03	36,51	63,49
	27	γ Virginis..... $= 10,88$	24,19	51,17	35,07	62,05
		γ Aquilæ..... $= 10,74$	24,19	51,17	34,93	61,91
	28	σ Leonis..... $= 14,24$	24,25	51,21	38,49	65,46
		ζ Aquilæ..... $= 13,87$	38,12	65,09
	29	a Virginis..... $= 14,17$	23,19	50,16	37,36	64,33
	30	σ Leonis..... $= 13,94$	23,23	37,17	64,14
		λ Virginis..... $= 13,40$	36,63	63,60
April	31	ϵ Pegasi..... $= 13,07$
	3	ϵ Leonis..... $= 15,70$	23,24	50,21	38,94	65,91
		a Aquilæ..... $= 15,06$	38,30	65,27
	4	ϵ Leonis..... $= 15,22$	22,58	49,54	37,80	64,76
	5	β Leonis..... $= 14,14$	27,21	49,19	36,35	63,33
		γ Aquilæ..... $= 14,07$	36,28	63,26
	6	σ Leonis Minoris..... $= 14,50$	21,61	48,57	36,11	63,07
		γ Aquilæ..... $= 14,37$	35,98	62,94
	11	σ Virginis..... $= 11,75$	22,04	49,00	33,79	60,75
		γ Aquilæ..... $= 11,32$	33,36	60,32
	12	β Corvi..... $= 12,98$	22,53	49,51	35,51	62,49
		γ Aquilæ..... $= 13,12$	35,65	62,63
	14	β Virginis..... $= 13,88$	22,02	48,98	35,90	62,86
	15	γ Aquilæ..... $= 14,18$	20,04	47,00	34,22	61,18
	16	a^2 Capricorni..... $= 13,05$	19,98	46,78	32,87	59,83
	17	γ Aquilæ..... $= 12,91$	20,68	47,65	33,59	60,56
	18	a Aquilæ..... $= 12,58$	20,39	47,36	32,97	59,94
	19	γ Aquilæ..... $= 12,70$	20,96	47,93	33,66	60,63
	20	γ Aquilæ..... $= 14,07$	20,17	47,15	34,24	61,22
	21	54 Leonis..... $= 16,22$	19,68	46,64	35,90	62,86
	22	γ Comæ Berenicis..... $= 14,20$	20,32	47,30	34,52	61,50
	23	γ Comæ Berenicis..... $= 12,60$	20,94	47,90	33,54	60,50
	24	π Virginis..... $= 12,11$	21,06	48,02	33,17	60,13
	25	m Virginis..... $= 10,50$	21,52	48,48	32,02	58,98
	26	π Virginis..... $= 14,23$	21,80	48,79	36,03	63,02
		μ Virginis..... $= 14,36$	36,16	63,15
	27	δ Virginis..... $= 14,92$	22,35	49,32	37,27	64,24
		a Virginis..... $= 14,86$	37,21	64,18
	28	a Virginis..... $= 14,87$	21,82	48,79	36,69	63,66
May	1	ϵ Virginis..... $= 13,94$	22,90	49,87	36,84	63,81
		a Virginis..... $= 14,02$	36,92	63,89
	4	β Corvi..... $= 16,06$	22,62	49,58	38,68	65,64
		a Virginis..... $= 16,02$	38,64
	5	ϵ Corvi..... $= 13,35$	21,96	48,92	35,31	62,27
		a Virginis..... $= 13,39$	35,35	62,31

Comparison of the observed transit of *Polaris* with

		NAMES OF STARS.	Deviation of the central wire from		Azimuth of			
			North mark.	South mark.	North mark.	South mark.		
1831		"			"	"		
May	6	<i>a</i> Comæ Ber.....	$x = 15,02$	20,40	47,37	35,42	62,39	
	13	γ Comæ Bor.....	$= 16,56$	19,49	46,46	36,05	63,02	
		<i>a</i> Virginis.....	$= 16,52$	36,01	62,98	
	14	δ Virginis.....	$= 17,00$	19,15	46,12	36,15	63,12	
		<i>a</i> Virginis.....	$= 17,04$	36,19	63,16	
	15	61 Virginis.....	$= 16,03$	19,65	46,62	35,68	62,65	
	16	η Corvi.....	$= 15,65$	20,72	47,69	36,37	63,34	
		ζ Virginis.....	$= 15,55$	36,27	63,24	
	18	γ Corvi.....	$= 14,56$	20,33	47,29	34,89	61,85	
	20	γ Comæ Bor.....	$= 14,26$	20,23	47,20	34,49	61,46	
	21	δ Virginis.....	$= 14,42$	19,57	46,54	33,99	60,96	
	22	<i>m</i> Virginis.....	$= 15,07$	20,03	47,00	35,10	62,07	
	23	λ Virginis.....	$= 14,97$	20,14	47,11	35,11	62,08	
	24	γ Comæ Bor.....	$= 15,97$	19,12	46,09	35,09	62,06	
		<i>v</i> Bootis.....	$= 15,87$	34,99	61,96	
	25	γ Corvi.....	$= 16,05$	20,34	47,31	36,39	63,36	
		<i>i</i> Virginis.....	$= 16,28$	36,62	63,59	
	26	<i>a</i> Canum Ven.....	$= 15,86$	17,50	44,47	33,36	60,33	
		<i>i</i> Virginis.....	$= 15,86$	33,36	60,33	
	28	γ Virginis.....	$= 15,38$	16,59	43,56	31,97	58,94	
		η Bootis.....	$= 15,18$	31,77	58,94	
	June	3	<i>v</i> Leonis.....	$= 19,59$	15,14	42,10	34,73	61,69
		4	κ Virginis.....	$= 19,42$	15,34	42,31	34,76	61,73
5		η Virginis.....	$= 19,89$	15,38	42,54	35,27	62,43	
		δ Libræ.....	$= 20,20$	35,58	62,74	
11		ζ Bootis.....	$= 18,48$	16,32	43,29	34,80	61,77	
15		<i>a</i> Comæ Ber.....	$= 19,39$	16,69	43,67	36,08	63,06	
		ζ Bootis.....	$= 19,09$	35,78	62,76	
		* <i>a</i> Cassiopeæ.....	$= 19,60$	36,29	63,27	
19		<i>a</i> Cassiopeæ.....	$= 19,20$	16,13	43,11	35,33	62,31	
28		<i>m</i> Virginis.....	$= 18,77$	15,73	42,69	34,50	61,46	
		* γ Pegasi.....	$= 18,64$	34,37	61,33	
		* <i>a</i> Arietis.....	$= 19,40$	35,13	62,09	
29		<i>i</i> Virginis.....	$= 18,97$	15,60	42,57	34,57	61,54	
	* γ Pegasi.....	$= 18,81$	34,41	61,38		
	* <i>a</i> Arietis.....	$= 19,11$	34,71	61,68		
July	1	* <i>a</i> Arietis.....	$= 19,99$	16,15	43,12	36,14	63,12	
	5	<i>a</i> ^a Libræ.....	$= 19,99$	14,88	41,85	34,87	61,84	
September	21	* <i>e</i> Ophiuchi.....	$= 20,25$	35,13	62,10	
		* Procyon.....	$= 12,47$	20,56	47,52	33,03	59,99	
	22	* <i>o</i> Sayitt.....	$= 14,44$	20,34	47,31	34,78	61,75	
	29	* ζ Cassiopeæ.....	$= 14,67$	20,39	47,89	35,60	62,56	
October	30	* γ Cassiopeæ.....	$= 14,68$	19,92	46,89	34,60	61,57	
	2	* ζ Cassiopeæ.....	$= 14,68$	21,10	48,96	35,78	62,74	
	4	* μ Andromedæ.....	$= 12,76$	20,33	47,30	33,09	60,06	
	6	* ζ Andromedæ.....	$= 14,74$	19,59	46,56	34,33	61,30	
	14	* <i>e</i> Piscium.....	$= 15,21$	19,91	46,87	35,12	62,08	
	20	* <i>i</i> Ceti.....	$= 14,28$	19,46	46,43	33,74	60,71	
	22	* Piscium.....	$= 14,51$	19,66	46,63	34,17	61,14	
	23	* <i>e</i> Piscium.....	$= 14,53$	19,81	46,78	34,34	61,31	
	24	* ζ Andromedæ.....	$= 14,25$	20,28	47,25	34,53	61,50	
	25	* <i>e</i> Piscium.....	$= 14,36$	20,33	47,30	34,69	61,66	
	31	* ζ Andromedæ.....	$= 16,15$	18,89	45,86	35,04	62,01	
November	1	* τ Andromedæ.....	$= 15,22$	18,43	45,40	33,65	50,62	

ON THE DEVIATION IN AZIMUTH.

Comparison of the observed transit of Polaris with

	NAMES OF STARS.		Deviation of the central wire from		Azimuth of	
			North mark.	South mark.	North mark.	South mark.
1831		"			"	"
November	2*	* μ Andromedæ..... $x = 14,21$	19,27	46,23	33,48	60,43
	4*	* β Ceti..... $= 15,06$	20,21	47,18	35,27	62,21
	5*	* β Ceti..... $= 15,27$	18,96	45,92	34,23	61,19
	6*	* β Ceti..... $= 15,27$	19,73	46,70	35,00	61,97
	8*	* β Ceti..... $= 15,30$	18,76	45,72	34,08	61,02
	19*	* ϵ Andromedæ..... $= 15,19$	18,86	45,83	34,05	61,02
December	21*	* δ Andromedæ..... $= 18,61$	17,52	44,49	36,13	63,10
	27*	* δ Andromedæ..... $= 19,42$	15,81	42,78	35,23	62,20
	3*	* ν Andromedæ..... $= 20,39$	18,12	45,09	38,51	65,48
	5*	* ν Andromedæ..... $= 16,71$	16,10	43,06	32,81	59,77
	8*	* μ Piscium..... $= 22,07$	16,90	43,87	38,07	65,94
	12*	* ν Andromedæ..... $= 24,53$	12,47	38,78	37,00	63,31
	14*	* ϵ Ceti..... $= 25,46$	12,14	39,11	37,00	64,57
	16*	* α Piscium..... $= 25,74$	12,89	39,67	38,63	65,61
	21*	* ζ Arietis..... $= 25,83$	12,99	39,97	38,82	65,80
	24*	* τ Ceti..... $= 21,23$	12,83	39,81	34,06	61,04

Taking the mean of 128 Observations we get
 the Azimuth of the North mark = 35",34 West of the Meridian
 South mark = 62",31 East of the Meridian
 differing only two tenths of a second from the numbers found at page 22.
 Now if N and S represent the deviation of the central wire from the North and South marks already given (page 14 et seq.) and C the error of Collimation we

$$\text{have } x = 35",34 \pm C - N.$$

$$x = 62",31 \mp C - S.$$

taking half the sum of these

$$x = \frac{97",65 - N - S}{2}$$

employing this with the values of N and S above named we get the

Deviation of the Transit Instrument in Azimuth.

	N - S.	97",65 - N - S. 2	REMARKS.		N - S.	97",65 - N - S. 2	REMARKS.
1831	"	"		1831	"	"	
March	6	77,40	10,12		March 10	78,43	9,61
	7	77,49	10,08		11	75,30	11,17
	8	80,38	8,63		12	76,43	10,61
	9	78,50	9,57		13	76,64	10,50

1831			REMARKS.	1831			REMARKS.
	N — S.	$\frac{97''_{65} - N - S}{2}$			N — S.	$\frac{97''_{65} - N - S}{2}$	
1831	"	"		1831	"	"	
March 14	75,80	10,92		May 3	72,05	12,66	
15	75,31	11,17	= 10'',23	4	72,20	12,72	
16	74,82	11,41	New lines.	5	70,88	13,38	
17	76,16	10,75		6	67,77	14,94	= 13'',42 wires broken
18	73,80	11,92		10	68,21	14,72	
19	71,53	13,06		11	67,85	14,90	
20	74,38	12,63		12	68,87	14,39	
21	73,74	11,96		13	65,95	15,85	
22	72,67	12,49	= 12'',03	14	65,27	16,19	
23	72,44	12,60	New lines.	16	68,41	14,62	
24	74,15	11,75		17	67,73	14,96	
25	72,03	12,81		18	67,62	15,01	= 15'',00
26	73,08	12,28		19	69,00	14,32	
27	75,36	11,15	= 12'',12 Instrument inverted.	26	61,97	17,84	
28	75,46	11,09	Instrument inverted.	27	60,33	18,66	
29	73,35	12,15		28	60,15	18,75	
30	= 11'',62	29	61,11	18,27	
April 2	72,29	12,68		30	60,91	18,37	
3	73,45	12,10		31	60,27	18,69	= 18'',55
4	72,12	12,76		June 1	59,19	19,23	
5	71,40	13,12		2	58,22	19,71	
6	70,18	13,73		3	57,24	20,20	
7	71,40	13,12	= 12'',92	4	57,65	20,00	
8	70,86	13,38		5	58,10	19,78	
9	70,81	13,42		6	58,14	19,75	
10	71,26	13,20		7	58,62	19,51	
11	71,04	13,30		8	57,80	19,93	
12	72,04	12,80		9	59,13	19,26	
13	69,71	13,97	= 13'',34 Instrument inverted.	10	58,82	19,42	
14	71,00	13,32		11	59,61	19,02	
15	67,04	15,31		12	58,68	19,49	= 19'',60
16	66,60	15,52		13	60,33	18,66	
17	68,33	14,66		14	60,23	18,71	
18	67,75	14,95		15	60,36	18,65	
19	68,89	14,38		16	61,02	18,31	
20	67,32	15,16	= 14'',76 wires broken	17	60,50	18,57	
21	66,32	15,66	Wires broken.	18	59,51	19,07	
22	67,62	15,03		19	59,24	19,20	
23	68,84	14,41		20	59,85	18,90	
24	69,08	14,28		21	59,34	19,15	
25	70,00	13,82		22	59,88	18,89	
26	70,59	13,53		23	60,19	18,73	= 18'',80
27	71,67	12,99		24	59,34	19,15	
28	70,61	13,52		25	59,47	19,09	
30	68,93	14,36	= 13'',99 wires broken	26	59,12	19,26	
				27	59,17	19,24	
				28	58,42	19,61	
				29	58,17	19,74	

ON THE DEVIATION IN AZIMUTH.

1831			REMARKS.	1831			REMARKS.	
	N - S.	$\frac{97''_{65} - N - S}{2}$			N - S.	$\frac{97''_{65} - N - S}{2}$		
June 30	59,67	18,99	= 19'',66	Aug. 21	70,12	13,76	= 13'',58	
July 1	59,27	19,19		22	68,97	14,34		
2	60,47	18,59		23	69,37	14,14		
3	57,14	20,25		24	69,89	13,88		
4	57,44	20,10		25	69,89	13,88		4th wire broken.
5	56,73	20,46		26	70,06	13,80		
6	58,86	19,35		27	70,01	13,82		
7	58,71	19,47		28	68,62	14,52		
8	56,96	20,33		29	67,90	14,87		
9	57,82	19,91		30	68,11	14,77		
10	56,74	20,45		31	68,82	14,41		
11	58,20	19,74		Sept. 1	70,16	13,75		
12	58,41	19,62		2	69,78	13,93		
13	56,77	20,44		3	67,42	15,11		
14	57,79	19,93		4	68,35	14,65		= 14'',32
15	58,55	19,55		7	73,80	11,92		This alteration unaccountable.
16	57,80	19,92		8	73,94	11,85		
17	56,71	20,47		9	77,25	10,20		
18	55,84	20,90		10	76,71	10,47		
19	57,00	20,32		11	75,69	10,98		= 11'',09
20	56,30	20,67		12	72,25	12,70		
21	56,01	20,82		13	71,95	12,85		
22	56,22	20,72		14	71,55	13,05		= 12'',87
23	55,01	21,32		15	70,17	13,74		
24	56,25	20,70		16	70,23	13,71		
25	55,94	20,85		17	69,58	14,03		
26	57,51	20,07		18	69,42	14,11		
27	57,80	19,92		20	69,59	14,03		= 13'',92
28	57,21	20,22		21	68,08	14,78		
29	57,11	20,27		22	67,65	15,00		
30	57,04	20,31		23	68,31	14,67		
Aug. 31	58,13	19,76	24	68,82	14,41			
1	57,07	20,29	25	68,62	14,51			
2	57,75	19,95	26	68,55	14,55			
3	57,35	20,15	27	69,65	14,00			
4	56,97	20,34	28	69,03	14,31			
5	56,97	20,34	29	68,82	14,41			
6	57,15	20,25	30	66,81	15,42			
7	56,04	20,80	Oct. 1	67,62	15,01			
8	55,15	21,25	2	69,16	14,25			
9	56,56	20,54	3	67,05	15,30			
10	55,67	20,99	4	67,63	15,01			
11	56,52	20,56	5	66,59	15,53			
12	56,42	20,61	6	66,15	15,75			
13	56,15	20,75	7	66,61	15,52			
14	56,98	20,33	8	67,05	15,30			
15	56,98	20,33	9	67,47	15,09			
16	71,68	12,98	10	66,53	15,56	= 14'',92		
17	72,26	12,69						
18	70,44	13,60						
19	70,85	13,40						
20	70,78	13,44						

The readings were reduced.

1831			REMARKS.	1831			REMARKS.
	N—S.	$\frac{97''\ 65'' - N - S}{2}$			N—S.	$\frac{97''\ 65'' - N - S}{2}$	
1831	"	"		1831	"	"	
Oct. 11	—	—		Nov. 20	61,25	18,20	
12	70,16	13,75		21	62,01	17,82	
13	69,17	14,24		22	62,63	17,51	
14	66,78	15,43		23	63,79	16,93	
15	66,77	15,44		26	60,40	18,62	
16	65,22	16,21		27	58,59	19,53	
17	67,12	15,26		Dec. 2	63,61	17,02	
18	65,50	16,07		3	63,21	17,22	
21	66,29	15,68		4	59,67	18,99	
22	66,29	15,68		5	59,16	19,25	
23	67,75	14,95		6	59,72	18,96	
24	67,53	15,06		7	60,29	18,68	
25	67,63	15,01		8	60,77	18,44	
27	65,68	15,98		9	61,18	18,23	
28	65,88	15,89		10	62,28	17,68	= 18'',21
29	65,85	15,90	= 15'',37	12	51,25	23,20	
30	65,05	16,30		13	51,99	22,83	
31	64,75	16,45		14	51,25	23,20	
Nov. 1	63,83	16,91		15	52,35	22,65	
2	65,50	16,08		16	52,76	22,45	
3	64,89	16,38		17	50,90	23,37	
4	67,39	15,13		18	52,76	22,45	
5	64,88	16,38		19	52,65	22,50	
6	66,43	15,61		20	53,14	22,25	
7	65,89	15,89		21	52,96	22,35	
8	64,48	16,58		22	51,76	22,95	
9	63,58	17,03		23	52,27	22,69	
10	63,17	17,24		24	52,64	22,50	
11	62,62	17,51		25	51,92	22,86	
12	64,00	16,82		26	52,18	22,73	
13	63,97	16,84		27	—	—	
14	64,86	16,40	= 16'',47	30	52,13	22,76	
				31	50,73	23,46	= 22'',78

The column "correction for Azimuth" is computed from the mean values of x set down above in the column of Remarks, where cor. in Azim. = $\frac{\cos. \text{Altitude } x}{15 \sin. \text{N.P.D.}}$ which (the Deviation being to the *West* of the North) is subtractive for observations between the Zenith and North Pole and additive in all other cases.

ON THE REDUCTIONS EMPLOYED.

The reduction of eight or ten thousand Observations which the books of this Observatory presented for the year 1831, was an undertaking attended

with considerable labor and anxiety, especially as the Native Assistants being very liable to blunder, I could expect from them but little assistance, at least I have thought it proper and found it necessary either to examine carefully every computation made by the Native Assistants or to procure from them a duplicate computation; with this in view I very gladly availed myself of those excellent tables by Mr. Baily, printed in the II Vol. of the Royal Astronomical Society's Memoirs, without which valuable assistance I could not possibly have undertaken the reduction of my observations; the formulæ from which these tables were constructed are.

$$\begin{aligned}
 A &= -18'',677 \cos. \odot \\
 B &= -20'',360 \sin. \odot \\
 C &= t - ,025 \sin. 2 \odot - 0'',344 \sin. \Omega + 0'',004 \sin. 2 \Omega \\
 D &= -0'',545 \cos. 2 \odot - 9'',250 \cos. \Omega + 0'',090 \cos. 2 \Omega \\
 a &= + \cos. a \sin. \delta & a' &= + \tan w \cos. \delta - \sin. a \sin. \delta \\
 b &= + \sin. a \sin. \delta & b' &= + \cos. a \sin. \delta \\
 c &= + 46'',021 + 20'',043 \sin. a \tan \delta & c' &= + 20'',043 \cos. a \\
 d &= + \cos. a \tan \delta & d' &= - \sin. a
 \end{aligned}$$

The total corrections for aberration precession and nutation.

$$\begin{aligned}
 \text{In. A.R.} &= a A + b B + c C + d D \\
 \text{In. N.P.D.} &= a' A + b' B + c' C + d' D
 \end{aligned}$$

In computing the quantities A and B which comprehend the correction for aberration, I have substituted for 20'',36 the more recently determined value 20'',50 from the Greenwich Observations so that we have $A = -18'',802 \cos. \odot$ and $B = -20'',500 \sin. \odot$ in all other respects I have strictly adhered to the tables except in one or two cases where errors had been committed.* The computations are made for 8h. 30m. mean time of each day (being about the middle of the times of observing) except for Polaris and δ Ursæ Minoris for which the computations have been adapted to the time of their passing the Meridian.

* *Errors in the Ast. Soc. Tables.*

No. 767	for $b' = -7,6504$	read $b' = +7,6504$
795	— $a' = 0,0000$	— $a' = -6,5682$
810	— $d = -7,6067$	— $d = +7,6067$
1167	— $b = +9,8912$	— $b = +8,8912$

ON THE CLOCK ERRORS AND CLOCK RATES.

As I have before stated, the greater part of the observations have been made by the Native Assistants during my residence at Madras, and were altogether made by them during my absence for five months on other duties in Bengal so that it will be proper to put the reader in possession of some of the early rough observations made by these Assistants in order that a confidence proportional to their merit may be established, accordingly the observations on two consecutive days as here annexed are given as much by way of specimen for this purpose as to furnish an example of the manner in which the Clock Error and Rates have been determined.

It will be proper to premise that the Assistants had been accustomed to observing transits with the Mural Circle for one month previously to the Transit Instrument being erected, during this time their progress being equal to what I could have expected I did not hesitate to commence observing with their assistance.

The names of the observers whose initials are attached to their respective observations are M for Mootosawny Moodeliar; A for Anuntacharyer Braminy and T for my own observations.

Observed Transits of the Sun Planets and fixed Stars over the Meridian of the Madras Observatory, extracted from the Transit book.

	I	II	III	IV	V	Mean.	Corrections for			Transit over the Meridian.	NAMES.	Observed by	
							Level	Azimuth	Coll.				
							+		+				
1831	s.	s.	m. s.	s.	s.	m. s.	s.	s.	s.	h. m. s.			
March 18	23,7	51,2	49 18,9	46,7	13,9	49 18,88	,235	+	197	,225	23 49 19,54	☉ 2 L.	M.
		59,0	3 38,9	17,7	57,6	3 38,51	,296	—	624	,321	5 3 38,51	a Aurigæ	
	52,8	20,3	22 47,9	15,0	42,9	22 47,78	,238	+	188	,225	5 22 48,43	δ Orionis	
L 3",65 W.	44,1	11,9	31 38,7	6,7	34,3	31 39,14	,234	+	208	,225	5 31 39,81	ξ Orionis	
	50,8	20,0	36 50,1	20,1	49,6	36 50,12	,213	+	504	,244	5 36 51,08	γ Leporis	
A 12",03 W.	16,2	54,6	46 33,7	12,8	51,9	46 33,84	,296	—	600	,318	5 46 33,85	β Aurigæ	
C 3",66 E.		24,0	50 57,8	31,4	5,0	50 57,65	,198	+	732	,276	5 50 58,86	γ Columbæ	
	24,0	52,8	57 21,0	49,9	17,8	57 21,10	,252	+	048	,228	5 57 21,63	ν Orionis	
	59,9	34,6	1 9,0	43,6	17,2	1 8,86	,198	+	773	,283	6 1 10,11	θ Columbæ	
	7,0	34,8	6 2,4	29,9	57,6	6 2,34	,230	+	266	,226	6 6 3,06	α Monocer	
	49,0	22,9	9 57,5	30,0	4,2	9 56,72	,198	+	728	,275	6 9 57,92	κ Columbæ	

		App. A.R. + Clock Er.	Aber- ration, &c.	Mean A.R. Jan. 1, 1831 + Clock Er.	Greenwich Mean A.R. Jan. 1, 1831.	Difference or Clock Error.	Error computed from Mean.	Mean Places Jan. 1, 1831.	Mean Errors of Clock.
1831		<i>h. m. s.</i>	<i>s.</i>	<i>h. m. s.</i>	<i>h. m. s.</i>	<i>s.</i>	<i>s.</i>	<i>h. m. s.</i>	
March	18	23 49 19,54					- 34,65		
M.	α Aurigæ.....	5 3 38,51	+ 0,05	5 3 38,56	5 4 13,02	- 34,46	34,32	5 4 12,88	
	δ Orionis.....	5 22 48,43	+ 0,04	5 22 48,47	5 23 22,63	34,16	34,30	5 23 22,77	
Daily Rate	ζ Orionis.....	5 31 39,81	- 0,01	5 31 39,80	5 32 14,08	34,28	34,29	5 32 14,09	M. 6h. 17m.
of Clock.	γ Leporis.....	5 36 51,08	- 0,00	5 36 51,08	5 37 25,17	34,09	34,29	5 37 25,37	
M. + 1",33	β Aurigæ.....	5 46 33,85	- 0,26	5 46 33,59	5 47 8,01	34,42	34,28	5 47 7,87	17 - 34",25
A. + 1",59	ν Columbae.....	5 50 58,86	- 0,08	5 50 58,78	5 51 32,74	33,96	34,27	5 51 33,05	A. 9h. 34m.
T. + 1",46	υ Orionis.....	5 57 21,63	- 0,16	5 57 21,47	5 57 55,43	33,96	34,27	5 57 55,74	
Mean.	θ Columbae.....	6 1 10,11	- 0,14	6 1 9,97			34,27	6 1 44,24	7 - 34",12
= + 1",46	α Monocerotis...	6 6 3,06	- 0,19	6 6 2,87	6 6 36,87	34,00	34,26	6 6 37,13	
	ε Columbae.....	6 9 57,92	- 0,26	6 9 57,66	6 10 32,61	34,93	34,26	6 10 31,94	
	ζ Canis Major...	6 13 15,78	- 0,27	6 13 15,51	6 13 49,71	34,20	34,25	6 13 49,76	T. 19h. 13m.
	ν Geminorum...	6 18 21,97	- 0,29	6 18 21,68	6 18 55,61	33,93	34,25	6 18 55,93	8 - 33",83
	δ Urs Min.....	6 26 7,68	+ 3,22	6 26 10,90	6 26 49,37*	38,47	34,24	6 26 45,14	
	55 Aurigæ.....	6 30 12,96	- 0,40	6 30 12,56			34,24	6 30 46,80	
	α Canis Maj....	6 36 7,87	- 0,41	6 36 7,88	6 37 41,91	34,03	34,23	6 36 42,11	
	χ Arg in pup....	6 41 0,86	- 0,50	6 41 0,86			34,23	6 41 34,59	
	Canis Maj....	6 44 10,08	- 0,51	6 44 9,57			34,23	6 44 43,80	
	ζ Geminor.....	6 53 31,35	- 0,45	6 53 30,90	6 54 4,86	33,96	34,22	6 54 5,12	
	63 Aurigæ.....	6 59 27,63	- 0,59	6 59 27,04			34,21	6 59 27,25	
	64 Aurigæ.....	7 5 42,88	- 0,57	7 5 42,31			34,20	7 6 16,51	
	60 Geminor.....	7 14 39,80	- 0,56	7 14 39,24	7 15 13,30	34,06	34,19	7 15 13,43	
	α Geminor.....	7 23 14,22	- 0,61	7 23 13,61	7 23 48,33	34,72	34,18	7 23 47,79	
	α Canis Min....	7 29 53,09	- 0,62	7 29 52,47	7 30 27,20	34,73	34,17	7 30 26,64	
	β Geminor.....	7 34 24,26	- 0,65	7 34 23,61	7 34 57,91	34,30	34,17	7 34 57,78	
	c Arg in pup....	7 38 40,54	- 0,96	7 38 39,58			34,17	7 39 13,75	
A.	γ ¹ Argus.....	8 3 44,82	- 1,35	8 3 43,47			34,21	8 3 43,68	
	γ ² Argus.....	8 3 47,02	- 1,35	8 3 45,67			34,21	8 3 45,88	
	q Arg in pup...	8 11 40,35	- 1,19	8 11 40,16			34,20	8 12 14,36	
	η Piscis Vol....	8 23 0,52	- 3,61	8 22 56,91			34,19	8 23 31,10	
	δ Hydræ.....	8 28 9,23	- 0,89	8 28 8,34	8 28 42,29	33,05	34,18	8 28 42,52	
	β Pixid Naut....	8 33 56,35	- 1,29	8 33 55,06			34,18	8 34 39,24	
	α Arg in Vel....	8 39 45,58	- 1,61	8 39 43,97			34,17	8 40 18,14	
	ζ Hydræ.....	8 45 54,50	- 0,95	8 45 53,55	8 46 27,34	33,79	34,16	8 46 27,71	
	Lyncis.....	8 55 12,33	- 1,03	8 55 11,30			34,15	8 55 45,45	
	c Arg in Vel....	8 57 47,28	- 1,78	8 57 45,50			34,15	8 58 19,65	
	λ Argus.....	9 1 14,75	- 1,68	9 1 13,07			34,15	9 1 47,22	
	μ Arg in Car...	9 5 59,54	- 2,45	9 5 57,09			34,14	9 6 31,23	
	ι Argus.....	9 12 2,71	- 2,53	9 12 0,18			34,14	9 12 34,32	
	α Hydræ.....	9 18 43,22	- 1,15	9 18 42,07	9 19 17,02	34,95	34,13	9 19 16,20	
	U ¹ Hydræ.....	9 42 48,37	- 1,31	9 42 47,06			34,11	9 43 21,17	
	π Leonis.....	9 50 43,71	- 1,14	9 50 42,57	9 51 16,72	34,15	34,10	9 51 16,67	
	η Centrum.....	9 55 28,62					34,10		
	Regulus.....	9 58 49,38	- 1,23	9 58 48,15	9 59 21,93	33,78	34,09	9 59 22,24	
	γ Leonis.....	10 10 5,59	- 1,16	10 10 4,43	10 10 38,66	34,23	34,08	10 10 38,51	
	γ Arg in Vel....	10 14 33,74	- 1,98	10 14 31,76			34,08	10 15 5,84	
	ρ Leonis.....	10 23 21,69	- 1,23	10 23 20,46	10 23 54,44	33,98	34,07	10 23 54,53	
	Polaris SP... ..	12 58 50,25	+ 25,18	12 59 15,43	12 59 45,70*	30,27	33,92	12 59 49,35	
T.	α Lyrae.....	18 30 39,08	+ 0,18	18 30 39,26	18 31 13,11	33,85	33,87	18 31 13,13	
	β Lyrae.....	18 43 16,57	+ 0,22	18 43 16,79	18 43 50,58	33,79	33,86	18 43 50,65	
	ζ Aquilæ.....	18 57 4,44	+ 0,26	18 57 4,70	18 57 38,61	33,91	33,84	18 57 38,54	
	δ Draconis.....	19 11 55,06	+ 0,80	19 11 55,86	19 12 29,69	33,83	33,83	19 12 29,69	
	δ Aquilæ.....	19 16 24,54	+ 0,36	19 16 24,90	19 16 58,58	33,68	33,83	19 16 58,73	
	γ Aquilæ.....	19 37 39,31	+ 0,45	19 37 39,76	19 38 13,55	33,79	33,80	19 38 13,56	
	α Aquilæ.....	19 41 57,98	+ 0,48	19 41 58,46	19 42 32,29	33,83	33,80	19 42 32,26	
	β Aquilæ.....	19 46 26,35	+ 0,50	19 46 26,85	19 47 0,79	33,94	33,79	19 47 0,64	

	NAMES.	App. A.R. + Clock Er.	Aber- ration, &c.	Mean A.R. Jan. 1, 1831 + Clock Er.	Greenwich Mean A.R. Jan. 1, 1831.	Difference or Clock Error.	Error computed from Mean.	Mean Places Jan. 1, 1831.	Mean Errors of Clock.
1831		<i>h. m. s.</i>	<i>s.</i>	<i>h. m. s.</i>	<i>h. m. s.</i>	<i>s.</i>	<i>s.</i>	<i>h. m. s.</i>	
March 19	⊙ 1 L.....	23 51 54,55					- 33,68		
M.	⊙ 2 L.....								
	♀	1 10 58,99					33,64		
	α Aurigæ.....	5 3 39,64	+ 0,07	5 3 39,71	5 4 13,02	33,31	33,51	5 4 13,22	
	β Tauri.....	5 15 3,66	+ 0,04	5 15 3,70	5 15 36,92	33,22	33,51	5 15 37,21	
	ε Orionis.....	5 27 4,82	+ 0,03	5 27 4,85	5 27 38,47	33,62	33,50	5 27 38,35	M. 6h. 9m.
	κ Orionis.....	5 39 10,88	- 0,02	5 39 10,86	5 39 44,56	33,70	33,50	5 39 44,36	12 = 33",48
M. + 0",77	ν Orionis.....	5 57 22,24	- 0,15	5 57 22,09	5 57 55,43	33,34	33,49	5 57 55,58	
A. + 0",76	θ Columbæ....	6 1 47,57	- 0,11	6 1 47,46			33,48	6 2 20,94	A. 9h. 30m.
	α Monocerotis..	6 6 3,82	- 0,17	6 6 3,65	6 6 36,87	33,22	33,48	6 6 37,13	3 = 33",36
	β Canis Maj....	6 14 42,27	- 0,23	6 14 42,04	6 15 15,57	33,53	33,48	6 15 15,52	
T. + 0",73	α Argus.....	6 19 38,28	- 0,34	6 19 37,94			33,48	6 20 11,42	T. 19h. 8m.
Mean.	f Monocer.....	6 23 11,25	- 0,30	6 23 10,95			33,47	6 23 44,42	6 = 33",10
= + 0",76	δ Urs Min. SP..	6 26 11,98	+ 2,87	6 26 14,85	6 26 49,37*	34,52	33,47	6 26 48,32	
	55 Aurigæ.....	6 30 13,32	- 0,35	6 30 12,97			33,47	6 30 46,44	
	α Canis Maj....	6 37 8,65	- 0,37	6 37 8,28	6 37 41,91	33,63	33,46	6 37 41,74	
	χ ⁺ Canis Maj....	6 42 58,64	- 0,47	6 42 58,17	6 43 31,89	33,72	33,46	6 43 31,63	
	ι Canis Maj....	6 48 3,24	- 0,44	6 48 2,80	6 48 36,09	33,29	33,46	6 48 36,26	
	ζ Geminor.....	6 53 31,84	- 0,43	6 53 31,41	6 54 4,86	33,45	33,45	6 54 4,86	
	63 Aurigæ.....	6 59 28,25	- 0,59	6 59 27,66			33,45	7 0 1,11	
	e ⁺ Canis Maj....	7 6 48,93	- 0,60	7 6 48,33	7 7 22,07	33,74	33,45	7 7 21,78	
A.	b ⁺ Arg in Car...	8 52 19,38	- 2,31	8 52 17,07			33,38	8 52 50,45	
	r Lyncis.....	9 10 12,42	- 1,25	9 10 11,17	9 10 44,43	33,26	33,37	9 10 44,54	
	κ Leonis.....	9 14 15,64	- 1,04	9 14 14,60			33,37	9 14 47,97	
	λ Leonis.....	9 21 31,83	- 1,04	9 21 30,79	9 22 4,01	33,22	33,36	9 22 4,15	
	N Arg in Car...	9 25 34,71	- 2,46	9 25 32,25			33,36	9 26 5,61	
	κ Hydræ.....	9 31 40,82	- 1,16	9 31 39,66			33,36	9 32 13,02	
	ν Argus.....	9 42 22,54	- 3,33	9 42 19,21			33,36	9 42 52,57	
	φ Argus.....	9 50 25,79	- 2,47	9 50 23,32			33,35	9 50 56,67	
	ψ Centrum.....	9 55 15,28					33,35		
	Regulus.....	9 58 49,54	- 1,22	9 58 48,32	9 59 21,93	33,61	33,34	9 59 21,66	
	q Arg in Vel....	10 7 8,91	- 1,85	10 7 7,06			33,34	10 7 40,40	
	γ Arg in Vel....	10 14 33,54	- 1,98	10 14 31,56			33,34	10 15 4,90	
	Polaris SP....	12 58 45,30	+ 25,39	12 59 10,69	12 59 45,70*	35,01	33,25	12 59 43,94	
T.	α Lyræ.....	18 30 39,75	+ 0,15	18 30 39,90	18 31 13,11	33,21	33,12	18 31 13,02	
	β Lyræ.....	18 43 17,51	+ 0,19	18 43 17,70	18 43 50,58	32,88	33,12	18 43 50,82	
	ζ Aquilæ.....	18 57 5,33	+ 0,23	18 57 5,56	18 57 38,61	33,05	33,11	18 57 38,67	
	δ Draconis.....	19 11 55,84	+ 0,74	19 11 56,58	19 12 29,69	33,11	33,10	19 12 29,68	
	α Aquilæ.....	19 41 58,58	+ 0,46	19 41 59,04	19 42 32,29	33,25	33,08	19 42 32,12	
	β Aquilæ.....	19 46 27,19	+ 0,48	19 46 27,67	19 47 0,79	33,12	33,08	19 47 0,75	

In the above the first column contains the day of the month, the Initials of the Observer's name placed opposite to his observations, and the daily Rate of the Clock as determined from the mean error set down in column 10 found by each separate Observer, compared with that found by the same on the preceding day, thus by the observer Mf.

	<i>d. h. m.</i>	<i>s.</i>
1831, March 18	6 17 from 17 obs. Clock too slow.....	34,25
	19 6 9 — 12	33,48
Difference...	23 52.....	+ 0,77

* These are omitted in taking the means.

Hence by	M. the Clock's Rate in 24h. = + 0",774 from 17 obs. and 12 obs.
similarly. —	A. = + 0",762 — 7 — and 3 obs.
—	T. = + 0",733 — 8 — and 6 obs.
	By the method of Minimum Squares the Rate = + 0",706

The second and third columns are copied from the Transit book.

The fourth column contains the corrections for aberration nutation and precession see Pages 29 and 30, which applied to the preceeding gives column five.

Column six is derived from the Greenwich approximate Catalogue for 1830 brought on by the annual precessions thereunto annexed. This Catalogue is called an *approximate* one as far as the Right Ascensions are concerned in consequence of the number of observations being limited generally to 5, but from Instruments such as those at Greenwich and from observers possessing the advantage of long continued practice this number of observations should furnish results as accurate as could be desired; this is I believe for the most part the case except in a few cases where errors have crept in which I have corrected;* at any rate by assuming the whole Catalogue of 720 Stars I may venture to hope to make up in part by quantity what this Catalogue may be deficient in quality.

The seventh column contains the difference between the two preceeding, or the error of the Clock; this column is divided by lines, and the means taken so as to leave each observer's result separate, which is placed in column ten; with this and the mean rate, the next or column eight is computed, which being applied to column five gives the mean place of each Star for January 1st 1831, column nine. The advantage and necessity of thus rendering each observer's results independent has been acknowledged and met with considerable attention by Professor Schumacher and other continental Astronomers who have bestowed considerable pains to ascertain the constant of time which one observer differs from another in estimating the time when a Star arrives at the wires of a Transit Instrument: That this quantity *does remain constant*, is a fact however not confirmed by my own experience, on the contrary I generally find that unpractised observers note the time in *excess* from three to seven tenths of a second when compared with those long accustomed to this species of observation.

* *Errors in Greenwich Approximate Catalogue.*

		<i>h. m. s.</i>	<i>h. m. s.</i>
δ Eridani.....	A.R. for	3 35 5,52	read 3 35 6,52
κ Cancri.....	— —	8 58 31,85	— 8 58 32,85
ο Sagittarii....	— —	18 54 30,59	— 18 54 29,59
ο Aquarii.....	— —	21 54 31,77	— 21 54 31,17

The A.R. of τ¹ τ² η¹ and η² Aquarii being doubtful have been omitted.

On taking the difference between Columns 7 and 8 Pages 34 and 35, we get the error of a single Observation as follows.

By observer M.		By observer A.		By observer T.	
March.....18th	19th	March.....18th	19th	March....18th	19th
s.	s.	s.	s.	s.	s.
0,14	0,20	0,23	0,11	0,02	0,09
,14	,29	,37	,14	,07	,24
,01	,12	,82	,27	,07	,06
,20	,20	,05		,00	,01
,14	,15	,31		,15	,17
,31	,26	,15		,01	,04
,31	,05	,09		,03	
,26	,17			,15	
,67	,26				
,05	,17				
,32	,00				
,20	,29				
,26					
,13					
,54					
,56					
,13					
Mean error of one } Observation.... }	0,225	—	0,254	—	0,079

The above mean errors cannot be considered large for observers with only two months experience, but finding towards the middle of the month of May that these numbers did not become as small as I had anticipated, I was induced to exchange these two observers for two others who had practiced observing transits with the Mural Circle during the three previous Months, and from whom I had reasons for expecting an increased degree of accuracy and attention. Accordingly the transit Observations to the end of the year are made by these observers as is explained in the transit book by the initials of their names, where R stands for Ragavacharyer Braminy and S for Soontharum Moodeliar; the rough observations made by these observers on two consecutive days which now follow are given in order to shew the progress they had made when their Observations were first recorded, and for the purpose of exhibiting the degree of accuracy which may be expected from the results.

Errors in Greenwich Approximate Catalogue.

	h. m.					
δ Sagittarii....	A.R.	17 50	from Madras Observation	appears	1"	in error.
π Virginis.....	—	11 52	—	—	—	0",8
κ Cephei.....	—	20 14	—	—	—	3"

	I	II	III	IV	V	Mean.	Corrections for			Transit over the Meridian.			NAMES.	Observed by		
							Level	Azimuth	Coll.	h.	m.	s.				
							-	.	+							
1831																
May	31	s. 2,2	s. 31,9	m. s. 28 1,9	s. 31,8	s. 0,9	m. s. 29 9,89	s. ,035	s. —	s. ,202	s. ,549	h. 4	m. 29	s. 10,21	⊙ 1 L. ⊙ 2 L.	S.
		18,8	48,3	30 17,9	47,7	17,5	0,5	9 49,66	,039	—	,701	,659	11	9 49,58	p Ursæ Majoris	
		38,9	14,2	9 49,9	24,8	0,5	22,0	27 19,28	,036	—	,382	,580	11	27 19,44	Ursæ Minoris	
		16,6	47,7	27 19,5	50,6	22,0	31	55,26	,035	—	,213	,552	11	31 55,56	92 Leonis	
		55,5	25,5	31 55,5	24,9	34,9	17	9,94	,018	+	2,570	1,090	12	17 13,59	a ² Crucis	
		12,0	11,1	17 9,9	9,0	7,7	38	8,46	,035	—	,101	,535	12	38 8,86	r Comæ Berenicis	
		10,2	39,5	38 8,8	37,9	5,9	8,5	39 10,64	,035	—	,118	,537	13	39 11,02	r Bootis	R.
		12,6	41,7	39 10,8	39,6	43,4	51	55,08	,020	+	2,317	1,007	13	51 58,39	β Centauri	
		6,7	1,0	51 55,3	49,0	46,0	3	50,34	,031	+	,480	,516	14	3 51,31	κ Virginis	
		54,7	22,5	3 50,4	18,1	15,3	9	55,08	,041	—	1,015	,748	14	9 54,77	λ Bootis	
		35,0	14,9	9 55,0	35,2	55,0	19	24,62	,042	—	1,298	,844	14	19 24,12	θ Bootis	
		54,2	39,5	19 24,6	9,8	34,1	24	30,14	,037	—	,449	,597	14	24 30,25	ρ Bootis	
		26,0	58,1	24 30,2	2,3	41,7	32	44,30	,035	—	,092	,535	14	32 44,71	π Bootis	
		47,0	15,6	32 44,2	13,0	31,5	43	33,14	,035	—	,150	,542	14	43 33,50	ξ Bootis	
		34,7	4,0	43 33,1	2,4	21,9	48	8,80	,026	+	1,341	,684	14	48 10,80	κ Centauri	
		55,8	32,2	48 8,9	45,2	45,5	55	32,62	,040	—	,773	,684	14	55 32,49	β Bootis	
		19,8	56,2	55 32,6	9,0	2,0	4	17,86	,020	+	2,218	,968	15	4 21,03	β Cricini	
		33,8	25,7	4 18,0	9,8	14,2	18	4,24	,038	—	,644	,649	15	18 4,21	μ Bootis	
		54,6	29,4	18 4,0	39,0	17,5	28	14,74	,028	+	,933	,585	15	28 16,23	40 Libræ	
		12,0	43,2	28 14,9	46,1	33,0	8	36,80	,031	+	,556	,523	20	8 37,85	a ² Capricorni	
		40,5	8,7	8 36,8	5,0	32,6	17	35,00	,030	+	,686	,538	20	17 36,19	π Capricorni	
		37,8	6,0	17 34,7	3,9	19,5	30	21,56	,030	+	,686	,538	20	30 22,75	o Capricorni	
		23,6	52,5	30 21,7	50,5	51,1	41	21,80	,030	+	,643	,536	20	41 22,95	δ 2 L.	
		23,2	52,5	41 21,8	51,1	41,0	54	42,62	,030	+	,729	,544	20	54 43,86	η Capricorni	
		44,2	13,4	54 42,7	11,8	15,5	0	19,30	,031	+	,534	,520	21	0 20,32	ν Aquarii	
		23,0	51,1	0 19,4	47,5	16,5	48	12,28	,028	+	,990	,592	22	48 13,83	Fomalhaut	S.
		8,5	40,0	48 12,4	44,0	13,7	56	16,84	,034	—	,028	,526	22	56 17,30	a Pegasi	
		20,0	48,2	56 16,8	45,5	38,8	59	36,10	,036	—	,369	,580	23	59 36,28	a Andromedæ	
		33,4	4,6	59 36,0	7,7	24,9	4	28,02	,034	—	,027	,526	00	4 28,49	γ Pegasi	
		31,0	59,8	4 28,0	56,4	30,9	30	54,12	,043	—	1,48	,902	0	30 53,50	a Cassiopeæ	
		17,0	5,5	30 54,4	42,8											
June	1	7,0	36,6	32 6,4	36,0	5,8	33	14,53	,047	—	,215	,634	4	33 14,90	⊙ 1 L. ⊙ 2 L.	S.
		23,3	53,0	34 22,7	52,5	22,0	19	52,90	,051	—	,280	,644	7	19 53,21	♀ Centrum	
		52,2	22,8	19 53,0	23,5	0,9	9	50,16	,051	—	,741	,759	11	9 50,13	p Ursæ Major	
		39,4	14,8	9 50,2	25,5	19,8	16	22,40	,040	+	,678	,615	11	16 23,65	γ Hyd. and Crat.	
		25,0	53,8	16 22,5	50,9	31,9	21	36,76	,043	+	,341	,589	11	21 37,65	c Leonis	
		41,5	9,1	21 36,9	4,4	57,7	28	2,48	,041	+	,492	,595	11	28 3,53	θ Hyd. and Crat.	
		7,0	34,8	28 2,9	30,0	1,5	37	6,22	,045	+	,127	,594	11	37 6,90	ν Virginis	
		10,9	38,7	37 6,1	33,9	14,4	46	18,70	,045	+	,080	,596	11	46 19,33	A Virginis	
		23,0	50,8	46 18,7	46,6	58,0	55	2,86	,045	+	,151	,593	11	55 3,56	* Virginis	
		7,7	35,2	55 2,9	30,5	14,9	2	46,72	,031	+	1,890	,944	12	2 49,52	ρ Centauri	
		18,8	2,6	2 46,9	30,4	6,6	11	11,72	,043	+	,290	,588	12	11 12,55	η Virginis	
		16,9	44,1	11 11,9	32,1	31,0	19	20,76	,036	+	1,290	,748	12	19 22,76	u Centauri	
		10,2	45,8	19 20,9	55,9	29,4	33	34,46	,043	+	,286	,588	12	33 35,29	γ ^a Virginis	
		39,4	7,2	33 34,4	1,9	53,9	44	54,42	,047	—	,222	,637	12	44 54,79	q Comæ Berenicis	
		55,0	24,8	44 54,4	24,0	3,6	55	8,52	,043	+	,357	,589	12	55 9,42	k ⁺ Virginis	
		13,3	40,9	55 8,8	36,0	1,7	4	5,34	,045	—	,014	,603	13	4 5,91	* Virginis	R.
		9,0	37,0	4 5,3	33,7	8,2	9	54,84	,053	—	,833	,785	13	9 54,74	h Can Ven	
		41,5	18,0	9 54,8	31,7	36,3	26	41,30	,042	+	,396	,591	13	26 42,24	l ³ Virginis	
		46,2	13,7	26 41,3	9,0	32,0	32	33,40	,047	—	,188	,530	13	32 33,70	1 Bootis	
		34,6	4,0	32 33,7	2,7	8,8	39	11,10	,047	—	,125	,620	13	39 11,55	r Bootis	
		13,3	42,1	39 11,2	40,1	18,8	44	59,30	,033	+	1,640	,853	13	45 1,76	ξ Centauri	
		39,8	19,5	44 59,3	39,1	27,0	51	9,90	,033	+	1,560	,825	13	51 12,25	μ ² Centauri	
		52,8	31,3	51 10,0	48,4	46,3	3	50,58	,041	+	,508	,596	14	3 51,64	κ Virginis	
		55,0	22,6	3 50,6	18,4											

L = 0", 50 E.
A = 19", 60 W.
C = 6", 56 E.

L = 0", 67 E.
A = 19", 60 W.
C = 7", 62 E.

	I	II	III	IV	V	Mean.	Corrections for			Transit over the Meridian.	NAMES.	Observed by			
							Level	Azimuth	Coll.						
							-		+						
1831															
June	1	s. 1,0	s. 39,6	m. 15	s. 18,5	s. 56,8	s. 35,5	m. 15	s. 18,28	,033	+ 1,550	,825	14 15 20,62	τ ^a Lupi	S.
		49,1	31,3	21 14,0	56,3	38,8	21 13,90	,031	+ 1,790	,910	14 21 16,57	σ Lupi			
		31,8	3,0	37 34,4	5,5	36,5	37 34,24	,049	- ,378	,666	14 37 34,48	ε Bootis			
		35,1	4,2	43 33,5	2,6	31,8	43 33,44	,047	- ,160	,625	14 43 33,86	ξ Bootis			
		59,4	26,8	51 54,7	22,5	50,2	51 54,72	,041	+ ,470	,595	14 51 55,75	δ Libræ			
		49,8	20,3	59 50,7	21,0	51,6	59 50,68	,031	+ 1,894	,943	14 59 53,48	ζ Lupi			
		26,0	6,1	6 46,4	26,8	7,2	6 46,50	,033	+ 1,870	,866	15 6 49,01	μ Lupi			
		55,0	29,7	18 4 6	39,5	14,2	18 4,60	,051	- ,702	,747	15 18 4,60	μ Bootis			
		58,8	26,7	25 54,6	22,5	50,1	25 54,54	,041	+ ,508	,597	15 25 55,60	f Libræ			
		12,7	41,6	32 10,9	39,7	8,8	32 10,74	,040	+ ,735	,623	15 32 12,06	κ Libræ			
		55,5	24,1	33 53,8	22,4	51,5	33 53,46	,040	+ ,621	,608	21 33 54,65	δ 2 L.			
		3,5	31,9	44 0,5	28,8	57,0	44 0,34	,040	+ ,621	,608	21 44 1,53	μ Capricorni			
		16,7	45,5	57 13,9	42,0	10,6	57 13,74	,040	+ ,627	,608	21 57 14,94	i Aquarii			
		9,0	40,5	48 12,8	44,9	16,5	48 12,74	,038	+ 1,050	,683	22 48 14,44	γ Fomalhaut			
		20,1	48,8	56 17,4	45,8	13,9	56 17,20	,046	- ,027	,607	22 56 17,73	α Pegasi			

The above column "Transit over the meridian" being transcribed into book No. 1, we have from the same.

	NAMES.	App. A.R. + Clock Er.	Aber- ration, &c.	Mean A.R. Jan. 1, 1831 + Clock Er.	Greenwich Mean A.R. Jan. 1, 1831.	Difference or Clock Error.	Error computed from Mean.	Mean Places Jan. 1, 1831.	Mean Errors of Clock.
		h. m. s.	s.	h. m. s.	h. m. s.	s.	s.	h. m. s.	
1831									
May	31	⊙ 1 L.							
	S.	⊙ 2 L.					- 4,76		
		p Ursæ Major...	11 9 49,58	- 0,41	11 9 49,17		4,66	11 9 53,83	
		Ursæ Min....	11 27 19,44	- 0,61	11 27 18,83		4,66	11 27 23,49	
		92 Leonis.....	11 31 55,56	- 0,70	11 31 54,86		4,66	11 31 59,52	
		a* Crucis.....	12 17 13,59	- 2,80	12 17 10,79		4,65	12 17 15,44	
		n Comæ Ber....	12 38 8,86	- 1,09	12 38 7,77		4,65	12 38 12,42	
	R.	τ Bootis.....	13 39 11,02	- 1,39	13 39 9,63		4,03	13 39 13,66	
		β Centauri....	13 51 58,39	- 3,57	13 51 54,82		4,03	13 51 58,85	
	R. + 0",48	κ Virginis....	14 3 51,31	- 1,76	14 3 49,55	14 3 53,61	- 4,06	4,03	14 3 53,58
	S. + 0",18	λ Bootis.....	14 9 54,77	- 1,51	14 9 53,26	14 9 57,24	3,98	4,03	14 9 57,29
	Mean.	θ Bootis.....	14 19 24,12	- 1,61	14 19 22,51	14 19 26,50	3,99	4,02	14 19 26,53
	= + 0",38	ρ Bootis.....	14 24 30,25	- 1,55	14 24 28,70	14 24 32,75	4,05	4,02	14 24 32,72
		π Bootis.....	14 32 44,71	- 1,61	14 32 43,10	14 32 47,13	4,03	4,02	14 32 47,12
		ξ Bootis.....	14 43 33,50	- 1,63	14 43 31,87	14 43 35,84	3,97	4,02	14 43 35,89
		κ Centauri....	14 48 10,80	- 2,69	14 48 8,11			4,02	14 48 12,13
		β Bootis.....	14 55 32,49	- 1,71	14 53 30,78	14 55 34,85	4,07	4,01	14 53 34,79
		β Cricini.....	15 4 21,03	- 3,82	15 4 17,21			4,01	15 4 21,22
		μ Bootis.....	15 18 4,21	- 1,77	15 18 2,44	15 18 6,50	4,06	4,01	15 18 6,45
		40 Libræ.....	15 28 16,23	- 2,37	15 28 13,86	15 28 18,02	4,16	4,01	15 28 17,87
		a* Capricorni...	20 8 37,85	- 1,46	20 8 36,39	20 8 40,34	3,95	3,94	20 8 40,33
		π Capricorni....	20 17 36,19	- 1,44	20 17 34,75			3,93	20 17 38,68
		ο Capricorni....	20 30 22,86	- 1,86	20 30 21,50			3,93	20 30 25,43
		δ 2 L.....	20 41 22,95					3,93	
		η Capricorni....	20 54 43,86	- 1,21	20 54 42,65	20 54 46,72	4,07	3,93	20 54 46,58
		ν Aquarii.....	21 0 20,32	- 1,16	21 0 19,16	21 0 22,90	3,74	3,93	21 0 23,09

R. at 15h. 20m.
12 = - 4",01

	NAMES.	App. A.R. + Clock Er.	Aber- ration, &c.	Mean A.R. Jan. 1, 1831 + Clock Er.	Greenwich Mean A.R. Jan. 1, 1831.	Difference or Clock Error.	Error computed from Mean.	Mean Places Jan. 1, 1831.	Mean Errors of Clock.
1831	S. Fomalhaut....	<i>h. m. s.</i> 22 48 13,83	— 0,32	<i>h. m. s.</i> 22 48 13,51	<i>h. m. s.</i> 22 48 17,65	— 4,14	— 4,38	<i>h. m. s.</i> 22 48 17,89	
	a Pegasi.....	22 56 17,30	— 0,67	22 56 16,63	22 56 20,96	4,33	4,38	22 56 21,01	S. at 23h. 40m.
	a Andromedæ..	23 59 36,28	— 0,37	23 59 35,91	23 59 40,12	4,21	4,37	23 59 40,28	
	γ Pegasi.....	0 4 28,49	— 0,28	0 4 28,21	0 4 32,62	4,41	4,37	0 4 32,58	5 = — 4",37
	a Cassiopeæ....	0 30 53,50	— 0,36	0 30 53,14	0 30 57,93	4,79	4,36	0 30 57,50	
June	1 ☉ 1 L.....	4 33 14,90					4,31		
	S. ☉ 2 L.....								
	☽.....	7 19 53,21					4,27		
	p Ursæ Major..	11 9 50,13	— 0,40	11 9 49,73			4,21	11 9 53,94	
	γ Hyd. and Crat.	11 16 23,65	— 0,94	11 16 22,71	11 16 26,78	4,07	4,20	11 16 26,91	
	e Leonis.....	11 21 37,65	— 0,84	11 21 26,81	11 21 40,97	4,16	4,20	11 21 31,01	
	θ Hyd. and Crat.	11 28 3,53	— 0,93	11 28 2,60	11 28 6,96	4,36	4,20	11 28 6,80	
	ν Virginis.....	11 37 6,90	— 0,83	11 37 6,07	11 37 10,34	4,27	4,20	11 37 10,27	S. at 11h. 35m.
	A Virginis.....	11 46 19,33	— 0,88	11 46 18,95			4,20	11 46 23,15	5 = — 4",20
	* Virginis.....	11 55 3,56	— 0,95	11 55 2,61			4,20	11 55 6,81	
	ρ Centauri.....	12 2 49,52	— 1,99	12 2 47,53			4,20	12 2 51,73	
	η Virginis.....	12 11 12,55	— 1,08	12 11 11,47	12 11 15,62	4,15	4,19	12 11 15,66	
	α Centauri.....	12 19 22,76	— 1,72	12 19 21,04			4,19	12 19 25,23	
	γ ^a Virginis.....	12 33 35,29	— 1,19	12 33 34,10			4,19	12 33 38,29	
	q Comæ Ber....	12 44 54,79	— 1,08	12 44 53,71			4,19	12 44 57,90	
	k ^a Virginis.....	12 55 9,42	— 1,35	12 55 8,07			4,18	12 55 12,25	
	R. * Virginis.....	13 4 5,91	— 1,25	13 4 4,66			3,63	13 4 8,29	
	h Can Ven.....	13 9 54,74	— 1,14	13 9 53,60			3,63	13 9 57,23	
	l ³ Virginis.....	13 26 42,24	— 1,53	13 26 40,71			3,62	13 26 44,33	
	l Bootis.....	13 32 33,70	— 1,33	13 32 32,37			3,62	13 32 35,99	
	τ Bootis.....	13 39 11,55	— 1,39	13 39 10,16			3,62	13 39 13,78	R. at 14h. 51m.
	ζ Centauri.....	13 45 1,76	— 2,61	13 44 59,15			— 3,62	13 45 2,77	6 = — 3",59
	μ ² Centauri.....	13 51 12,25	— 2,58	13 51 9,67			3,61	13 51 13,28	
	κ Virginis.....	14 3 51,64	— 1,75	14 3 49,89	14 3 53,61	— 3,72	3,61	14 3 53,50	
R. + 0",43	τ ^a Lupi.....	14 15 20,62	— 2,70	14 15 17,92			3,61	14 15 21,53	
S. + 0",49	σ Lupi.....	14 21 16,57	— 2,98	14 21 13,59			3,60	14 21 17,19	
Mean.	e Bootis.....	14 37 34,48	— 1,59	14 37 32,89	14 37 36,36	3,47	3,60	14 37 36,49	
= + 0",45	ξ Bootis.....	14 43 33,86	— 1,63	14 43 32,23	14 43 35,84	3,61	3,59	14 43 35,82	
	δ Libræ.....	14 51 55,75	— 1,88	14 51 53,87	14 51 57,28	3,41	3,59	14 51 57,46	
	ζ Lupi.....	14 59 53,48	— 3,26	14 59 50,22			3,59	14 59 53,81	
	μ Lupi.....	15 6 49,01	— 3,02	15 6 45,99			3,59	15 6 49,58	
	μ Bootis.....	15 18 4,60	— 1,76	15 18 2,84	15 18 6,50	3,66	3,58	15 18 6,42	
	f Libræ.....	15 25 55,70	— 2,00	15 25 53,70			3,58	15 25 57,28	
	κ Libræ.....	15 32 12,06	— 2,20	15 32 9,86	15 32 13,54	3,68	3,58	15 32 13,44	
	S.) 2 L.....	21 33 54,65							
	μ Capricorni....	21 44 1,53	— 0,92	21 44 0,61	21 44 4,64	4,03	3,93	21 44 4,54	
	i Aquarii.....	21 57 14,94	— 0,84	21 57 14,10	21 57 18,19	4,09	3,93	21 57 18,03	S. at 22h. 21m.
	Fomalhaut....	22 48 14,44	— 0,35	22 48 14,09	22 48 17,55	3,56	3,91	22 48 18,00	4 = — 3",90
	a Pegasi.....	22 56 17,73	— 0,70	22 56 17,03	22 56 20,96	3,93	3,91	22 56 20,94	

Taking the difference between Columns 7 and 8, we get the errors of each single Observation by

Observer R.			Observer S.		
May 31	June 1		May 31	June 1	
s.	s.		s.	s.	
0,03	—	0,11	—	0,24	—
,05	—	,13	—	0,05	—
,03	—	,02	—	0,16	—
,03	—	,18	—	0,04	—
,01	—	,08	—	0,43	—
,05	—	,10	—		
,06					
,05					
,15					
,01					
,14					
,19					
Mean error of one } Observation.... }	,079			,142	

On inserting the Clock errors at pages 34 and 35, it is found that little or no difference exists between the times at which the observers M, A and T note the passage of a Star, this however was not always the case, with the first observations a difference of from three to five tenths of a second of time in excess invariably occurred between M and A when compared with T. Comparing the Clock errors by observers R and S as last found it appears from the observations of 31st May that observer R noted the time in excess above observer S = 0,47 and from the observations of 1st June that R noted the time in excess 0,55; we will now see if this difference continues constant, for which purpose we will examine the observations on two days near the end of the year as follows.

	I	II	III	IV	V	Mean.	Corrections for			Transit over the Meridian.	NAMES.	Observed by
							Level	Azimuth	Coll.			
							—		+			
1831												
December 24	s.	s.	m. s.	s.	s.	m. s.	s.	s.	s.	h. m. s.		
	42,2	12,8	9 42,5	12,7	42,7	10 53,58	,257	+	,986	,681	18 10 54,99	⊙ 1 L.
		34,4	12 4,5	34,7	4,7						18 10 54,99	⊙ 2 L.
			4 10,9				2,70	—	53,17	20,38	1 3 35,41	Polaris
	19,6	13,7	18 7,7	1,7	55,5	18 7,64	,403	—	2,15	1,227	1 18 6,31	δ Cassiopeæ
	40,1	7,7	24 35,5	2,9	30,5	24 35,34	,292	+	,207	,625	1 24 35,88	μ Piscium
	32,51,8		34 35,7	35,27,7	36,19	34 35,71	,182	+	2,72	1,186	1 34 39,43	α Eridani
	29,8	58,8	39 27,8	55,9	24,6	39 27,38	,266	+	,793	,652	1 39 28,56	τ Ceti
	33,6	2,9	47 32,5	0,7	29,8	47 31,90	,310	—	,150	,659	1 47 32,10	γ Arietis
		51,4,0	52 31,6	53,58,6	55,25,5	52 31,47	,487	—	4,108	1,954	1 52 28,83	50 Cassiopeæ
	44,4	42,0	4 9,5	37,0	4,3	4 9,44	,292	+	,255	,625	2 4 10,03	62 Ceti
	4,3	33,4	12 2,2	31,2	0,1	12 2,24	,310	—	,164	,661	2 12 2,43	θ' Arietis
	51,2	19,1	18 47,0	14,6	42,4	18 46,86	,297	+	,109	,631	2 18 47,30	* Ceti
	38,5	20,3	24 2,0	43,0	24,8	24 1,72	,213	+	2,01	,943	2 24 4,46	κ Eridani
	11,5	39,4	30 7,0	34,8	2,3	30 7,00	,297	+	,186	,627	2 30 7,52	* Ceti

	I	II	III	IV	V	Mean.	Corrections for			Transit over the Meridian.	NAMES.	Observed by		
							Level	Azimuth	Coll.					
							-		+					
1831	s.	s.	m.	s.	s.	m.	s.	s.	s.	h.	m.	s.		
December 24	43,9	11,2	34 40,3	8,2	36,6	34 40,04	,275	+	,672	,640	2 34	41,08	c Ceti	R.
	24,9	53,1	39 21,7	50,9	18,2	39 21,60	,270	+	,731	,645	2 39	22,71	π Ceti	
	20,2	51,0	43 21,9	52,6	23,1	43 21,76	,323	-	,396	,696	2 43	21,74	c Arietis	
	53,3	22,9	52 52,2	21,7	50,9	52 52,20	,315	-	,309	,666	2 52	52,34	c Arietis	
	46,2	14,1	57 41,9	9,9	37,6	57 41,94	,279	+	,562	,630	2 57	42,85	ρ ^a Eridani	
		27,0	10 55,0	22,8		10 54,93	,275	+	,588	,640	3 10	55,88	ζ Eridani	
	6,0	33,8	19 1,8	29,7	57,0	19 1,66	,297	+	,123	,630	3 19	2,12	o Tauri	
		21,8	22 2,2	43,0		22 2,33	,363	-	,126	,932	3 22	1,64	σ Persei	
L = 4",43 E.	13,9	51,0	27 27,8	4,9	41,7	27 27,86	,226	+	,167	,841	3 27	30,15	z Eridani	
A = 22",78 E.		5,4	31 32,9	0,3		31 32,87	,288	+	,346	,631	3 31	33,56	E Tauri	
C = 8",45 E.		50,2	27,4 37 4,6	41,9	19,2	37 4,66	,350	-	,104	,842	3 37	4,11	v Persei	
		26,2	56,3 42 26,4	51,5	26,3	42 26,34	,319	-	,303	,681	3 42	26,40	f Pleiadum	
		46,3	18,8 46 51,0	23,0	55,1	46 50,84	,328	-	,560	,731	3 46	50,68	ζ Persei	
		24,9	5,5 59 46,0	26,8	7,0	59 46,04	,363	-	,125	,922	3 59	45,35	c Persei	
		32,8	1,8 16 30,6	59,2	28,0	16 30,48	,306	-	,112	,654	4 16	30,72	δ ^a Tauri	
		22,7	51,2 22 20,0	48,7	17,1	22 19,94	,306	-	,056	,648	4 22	20,23	o ^a Tauri	
		45,0	13,0 29 40,9	8,8	36,6	29 40,86	,297	-	,089	,632	4 29	41,11	d Tauri	
		1,2	29,5 33 57,8	26,0	53,8	33 57,66	,301	-	,032	,638	4 33	57,96	c ^a Tauri	
		12,0	46,7 39 21,4	56,0	30,4	39 21,30	,239	+	,147	,787	4 39	23,32	β Ceti Sculp	
		20,7	49,0 46 17,2	45,7	14,0	46 17,32	,306	+	,026	,743	4 46	17,78	o ^a Orionis	
		57,2	53,0 51 48,0	Cloudy	38,5	51 48,14	,403	-	,224	1,255	4 51	46,75	d ^a Camelopardi	
		48,0	24,6 58 1,1	37,4	13,8	58 0,98	,340	-	,045	,828	4 58	0,52	η Aurigæ	
		50,5	18,2 7 45,8	13,3	40,7	7 45,70	,292	+	,278	,625	5 7	46,31	ρ Orionis	
		12,2	39,9 55 7,5	35,1	2,5	55 7,44	,292	+	,227	,625	10 55	8,00	d Leonis	
		54,2	22,9 33 50,8	18,9	47,0	33 50,76	,297	+	,174	,628	11 33	51,26	γ 2 L.	
		56,0	23,9 39 51,6	19,7	47,2	39 51,68	,297	+	,102	,632	11 39	52,12	ε ^a Virginis	
		15,9	43,4 45 11,0	38,3	6,0	45 10,92	,292	+	,278	,625	11 45	11,53	β Virginis	
		34,5	2,2 54 30,0	57,8	25,5	54 30,00	,297	+	,145	,629	11 54	30,48	π Virginis	
		57,6	25,4 59 53,5	21,1	48,9	59 53,30	,297	+	,094	,633	11 59	53,73	o Virginis	
25			14	42,1	12,1						18		⊙ 1 L.	
		34,7	4,9 16 34,9	5,0	34,8	16 34,86	,256	+	,985	,681	18 16	36,27	⊙ 2 L.	R.
		14,5	42,0 19 9,7	37,2	4,8	19 9,64	,282	+	,426	,625	0 19	10,41	* Ceti	
		23,8	1,1 24 32,8	4,0	35,0	24 34,34	,322	-	,473	,713	0 24	34,26	28 Andromeda	
		21,7	49,9 31 18,2	46,7	15,0	31 18,30	,304	-	,032	,644	0 31	18,61	53 Piscium	
		3,6	32,0 37 0,3	28,3	56,5	37 0,14	,260	+	,913	,669	0 37	1,46	* Ceti	
		7,1	38,0 44 8,9	39,8	10,5	44 8,86	,322	-	,394	,700	0 44	8,84	k Piscium	
		15,6	45,5 49 15,1	45,0	14,9	49 15,22	,318	-	,386	,696	0 49	15,21	k Piscium	
		37,9	6,0 53 34,0	2,1	30,0	53 34,00	,273	+	,662	,639	0 53	35,03	φ ^a Ceti	
		58,6	28,0 59 57,3	26,8	56,0	59 57,34	,313	-	,209	,667	0 59	57,49	ψ ^a Piscium	
		24,9	52,5 11 20,1	47,6	15,0	11 20,02	,282	+	,428	,625	1 11	20,79	39 Ceti	
		23,9	18,2 18 12,0	5,8	59,8	18 11,94	,401	-	2,150	1,227	1 18	10,62	δ Cassiopeæ	
		43,6	11,6 24 39,1	6,8	34,3	24 39,12	,291	+	,227	,625	1 24	39,68	μ Piscium	
		32,0	0,1 31 28,2	56,2	24,0	31 28,10	,300	+	,045	,637	1 31	28,48	π Piscium	
		41,1	10,5 36 39,8	9,0	37,9	36 39,66	,309	-	,176	,663	1 36	39,84	107 Piscium	
		33,1	2,1 39 31,0	59,8	28,5	39 30,90	,265	+	,793	,652	1 39	32,08	τ Ceti	
		37,7	6,9 47 35,8	4,7	33,5	47 35,72	,309	-	,150	,659	1 47	35,92	γ Arietis	
			51,9,2 52 36,0	54,3,2		52 36,13	,485	-	4,108	2,000	1 52	33,54	50 Cassiopeæ	
			55,4,15 56 40,8	57,4,0		56 40,77	,159	+	,313	,136	1 56	45,10	α Hydri	
		16,2	43,7 4 11,2	38,9	6,1	4 11,22	,291	+	,255	,625	2 4	11,81	62 Ceti	
		5,9	35,0 12 4,1	33,1	2,0	12 4,02	,309	-	,164	,661	2 12	4,21	θ ^a Arietis	
		40,7	22,0 24 3,6	45,0	26,3	24 3,52	,212	+	2,010	,943	2 24	6,26	κ Eridani	
L = 4",41 E.		48,9	17,4 30 45,0	13,1	41,0	30 45,08	,300	+	,035	,638	2 30	45,45	U Arietis	
A = 22",78 E.		45,7	13,9 34 41,9	10,1	38,2	34 41,96	,273	+	,672	,640	2 34	43,00	ε Ceti	
C = 8",45 E.		37,40,0	38,57,0 40 14,0	41,31,5	42,48,0	40 14,10	,123	+	4,067	1,690	2 40	19,73	ε Hydri	
		15,0	42,8 56 10,7	38,5	6,0	56 10,60	,278	+	,562	,630	2 56	11,51	ρ ^a Eridani	

	I	II	III	IV	V	Mean.	Corrections for			Transit over the Meridian.	NAMES.	Observed by	
							Level	Azimuth	Coll.				
							-		+				
1831 December 25	s.	s.	m. s.	s.	s.	m. s.	s.	s.	s.	h. m. s.		S.	
	21,5	57,6	0 33,5	9,5	45,3	0 33,48	,344	-	,913	,818	3 0 33,04		β Persei
	33,4	2,9	8 32,1	1,5	30,9	8 32,16	,309	-	,208	,666	3 8 32,31		ξ Arietis
	7,5	5,0	31 32,5	0,1	27,7	31 32,56	,287	+	,346	,624	3 31 33,24		E Tauri
	49,5	26,9	37 4,0	40,8	17,8	37 3,80	,348	-	,104	,842	3 37 4,19		ν Persei
	25,8	55,9	42 25,7	55,9	25,6	42 25,78	,318	-	,729	,681	3 42 25,41		f Pleiadum
	45,8	17,8	46 50,4	22,7	54,5	46 50,24	,326	-	,560	,731	3 46 50,09		ζ Persei
		4,5	59 44,8	25,5	6,5	59 45,12	,362	-	1,250	,829	3 59 44,34		c Persei
	28,7	10,0	5 50,8	32,3	12,9	5 50,94	,362	-	1,300	,934	4 5 50,21		μ Persei
	23,7	1,7	11 39,0	16,8	53,4	11 38,92	,225	+	1,730	,844	4 11 41,27		a Horologii
	32,5	1,2	16 30,0	58,8	27,6	16 30,02	,304	-	,112	,654	4 16 30,26		δ ¹ Tauri
		34,7	22 3,7	33,0	1,5	22 3,74	,309	-	,159	,595	4 22 3,87		e Tauri
	44,5	12,8	29 40,5	8,5	36,4	29 40,54	,295	+	,089	,633	4 29 40,97		d Tauri
	25,5	55,5	35 25,4	55,5	25,3	35 25,44	,313	-	,270	,676	4 35 25,53		τ Tauri
	25,4	53,3	40 20,3	47,8	15,8	40 20,52	,282	+	,428	,625	4 40 21,29		μ Eridani
	20,0	48,6	46 16,6	45,5	13,4	46 16,82	,304	-	,026	,743	4 46 17,23		o ¹ Orionis
	49,56,6	50,52,5	51 48,3	52,42,6	53,37,7	51 47,54	,401	-	2,240	1,255	4 51 46,15		d ¹ Camelopardi
	46,9	23,5	57 59,8	36,8	12,5	57 59,90	,344	-	,945	,828	4 57 59,44		η Aurigæ
	49,7	17,7	7 45,7	12,4	40,0	7 45,10	,291	+	,278	,625	5 7 45,71		ρ Orionis
	29,5	57,6	15 25,6	53,8	21,9	15 25,68	,273	+	,670	,640	5 15 26,72		ν Leporis
	26,6	54,5	19 21,6	49,7	17,7	19 22,02	,384	+	,184	,628	5 19 22,45		γ Orionis
		30,5	25 3,4	35,5	7,7	25 3,09	,326	-	,585	,664	5 25 2,84		κ Aurigæ
	31,5	59,6	30 27,0	54,8	22,5	30 27,08	,278	+	,503	,627	5 30 27,93		ι Orionis
	24,4	54,4	38 23,8	53,8	23,6	38 24,00	,260	+	,957	,676	5 38 25,37		12 Leporis
	37,7	6,6	42 34,6	3,4	31,3	42 34,72	,209	+	,731	,646	5 42 35,83		ξ Leporis
	8,6	41,5	48 16,3	49,9	24,2	48 16,10	,218	+	1,410	,770	5 48 18,06		β Columbæ
	2,6	31,2	51 58,9	28,0	56,6	51 59,46	,269	+	,728	,644	5 52 0,56		η Leporis
	12,5	42,4	57 11,6	40,6	9,8	57 11,38	,309	-	,192	,665	5 57 11,54		χ ² Orionis
	46,0	43,9	5 42,2	39,3	36,9	5 41,66	,410	-	2,390	1,309	6 5 40,17		α Lyncis
	48,5	30,3	15 Hazy	54,5	36,7	15 12,61	,366	-	1,390	,867	6 15 11,72		d Aurigæ
	15,5	44,9	22 14,4	43,5	13,2	22 14,30	,198	+	2,280	1,031	6 22 17,41	a Argus	
	14,6	44,5	11 13,5	42,3	12,1	11 13,40	,309	-	,192	,664	14 11 13,56	a Bootis	

The above column "Transit over the meridian" being transcribed into book No. 1, we have from the same.

	NAMES.	App. A.R. + Clock Er.	Aber- ration, &c.	Mean A.R. Jan. 1, 1831 + Clock Er.	Greenwich Mean A.R. Jan. 1, 1831.	Difference or Clock Error.	Error computed from Mean.	Mean Places Jan. 1, 1831.	Mean Errors of Clock.
1831 Dec. 24	⊙ 1 L.....	h. m. s.	s.	h. m. s.	h. m. s.	m. s.	m. s.	h. m. s.	
S.	⊙ 2 L.....	18 10 54,99					3 12,20		
	Polaris.....	1 3 35,41	- 38,96	1 2 56,45	0 59 45,70	*3 10,75	3 12,67		
	δ Cassiopeæ....	1 18 6,31	- 4,32	1 18 1,99	1 14 49,61	3 12,38	3 12,70	1 14 49,29	
	μ Piscium.....	1 24 35,88	- 2,75	1 24 33,13			3 12,70	1 21 20,43	S. at 1h. 31m.
	α Eridani.....	1 34 39,43	- 1,85	1 34 37,58			3 12,71	1 31 24,87	3 = 3' 12", 71
	τ Ceti.....	1 39 28,56	- 2,44	1 39 26,12	1 36 13,23	3 12,89	3 12,72	1 36 13,40	
	γ Arietis,.....	1 47 32,10	- 2,93	1 47 29,17	1 44 16,32	3 12,85	3 12,73	1 44 16,44	
	50 Cassiopeæ..	1 52 28,83	- 6,62	1 52 22,21	1 49 10,55	*3 11,66	3 12,73	1 49 9,48	

* These are omitted in taking the mean.

		NAMES.	App. A.R. + Clock Er.	Aber- ration, &c.	Mean A.R. Jan. 1, 1831 + Clock Er.	Greenwich Mean A.R. Jan. 1, 1831.	Difference or Clock Error.	Error computed from Mean.	Mean Places Jan. 1, 1831.	Mean Errors of Clock.
1831			<i>h. m. s.</i>	<i>s.</i>	<i>h. m. s.</i>	<i>h. m. s.</i>	<i>m. s.</i>	<i>m. s.</i>	<i>h. m. s.</i>	
Dec.	24	62 Ceti.....	2 4 10,03	—	2,94	2 4 7,09		3 14,82	2 0 52,27	
R.		θ ¹ Arietis.....	2 12 2,43	—	3,37	2 11 59,06	2 8 44,48	3 14,58	2 8 44,24	
		* Ceti.....	2 18 47,30	—	3,12	2 18 44,18		3 14,83	2 15 29,35	
		κ Eridani.....	2 24 4,46	—	2,30	2 24 2,16		3 14,83	2 20 47,33	
		* Ceti.....	2 30 7,52	—	3,12	2 30 4,40		3 14,81	2 26 49,59	
		ε Ceti.....	2 34 41,08	—	2,85	2 34 38,23	2 31 23,70	3 14,53	2 42 23,42	
		π Ceti.....	2 39 22,71	—	2,85	2 39 19,86	2 36 4,94	3 14,92	2 36 5,01	
		c Arietis.....	2 43 21,74	—	3,67	2 43 18,07	2 40 3,26	3 14,81	2 40 3,21	
R. + 1 ^h , 65		ε Arietis.....	2 52 52,34	—	3,57	2 52 48,77	2 49 33,86	3 14,91	2 49 33,90	
		ρ ² Eridani.....	2 57 42,85	—	3,01	2 57 39,84		3 14,88	2 54 24,96	
		ζ Eridani.....	3 10 55,88	—	3,05	3 10 52,83	3 7 37,76	3 15,07	3 7 37,94	
		ο Tauri.....	3 19 2,12	—	3,40	3 18 58,72	3 15 43,78	3 14,94	3 15 43,82	R. at 3h. 19m.
		σ Persei.....	3 22 1,64	—	4,88	3 21 56,76		3 14,90	3 18 41,86	12 = 3' 14", 90
		z Eridani.....	3 27 30,15	—	2,91	3 27 27,24		3 14,91	3 24 12,33	
		E Tauri.....	3 31 33,56	—	3,00	3 31 30,56		3 14,91	3 28 15,65	
		ν Persei.....	3 37 4,11	—	4,64	3 36 59,47	3 33 44,39	3 15,08	3 33 44,55	
		f Pleiadum...	3 42 26,40	—	3,88	3 42 22,52		3 14,92	3 39 7,60	
		ζ Persei.....	3 46 50,68	—	4,24	3 46 46,44	3 43 31,61	3 14,83	3 43 31,51	
		c Persei.....	3 59 45,35	—	5,09	3 59 40,26		3 14,95	3 56 25,31	
		δ ¹ Tauri.....	4 16 30,72	—	3,82	4 16 26,90	4 13 11,87	3 15,03	4 13 11,93	
		θ ² Tauri.....	4 22 20,23	—	3,79	4 22 16,44	4 19 1,36	3 15,08	4 19 1,46	
		d Tauri.....	4 29 41,11	—	3,67	4 29 37,44		3 14,99	4 26 22,45	
		c ² Tauri.....	4 33 57,96	—	3,73	4 33 54,23		3 14,99	4 30 39,24	
		β Cæli Scalp...	4 39 23,32	—	3,00	4 39 20,32		3 15,00	4 36 5,32	
		ο ¹ Orionis.....	4 46 17,78	—	3,82	4 46 13,96		3 15,01	4 42 58,05	
		d ¹ Camelopardi.	4 51 46,75	—	6,70	4 51 40,05		3 15,01	4 48 25,04	
		γ Aurigæ.....	4 58 0,52	—	4,88	4 57 55,64	4 54 40,57	3 15,07	4 54 40,62	
		ρ Orionis.....	5 7 46,31	—	3,62	5 7 42,69		3 15,03	5 4 27,66	
		d Leonis.....	10 55 8,00	—	2,67	10 55 5,33	10 51 49,79	3 15,54	3 15,53	10 51 49,80
) 2 L.....	11 33 51,26	—						R. at 11h. 31m.
		ξ ¹ Virginis.....	11 39 52,12	—	2,38	11 39 49,74	11 36 34,10	3 15,64	3 15,51	11 36 34,23
		β Virginis.....	11 45 11,53	—	2,22	11 45 9,31	11 41 53,78	3 15,53	3 15,58	11 41 53,73
		π Virginis.....	11 55 30,48	—	2,28	11 55 28,20	11 52 12,08	3 16,12	3 15,69	11 52 12,51
		ο Virginis.....	11 59 53,73	—	2,23	11 59 51,50	11 56 35,95	3 15,55	3 15,60	11 56 35,90
25		⊙ 2 L.....	18 16 36,27							
R.		* Ceti.....	0 19 10,41	—	2,23	0 19 8,18		3 16,45	0 15 51,73	
		28 Andromedæ..	0 24 34,26	—	2,62	0 24 31,68		3 16,46	0 21 15,22	
		53 Piscium.....	0 31 18,61	—	2,50	0 31 16,11		3 16,46	0 27 59,65	
		* Ceti.....	0 37 1,46	—	2,25	0 36 59,21		3 16,47	0 33 42,74	
		i Piscium.....	0 44 8,84	—	2,76	0 44 6,08		3 16,48	0 40 49,60	
		k Piscium.....	0 49 15,21	—	2,80	0 49 12,41		3 16,49	0 45 55,92	
		φ ⁴ Ceti.....	0 53 35,03	—	2,32	0 53 32,71		3 16,49	0 50 16,21	
		ψ ¹ Piscium.....	0 59 57,49	—	2,76	0 59 54,73		3 16,50	0 56 38,23	
		39 Ceti.....	1 11 20,79	—	2,55	1 11 18,24		3 16,51	1 8 1,73	
		δ Cassiopeæ...	1 18 10,62	—	4,29	1 18 6,33	1 14 49,61	3 16,72	3 16,52	1 14 49,81
		μ Piscium.....	1 24 39,68	—	2,73	1 24 36,95	1 21 20,21	3 16,74	3 16,53	1 21 20,42
		π Piscium.....	1 31 28,48	—	2,87	1 31 25,61	1 28 9,07	3 16,54	3 16,53	1 28 9,08
		107 Piscium.....	1 36 39,84	—	3,04	1 36 36,80		3 16,54	1 33 20,26	
		τ Ceti.....	1 39 32,08	—	2,43	1 39 29,65	1 36 13,23	3 16,42	3 16,54	1 36 13,11
		γ Arietis.....	1 47 35,92	—	2,96	1 47 32,96	1 44 16,32	3 16,64	3 16,55	1 44 16,41
		50 Cassiopeæ...	1 52 33,54	—	6,51	1 52 27,03	1 49 10,55	3 16,48	3 16,56	1 49 10,47

† The Madras Observations indicate an error of 0,6 in the Greenwich determination of the A.R. of this Star it is therefore omitted.

		App. A.R. + Clock Er.	Aber- ration, &c.	Mean A.R. Jan. 1, 1831 + Clock Er.	Greenwich Mean A.R. Jan. 1, 1831.	Difference or Clock Error.	Error computed from Mean.	Mean Places Jan. 1, 1831.	Mean Errors of Clock.
1831									
Dec.	25 ^a	<i>h. m. s.</i>	<i>s.</i>	<i>h. m. s.</i>	<i>h. m. s.</i>	<i>m. s.</i>	<i>m. s.</i>	<i>h. m. s.</i>	R. at 1h. 52m.
		1 56 45,10	— 1,84	1 56 43,26			3 16,56	1 53 26,70	9 = 3' 16",56
	62	2 4 11,21	— 2,93	2 4 8,28			3 16,57	2 0 51,71	
	θ^1	2 12 4,21	— 3,28	2 12 0,93	2 8 44,48	16,45	3 16,58	2 8 44,34	
	κ	2 24 6,26	— 2,28	2 24 3,98			3 16,59	2 20 47,39	
	U	2 30 45,45	— 3,22	2 30 42,23			3 16,60	2 27 25,63	
	e ¹	2 34 43,00	— 2,84	2 34 40,16	2 31 23,70	16,46	3 16,60	2 31 23,56	
	ϵ	2 40 19,73	— 1,96	2 40 17,77			3 16,61	2 37 1,16	
	ρ^1	2 56 11,51	— 3,00	2 56 8,51			3 16,63	2 52 51,88	
	β	3 0 33,04	— 4,29	3 0 28,75	2 57 12,11	16,64	3 16,64	2 57 12,11	
	ζ	3 8 32,31	— 3,64	3 8 28,67			3 16,65	2 5 12,02	
	S. E	3 31 33,24	— 3,27	3 30 29,97			3 14,25	3 27 15,72	
	ν	3 37 3,25	— 4,64	3 36 58,61	3 33 44,39	3 14,22	3 14,25	3 33 44,36	
	f	3 42 25,41	— 3,88	3 42 21,53			3 14,26	3 30 7,27	
	ζ	3 46 50,09	— 4,18	3 46 45,91	3 43 31,61	3 14,30	3 14,26	3 43 31,65	
	c	3 59 44,34	— 5,09	3 59 39,25			3 14,28	3 56 24,97	
	μ	4 5 50,21	— 5,18	4 5 45,03	4 2 31,09	3 13,94	3 14,29	4 2 30,74	
R. + 1",76	α	4 11 41,27	— 2,86	4 11 38,41			3 14,30	4 8 24,11	
S. + 1",42	δ	4 16 30,26	— 3,82	4 16 26,44	4 13 11,87	3 14,57	3 14,31	4 13 12,13	
Mean.	ϵ	4 22 3,87	— 3,89	4 21 59,98	4 18 45,45	3 14,53	3 14,32	4 18 45,66	S. at 4h. 46m.
= + 1",64	d	4 29 40,97	— 3,96	4 29 37,01			3 14,32	4 26 22,69	13 = 3' 14",34
	τ	4 35 25,53	— 4,04	4 35 21,49	4 32 6,82	3 14,67	3 14,33	4 32 7,16	
	μ	4 40 21,29	— 3,44	4 40 17,85			3 14,33	4 37 3,52	
	σ^1	4 46 17,23	— 3,82	4 46 13,41			3 14,34	4 42 59,07	
	d ¹	4 51 45,75	— 6,70	4 51 39,05	4 48 25,16	3 13,89	3 14,34	4 48 24,71	
	η	4 57 59,44	— 4,89	4 57 54,55	4 54 40,57	3 13,98	3 14,35	4 54 40,20	
	ρ	5 7 45,71	— 3,63	5 7 42,08			3 14,36	5 4 27,72	
	ν	5 15 26,72	— 3,38	5 15 23,34			3 14,37	5 12 8,97	
	γ	5 19 22,45	— 3,70	5 19 18,75	5 16 4,20	3 14,55	3 14,37	5 16 4,38	
	χ	5 25 2,84	— 4,48	5 24 58,36			3 14,38	5 21 43,98	
	ι	5 30 27,93	— 3,49	5 30 24,44	5 27 10,22	3 14,22	3 14,39	5 27 10,05	
	12	5 38 25,37	— 3,29	5 38 22,08			3 14,40	5 35 7,68	
	ζ	5 42 35,83	— 3,38	5 42 32,45	5 39 18,13	3 14,32	3 14,40	5 39 18,05	
	β	5 48 18,06	— 3,20	5 48 14,86	5 45 0,21	3 14,65	3 14,41	5 45 0,45	
	η	5 52 0,56	— 3,37	5 51 57,19	5 48 42,67	3 14,52	3 14,42	5 48 42,77	
	χ^3	5 57 11,54	— 4,06	5 57 7,48			3 14,42	5 53 53,06	
	a	6 5 40,17	— 6,93	6 5 33,24			3 14,43	6 2 18,81	
	d	6 15 11,72	— 5,44	6 15 6,28			3 14,44	6 11 51,84	
	α	6 22 17,41	— 3,27	6 22 14,14			3 14,45	6 18 59,69	
	α	14 11 13,56	— 1,16	14 11 12,40	14 7 57,37	3 15,03	3 15,02	14 7 57,38	

The above errors of the Clock shew that differences to the astonishing amount of 2",07 and 2",39 existed between the times at which the observers R and S noted the passage of Stars on the 24th and 25th December respectively, whereas we have found at page 41 that a difference of only half a second occurred in the observations of these observers on the 31st May and 1st June, it will consequently be interesting to inspect the differences for each day which are here subjoined.

CLOCK ERRORS AND CLOCK RATES.

Observer R noted the time in excess above observer S.

1831		REMARKS.	1831		REMARKS.
April	18	+ 0,11	Sept.	3	+ 1,11
	19	+ 0,31		4	
	20	- 0,11		9	+ 1,05
	21	- 0,17		11	+ 1,00
	22	+ 0,10		13	+ 1,05
	23	+ 0,30		14	+ 1,06
	24	+ 0,17		15	+ 0,90
May	15	+ 0,56		20	+ 1,26
	16	+ 0,46		21	+ 1,53
	17	+ 0,26		22	+ 1,69
	18	+ 0,29		24	+ 1,89
	19	+ 0,28		29	+ 2,70
	20	+ 0,79		30	+ 2,12
	21	+ 0,59	Oct.	2	+ 1,98
	22	+ 0,56		3	+ 1,70
	23	+ 0,28		5	+ 2,52
	26	+ 0,53		6	+ 2,56
	27	+ 0,52		8	+ 2,32
	29	+ 0,62		14	+ 2,74
	30	+ 0,56		17	+ 2,77
	31	+ 0,47		21	+ 2,36
June	1	+ 0,55		22	+ 2,53
	3	+ 0,77		23	+ 2,68
	4	+ 0,57		25	+ 2,70
	5	+ 0,60		30	+ 3,21
	6	+ 0,60		31	+ 2,53
	7	+ 0,87	Nov.	1	+ 2,93
	8	+ 0,97		2	+ 2,91
	15	+ 0,98		3	+ 2,62
	20	+ 0,89		4	+ 2,65
	27	+ 1,52		5	+ 2,92
	28	+ 0,68		6	+ 2,55
	29	+ 1,45		8	+ 2,74
	30	+ 1,10		9	+ 2,61
July	3	+ 1,20		10	+ 2,60
	7	+ 1,02		20	+ 2,48
	10	+ 0,83		21	+ 2,35
	12	+ 0,69		27	+ 2,18
	13	+ 0,75	Dec.	2	+ 2,26
	14	+ 0,54		3	+ 2,26
	17	+ 0,60		5	+ 3,36
	22	+ 0,72		6	+ 3,43
	25	+ 0,58		7	+ 2,98
	28	+ 0,69		8	+ 2,73
Aug.	1	+ 0,86		12	+ 2,63
	9	+ 0,93		13	+ 2,58
	10	+ 0,51		15	+ 2,28
	11	+ 0,38		16	+ 1,99
	12	+ 0,91		19	+ 1,96
	23	+ 0,57		20	
	26	+ 0,51		21	+ 2,05
	28	+ 0,45		24	+ 2,07
	29	+ 0,61		25	+ 2,39
	30	+ 1,45		29	+ 2,56
Sept.	1	+ 1,09			

5 observ. by S and 10 by R.

one observation by S.

only one Star observed.

one observation.

M - R = 0,16.

On comparing recent observations I find that observers R, T and M very nearly agree in estimating the instant of the passage of a Star, this it will be recollected was found to be the case with the observations at the earlier part of the year, hence it is fair to presume that the change which has taken place between R and S is chargeable on the latter observer only : to account for this difference which is found between one observer and another we are led to suppose an error either in the ear or eye, or possibly in both of these organs ; now any error arising from the former would be constant with the same observer for every Star, whereas the latter would introduce an error depending on the nature and magnitude of the body observed and on the apparent rapidity of its motion through the field of the telescope ; thus, if from an over anxiety peculiar to young observers, the first edge of the wire or the first edge of the Star should be observed instead of the centre then in the former case an error would be committed (with the lines now in use) to the amount of $\frac{0^{\circ},073}{\sin. N.P.D.}$ and in the latter (for bright Stars) to probably five times this amount, but these are wholly insufficient to account for the enormous differences above found, it will however be desirable to ascertain if the Clock errors vary with the declination of the Star, for this purpose I have reduced each separate error to the same instant of time by means of the daily Rate for the sake of comparison as follows.

Comparison of the Clock errors as found by observer S.

Days.	NAMES.	A. R.	Error of Clock at		Declina- tion.
			Passage.	<i>h. m.</i> I 31	
1831		<i>h. m.</i>	<i>m. s.</i>	<i>m. s.</i>	<i>° ' "</i>
December 24	τ Ceti.....	1 39	+ 3 12,89	+ 3 12,88	- 16 50
	γ Arietis.....	1 47	12,85	12,83	+ 18 27
	ϵ Cassiopeæ.....	1 18	12,38	12,40	+ 59 21
	β Cassiopeæ.....	1 52	11,66	11,64	+ 71 36
				<i>h. m.</i> Passage. at 4 46	
25	ι Orionis.....	5 30	+ 3 14,22	+ 3 14,17	- 6 1
	γ Orionis.....	5 16	14,55	14,52	+ 6 11
	η Leporis.....	5 52	14,52	14,44	- 14 12
	ζ Leporis.....	5 42	14,32	14,25	- 14 53
	α Tauri.....	4 16	14,57	14,60	+ 17 8
	ϵ Tauri.....	4 22	14,53	14,56	+ 18 48
	α Bootis.....	14 11	15,03	14,38	- 20 9
	τ Tauri.....	4 35	14,67	14,68	+ 22 37
	ζ Persei.....	3 47	14,30	14,37	+ 31 22
	β Columbæ.....	5 48	14,65	14,59	- 35 50
	η Aurigæ.....	4 54	13,98	13,97	+ 41 0
	ν Persei.....	3 37	14,22	14,30	+ 42 2
	μ Persei.....	4 5	13,94	13,99	+ 47 58
	δ Camelopard.....	4 51	13,89	13,88	+ 60 11

The above two cases shew that an error varying with the declination of the Star really does exist; to ascertain its value let x represent the true error of the Clock, and e the error committed, and we have the following equations of conditions.

December 24.	December 25.
<i>m. s.</i>	<i>m. s.</i>
$x - 3 \ 12,88 = \frac{e}{,957}$	$x - 3 \ 14,17 = \frac{e}{,994}$
$x - 3 \ 12,83 = \frac{e}{,949}$	$x - 3 \ 14,52 = \frac{e}{,994}$
$x - 3 \ 12,40 = \frac{e}{,510}$	$x - 3 \ 14,44 = \frac{e}{,969}$
$x - 3 \ 11,64 = \frac{e}{,316}$	$x - 3 \ 14,25 = \frac{e}{,966}$
	$x - 3 \ 14,60 = \frac{e}{,956}$
	$x - 3 \ 14,56 = \frac{e}{,947}$
	$x - 3 \ 14,38 = \frac{e}{,939}$
	$x - 3 \ 14,68 = \frac{e}{,923}$
	$x - 3 \ 14,37 = \frac{e}{,854}$
	$x - 3 \ 14,59 = \frac{e}{,811}$
	$x - 3 \ 13,97 = \frac{e}{,755}$
	$x - 3 \ 14,30 = \frac{e}{,743}$
	$x - 3 \ 13,99 = \frac{e}{,670}$
	$x - 3 \ 13,88 = \frac{e}{,497}$

eliminating e so that the sum of the squares of the errors may be a minimum from the observations of the 24th we have $e = ^s,58$

 25th $= ^s,50$

Hence to turn the above observations to account it becomes necessary to apply to each *mean error* of the Clock a correction

$\frac{1}{n} \left(\frac{e}{\sin. P.} + \frac{e}{\sin. P.^1} + \frac{e}{\sin. P.^2} + \&c. \frac{e}{\sin. P.^n} \right)$ where P. P.¹ express the

Polar distance of the several Stars which are employed in determining the Clock's error; and in computing the column δ from this *corrected mean error*,

a correction $\frac{e}{\sin. P.}$ must be employed. I have here supposed the error to

be committed by the observer; but it is plain that an error of the Instrument

either in Azimuth or Collimation would equally well reconcile all the above observations; to decide upon this point we will now examine the observations by R on these two days; arranging them in order of declination we have

December 24.					December 25.				
NAMES.	A. R.	Error of the Clock at		Declination.	NAMES.	A. R.	Error of the Clock at		Declination.
		Passage.	<i>h. m.</i> 3 19				Passage.	<i>h. m.</i> 1 52	
	<i>h. m.</i>	<i>m. s.</i>	<i>m. s.</i>	'		<i>h. m.</i>	<i>m. s.</i>	<i>m. s.</i>	'
θ Tauri...	3 19	+ 3 14,94	3 14,94	+ 8 26	μ Piscium..	1 21	- 3 16,74	3 16,77	+ 5 16
ζ Eridani..	3 7	15,07	15,08	- 9 27	π Piscium..	1 31	16,54	16,57	+ 11 16
ϵ Ceti.....	2 31	14,53	14,58	- 12 36	ϵ^2 Ceti... .	2 34	16,46	16,41	- 12 36
π Ceti.....	2 36	14,92	14,97	- 14 34	τ Ceti... .	1 36	16,42	16,43	- 16 50
θ^2 Tauri...	4 19	15,08	15,02	+ 15 29	ν Arietis...	1 44	16,64	16,65	+ 18 28
δ^1 Tauri...	4 13	15,03	15,97	+ 17 80	θ^1 Arietia..	2 12	16,45	16,43	+ 19 70
θ^1 Arietis..	2 52	14,58	14,65	+ 19 70	β Persei... .	2 57	16,64	16,57	+ 40 58
ϵ Arietis... .	2 52	14,91	14,94	+ 20 40	δ Cassiop... .	1 14	16,72	16,76	+ 59 21
ϵ Arietis... .	2 40	14,81	14,85	+ 26 34	50 Cassiop... .	1 52	16,48	16,48	+ 71 36
ζ Persei... .	3 43	14,83	14,81	- 31 22					
η Aurigæ... .	4 54	15,07	14,98	- 41 00					
ν Persei... .	3 33	15,08	15,07	- 42 20					

Here we find that the Clock errors are in no wise affected by the declination of the Star; and further, I may remark that this solitary instance is confirmed by the totality of the observations during the year; whence it appears that by observers R, T and M the Clock errors given by Stars wherever situated agree among themselves; but that the observations by S with but one or two exceptions exhibit an error ϵ , of the nature above found as follows.

1831	Value of ϵ	REMARKS.	1831	Value of ϵ	REMARKS.
	<i>s.</i>			<i>s.</i>	
April 22	0,29		June 6	0,41	one observation only
23	0,62		7	0,41	
24	0,00		8	0,28	one observation
25	0,50		21	0,26	
26	0,44		July 7	0,36	
May 1	0,36		10	0,32	
2	0,61		11	0,99	
6	1,20		12	0,90	
12	0,83		13	0,80	
16	0,68		13	0,37	
18	1,10		22	0,71	
20	0,42		25	0,21	
26	0,45		28	0,28	
31	0,73		Aug. 9	0,93	
June 2	0,18		10	0,90	} one observation affording but a small coefficient I have used
4	0,60		11	1,75	
5	0,40		23	1,31	

1831	Value of e	REMARKS.	1831	Value of e	REMARKS.
	<i>s.</i>			<i>s.</i>	
Aug. 26	1,47	} only one observ. the observations by R indicate $e = 0^s,27$ from 3 observ. on the 29th & 30th.	Oct. 22	1,29	} only one observation
29	2,21		23	1,60	
30	2,10		25	1,78	
31	0,80		30	1,42	
Sept. 1	1,06	} δ Urs Min gives $0^s,30$ for e observations by R $0^s,35$	31		
2			Nov. 2	1,70	
3	1,98	} δ Urs Min by R gives $- 0^s,03$ other Stars — — + $0^s,30$	8	1,64	
4			9	0,90	
9	0,30		12	1,25	
13	0,93		20	0,55	
14	1,00	} δ Urs Min $e = 0^s,00$ δ Urs Min $e = 0^s,30$	21	0,95	
15	1,20		30	0,33	
16	0,30	} δ Urs Min	Dec. 1	0,00	
20	1,31		3	0,93	
22	1,20	} A very good set of observations by R give $e = 0^s,03$ R gives $e = 0^s,00$ Morning observations	5	1,00	
24	1,08		7	0,32	
28	1,28		8	0,73	
29	1,25		12	1,23	
Oct. 2	1,45		16	0,73	
2	0,75		18	0,66	
3	0,94		19	0,76	
6	0,76		24	0,60	
7	0,89	25	0,52		
12	1,43	26	1,70		
14	1,34	28	1,08		
17	1,67	29	0,68		
21	0,71	30	1,12		

Two observations of δ Urs Min by R on 10th and 11th of August give $e = 0,04$ and $0,00$ respectively.
One ————— by T on the 27th ————— give $e = 0,33$

On employing with the observations by S the value of e found above, the agreement is such as to forbid their being rejected, which at the commencement of the enquiry I feared would be a necessary measure; in the case of the Sun, Moon and Planets, and of Stars situated above 30° from the Pole the observations by observer S will I estimate be doubtful to less than one tenth of a second; I have therefore reduced all the observations by S with these values rejecting only the observations of those Stars which are situated within this limit, but these I am happy to state are few and far between. The observations by R indicate an error $e = 0^s,30$ between the 27th August and 4th September, but since this does not occur at any other period during the year, (in the absence of any cause to explain why it should thus happen) in cases where its effect is less than one tenth of a second I have preferred not to allow for it, and to reject the remaining observations.

The Table which now follows exhibits the daily rate of the Clock as determined by the Sun and by Stars; from the column of differences which is annexed, some sort of estimate can be formed of the dependance to be placed in a single observation of the Sun.

Comparison of the Rate of the Transit Clock as determined by the Sun with that determined by the Stars.

1831	Clock Rate by		Difference.	REMARKS.	1831	Clock Rate by		Difference.	REMARKS.
	Sun.	Stars.				Sun.	Stars.		
Feb. 18 & 19	s.	s.	s.		April	s.	s.	s.	
20		-4,49			6	+ 0,02	+ 0,03	0,01	
21	-3,45	-3,52	0,07		7	- 0,23	+ 0,02	0,25	
22				The Clock stopt for 23 seconds without any apparent cause.	8		- 0,23		
23		-4,36			9	+ 0,20			
24	-3,95	-3,75	0,20		10		+ 0,16		
25	-3,49	-3,80	0,31		11		- 0,25		
26		-3,96			12		*		
27	-4,27	-4,13	0,14		13	+ 2,21	+ 2,46	0,25	
28		-4,38			14	+ 2,31	+ 2,57	0,26	
March		-4,51		Stopt 10s. in winding.	15		+ 1,47		
1				Regulated.	16	+ 1,29	+ 1,05	0,24	
2				Stopt 6 seconds.	17		+ 1,07		
3				Stopt 10 seconds.	18	- 0,18	+ 0,75		
4				I applied oil to the escapement.	19	+ 1,11	+ 0,82	0,29	
5		+ 7,88			20	+ 0,65	+ 0,85	0,20	
6		+ 7,77			21				The Clock stopt about 10s. without any apparent cause.
7	-7,30	+ 7,70	0,40		22		+ 1,48		
8				Regulated.	23		+ 1,08		
9		- 0,30			24		+ 0,61		
10	- 0,39	- 0,38	0,01		25	+ 0,19	+ 0,56	0,36	
11	+ 0,04	- 0,00	0,40		26		+ 0,81		
12	- 0,26	- 0,00	0,26		27		+ 0,50		
13	+ 0,05	- 0,01	0,06		28		+ 0,58		
14	- 0,02	- 0,25	0,23		29	+ 0,99	+ 0,85	0,14	
15	- 0,17				30	+ 0,59	+ 0,53	0,06	
16		- 0,19			May	1		+ 0,65	
17	+ 0,67	+ 1,20		The Clock had stopt 1s. in the night.	2	+ 0,63	+ 0,82	0,19	
18	+ 0,62	+ 0,55	0,07		3	+ 1,20	+ 1,19	0,01	
19		+ 0,78			4		+ 1,15		
20	+ 0,55	+ 0,75	0,20		5		+ 1,96		
21		+ 0,88			6		+ 1,22		
22	+ 0,68	+ 0,35	0,33		7	+ 0,77	+ 0,95	0,18	
23	+ 0,03	+ 0,10	0,07		10	+ 0,86	+ 0,99		
24	+ 0,28	+ 0,23	0,05		11		+ 0,91		
25	+ 0,17	+ 0,55	0,38		12	+ 0,69			
26	+ 0,75	+ 0,42	0,33		13	+ 0,53	+ 0,58	0,05	
27		- 0,10			14	+ 0,93	+ 0,63	0,30	
28	+ 0,22	+ 0,45	0,23		15		+ 0,51		
29	+ 0,20	- 0,05	0,25		16	+ 0,22	+ 0,30	0,08	
30	- 0,23	- 0,07	0,16		17	+ 0,40	+ 0,22	0,18	
31	- 0,18	- 0,30	0,12		18	+ 0,52	+ 0,32	0,20	
April		+ 0,37			19	+ 0,28	+ 0,26	0,02	
1		+ 0,29	0,06		20	+ 0,25	+ 0,55	0,30	
2	+ 0,23				21	+ 0,57	+ 0,26	0,30	
3		+ 0,40			22	+ 0,22	+ 0,26	0,04	
4		- 0,24			23	+ 0,47	+ 0,25	0,22	
5		+ 0,29							

* Evening observations from 4 to 8 P.M. give the Rate - 0,13

— 8 to 12 P.M. — — — + 0,09

Next morning — 6 to 9 A.M. — — — + 0,82

1831	Clock Rate by			REMARKS.	1831	Clock Rate by			REMARKS.
	Sun.	Stars.	Difference.			Sun.	Stars.	Difference.	
May	24	<i>s.</i> + 0,24	<i>s.</i> + 0,11	<i>s.</i> 0,13	July	24	<i>s.</i> + 0,52	<i>s.</i> + 0,42	<i>s.</i> 0,10
	25	+ 0,11	+ 0,13	0,02		25		+ 0,42	
	26	- 0,24	- 0,10	0,14		26	+ 0,50	+ 0,57	0,07
	27	- 0,14	+ 0,40	0,17		27		+ 0,24	
	28		+ 0,60			28	+ 0,46		
	29		+ 0,67			29		+ 0,27	
	30		+ 0,22			30	+ 0,30	+ 0,80	0,50
	31	+ 0,73	+ 0,38	0,35	August	31	+ 0,57	} + 0,56	Only two Stars observed the rate by the Sun is therefore entitled to the same weight as that by the Stars.
June	1	- 0,10	+ 0,45	0,55		1			
	2	+ 0,61	+ 1,02	0,41	2	+ 0,51			
	3	+ 1,45	+ 0,96	0,49	3				
	4	+ 0,73	+ 1,01	0,28	4	+ 0,64			
	5	+ 0,26	+ 0,52	0,26	5	+ 0,69	+ 0,63	0,06	
	6	+ 0,74	+ 0,81	0,07	6	+ 0,81			
	7	+ 1,28	+ 0,74	0,54	7	+ 0,36			
	8	+ 0,45	+ 0,88	0,43	8	+ 0,93	+ 0,66	0,27	
	9	+ 0,84	- 0,73	0,11	9				
	10	+ 0,32	} - 0,60		10	+ 0,55	+ 0,63	0,08	
	11	+ 1,23				11		+ 0,66	
	12	+ 0,68	} - 0,73		12	+ 0,45	+ 0,65	0,20	
	13	+ 1,02				13	+ 0,69		
	14	+ 0,87	} - 0,87		14	+ 0,23	+ 0,69		
	15	} + 0,86				18	+ 0,83		
	16					20	+ 0,69		
	19		} - 0,78		21		} + 0,82		
	20	+ 0,64				22		+ 0,56	} + 0,46
	21		+ 0,68		23				
	22	+ 0,52	+ 0,77	0,25	24	+ 0,33	+ 0,29	0,04	
	23	+ 0,47	} + 1,06	... The Clock has probably tript 1s. between the 24th and 25th.	25	+ 0,27	+ 0,47	0,20	
	24	+ 0,50					26	} + 0,54	+ 0,55
	25	+ 1,67				27	+ 0,58		
	26		+ 0,50			28	+ 0,56		
	27	+ 0,64			29	+ 0,89	+ 0,88	0,01	
	28	+ 0,61	+ 0,71	0,10	30		+ 0,87		
	29	+ 0,34	+ 0,41	0,07	31		+ 1,33		
	30	+ 0,65	+ 0,63	0,02	September	1	+ 1,06	+ 0,55	0,49
July	1		+ 0,58			2		+ 0,95	
	2	+ 0,31	+ 0,49	0,18	3		+ 1,01		
	3	+ 0,16			4		+ 1,17		
	4		+ 1,07		5				
	5				7	+ 0,64	+ 0,77	0,13	
	6				8	+ 1,14	} + 0,73		
	7	+ 0,45	+ 0,60	0,15	9				
	8	+ 0,25	+ 0,43		10				
	9				11	+ 0,98	} + 0,82		
	10	+ 0,55	+ 0,56	0,01	12	+ 1,04			
	11		+ 0,24		13	+ 0,39	+ 0,79	0,40	
	12		+ 0,33		14	+ 0,78	+ 0,56	0,22	
	13	+ 0,21	+ 0,07	0,14	15	+ 0,53	+ 0,85	0,32	
	14		+ 0,22		16	+ 1,18	+ 0,70	0,48	
	15				17	+ 0,62	+ 0,80	0,18	
	16	+ 0,56	+ 0,62		19	+ 0,90			
	17		+ 0,48		20		+ 0,93		
	21	+ 0,83	+ 0,80	0,03	21	} + 1,14	+ 0,99		
	22				22		+ 0,54	+ 1,04	0,50

1831	Clock Rate by		Difference.	REMARKS.	1831	Clock Rate by		Difference.	REMARKS.
	Sun.	Stars.				Sun.	Stars.		
September 24	s. + 1,24	s. + 1,07	s. 0,17		November 18	s.	s.	s.	
25	+ 0,67				19	+ 1,32			
26	+ 1,07	+ 0,70	0,37		20		+ 1,17		
27	+ 0,52				21	+ 1,64	+ 1,19	0,45	
28	+ 0,48	+ 0,69	0,21		22	+ 1,53	+ 1,62	0,09	
29	+ 0,35	+ 0,28	0,07		27		+ 1,49		
30		+ 0,96			28		+ 1,18		
October 2		+ 0,94			29				
3		+ 1,25			30	+ 1,09			
4		+ 0,89			December 1		+ 1,49		
5	+ 0,72	+ 0,93	0,21		2	+ 1,32	+ 1,12	0,20	
6	+ 0,84	+ 0,97	0,13		3	+ 1,10	+ 1,41	0,31	
7	+ 0,99	+ 1,02	0,03		4	+ 1,17	+ 1,11	0,06	
8	+ 1,02	+ 1,09	0,07		5	+ 1,24			
12	+ 1,13	+ 1,04	0,09		6	+ 1,07	+ 1,44	0,37	
13	+ 1,19	} + 1,02			7	+ 1,39	+ 1,30	0,09	
14	+ 1,18				8	+ 1,22	+ 1,68	0,46	
15	+ 0,75	} + 0,98			9	} + 1,69	+ 1,75		
16					10		+ 1,56		
17	+ 1,08				11	} + 1,97	+ 1,79		
20		+ 1,15		12					
21	+ 1,24	+ 1,13	0,11		13	+ 1,67	+ 1,45	0,22	
22	+ 1,08	+ 1,15	0,07		14	+ 1,54	+ 1,64	0,10	
23	+ 0,77	+ 1,03	0,26		15	+ 1,64	+ 1,94	0,30	
24	} + 1,29	+ 1,01			16		+ 1,50		
25			+ 1,32	17	+ 1,48	} + 1,51		
29		+ 1,31		18	} + 1,56				
30		+ 1,19		19					
31		+ 0,99		20	} + 1,35	+ 1,29			
November 1	+ 1,39	+ 1,19	0,20	21			+ 1,30		
2	+ 1,22	+ 1,10	0,12		22	+ 1,26	} + 1,33		
3	+ 1,43	+ 1,36	0,07		23	+ 1,57			
4		+ 1,20			24	+ 1,43	+ 1,65	0,22	
5		+ 1,45			25		+ 1,66		
6		+ 1,26			26	+ 1,51	+ 1,82	0,31	
8		+ 1,18			28 to 29	} + 1,75	+ 1,73		Only two observations
9	+ 1,38	+ 1,22	0,16		30			+ 1,31	0,44
10	+ 1,42	+ 1,30	0,12		31				
12	+ 1,26	+ 1,21	0,05						

Taking the mean it appears that the mean error of each rate by the Sun = 0,199 or the mean uncertainty of a single observation of the Sun may be stated at this amount, whereas for Stars the uncertainty does not I imagine much exceed the half of this.

OF THE MURAL CIRCLE.

This Instrument which is represented by Plate III, was constructed by Dollond on purpose for this Observatory; it consists of a circular brass ring 48 Inches diameter, firmly attached by sixteen conical spokes to the base of a hollow but strong conical axis; this axis is 3 feet long and from $5\frac{3}{4}$ Inches diameter at the end next the circle tapers down to a diameter of $2\frac{1}{2}$ Inches; the two ends E, e, Plate IV fig. 1. are accurately ground to admit of their working into collars *ab*, *ac*, attached to the ends of a hollow conical frustrum through which the axis passes; this conical frustrum is strengthened at its base *b* by the metal forming the collar in which the axis of the circle works, and is likewise strengthened at the smaller end *c* by the collar in which the other end of the axis works; the axis of the circle is made to fit the collars and retained in its place to any required degree of tightness by a plate of brass fastened by six screws to the smaller end of the axis, working upon the collar fig. 3; the whole frustrum passing through an aperture in the pier, reposes at the end *b* next the circle upon a flat brass plate *a*, *a*, fig. 2. which is firmly cemented and screwed to the stone pier; and is kept steady by a steel pin working into the collar at *b*; the further or smaller end is secured by four antagonizing screws *s*, *s*, *s*, *s*, fig. 3. working into a strong brass frame likewise very securely cemented and screwed to the pier; these screws are for the purpose of communicating small motions to the axis in order to level it or bring the circle into the plane of the meridians.

The Telescope is attached to the circular ring, at each end by appropriate braces each secured by four strong screws; and is further supported in the middle by an axis (represented by dotted lines fig. 1.) which passes through the axis of the circle and is secured by a screw C affixed to its smaller end; the divisions of the circle (which are drawn upon a slip of gold let into its circumference to every five minutes of a degree) were effected upon an entirely new plan invented by Mr. Dollond, the principles of which have not been made public; I have not yet found time to examine into the accuracy attained on a large scale, but form a few divisions which *I have* inspected. I am able to state their accuracy is such as to place them in a class second to none. The circle is furnished with four spring micrometer microscopes which are firmly screwed to the pier Plate III, these (as they are of the ordinary construction) require no further description than a general one: the power of the microscopes is about 12; the micrometer screw acts upon a frame carrying a pair of wires crossing one another at an angle of 25° for the purpose of

reading off the circle; the value of the divisions of the micrometers were rendered equivalent to seconds by adjusting the distance of the object glass until five revolutions of the screw corresponded with the image of the interval between any two divisions of the circle, the centre of which was for this purpose placed opposite to the Zero point of the microscope; in reading off the circle that division which is nearest to the Zero point of the microscope is the one at which the bisection is always made. The object glass is 49 Inches focal length with a clear aperture of $3\frac{3}{4}$ Inches; the optical powers of this glass though inferior to those of the Transit Instrument, are such as do the highest possible credit to the maker and leave me nothing to desire on this head; the eye piece is made to slide and contains a power of about 140; the eye end is furnished with one horizontal, and five vertical fixed wires which are adjustable to horizontal and vertical positions by an arm acted upon by two opposing screws *a, a* fig. 4, and for Collimation by the screws *b, b*; to this is added a horizontal wire moveable by a screw micrometer C, adapted for measuring the difference of declination of Stars or the diameters of Planets, the circle is relieved from wear by two friction wheels which by appropriate weights are brought to bear upon a grooved circular plate, fig. 1. attached to the circle; by this means I estimate that one half or more of the weight of the circle is supported whereby the motion on its axis is rendered exceedingly pleasant and easy. The circle is furnished with four spring clamps of a very superior construction which are screwed to the stone pier (Plate III,) whereby the observer has always one conveniently situated within his reach; by a particular contrivance (due I believe to Mr. Dollond) the two pieces forming the clamp are brought at once to embrace the circle equally on both sides on the clamping screw being tightened thus leaving the circle free from strain and at the same time secure.

In the measurement of North Polar Distance with the Mural Circle the variation produced by deviations of the Instrument (to the amount of one or two minutes) in Level Azimuth or Collimation is so insensible that it is quite unnecessary to perform the adjustment or allow for its effect when within this limit; having this in view my first care on the erection of the Mural Circle was to adjust it to such accuracy as would render corrections unnecessary. The error of Level was discovered by observing the transit of any Star situated near the Zenith by direct vision over the 1st and 2d wires, these by means of the Equatoreal Interval were reduced to give the time of passing the centre wire; in a similar manner observing the transit of the reflected image of the same Star in a basin of quicksilver over the 4th and 5th wires gave also the time of passing the centre wire; the difference $\times 15 \sin. N.P.D. \div 2 \sin. \text{Altitude} = \text{error of Level.}$

The error for Azimuth was obtained from comparing the transits of known Stars situated at from 55° to 60° of N.P.D. with those at from 125° to 130° of N.P.D. by selecting these limits large coefficients are obtained in Azimuth and but small ones for Collimation, for if E, E^1 , represent the errors of the Clock by any two Stars. P, P^1 , their N.P.D. A, A^1 , their Altitude. $x =$ error in Azimuth. $y =$ error in Collimation, we have

$$E - E^1 = \frac{x \cos. A}{15 \sin. P.} + \frac{x \cos. A^1}{15 \sin. P^1} + \frac{y}{15 \sin. P.} - \frac{y}{15 \sin. P^1}$$

In the above case where the sines of P and P^1 are nearly alike

$$x = E - E^1 \times \frac{15 \sin. P. \sin. P^1}{\cos. A. \sin. P^1 + \cos. A^1 \sin. P.} \text{ nearly.}$$

The Azimuthal error being corrected, the Collimation error was determined by means of the Clock error found from the transits of known Stars situated near the Pole, compared with that given by Stars near the equator, for we have

$$y = E - E^1 \times \frac{15 \sin. P. \sin. P^1}{\sin. P^1 - \sin. P.}$$

In this way the amount of deviation for Level, Azimuth, and Collimation, was at first determined and the proper adjustment made; to discover if the deviation remained constant, or within allowable limits, I have from time to time compared the transits of Stars over the meridian wire, with those observed at the same time with the Transit Instrument; the result of these comparisons shew that the circle when left undisturbed has never erred in any way to the amount of $1'$; this introducing but an insensible error in the determination of N.P.D. has been disregarded.

OF THE METEOROLOGICAL INSTRUMENTS.

Of two Barometers which were sent out to this country by Dollond with the Instrument above described, one only had arrived in safety; this on being compared with two others which had been formerly supplied, exhibited differences of two or three tenths of an Inch; to decide which among the three

was most entitled to credit, I set to work to fill a tube myself; after a little experience I succeeded in refilling a tube three times as follows.

From the mean of 10 observations D — T = 0,1114 Inches.			
Emptied and refilled 10	—	=	,1153 „
Do. do. 10	—	=	,1007 „

The near agreement of these differences among themselves led me to suppose that my Barometer was not far from a good one on the one hand, whereas on the other hand, neither the difference of bore of the tubes nor difference of the specific gravities of the quicksilver accounting for the difference itself, I was assured that one or both of these were in error; in this dilemma I began to observe using Dollond's Barometer, and in the mean time applied through Government to the Surveyor General at Calcutta for the loan of the Meteorological Instruments which had recently been supplied to his office for the sake of comparison, this through the zeal which has always distinguished that Officer in affairs connected with scientific research, was most readily and liberally granted, for on my return to this Presidency from Calcutta in February last, I had the pleasure to find myself equipped with two Barometers, and two Thermometers, of the best possible construction, which at the recommendation of the Surveyor General the Supreme Government had allowed to be transferred to this Establishment.

The Barometers furnished to the Surveyor General's Office consisted of Nos. 3, 5, and 6, by Gilbert, constructed with glass cisterns for the adjustment of the Zero point; a Barometer by Troughton, known by the name of Colonel Blacker's standard, and another by the same maker constructed for the Pendulum experiments, both likewise fitted up with glass cisterns: from the mean of 5 careful readings on 27th January 1832, at the Surveyor General's Office.

	Reading Inches.	Cor. for Cap. Act.	Corrected Reading.	Specific Gravity of Quicksilver.
No. 3 by Gilbert.....	30,166	+ ,027	30,193	13,656
No. 5 —————	,135	,047	,182	13,656
No. 6 —————	,148	,051	,199	13,656
Colonel Blacker's Standard by Troughton....	,149	,039	,188	not given.
Barometer employed in Pendulum experiments } by Troughton.	,169	,041	,210	not given.
Mean.....	—	—	30,194	

from the near agreement of these five Barometers *inter se*, little doubt can be entertained of the accuracy of one and all; the two which I was allowed to select were Nos. 3, and 6, by Gilbert; on my arrival at Madras my first care was to compare these Instruments with those which had heretofore been used;

that by Dollond had unfortunately been broken during my absence by a gale of wind on the 24th November, the comparison with that made by myself is as follows.

		Gilbert.		
		No. 3	No. 6	T
		Inches.	Inches.	Inches.
1832, February 25	Mean of 10 readings.....	30,028	30,013	29,921
March 1	Mean of 10 readings.....	30,086	30,076	29,984
Mean.....		<hr/> 30,057	<hr/> 30,044	<hr/> 29,952
Capillary Action.....		+ ,027	,051	,058
Corrected Reading.....		<hr/> 30,084	<hr/> 30,095	<hr/> 30,010
D — T.....				+ ,1007
D.....				<hr/> 30,1107

In the first place the small difference between Nos. 3 and 6, so nearly corresponding with that found on 27th January in Calcutta, affords a presumptive proof that neither of them have sustained any injury on the voyage, and the further agreement of these two with Dollond's Barometer (for we have $D - G = ,027$ Inches and $0,016$ Inches respectively) will I apprehend justify the use of Barometers Nos. 3 and 6 by Gilbert as being sufficiently accurate for every practical purpose. At page 58 it appears that No. 3 differed only ,001 from the mean of five Barometers, I therefore propose in future to consider this the standard, hence it appears that when Dollond's Barometer was in use, its indications as set down in the Circle book were *too great ,027 but having neglected to employ the correction for capillary action*, the registered indications are *too small* ($,058 - ,027 = ,031$); this holds good up to the 24th November, from this time to the end of the year a Barometer by Cary was employed; which on being compared with No. 3, gave

	Gilbert No. 3.	Cary.
1832, February 25..	29,990	29,864
27..	30,016	29,875
29..	30,077	29,959
March 1..	30,111	29,986
2..	30,096	29,979
Mean.....	30,058	29,933
Capillary Action.... +	,027	
	<hr/> 30,085	

hence it appears that the indications of the Barometer as registered in the

N.P.D.			N.P.D.		
23° add to Greenwich Catalogue	0,38	63° add to Greenwich Catalogue	0,92
28	do.0,44	68	do.0,99
33	do.0,51	73	do.1,06
38	do.0,58	78	do.1,16
43	do.0,65	83	do.1,29
48	do.0,72	88	do.1,42
53	do.0,78	90	do.1,50
58	do.0,86			

By way of example we will now compute the Index Error for one day: from the observation Book we have

	Baro- meter.	Thermometer.		Instrumental Reading of				Mean.	NAMES.
		In.	Out.	A	B	C	D		
1831 January 21	Inches. 30,102	75,3	75,2	67 22 30,0	30,2	38,5	19,7	29,6	a Arietis
				82 0 3,0	2,7	16,5	50,5	3,2	δ S. L.
				159 26 43,2	40,0	56,5	26,1	41,4	δ Hydri
				231 18 12,0	12,8	57,1	27,6	12,4	ε Ceti
				102 37 25,1	19,6	33,5	6,4	21,2	ζ Ceti
				80 38 18,5	17,7	31,0	3,2	17,6	μ Ceti
				158 16 19,8	16,8	26,8	59,8	15,8	* Hydri
				234 19 18,5	19,7	30,8	0,6	17,4	R. { 7 Eridani
				99 36 22,8	21,3	38,3	6,7	22,3	D. { 7 Eridani
	30,129	74,3	72,7	86 36 37,8	36,5	48,3	24,8	36,8	α Ceti
				69 37 25,3	23,4	33,8	15,0	24,4	ζ Arietis
				40 47 28,8	29,2	39,6	14,0	27,9	ω Persei
				80 53 46,0	45,2	58,8	32,0	45,5	ε Tauri
	30,124	74,3		233 51 37,5	35,5	52,2	22,8	37,0	R. { ε Eridani
				100 4 0,9	56,8	12,8	43,9	58,6	D. { ε Eridani
				58 17 32,8	32,7	42,0	19,0	31,6	* Persei
				231 16 41,0	39,1	57,9	23,8	40,4	R. { π Eridani
				102 39 0,9	57,0	9,9	40,6	57,1	D. { π Eridani
				283 23 56,5	57,9	12,3	39,0	56,4	R. { ε Persei
	30,126	74,0	71,4	50 31 38,0	39,6	54,2	24,7	39,1	D. { ε Persei
				151 51 43,7	44,0	3,4	25,3	44,1	δ Reticuli
				81 34 52,9	50,1	5,6	37,2	51,4	μ Tauri
	30,138	73,2	70,6	124 14 18,8	10,4	26,6	59,7	13,9	χ Eridani
				75 42 44,5	40,5	55,5	26,0	41,6	π Tauri
				73 52 25,5	20,4	38,0	6,7	22,6	Aldeburan
				223 53 51,8	48,3	5,0	30,1	48,8	R. { 54 Eridani
				110 1 49,7	45,2	2,0	34,3	47,8	D. { 54 Eridani
				24 0 33,7	35,2	44,9	19,9	33,4	* Camelopardi
				76 4 29,0	24,0	38,5	9,1	25,1	ο Orionis
	30,144	73,0	68,9	281 29 41,6	48,3	55,2	25,1	42,6	R. { ω Aurigæ
				52 25 52,6	56,7	5,2	35,9	52,6	D. { ω Aurigæ
				68 41 40,0	40,4	56,8	29,2	41,6	ι Tauri
				74 39 45,8	41,7	59,5	29,9	44,2	γ* Orionis
	30,144	72,2	68,7	125 5 16,8	10,9	25,9	58,2	12,9	ο Columbæ
				97 26 3,0	58,7	16,6	44,8	0,8	U Orionis
	30,150	72,0	68,5	124 11 26,2	28,1	36,2	8,0	24,6	α Columbæ
				30 12 48,8	47,0	58,1	34,7	47,1	31 Camelopardi
				82 40 56,1	53,7	10,4	40,6	55,2	α Orionis
	30,186	72,1	68,2	66 46 23,2	21,8	37,2	9,2	22,8	H Geminor

	Baro- meter.	Thermometer.		Instrumental Reading of			D	Mean.	NAMES.
		In.	Out.	A	B	C			
1831 January 21	Inches.	°	°	°	"	"	"	"	
				20 41 32,6	32,4	47,2	22,5	33,7	* Camelopardi
				96 15 40,0	37,5	59,0	25,5	40,5	a Monocerotis
				107 54 19,8	15,6	35,8	58,6	17,4	β Canis Major
				142 36 35,2	32,0	47,0	13,9	32,0	a Argo Navis
	30,132	72,6	68,3	22 18 45,0	47,5	5,4	22,2	45,0	42 Camelopardi
				31 25 5,0	4,8	16,5	48,6	3,7	c Lynxis
				111 51 4,5	0,2	15,8	44,6	1,3	θ Canis Major
				69 13 37,2	36,0	50,8	21,1	36,3	ζ Geminor
	30,130	71,8	67,0	116 9 15,5	12,2	29,5	56,2	13,3	δ Canis Major
				73 11 54,2	51,0	11,3	33,5	52,5	λ Geminor
				61 54 50,5	48,6	7,2	40,4	51,7	60 Geminor
	30,128	70,9	66,9	57 47 25,3	22,0	40,2	9,5	24,2	a Geminor
				84 22 57,5	54,2	8,2	45,6	56,4	a Canis Min
	30,126	72,0	66,3	118 34 38,1	34,2	49,3	17,3	34,7	3 Argus

Selecting those Stars which are situated within the prescribed limits and whose places are given in the Greenwich Catalogue we have

NAMES.	N.P.D.	Refraction.	4 Equation.	N.P.D. Jan. 1, 1831 + I.			Greenwich Catalogue reduced by Atkinson's Refraction.	Index Error No. 1.	Madras Catalogue Jan. 1, 1831 Atkinson's Refraction.			Index Error No. 2.
				°	'	"			°	'	"	
1831, Jan. 21.												
a Arietis.....	67 22 29,6	—	9,33	1,76	67 22 18,51	67 20 26,17	* 1 52,34	67 20 24,86	* 1 53,65			
μ Ceti.....	80 38 17,6	+	3,52	7,54	80 38 13,58	80 36 14,43	1 59,15	80 36 17,64	1 55,91			
a Ceti.....	86 36 36,8	+	9,39	10,06	86 36 36,13	86 34 42,30	1 53,83	86 34 40,10	1 56,03			
ζ Arietis.....	69 37 24,4	—	7,13	4,54	69 37 12,73	69 35 15,60	1 57,13	69 35 16,46	1 56,27			
a Persei.....	40 47 27,9	—	40,55	4,16	40 46 51,51	40 44 52,31	1 59,20	40 44 54,51	1 57,00			
ε Tauri.....	80 53 45,5	+	3,80	8,68	80 53 40,62	80 51 42,75	1 57,87	80 51 44,84	1 55,78			
e Persei.....	50 31 39,1	—	27,62	0,03	50 31 11,45	50 29 11,65	1 59,80	50 29 13,49	1 57,96			
a Tauri.....	73 52 22,6	—	3,00	8,07	73 52 11,53	73 50 15,77	1 55,75	73 50 13,10	1 58,43			
Camelopardi	24 0 33,4	— 1	13,60	3,71	23 59 23,41	23 57 29,19	1 54,22	23 57 25,17	1 58,24			
t Tauri.....	68 41 41,6	—	8,08	7,33	68 41 26,19	68 39 33,38	1 52,81	68 39 29,03	1 57,16			
a Orionis....	82 40 55,2	+	5,58	11,34	82 39 49,44	82 37 54,90	1 54,54	82 37 53,49	1 55,95			
H Geminorum.	66 46 22,8	—	10,03	8,40	66 46 4,37	66 44 9,27	1 55,10	66 44 8,14	1 56,23			
ζ Geminorum.	69 13 36,3	—	7,58	9,90	69 13 18,82	69 11 22,69	1 56,13	69 11 22,36	1 56,46			
λ Geminorum.	73 11 52,5	—	3,65	10,45	73 11 38,40	73 9 48,66	1 54,74	73 9 38,80	1 59,60			
60 Geminorum	61 54 51,7	—	15,03	9,58	61 54 27,09	61 52 26,41	2 0,68	61 52 28,47	1 58,62			
a Geminorum.	57 47 24,2	—	19,45	9,41	57 46 55,34	57 44 57,06	1 58,28	57 44 57,08	1 58,26			
a Canis Minor	84 22 56,4	+	7,27	10,88	84 22 52,79	84 20 54,24	1 58,55	84 20 55,45	1 57,74			
							Mean of 16 =	1 56,76	—	1 57,23		

The large differences between the results of column 7 (entitled "Index error No. 1,") can only be accounted for, by supposing errors in the observations, in the divisions of the Instrument, or in the Greenwich determination of the N.P.D. the magnitude of the error of observation can best be determined by consulting

* This is omitted in taking the mean.

column 9, "Index error No. 2," whence it appears that the mean error of each observation is 1',03 and the largest deviation from the mean = 2',37: these errors are not larger than what may reasonably be expected, since errors to this amount occur with the Greenwich observations; hence the disagreement in question can only be explained by supposing errors of division in the Madras Mural Circle, or discrepancies in the N.P.D. assumed from the Greenwich Catalogue; to which of these however, the disagreement is attributable, I am at present unable to decide.

In the reduction of the observations, I have of necessity in the first instance been obliged to use Index error No. 1, but the difference which would arise from the use of No. 2, is so small that I have not thought it necessary to recompute them.

We will now compute the Index error from the observations by Reflection; assuming the latitude as determined by Mr. Goldingham 13° 4' 9" (which agrees very nearly with my own determination); putting D for the direct vision observation, R the observation by Reflection; I, the Index error; A the Altitude reckoned from the South horizon, we have

$$D = P + I \text{ and } R = P + 2A + I$$

whence $\frac{D + R}{2} = P + A + I = P + 180^\circ - L - P + I$

$$\therefore I = \frac{D + R}{2} - 176^\circ 55' 51''$$

applying this in the case of the above observations we have

		$\frac{D + R}{2}$		I
	• / "	• / "		/ "
γ Eridani.....	{ R 234 19 17,4 } { D 99 36 22,3 }	166 57 49,85	—	1 58,85
ε Eridani.....	{ R 233 51 37,0 } { D 100 3 58,6 }	47,80	—	1 56,80
π Eridani.....	{ R 231 16 40,4 } { D 102 38 57,1 }	48,75	—	1 57,75
ε Persei.....	{ R 283 23 56,4 } { D 50 31 39,1 }	47,75	—	1 56,75
54 Eridani.....	{ R 223 53 48,8 } { D 110 1 47,8 }	48,30	—	1 57,30
ω Aurigæ.....	{ R 281 29 42,6 } { D 52 25 52,6 }	47,60	—	1 56,60

Mean of 6 = — 1 57,34

The near agreement of the result from reflection observations, with No. 1 and 2, on this and other occasions, speaks in favor of the Madras Mural Circle, but from the few observations which have been made in this way, this testimony is necessarily but a slight one I have not been able to come to any certain conclusion on the subject.

INDEX ERROR OF THE MADRAS MURAL CIRCLE FOR THE YEAR 1831.

Date.	No. of Observations.	Index Error No. 1.	Index Error No. 2.	REMARKS.	Date.	No. of Observations.	Index Error No. 1.	Index Error No. 2.	REMARKS.
1831		<i>m. s.</i>	<i>m. s.</i>		1831		<i>m. s.</i>	<i>m. s.</i>	
January	1 15	1 53,74	1 52,80		March	17 4	2 25,73	1 24,67	Found the horizontal wire broken, the micrometer wire was used on this day.
	2 20	1 52,74	1 53,19			18 10	2 2,92	2 1,96	
	3 15	1 52,25	1 52,51			18 6	2 21,71	2 21,55	Put in a new wire.
	4 8	1 52,14	1 52,92			19, 20 25	2 25,37	1 25,05	The horizontal wire was not so uniform as I could wish, I replaced it with a better one.
	5 21	1 52,03	1 52,36			21, 22 19	1 15,50	1 15,36	
	6 6	1 52,90	1 52,87			23, 24 26	1 17,36	1 17,04	
	7 7	1 53,31	1 53,17			25, 26 24	1 17,13	1 16,87	
	8 15	1 53,99	1 54,52			27, 28 25	1 17,28	1 16,57	
	9, 10 12	1 55,31	1 55,04			29, 30 23	1 16,93	1 16,57	
	11, 12 12	1 54,85	1 54,97		31 Apr. 1 18	1 17,44	1 16,76		
	13 9	1 54,80	1 54,59		2, 3 19	1 16,70	1 16,60		
	14 10	1 54,92	1 54,84		4, 5 20	1 17,08	1 16,78		
	15 13	1 55,08	1 54,67		6, 7 14	1 16,35	1 16,67		
	16, 17 19	1 55,12	1 56,12		9, 10 15	1 18,10	1 17,41		
	18 17	1 54,69	1 55,08		11, 12 19	1 11,10	1 10,82		
	19 18	1 56,16	1 56,46		13, 14 19	1 10,28	1 10,21		
	20 9	1 56,38	1 56,04		15, 18 15	1 10,03	1 10,52		
	21 16	1 56,76	1 51,23	This change is quite unaccountable.	18, 22 16	1 5,91	1 5,00		
	22 15	1 56,49	1 56,97		23, 24 9	1 5,66	1 4,38		
	23, 24 11	1 45,64	1 45,70		25, 26 14	1 4,13	1 3,80		
	25 14	1 46,23	1 45,81		27, 30 19	1 1,27	1 0,89		
	26 14	1 45,68	1 45,54		May 1, 3 13	1 0,22	0 59,97		
	27 9	1 44,49	1 45,53		4, 5 12	1 0,24	1 0,01		
	28 14	1 44,90	1 45,04		6, 7 11	0 55,22	0 55,04		
	29 29	1 45,48	1 45,60		8, 9 19	0 57,44	0 57,37		
	30 9	1 46,75	1 46,44		10, 12 16	0 57,10	0 56,52		
	31 13	1 45,94	1 45,76		13, 15 13	0 57,57	0 56,95		
February	1 10	1 45,85	1 45,77		16, 17 9	0 46,29	0 46,55	Alteration unaccountable.	
	2 10	1 45,76	1 45,47		18, 21 14	0 44,89	0 43,82		
	3, 4 13	1 46,73	1 47,29		22, 24 12	0 40,42	0 41,51		
	5, 6 18	1 48,32	1 47,95		25, 26 10	0 39,01	0 39,36	The circle on this day moved very stiffly which accounts for this sudden change; it originated in the clamp situated between Band D.	
	7, 8 16	1 47,51	1 46,58		27, 28 8	1 10,62	1 10,41		
	9, 10 24	1 46,86	1 46,80		29, 30 10	1 12,05	1 11,64		
	11, 12 11	1 47,25	1 46,96		31 13	1 11,37	1 11,93		
	13, 14 16	1 46,27	1 47,14		June 1, 3 21	1 10,25	1 11,25		
	15, 16 15	1 46,59	1 46,50		4, 6 35	1 10,72	1 10,91		
	17, 18 11	1 47,07	1 46,57		7, 13 22	1 12,03	1 11,98		
	19, 20 12	1 47,86	1 47,16		15, 20 17	1 15,99	1 16,36		
	21, 22 17	1 47,31	1 46,95		21, 27 12	1 17,41	1 17,60		
	23, 24 28	1 47,06	1 46,89		28 6	1 16,79	1 17,38		
	25, 26 15	1 48,66	1 47,66		29, 30 10	1 10,01	1 9,71	The horizontal wire appeared bent, I removed it and introduced silk lines.	
	27, 28 19	1 48,23	1 48,03		July 1, 3 7	1 8,26	1 8,83		
Mar.	1, 2 23	1 47,64	1 47,49		5, 7 11	3 11,14	3 10,04		
	3, 4 34	1 46,85	1 46,83		9, 12 31	3 10,85	3 10,58		
	5, 6 48	1 47,26	1 47,67		13, 15 10	3 10,01	3 10,66		
	7, 8 31	1 46,96	1 46,98		16, 17 15	3 4,87	3 4,70		
	9, 10 27	1 47,35	1 47,23		22 2	3 3,70	3 3,27		
	11, 12 31	1 48,25	1 47,76						
	13, 14 31	1 47,41	1 47,14						
	15, 16 29	1 46,75	1 46,42						
	17 5	1 45,73	1 45,72						

Date.	No. of Observations.	Index Error No. 1.	Index Error No. 2.	REMARKS.	Date.	No. of Observations.	Index Error No. 1.	Index Error No. 2.	REMARKS.
1831		<i>m. s.</i>	<i>m. s.</i>		1831		<i>m. s.</i>	<i>m. s.</i>	
July 24, 26	15	3 14,30	3 15,20		Oct. 4, 5	13	3 0,30	3 0,20	
28, 31	19	3 15,12	3 15,61		6, 12	21	3 1,50	3 1,88	
Aug. 1, 9	20	3 15,13	3 15,93		14, 21	21	3 0,52	3 0,68	
10, 11	27	3 15,67	3 16,38		22, 23	21	3 2,04	3 1,97	
12, 13	8	3 16,66	3 16,78		24, 29	23	2 2,26	3 2,06	
20, 22	6	3 15,68	3 16,43		30, 31	26	3 1,62	3 2,18	
23, 24	23	3 15,75	3 15,51		Nov. 1	15	3 1,78	3 1,51	
25, 27	22	3 15,96	3 15,55		2, 3	25	3 2,59	3 2,15	
28, 29	32	3 16,04	3 15,36		4, 5	13	3 2,37	3 2,64	
30, 31	17	3 14,71	3 14,90		6, 8	17	3 3,49	3 2,97	
Sept. 1, 2	21	3 13,10	3 13,24		9, 19	20	3 2,28	3 2,38	
3, 4	17	3 12,62	3 13,27		21, 30	18	3 3,18	3 3,66	
7, 13	13	3 10,19	3 10,61		Dec. 1, 6	8	3 4,86	3 4,24	
14	11	3 10,26	3 10,75		7, 9	13	3 5,11	3 6,05	
15	13	3 10,48	3 10,57		10, 15	10	3 7,89	3 8,46	
16, 21	22	3 10,27	3 10,24		16, 23	18	3 7,98	3 8,13	
22, 27	17	3 10,72	3 11,05		24, 25	9	3 9,18	3 9,29	
28, 30	23	3 3,16	3 2,59		28, 30	9	3 0,85	3 0,74	Unaccountable.*
Oct. 1, 3	28	3 3,87	3 2,69						

The column of Index error No. 1, is as I have before stated that which has been employed in reducing the places of the Stars, but in the case of the Sun, Planets, and Moon, when the *limb* is observed, (the *edge* of the wire being made a tangent to the limb) it is the Index error of the North or South edge (according to which limb is observed) which should be employed; the simplest way to apply the correction is to increase the semi-diameter of the object observed by half the thickness of the wire, whereby the consideration of which edge has been observed is evaded: I estimate the diameter of the wire first employed at 2",42; that put in on the 20th March at 2",5, and the silk line put in on the 5th July at 2",1, I have employed throughout 1",2 for the semi-diameter.

* With reference to this remark I am under the painful obligation to observe that unless the cause of alteration happened when using the Instrument *myself* or superintending its use, I stand no chance of being informed of it.

RESULTS DERIVED FROM THE OBSERVATIONS MADE WITH THE TRANSIT INSTRUMENT AND MURAL CIRCLE IN THE YEAR 1831.

In the first place we will examine the observations of the Sun

The observation of the Sun's first and second limb over the five wires of the Transit Instrument affords a ready means of determining the diameter, for we have Sun's Apparent Semi-d. = $\left(\frac{\odot 2 L - \odot 1 L}{30}\right) \left(\frac{a' - a}{48h.}\right) \sin. N.P.D.$
 a' and a are the Right Ascensions of the Sun at the noon preceding, and following the day of observation, as found in the Nautical Almanac; from this the mean semi-diameter or the semi-diameter seen from the Earth when at its mean distance has been found, by dividing the above by the distance $\odot - \oplus$, whose logarithm is given in the Nautical Almanac; thus we have

Interval in time occupied by the Sun's diameter in passing the Meridian applied to the determination of its Mean semi-diameter.

1831	☉ 2 L — ☉ 1 L.	Reduced Apparent Semi.	Mean Semi-dia- meter ☉ Dist. 1.	1831	☉ 2 L — ☉ 1 L.	Reduced Apparent Semi.	Mean Semi-dia- meter ☉ Dist. 1.		
	<i>m. s.</i>	<i>m. s.</i>	<i>m. s.</i>		<i>m. s.</i>	<i>m. s.</i>	<i>m. s.</i>		
February	19	2 12,62	16 12,0	16 1,1	March	28	2 8,98	16 4,7	16 3,4
	21	2 12,72	16 15,0	16 4,6		30	2 9,12	16 3,7	16 3,2
	22	2 14,08	16 26,3	16 16,0	April	1	2 9,08	16 2,9	16 2,8
	23	2 12,28	16 14,1	16 4,2		2	2 9,12	16 7,8	16 8,1
	24	2 11,82	16 11,8	16 2,1		3	2 9,84	16 8,1	16 8,6
	25	2 11,22	16 8,4	15 59,0		5	2 9,94	16 7,3	16 8,3
	27	2 11,04	16 9,0	16 0,1		6	2 10,00	16 2,0	16 3,3
March	3	2 10,86	16 11,4	16 3,6		8	2 9,38	16 3,8	16 5,9
	4	2 10,92	16 12,6	16 5,1		10	2 9,84	15 58,9	16 1,1
	5	2 10,70	16 11,6	16 4,4	13	2 9,38	16 0,9	16 4,2	
	8	2 10,40	16 11,5	16 5,1	15	2 10,02	16 0,9	16 4,9	
	10	2 9,80	16 8,0	16 2,1	16	2 10,28	15 59,9	16 3,9	
	11	2 9,62	16 7,1	16 1,5	17	2 9,36	15 52,0	15 56,3	
	12	2 9,50	16 6,7	16 1,3	18	2 7,84	15 39,8	15 44,3	
	13	2 9,12	16 4,3	15 59,1	19	2 10,50	15 58,3	16 3,1	
	14	2 9,50	16 7,5	16 2,6	20	2 10,66	15 58,3	16 3,4	
	15	2 9,00	16 4,0	15 59,4	23	2 10,38	15 52,7	15 58,5	
	17	2 9,12	16 5,4	16 1,4	25	2 8,88	15 39,2	15 45,5	
	19	2 8,96	16 4,5	16 0,9	26	2 8,36	15 34,2	15 40,7	
	20	2 9,00	16 4,9	16 1,6	27	2 10,24	15 46,7	15 53,4	
	22	2 8,98	16 4,7	16 1,9	28	2 11,78	15 53,6	15 53,2	
24	2 9,58	16 9,1	16 6,7	29	2 11,36	15 52,2	15 59,5		
25	2 9,10	16 5,3	16 3,2	30	2 11,56	15 52,3	15 59,9		
26	2 9,18	16 5,7	16 3,9	May	1	2 9,72	15 37,5	15 45,1	
27	2 8,76	16 2,3	16 0,7		2	2 11,74	15 50,7	15 58,7	

RESULTS FROM OBSERVATIONS, 1831.

1831	☉ 2 L.— ☉ 1 L.	Reduced Apparent Semi.	Mean Semi-dia- meter ⊖ Dist. 1.	1831	☉ 2 L.— ☉ 1 L.	Reduced Apparent Semi.	Mean Semi-dia- meter ⊖ Dist. 1.			
May	<i>m. s.</i>	<i>m. s.</i>	<i>m. s.</i>	July	<i>m. s.</i>	<i>m. s.</i>	<i>m. s.</i>			
	5	2 12,78	15 54,1		16 2,9	11	2 16,58	15 45,2	16 1,0	
	6	2 11,52	15 43,6		15 52,6	14	2 16,50	15 47,4	16 3,3	
	7	2 13,32	15 55,2		16 4,5	17	2 15,62	15 44,4	15 59,8	
	11	2 14,02	15 54,6		16 4,8	21	2 14,56	15 41,4	15 56,5	
	12	2 12,24	15 40,5		15 50,7	25	2 14,28	15 44,3	15 58,9	
	13	2 12,16	15 38,5		15 48,8	27	2 14,02	15 45,0	15 59,5	
	14	2 13,64	15 47,8		15 58,5	29	2 13,88	15 44,7	16 0,9	
	15	2 15,00	15 56,1		16 7,0	30	2 13,74	15 47,0	16 1,1	
	17	2 15,26	15 55,3		16 6,6	August 4	2 12,66	15 41,6	15 55,0	
	18	2 13,70	15 42,9		15 54,3	6	2 12,34	15 46,6	15 59,9	
	19	2 14,72	15 48,8		16 0,5	7	2 12,32	15 47,9	16 1,0	
	20	2 14,88	15 48,6		16 0,5	8	2 12,20	15 48,4	16 1,3	
	21	2 14,94	15 47,8		15 59,8	11	2 11,14	15 44,8	15 57,2	
	22	2 15,66	15 51,4		16 3,6	13	2 11,06	15 46,9	15 59,0	
	23	2 15,38	15 58,3		16 0,8	14	2 11,22	15 49,4	16 1,4	
	24	2 15,94	15 51,2		16 3,6	18	2 10,62	15 50,2	16 1,4	
	25	2 15,46	15 46,6		15 59,3	20	2 10,34	15 50,6	16 1,5	
	26	2 15,40	15 45,1		15 57,8	22	2 10,10	15 51,3	16 1,8	
	27	2 16,22	15 49,7		16 2,7	23	2 9,82	15 52,4	16 0,6	
	29	2 15,64	15 43,5		15 56,7	24	2 9,72	15 52,9	16 0,8	
	30	2 16,62	15 49,4		16 2,7	25	2 9,64	15 52,4	16 1,2	
	31	2 16,30	15 46,1		15 59,6	27	2 9,60	15 53,3	16 2,7	
	June	1	2 16,34		15 45,3	15 59,0	31	2 8,86	15 55,8	16 0,3
		2	2 16,66		15 46,6	16 0,4	September 1	2 8,84	15 56,8	16 1,1
		3	2 16,50		15 44,6	15 58,5	5	2 8,28	15 55,0	15 59,3
		4	2 16,60		15 44,5	15 58,5	7	2 8,18	15 52,7	15 59,6
		5	2 17,14		15 47,4	16 1,7	8	2 8,18	15 53,4	16 0,0
		6	2 17,98		15 52,4	16 6,9	11	2 8,22	15 55,5	16 1,4
		7	2 17,26		15 46,7	16 1,2	13	2 8,18	15 56,2	16 1,6
		8	2 17,10		15 45,0	15 59,5	14	2 7,98	15 55,1	16 0,3
9		2 17,54	15 47,3	15 58,5	15	2 7,58	15 52,5	15 57,4		
10		2 17,34	15 45,4	16 0,1	16	2 8,10	15 56,8	16 1,3		
11		2 17,66	15 47,0	16 1,9	17	2 7,74	15 54,4	15 58,6		
12		2 17,62	15 46,3	16 1,2	19	2 8,04	15 57,1	16 0,8		
13		2 17,40	15 44,3	15 59,3	21	2 7,96	15 56,9	16 0,1		
14		2 17,70	15 45,9	16 1,0	22	2 8,16	15 58,4	16 1,4		
16		2 17,94	15 46,9	16 2,1	24	2 7,80	15 55,8	15 58,1		
19		2 17,54	15 46,6	15 58,8	25	2 8,14	15 58,3	16 0,4		
21		2 17,16	15 40,7	15 56,0	26	2 8,02	15 57,3	15 59,1		
22		2 17,78	15 44,9	16 3,5	27	2 8,50	16 0,8	16 2,3		
23		2 18,12	15 47,3	16 2,9	28	2 8,04	15 57,1	15 58,4		
24		2 17,76	15 44,9	16 0,6	29	2 8,66	16 1,6	16 2,6		
25		2 17,94	15 46,2	16 1,9	October 4	2 8,98	16 2,2	16 1,9		
26		2 18,04	15 47,1	16 2,9	5	2 8,92	16 1,3	16 0,7		
27		2 17,64	15 44,6	16 0,4	6	2 9,36	16 4,0	16 3,1		
28		2 18,12	15 48,2	16 4,0	7	2 9,28	16 3,1	16 2,0		
29		2 17,98	15 47,5	16 3,4	8	2 9,60	16 4,6	16 3,2		
30		2 17,58	15 45,1	16 1,0	13	2 10,02	16 4,0	16 1,2		
July		1	2 17,82	15 47,2	16 3,1	14	2 10,08	16 3,6	16 0,5	
		2	2 17,96	15 48,6	16 4,6	15	2 10,70	16 7,4	16 4,0	
		3	2 17,20	15 43,8	15 59,7	17	2 10,68	16 5,3	16 1,4	
		4	2 17,64	15 47,5	16 3,4	22	2 11,62	16 6,9	16 1,6	
	5	2 17,18	15 44,9	16 0,8	23	2 11,40	16 4,1	15 58,6		
	6	2 17,10	15 44,9	16 0,8	31	2 13,56	16 9,7	16 2,0		
	7	2 16,96	15 44,6	16 0,5	November 1	2 13,38	16 6,3	15 59,0		
	8	2 16,94	15 45,3	16 1,1	2	2 14,26	16 11,9	16 3,8		

1831				1831			
	☉ 2 L.— ☉ 1 L.	Reduced Apparent Semi.	Mean Semi-dia- meter ⊖ Dist. 1.		☉ 2 L.— ☉ 1 L.	Reduced Apparent Semi.	Mean Semi-dia- meter ⊖ Dist. 1.
	<i>m. s.</i>	<i>m. s.</i>	<i>m. s.</i>		<i>m. s.</i>	<i>m. s.</i>	<i>m. s.</i>
November 3	2 14,02	16 8,7	16 0,3	December 10	2 21,96	16 17,9	16 2,7
8	2 14,38	16 4,0	15 54,6	12	2 22,00	16 16,8	16 1,4
9	2 15,40	16 9,6	16 0,0	13	2 22,10	16 17,0	16 1,5
10	2 15,88	16 11,7	16 1,7	14	2 22,24	16 17,5	16 1,9
12	2 16,28	16 11,6	16 1,1	17	2 22,42	16 17,5	16 1,7
13	2 16,24	16 9,9	15 59,2	20	2 22,42	16 16,9	16 0,1
18	2 17,66	16 13,0	16 0,4	21	2 22,34	16 16,0	16 0,0
27	2 19,54	16 14,2	16 0,8	22	2 22,80	16 19,3	16 3,0
December 1	2 20,48	16 15,8	16 1,7	26	2 22,50	16 17,7	16 1,3
2	2 20,84	16 17,1	16 2,9	29	2 22,12	16 15,9	15 59,5
7	2 21,14	16 14,5	15 59,7				

Mean of 184 gives for the Sun's semi-diameter when viewed at the Earth's Mean Distance..... 16' 0',742
 Rejecting the observations between the 23d April and 2d of May which were made by an inexperienced Assistant, we get..... 16' 0',15

Thus much for a rough approximation, for the value here assigned can only be thus regarded from certain uncontrollable circumstances which it may be as well here to enumerate. In the first place it is a generally received opinion that all telescopes do not give the same value for the diameter of the Sun and Planets; be this as it may, in the absence of proof, probability seems strongly to favor the notion; in the next place an *irradiation* as it is called, will I apprehend be found with every observer of a nature much more serious than the above, for on comparing several hundred observations made at Greenwich with the old 8 feet transit Instrument with 3 Inches aperture I found a difference of 6 or 8 seconds between the value assigned for the Sun's diameter by different observers*; whereas on this Instrument being superceded by the present 10 feet with 5 Inches aperture, no appreciable difference resulted in the value it gave for the diameter of the Sun, when compared with that given by the old Instrument.

Again, I am aware of no observation sufficiently nice having been made whereby we are justified in assuming the Sun to be perfectly Spherical an assumption indeed which would be at variance with the received laws of

* So certainly did a change occur in the value for the Sun's diameter determined in this way, in the days of Dr. Marklyne on his engaging a new Assistant, that is no difficult matter to discover the day when a new Assistant came from this circumstance.

gravity ; as the matter now stands (in consequence of the small inclination of the Solar axis to the axis of the ecliptic) our measure is very nearly the Sun's Equatorial diameter, and being the mean diameter from three or four different observers I propose for the present (notwithstanding the imperfections to which it is liable,) for want of a better, to employ this result in any case where the Sun's diameter is required.

We will now compare the observed Right Ascension and North Polar Distance of the Sun with its place as interpolated from the Nautical Almanac.

	Observed A.R.			A.R. from Nautical Almanac.		Error of Tables.	Observed N.P.D.			N.P.D. from Nautical Almanac.		Error of Tables.	
	h.	m.	s.	m.	s.		°	m.	s.	m.	s.		
1831													
January	28						108	21	56,49	22	2,00	+	5,5
	29						108	6	13,47	6	15,00	+	1,5
	30						107	50	5,55	50	10,00	+	4,5
February	1						107	16	53,95	17	1,00	+	7,0
	2						106	59	51,73	59	58,00	+	6,3
	3						106	42	34,20	42	38,00	+	3,8
	4						106	24	58,40	25	0,00	+	1,6
	5						106	7	1,77	7	5,00	+	3,2
	6						105	48	50,63	48	53,00	+	2,4
	7						105	30	27,28	30	25,00	-	2,3
	8						105	11	38,30	11	41,00	+	2,7
	11						104	13	57,79	13	59,00	+	1,2
	12						103	54	7,91	54	16,00	+	8,1
	13						103	34	13,64	34	19,00	+	5,4
	14						103	14	4,77	14	9,00	+	4,2
	15						102	53	40,05	53	47,00	+	6,9
	16						102	33	9,32	33	12,00	+	2,7
	17						102	12	19,59	12	25,00	+	5,4
	18						101	51	21,19	51	26,00	+	4,8
	19						101	30	11,38	30	16,00	+	4,6
	20	22	11	58,48	11	58,10	101	8	54,14	8	55,00	+	0,9
	21	22	15	48,37	15	47,90	100	47	29,15	47	25,00	-	4,1
	22	22	19	37,10	19	37,10							
	23	22	23	25,62	23	25,70							
	24	22	27	13,78	27	13,70	100	3	50,81	3	55,00	+	4,2
	25	22	31	1,30	31	1,10	99	41	57,63	41	55,00	-	2,6
	26						99	19	40,51	19	47,00	+	6,5
	27	22	38	34,89	38	34,00	98	57	29,42	57	31,00	+	1,6
	28						98	35	1,51	35	7,00	+	5,5
March	2	22	49	49,81	49	49,40	98	12	33,27	12	35,00	+	1,7
	3	22	53	33,04	53	33,40	97	27	2,25	27	10,00	+	7,7
	4	22	57	16,86	57	17,00	97	4	13,18	4	18,00	+	4,8
	5	23	1	1,41	1	0,10	96	41	14,55	41	21,00	+	6,4
	6						96	18	6,27	18	17,00	+	10,7
	7						95	55	2,69	55	8,00	+	5,3
	8	23	12	7,40	12	7,10	95	31	45,07	31	54,00	+	8,9
	9	23	15	49,23	15	48,70	95	8	34,53	8	35,00	+	0,5
	10	23	19	30,46	19	29,80	95	45	12,92	45	13,00	+	0,1
	11	23	23	11,53	23	10,70	94	21	38,54	21	46,00	+	7,5
	12	23	26	51,98	26	51,30	93	58	12,89	58	16,00	+	3,1
							93	34	35,62	34	44,00	+	8,4

		Observed A.R.			A.R. from Nautical Almanac.		Error of Tables.	Observed N.P.D.			N.P.D. from Nautical Almanac.		Error of Tables.	
1831		<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>m.</i>	<i>s.</i>	<i>s.</i>	<i>°</i>	<i>m.</i>	<i>s.</i>	<i>m.</i>	<i>s.</i>	<i>s.</i>	
March	13	23	30	32,11	30	31,50	— 0,61	93	11	1,71	11	8,00	+ 6,3	
	14	23	34	12,31	34	11,50	— 0,81	92	47	23,76	47	31,00	+ 7,2	
	15	23	37	52,15	37	51,20	— 0,95	92	23	45,89	23	51,00	+ 5,1	
	16							92	0	7,81	0	10,00	+ 2,2	
	17	23	45	10,01	45	9,80	— 0,21	91	36	27,07	36	29,00	+ 1,9	
	18													
	19	23	52	28,23	52	27,50	— 0,73	90	48	59,36	49	4,00	+ 4,6	
	20	23	56	6,47	56	6,10	— 0,37	90	25	21,09	25	22,00	+ 0,9	
	21	23	59	44,70	59	44,50	— 0,20							
	22	0	3	23,67	3	22,70	— 0,97	89	37	55,28	38	0,00	+ 4,7	
23	0	7	2,15	7	0,90	— 1,25	89	14	20,76	14	21,00	+ 0,2		
24	0	10	39,89	10	38,90	— 0,99								
25	0	14	17,28	14	16,90	— 0,38	88	27	4,51	27	8,00	+ 3,5		
26	0	17	55,70	17	54,80	— 0,90								
27	0	21	33,42	21	32,60	— 0,82	87	39	58,84	40	3,00	+ 4,2		
28	0	25	11,25	25	10,50	— 0,75	87	16	23,75	16	35,00	+ 6,2		
29							86	53	8,04	53	11,00	+ 3,0		
30	0	32	27,00	32	26,20	— 0,80	86	29	41,22	29	50,00	+ 8,0		
31	0	36	5,06	36	4,20	— 0,86	86	6	31,75	6	32,00	+ 0,3		
April	1	0	39	43,24	39	42,30	— 0,94	85	43	13,33	43	19,00	+ 5,7	
	2	0	43	21,00	43	20,40	— 0,60	85	20	8,81	20	11,00	+ 2,2	
	3							84	57	2,59	57	8,00	+ 5,4	
	5	0	54	16,60	54	15,80	— 0,80	84	11	15,69	11	16,00	+ 0,3	
	6	0	57	55,17	57	54,50	— 0,67	83	48	23,03	48	30,00	+ 7,0	
	7							83	25	44,47	25	50,00	+ 5,5	
	8	1	5	13,61	5	12,30	— 0,31	83	3	8,78	3	16,00	+ 7,2	
	10	1	12	32,80	12	31,90	— 0,90	82	18	22,10	18	30,00	+ 7,9	
	11							81	56	16,78	56	18,00	+ 1,2	
	12							81	34	4,58	34	14,00	+ 9,4	
	13	1	23	33,51	23	33,00	— 0,51	81	12	19,29	12	18,00	— 1,3	
	14							80	50	30,31	50	32,00	+ 1,7	
	15	1	30	55,73	30	55,10	— 0,63	80	28	56,48	28	55,00	— 1,5	
	16	1	34	37,18	34	36,50	— 0,68	80	7	29,46	7	27,00	— 2,5	
	17	1	38	19,17	38	18,30	— 0,87							
	18	1	42	0,58	42	0,50	— 0,08	79	24	58,24	25	1,00	+ 2,8	
	19	1	45	43,41	45	43,20	— 0,21	79	4	0,33	4	4,00	+ 3,7	
	20	1	49	26,35	49	26,20	— 0,15	78	43	11,35	43	17,00	+ 5,7	
	21							78	22	33,45	22	41,00	+ 7,5	
22							78	2	12,26	2	17,00	+ 4,7		
23	1	0	38,07	0	37,60	— 0,47	77	41	56,56	42	5,00	+ 8,4		
24							77	21	59,27	22	5,00	+ 5,7		
25	1	8	7,57	8	7,30	— 0,27	77	2	16,12	2	18,00	+ 1,9		
26							76	42	35,98	42	44,00	+ 8,0		
27							75	23	21,54	23	23,00	+ 1,5		
28	1	19	26,18	19	25,50	— 0,68	76	4	10,13	4	14,00	+ 3,9		
29	1	23	13,55	23	12,60	— 0,95	75	45	18,90	45	19,00	+ 0,1		
30	1	27	1,21	27	0,20	— 1,01	75	26	34,41	26	38,00	+ 3,6		
May	1	1	30	49,14	30	48,20	— 0,94	75	8	8,19	8	11,00	+ 2,8	
	2	1	34	37,37	34	36,90	— 0,47							
	3							74	32	1,03	32	4,00	+ 3,0	
	4	1	42	16,90	42	16,00	— 0,90	74	14	16,48	14	22,00	+ 5,5	
	5	1	46	7,67	46	6,40	— 1,27	73	56	52,43	56	56,00	+ 3,6	
	6	1	49	58,55	49	57,30	— 1,25	73	39	44,54	39	45,00	+ 0,5	
	7	1	53	49,87	53	48,90	— 0,97	73	22	50,74	22	52,00	+ 1,3	
	8							72	49	46,20	49	53,00	+ 6,8	

RESULTS FROM OBSERVATIONS, 1831.

		Observed A.R.			A.R. from Nautical Almanac.		Error of Tables.	Observed N.P.D.			N.P.D. from Nauti- cal Almanac.		Error of Tables.		
1831		<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>m.</i>	<i>s.</i>	<i>s.</i>	<i>°</i>	<i>m.</i>	<i>s.</i>	<i>m.</i>	<i>s.</i>	<i>s.</i>		
May	10							72	33	50,36	33	49,00	—	1,4	
	11	3	9	21,68	9	21,10	—	0,58	72	18	0,60	18	3,00	+	2,4
	12	3	13	15,96	13	15,60	—	0,36	72	2	37,19	2	34,00	—	3,2
	13	3	17	10,91	17	10,60	—	0,31	71	47	20,31	47	23,00	—	2,7
	14	3	21	6,91	21	6,30	—	0,61	71	32	27,67	32	31,00	+	3,3
	15	3	25	3,37	25	2,50	—	0,87	71	17	54,16	17	58,00	+	3,8
	16	3	28	59,39	28	59,20	—	0,19	71	3	41,76	3	43,00	+	1,2
	17	3	32	56,79	32	56,50	—	0,29	70	49	42,30	49	47,00	+	4,7
	18	3	36	54,70	36	54,30	—	0,40	70	36	13,04	36	11,00	—	2,0
	19	3	40	53,35	40	52,80	—	0,55	70	22	51,05	22	54,00	+	2,9
	20	3	44	52,38	44	51,70	—	0,68	70	10	3,26	9	58,00	—	5,3
	21	3	48	51,86	48	51,20	—	0,66	69	57	25,58	57	23,00	—	2,6
	22	3	52	51,67	52	51,30	—	0,37	69	45	7,72	45	8,00	+	0,3
	23	3	56	52,36	56	51,70	—	0,66	69	33	10,88	33	13,00	+	2,1
	24	4	0	53,47	0	52,80	—	0,67	69	21	38,75	21	39,00	+	0,2
	25	4	4	54,99	4	54,30	—	0,69	69	10	24,61	10	26,00	+	1,4
	26	4	8	56,58	8	56,30	—	0,28	68	59	31,15	59	35,00	+	3,8
	27	4	12	59,23	12	58,80	—	0,43	68	48	59,83	49	5,00	+	5,2
	29	4	21	5,88	21	5,30	—	0,58							
	30	4	25	10,17	25	9,30	—	0,87	68	19	41,32	19	49,00	+	7,7
31	4	29	14,68	29	13,80	—	0,88								
June	1	4	33	19,27	33	18,60	—	0,67	68	2	12,49	2	10,00	—	2,5
	2	4	37	24,71	37	24,10	—	0,61	67	53	50,07	53	55,00	+	4,9
	3								67	45	57,42	46	2,00	+	4,6
	4	4	45	36,71	45	36,00	—	0,71	67	38	27,39	38	33,00	+	5,6
	5	4	49	43,01	49	42,50	—	0,51	67	31	31,87	31	28,00	—	3,9
	6	4	53	49,82	53	49,30	—	0,52	67	24	44,60	24	45,00	—	0,4
	7	4	57	57,41	57	56,60	—	0,81	67	19	27,93	18	26,00	—	1,9
	8	5	2	5,04	2	4,20	—	0,84	67	12	27,00	12	32,00	+	4,1
	9	5	6	13,05	6	12,10	—	0,95	67	7	2,31	7	1,00	—	1,3
	10	5	10	21,24	10	20,40	—	0,84	67	1	55,96	1	55,00	—	1,0
	11	5	14	29,35	14	28,90	—	0,45	66	57	14,54	57	12,00	—	2,5
	12	5	18	37,90	18	37,50	—	0,40	66	52	57,91	52	53,00	—	4,9
	13	5	22	47,02	22	46,30	—	0,72	66	49	4,12	49	0,00	—	4,1
	14	5	26	56,03	26	55,30	—	0,73							
	16	5	35	14,32	35	13,80	—	0,52	66	39	49,73	39	46,00	—	3,7
	19	5	47	42,23	47	42,10	—	0,13	66	34	9,04	34	15,00	+	6,0
	21	5	56	1,77	56	1,10	—	0,67	66	32	38,75	32	38,00	—	0,7
	22	6	0	11,02	0	10,60	—	0,42	66	32	30,67	32	26,00	—	4,7
	23	6	4	20,67	4	20,10	—	0,47	66	32	37,94	32	40,00	+	2,1
	24								66	33	20,66	33	18,00	—	2,7
	25	6	12	39,22	12	38,80	—	0,42	66	34	24,83	34	21,00	—	3,8
	26	6	16	48,66	16	48,00	—	0,66	66	35	50,95	35	49,00	—	1,9
	27	6	20	57,79	20	57,00	—	0,79	66	37	48,33	37	42,00	—	6,3
	28	6	25	6,83	25	6,00	—	0,83	66	39	58,57	39	59,00	—	0,4
29	6	29	15,57	29	14,90	—	0,67	66	42	43,86	42	40,00	—	3,9	
30	6	33	24,45	33	23,50	—	0,95	66	45	44,26	45	47,00	+	1,2	
July	1	6	37	32,97	37	32,00	—	0,97	66	49	20,65	49	16,00	+	4,6
	2	6	41	40,99	41	40,20	—	0,79	66	53	16,37	53	11,00	—	5,4
	3	6	45	48,40	45	48,20	—	0,20	66	57	34,25	57	30,00	—	4,2
	4	6	49	56,93	49	55,90	—	1,03							
	5	6	54	4,45	54	3,40	—	1,05							
	6	6	58	11,47	58	10,60	—	0,87	67	12	55,05	12	52,00	—	3,0
	7	7	2	18,45	2	17,60	—	0,85	67	18	48,93	18	48,00	—	0,9
	8	7	6	24,67	6	24,20	—	0,47	67	25	8,24	25	7,00	—	1,2
	11	7	18	41,63	18	41,40	—	0,23	67	46	22,42	46	23,00	+	0,6

RESULTS FROM OBSERVATIONS, 1831.

		Observed A.R.			A.R. from Nautical Almanac.		Error of Tables.	Observed N.P.D.			N.P.D. from Nautical Almanac.		Error of Tables.
1831		<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>m.</i>	<i>s.</i>	<i>s.</i>	<i>°</i>	<i>m.</i>	<i>s.</i>	<i>m.</i>	<i>s.</i>	<i>s.</i>
July	14	7	30	55,52	30	54,70	— 0,82	68	11	6,01	11	7,00	+ 1,0
	17	7	43	4,45	43	3,60	— 0,85	68	39	10,43	39	11,00	+ 0,6
	21	7	59	8,92	59	7,90	— 1,02	69	21	45,35	21	41,00	— 4,4
	25	8	15	3,90	15	3,20	— 0,70	70	4	48,04	9	42,00	— 6,0
	27	8	22	58,16	22	57,30	— 0,86	70	35	49,23	35	43,00	— 6,2
	29	8	31	50,44	30	49,20	— 1,24	71	3	7,19	3	1,00	— 6,2
	30	8	35	45,24	34	44,30	— 0,94	71	17	12,01	17	8,00	— 4,0
August	31	8	39	39,65	38	38,70	— 0,95	71	31	34,79	31	34,00	— 0,8
	3	8	50	19,49	50	18,50	— 0,99	72	16	44,58	16	40,00	— 4,6
	4	8	54	11,51	54	10,60	— 0,91	72	32	22,80	32	17,00	— 5,8
	5	8	58	3,07	58	2,10	— 0,97	72	48	16,31	48	12,00	— 4,3
	6	9	1	54,25	1	53,10	— 1,15	73	4	25,81	4	24,00	— 1,8
	7	9	5	44,19	5	43,40	— 0,79	73	20	49,07	20	52,00	+ 2,9
	8	9	9	34,29	9	33,20	— 1,09	73	37	41,20	37	36,00	— 5,2
	11	9	20	59,70	20	59,00	— 0,70	74	29	23,73	29	21,00	— 2,7
	12	9	24	46,96	24	46,40	— 0,56	74	47	11,87	47	7,00	— 4,9
	13	9	28	34,14	28	33,20	— 0,94	75	5	11,77	5	7,00	— 4,8
	14	9	32	19,98	32	19,50	— 0,48	75	23		23	21,00	
	18	9	47	20,03	47	19,00	— 1,03	76	38	31,71	38	34,00	+ 2,3
	20	9	54	46,38	54	45,60	— 0,78	77	17	27,79	17	27,00	— 0,8
	22	10	2	10,65	2	10,10	— 0,55	77	57	7,73	57	10,00	+ 2,3
23	10	5	52,00	5	51,70	— 0,30	78	17	15,76	17	18,00	+ 2,2	
24	10	9	33,32	9	32,80	— 0,52	78	37	39,65	37	37,00	— 2,6	
25	10	13	13,87	13	13,60	— 0,27	78	58	5,12	58	7,00	+ 1,9	
27	10	20	34,08	20	33,80	— 0,28	79	39	48,67	39	39,00	— 9,7	
31	10	35	10,85	35	9,90	— 0,95	81	4	43,70	4	38,00	— 5,7	
Sept.	1	10	38	48,72	38	48,00	— 0,72	81	26	19,46	26	15,00	— 4,5
	5	10	53	18,77	53	18,10	— 0,67	82	54	5,57	54	7,00	+ 1,4
	7	11	0	32,21	0	31,80	— 0,41	83	38	47,30	38	45,00	— 2,3
	8	11	4	9,12	4	8,30	— 0,82	84	1	13,07	1	14,00	+ 0,9
	10	11	11	21,02	11	20,60	— 0,42	84	46	38,32	46	31,00	— 7,3
	11	11	14	56,98	14	56,40	— 0,58	85	9	24,86	9	17,00	— 7,9
	12	11	18	33,17	18	32,40	— 0,77	85	32	11,87	32	8,00	— 3,9
	13	11	22	8,57	22	8,20	— 0,37	85	55	7,08	56	3,00	— 4,1
	14	11	25	44,29	25	43,70	— 0,59	86	18	8,00	18	2,00	— 6,0
	15	11	29	19,82	29	19,20	— 0,62	86	41	5,93	41	5,00	— 0,9
	16	11	32	5,70	32	5,40	— 1,10	87	4	14,31	4	12,00	— 2,3
	17	11	36	30,82	36	29,90	— 0,92	87	27	27,62	27	23,00	— 4,6
	19							88	13	53,74	13	51,00	— 2,7
	21	11	50	52,24	50	41,30	— 0,94	89	0	32,99	0	29,00	— 4,0
22	11	54	27,86	54	26,80	— 1,06	89	23	58,32	23	51,00	— 7,3	
24	12	1	39,09	1	38,00	— 1,09	90	10	44,58	10	38,00	— 6,6	
25	12	5	14,41	5	13,80	— 0,61	90	34	4,23	34	2,00	— 2,2	
26	12	8	50,68	8	49,70	— 0,98	90	57	30,43	57	27,00	— 3,4	
27	12	12	26,95	12	25,80	— 1,15	91	20	57,14	20	52,00	— 5,1	
28	12	16	3,20	16	2,30	— 0,90	91	44	17,17	44	16,00	— 1,2	
29	12	18	39,83	19	39,00	— 0,83	92	7	42,70	7	40,00	— 2,7	
October	4	12	37	47,36	37	46,20	— 1,16	94	4	17,70	4	20,00	+ 2,3
	5	12	41	25,86	41	24,70	— 1,16	94	27	32,17	27	33,00	+ 0,8
	6	12	45	4,57	45	3,40	— 1,17	94	50	44,85	50	42,00	— 2,8
	7	12	48	43,48	48	42,60	— 0,88	95	13	56,28	13	48,00	— 8,3
	8	12	52	22,75	52	22,20	— 0,55	95	36	52,11	36	50,00	— 2,1
	12	13	7	5,70	7	4,60	— 1,10	97	8	11,00	8	12,00	+ 1,0
	13	13	10	47,47	10	46,30	— 1,17	97	30	48,88	30	48,00	— 0,9
	14	13	14	29,80	14	28,60	— 1,20	97	53	16,97	53	17,00	+ 0,0
15	13	18	12,33	18	11,40	— 0,93	98	15	40,26	15	40,00	— 0,3	

		Observed A.R.			A.R. from Nautical Almanac.		Error of Tables.	Observed N.P.D.		N.P.D. from Nauti- cal Almanac.	Error of Tables.	
1831		<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>m.</i>	<i>s.</i>	<i>s.</i>	<i>°</i>	<i>m.</i>	<i>s.</i>	<i>m.</i>	<i>s.</i>
October	17	13	25	39,52	25	38,50	— 1,02	99	0	3,95	0	7,00
	21	13	40	40,70	40	39,70	— 1,00	100	27	22,74	27	19,00
	22	13	44	27,55	44	26,50	— 1,05	100	48	50,54	48	46,00
	23	13	48	15,10	48	14,10	— 1,00	101	10	1,26	10	2,00
	25	13	55	52,39	55	51,30	— 1,09	101	52	6,20	52	4,00
	30	14	15	7,99	15	7,50	— 0,49	103	33	57,74	33	56,00
Nov.	31	14	19	1,42	19	1,00	— 0,42	103	53	39,44	53	43,00
	1	14	22	56,61	22	55,40	— 1,21	104	13	16,84	13	15,00
	2	14	26	51,44	26	50,50	— 0,94	104	32	33,34	32	34,00
	3	14	30	47,49	30	46,40	— 1,09	104	51	42,87	51	38,00
	8	14	50	39,43	50	38,70	— 0,73	106	23	10,39	23	16,00
	9	14	54	40,09	54	39,60	— 0,49	106	40	49,01	40	47,00
	10	14	58	42,11	58	41,40	— 0,71	106	58	2,44	58	1,00
	12	15	6	48,06	6	47,30	— 0,76	107	31	37,85	31	38,00
	13	15	10	51,95	10	51,60	— 0,35	107	48	6,76	47	59,00
	18	15	31	26,05	31	25,40	— 0,65	109	4	56,01	4	55,00
	19	15	35	34,74	35	34,60	— 0,14	109	19	18,39	19	19,00
	20	15	39	45,55	39	44,60	— 0,95	109	33	22,84	33	21,00
	21	15	43	56,45	43	55,40	— 1,05	109	47	2,57	47	1,00
	22	15	48	8,07	48	7,20	— 0,87	110	0	22,31	0	21,00
Dec.	27	16	9	18,20	9	17,60	— 0,60	111	1	18,15	1	22,00
	29	16	17	52,19	17	51,20	— 0,99	111	23	3,48	23	2,00
	1	16	26	28,15	26	27,70	— 0,45	111	43	6,28	43	7,00
	2	16	30	47,72	30	47,00	— 0,72	111	52	28,75	52	32,00
	3	16	35	7,18	35	6,80	— 0,38	112	1	31,69	1	32,00
	4	16	39	27,74	39	27,30	— 0,44	112	10	4,13	10	6,00
	5	16	43	49,14	43	48,40	— 0,74	112	18	9,56	18	15,00
	6	16	48	10,47	48	10,10	— 0,37					
	7	16	52	33,00	52	32,30	— 0,70	112	33	12,14	33	12,00
	8	16	56	55,51	56	55,10	— 0,41	112	39	59,79	40	2,00
	10	17	5	42,17	5	41,90	— 0,27	112	52	19,09	52	21,00
	12	17	14	31,14	14	30,20	— 0,94	113	2	45,82	2	50,00
	13	17	18	56,06	18	55,00	— 1,06	113	7	21,50	7	23,00
	14	17	23	21,06	23	20,10	— 0,96	113	11	24,33	11	30,00
	15	17	27	46,13	27	45,50	— 0,63	113	15	6,63	15	9,00
	16	17	32	12,26	32	11,10	— 1,16	113	18	17,11	18	20,00
	17	17	36	37,94	36	36,80	— 1,14	113	20	58,32	21	2,00
	19	17	45	30,27	45	29,00	— 1,27	113	24	57,30	25	3,00
	20	17	49	56,22	49	55,40	— 0,82	113	26	18,21	26	21,00
21	17	54	23,11	54	21,90	— 1,21	113	27	7,54	27	11,00	
22	17	58	49,56	58	48,50	— 1,06	113	27	31,33	27	33,00	
23	18	3	16,20	3	14,90	— 1,30	113	27	21,84	27	27,00	
24	18	7	42,62	7	41,50	— 1,12	113	26	41,37	26	52,00	
26							113	24	10,09	24	17,00	
29	18	29	54,80	29	53,80	— 1,00	113	16	47,34	16	53,00	
31	18	38	46,41	38	45,30	— 1,11	113	9	31,19	9	36,00	

We can now compute from the above observations the value of ω the apparent obliquity of the Ecliptic: for this purpose we have at once, $\tan.$

Note. The above columns of A.R. and N.P.D. are necessarily due to the instant of the Sun's centre transiting the Meridian of the Madras Observatory, which according to the observations of my predecessor is situated in longitude $5h. 21m. 9,00s.$ East of the Greenwich Royal Observatory.

$w = \tan. D \times \text{cosect. A.R.}$ or we may compute the reduction to the Solstice R by the formulæ $R = 2 \sin. 2\frac{1}{2} d \text{ A.R.} \times \cos. D \times \frac{\sin. w'}{\sin. 1''}$. where d A.R. represents the distance of the Sun's true Right Ascension, from the Solstice, and w' , the apparent obliquity as near as it is known, for an error of 3 or 4 seconds in the assumption does not perceptibly alter the value of R. In each of these cases we determine the obliquity with reference to the place of the Sun, which (by reason of the action of the Moon and Planets upon the Earth,) does not appear accurately to describe the Ecliptic; account must consequently be taken of the Sun's latitude. Making use of the second formulæ, assuming $w' = 23^\circ 27' 34''$, and computing the Sun's latitude from Vince's Solar Tables, we have from the observations about the time of Summer Solstice.

	1831	N.P.D. of the Sun at the Summer Solstice.			R	Sun's Latitude.	Solstitial N.P.D.	Correction for		REMARKS.
		°	'	"				D Nut.	☉ Nut. t. 0,46 + 365	
May	30	68	19	41,32	1 47 20,88	+ 0,48	*66 32 20,92	+ 7,68	+ 0,59	
June	1	68	2	12,49	1 29 42,43	+ 0,18	30,24	+ 7,68	+ 0,61	
	2	67	53	50,07	1 21 27,12	+ 0,04	22,99	+ 7,67	+ 0,62	
	3	67	45	57,42	1 13 33,38	— 0,10	23,94	+ 7,67	+ 0,63	
	4	67	38	27,39	1 6 5,60	— 0,25	* 21,54	+ 7,66	+ 0,65	
	5	67	31	31,87	0 59 0,24	— 0,39	31,24	+ 7,65	+ 0,66	
	6	67	24	44,60	0 52 19,15	— 0,49	21,96	+ 7,65	+ 0,67	
	7	67	18	27,93	0 45 59,96	— 0,56	27,41	+ 7,64	+ 0,68	
	8	67	12	27,90	0 40 4,20	— 0,60	23,10	+ 7,64	+ 0,69	
	9	67	7	2,31	0 34 33,26	— 0,62	28,43	+ 7,63	+ 0,70	
	10	67	1	55,96	0 29 27,00	— 0,59	28,37	+ 7,62	+ 0,71	
	11	66	57	14,54	0 24 42,75	— 0,53	31,26	+ 7,62	+ 0,72	
	12	66	52	57,91	0 20 26,97	— 0,43	30,51	+ 7,61	+ 0,73	
	13	66	49	4,12	0 16 33,00	— 0,30	30,82	+ 7,60	+ 0,73	
	14	66	45	29,31	0 13 3,95	— 0,16	25,20	+ 7,60	+ 0,73	
	16	66	39	49,73	0 7 19,62	+ 0,14	30,25	+ 7,59	+ 0,74	
	19	66	34	9,04	0 1 48,58	+ 0,58	* 21,04	+ 7,58	+ 0,76	
	21	66	32	38,75	0 0 11,38	+ 0,74	28,11	+ 7,58	+ 0,76	
	22	66	32	30,67	0 0 0,02	+ 0,77	31,42	+ 7,57	+ 0,76	
	23	66	32	37,94	0 0 13,50	+ 0,77	25,21	+ 7,56	+ 0,76	
	24	66	33	20,66	0 0 51,63	+ 0,74	29,77	+ 7,56	+ 0,76	
	25	66	34	24,83	0 1 54,74	+ 0,66	30,75	+ 7,55	+ 0,76	
	26	66	35	50,95	0 3 22,64	+ 0,58	28,89	+ 7,54	+ 0,76	
	27	66	37	48,33	0 5 15,12	+ 0,43	33,64	+ 7,54	+ 0,75	
	28	66	39	58,57	0 7 32,22	+ 0,29	26,64	+ 7,53	+ 0,75	
	29	66	42	43,86	0 10 13,83	+ 0,14	30,17	+ 7,53	+ 0,75	
	30	66	45	44,26	0 13 20,22	+ 0,02	24,06	+ 7,52	+ 0,75	
July	1	66	49	20,65	0 16 50,87	— 0,13	29,65	+ 7,52	+ 0,74	
	2	66	53	16,37	0 20 45,68	— 0,26	30,43	+ 7,51	+ 0,74	
	3	66	57	34,25	0 25 4,28	— 0,36	29,61	+ 7,51	+ 0,73	

* These are omitted in taking the mean.

RESULTS FROM OBSERVATIONS, 1831.

	1831	N.P.D. of the Sun at the Summer Solstice.			R	Sun's Latitude.	Solstitial N.P.D.	Correction for		REMARKS.			
		°	'	"				☉ Nut.	☉ Nut. t. 0,46 + 365				
July	6	67	12	55,05	0 40	27,24	—	0,50	27,31	+ 7,49	+ 0,71		
	7	67	18	48,93	0 46	22,70	—	0,48	25,75	+ 7,48	+ 0,70		
	8	67	25	8,24	0 52	41,10	—	0,44	26,70	+ 7,48	+ 0,69		
	11	67	46	22,42	1 13	57,38	—	0,09	24,95	+ 7,46	+ 0,67		
	14	68	11	6,01	1 38	42,18	+	0,36	24,19	+ 7,45	+ 0,64		
	17	68	39	10,43	2 0	47,20	+	0,78	24,01	+ 7,44	+ 0,60		
Mean of 33 = 66 32 27,87													
90													
Mean value of ϖ = 23 27 32,13											+ 7,57	+ 0,71	

Similarly we have from the observations about the time of the Winter Solstice.

	1831	N.P.D. of the Sun at the Winter Solstice.			R	Sun's Latitude.	Solstitial N.P.D.	Correction for		REMARKS.		
		°	'	"				☉ Nut.	☉ Nut. t. 0,46 + 365			
Jan.	28	108	21	56,49	5 5	31,86	—	0,45	113 27 27,90	+ 8,23	+ 0,16	
	29	108	6	13,47	5 21	18,90	—	0,31	32,06	+ 8,23	+ 0,14	
	30	107	50	5,55	5 37	25,72	—	0,17	31,10	+ 8,22	+ 0,12	
Feb.	1	107	16	53,95	6 10	35,34	+	0,14	29,44	+ 8,21	+ 0,08	
	2	106	59	51,73	6 27	37,72	+	0,29	29,74	+ 8,20	+ 0,06	
Nov.	12	107	31	37,85	5 55	55,38	+	0,27	33,50	+ 6,79	+ 0,49	
	13	107	48	6,76	5 39	36,83	+	0,10	33,69	+ 6,78	+ 0,51	
	18	109	4	56,01	4 22	38,30	—	0,42	33,89	+ 6,74	+ 0,61	
	19	109	19	18,39	4 8	17,28	—	0,44	35,22	+ 6,74	+ 0,63	
	20	109	33	22,84	3 54	12,45	—	0,44	34,85	+ 6,73	+ 0,64	
	21	109	47	2,57	3 40	31,22	—	0,38	33,41	+ 6,73	+ 0,66	
	22	110	0	22,31	3 27	12,07	—	0,31	34,07	+ 6,72	+ 0,68	
	27	111	1	18,15	2 26	11,78	+	0,40	30,33	+ 6,69	+ 0,77	
	29	111	23	3,43	2 4	29,45	+	0,68	33,61	+ 6,68	+ 0,80	
Dec.	1	111	43	6,28	1 44	26,36	+	0,84	33,48	+ 6,67	+ 0,82	
	2	111	52	28,75	1 35	1,05	+	0,87	30,67	+ 6,66	+ 0,83	
	3	112	1	31,69	1 26	2,20	+	0,87	34,76	+ 6,65	+ 0,84	
	4	112	10	4,13	1 17	28,07	+	0,84	33,04	+ 6,64	+ 0,85	
	5	112	18	9,56	1 9	19,22	+	0,77	29,55	+ 6,63	+ 0,87	
	7	112	33	12,14	0 54	21,03	+	0,53	33,70	+ 6,62	+ 0,90	
	8	112	39	59,79	0 47	31,86	+	0,38	32,03	+ 6,61	+ 0,91	
	10	112	52	19,09	0 35	13,50	+	0,09	32,68	+ 6,60	+ 0,93	
	12	113	2	45,82	0 24	43,05	—	0,23	28,64	+ 6,59	+ 0,94	
	13	113	7	21,50	0 20	9,17	—	0,35	30,32	+ 6,58	+ 0,95	
	14	113	11	24,33	0 16	3,08	—	0,45	26,96	+ 6,57	+ 0,96	
	15	113	15	6,63	0 12	24,86	—	0,51	30,98	+ 6,57	+ 0,96	
	16	113	18	17,11	0 9	13,97	—	0,52	30,56	+ 6,56	+ 0,97	
	17	113	20	58,32	0 6	34,60	—	0,52	32,40	+ 6,56	+ 0,98	
	19	113	24	57,30	0 2	30,67	—	0,40	27,57	+ 6,54	+ 0,98	
	20	113	26	18,21	0 1	12,63	—	0,30	30,54	+ 6,53	+ 0,98	
	21	113	27	7,54	0 0	22,60	—	0,15	29,99	+ 6,52	+ 0,99	
	22	113	27	31,33	0 0	1,00	—	0,01	32,32	+ 6,52	+ 0,99	

	N.P.D. of the Sun at the Summer Solstice.	R	Sun's Latitude.	Solstitial N.P.D.	Correction for		REMARKS.
					☽ Nut.	☉ Nut. t. 0,46 + 365	
1831	° ' "	° ' "	"	° ' "	"	"	
Dec. 23	113 27 21,84	0 0 7,67	+ 0,13	29,64	+ 6,51	+ 1,00	
24	113 26 41,37	0 0 42,65	+ 0,28 *	24,30	+ 6,50	+ 0,99	
26	113 24 10,09	0 3 17,36	+ 0,55	28,00	+ 6,49	+ 0,98	
29	113 16 47,34	0 10 41,64	+ 0,76	29,74	+ 6,47	+ 0,97	
31	113 9 31,19	0 17 57,96	+ 0,74	29,89	+ 6,46	+ 0,96	
Mean of 36 = 113 27 31,397							
90							
Mean value of ω = 23 27 31,397					+ 6,84	+ 0,74	

In the above column "☉. Nut. + $\frac{t. 0'',46''}{365}$ ". t represents the number of days from January 1, so that we have the mean obliquity for January 1, 1831, from observations near the Summer Solstice = $23^\circ 27' 40'',41$
 ----- Winter Solstice = $38'',98$.

Now at the time of Summer Solstice, the Sun is situated $10^\circ 23'$ to the North, and at the Winter Solstice, $36^\circ 31'$ to the South of the Zenith; hence the former (on account of the small uncertainty in the refraction) is preferable to the latter, and cannot on this account err to the amount of one tenth of a second; in the case of the Winter Solstice, an error exceeding four or five tenths of a second need not be expected, since the several tables of refraction do not disagree, (at this distance from the Zenith) to a larger amount; the disagreement above found is consequently to be accounted for, by errors of observation, and a wrong assumption of the latitude; in the absence of data to show the probable amount of the error of observation, we will suppose the latitude to be the sole cause of disagreement, hence it appears that instead of $13^\circ 4' 9'',0$ (which we have assumed for the latitude of Madras,) we must use $13^\circ 4' 8'',29$ in order to reconcile the observations of the Summer and Winter Solstice, and we have the mean obliquity for January 1, 1831, = $23^\circ 27' 39'',70$.

We will now from the observations near the time of the Equinoxes compare the Right Ascension of the Sun as determined from Stars by the Transit Instrument, with the A.R. computed from the observations of N.P.D. we have as follows.

* This is omitted in taking the mean.

RESULTS FROM OBSERVATIONS, 1831.

Observations of the Sun near the Vernal Equinox.

	Observed N.P.D. of the Sun.	Correc- tions.	Reduced N.P.D.	Computed A.R.	Observed A.R.	Error of Eq. Point.	REMARKS.
1831	° m. s.	s.	° m. s.	h. m. s.	m. s.	s.	
May 9	94 45 12,92	+ 0,36	94 45 13,28	23 15 48,53	15 49,23	+ 0,70	
10	94 21 38,54	+ 0,24	94 21 38,78	21 19 31,04	19 30,46	- 0,58	
11	93 58 12,89	+ 0,11	93 58 13,00	23 23 11,27	23 11,53	+ 0,26	
12	93 34 35,62	- 0,03	93 34 35,59	23 26 52,54	26 51,98	- 0,56	
13	93 11 1,71	- 0,15	93 11 1,56	23 30 32,58	30 32,11	- 0,47	
14	92 47 23,76	- 0,30	92 47 23,46	23 34 12,63	34 12,31	- 0,32	
15	92 23 45,89	- 0,44	92 23 45,45	23 37 52,16	37 52,15	+ 0,04	
17	91 36 27,07	- 0,63	91 36 26,44	23 45 10,22	45 10,01	- 0,21	
19	90 48 59,36	- 0,70	90 48 58,66	23 52 28,45	52 28,23	- 0,22	
20	90 25 21,09	- 0,70	90 25 20,39	23 56 6,43	56 6,47	+ 0,04	
22	89 37 55,28	- 0,60	89 37 54,68	0 3 23,63	3 23,67	+ 0,04	
23	89 14 20,76	- 0,50	89 14 20,26	0 7 0,98	7 2,15	+ 1,17	
25	88 27 4,51	- 0,22	88 27 4,29	0 14 17,32	14 17,28	- 0,04	
27	87 39 58,84	- 0,05	87 39 58,79	0 21 33,23	21 33,42	+ 0,19	
28	87 16 28,75	+ 0,18	87 16 28,93	0 25 11,37	25 11,25	- 0,12	
30	86 29 41,22	+ 0,41	86 29 41,63	0 32 27,38	32 27,00	- 0,38	
31	86 6 31,75	+ 0,50	86 6 32,25	0 36 4,14	36 5,06	+ 0,92	
April 1	85 43 13,33	+ 0,54	85 43 13,87	0 39 43,04	39 43,24	+ 0,20	
2	85 20 8,81	+ 0,55	85 20 9,36	0 43 20,60	43 21,09	+ 0,40	
Mean of 19 = + 0,055							

Observations of the Sun near the Autumnal Equinox.

	Observed N.P.D. of the Sun.	Correc- tions.	Reduced N.P.D.	Computed A.R.	Observed A.R.	Error of Eq. Point.	REMARKS.
1832	° m. s.	s.	° m. s.	h. m. s.	m. s.	s.	
Sept. 11	85 9 24,86	+ 0,36	85 9 25,72	11 14 57,88	14 56,98	- 0,90	
12	85 32 11,87	+ 0,88	85 32 12,75	11 18 33,25	18 33,17	- 0,80	
13	85 55 7,08	+ 0,87	85 55 7,95	11 22 8,89	22 8,57	- 0,32	
14	86 18 8,00	+ 0,83	86 18 8,83	11 25 44,67	25 44,29	- 0,38	
15	86 41 5,93	+ 0,74	86 41 6,67	11 29 19,28	29 19,82	+ 0,54	
16	87 4 14,31	+ 0,62	87 4 14,93	11 32 54,89	32 55,70	+ 0,81	
17	87 27 27,62	+ 0,49	87 27 28,11	11 36 30,71	36 30,82	+ 0,11	
19	88 13 53,74	+ 0,22	88 13 53,96	11 43 40,89	43 41,46	+ 0,57	
21	89 0 32,99	- 0,02	89 0 32,97	11 50 51,84	50 52,24	+ 0,40	
22	89 23 58,32	- 0,17	89 23 58,15	11 54 27,85	54 27,86	+ 0,01	
24	90 10 44,58	- 0,33	90 10 44,25	12 1 38,97	1 39,09	+ 0,12	
25	90 34 4,23	- 0,36	90 34 3,87	12 5 14,01	5 14,41	+ 0,40	
26	90 57 30,43	- 0,35	90 57 30,08	12 8 50,17	8 50,68	+ 0,51	
27	91 20 57,14	- 0,32	91 20 56,82	12 12 26,60	12 26,95	+ 0,35	
28	91 44 17,17	- 0,25	91 44 16,92	12 16 2,26	16 3,20	+ 0,94	
29	92 7 42,70	- 0,16	92 7 42,54	12 19 39,09	19 39,83	+ 0,74	
Oct. 6	94 50 44,85	+ 0,74	94 50 45,59	12 45 3,85	45 4,57	+ 0,72	
Mean of 17 = + 0,267							

Taking the mean between 19 observations in the Spring and 17 in the Autumn, we have for the error of the assumed Equinoctial point + 0",16. Now the Equinoctial point we have assumed is Dr. Marklyne + 0",21, hence the true point of the Equinox from the observations appears to be Dr. Marklyne + 0",05. To account for the disagreement between the results of Spring and Autumn we will again suppose the latitude in error, for this purpose, at the time of the Equinoxes ; we have

$$d \text{ Declination} = \frac{d \text{ A.R.}}{\cot. w}$$

this being applied in the above case gives the latitude of Madras (which will reconcile the observations of the Sun at the Vernal and Autumnal Equinox,) 13° 4' 8",20 agreeing very nearly with that found at Page 77.

We now come to the Planetary Observations ; these have been reduced only to their *apparent* place, as would be viewed by an observer situated at the centre of the Earth, and consequently require to be corrected for aberration; the Parallaxes have been computed from the numbers given in the Supplement to the Nautical Almanac which are stated to be computed with the assistance of Professor SCHUMACHER'S Tables.

Apparent Right Ascension and North Polar Distance of MERCURY.

Date.	Mean Time of Observation.	Point Observed.	A.R.	Point Observed.	N.P.D.	REMARKS.
1831	<i>h. m. s.</i>		<i>h. m. s.</i>		<i>m. s.</i>	
February 18	22 26 6,1	Centre.	20 19 42,93	Centre.	109 7 29,06	
20	22 28 50,8	—	20 30 21,34	—	108 59 27,52	
23	—	—	—	—	108 37 33,61	
26	22 34 1,0	—	20 59 11,27	—	108 3 37,27	
27	22 35 17,6	—	21 4 24,65	—	107 49 46,64	
March 1	22 38 6,0	—	21 15 6,58	—	107 18 1,81	
2	—	—	—	—	107 0 6,07	
3	22 43 16,4	—	21 28 11,10	—	107 40 53,22	
5	—	—	—	—	105 58 33,29	
6	22 46 23,5	—	21 43 8,36	—	105 35 27,14	
7	22 48 3,0	—	21 48 54,69	—	105 10 59,85	
8	22 50 6,9	—	21 54 45,41	—	104 45 15,29	
10	22 54 4,0	—	22 6 36,35	—	103 49 51,26	
11	22 56 6,4	—	22 12 35,78	—	103 20 14,49	
12	22 58 21,0	—	22 18 47,19	—	102 49 22,39	
16	23 7 0,1	—	22 43 13,97	—	100 33 20,39	
19	—	—	—	—	98 38 11,70	
21	23 19 4,1	—	23 15 2,56	—	97 15 31,28	
24	23 26 54,7	—	23 34 43,61	—	—	
28	23 38 7,6	—	0 1 45,23	—	91 49 32,43	
30	23 44 7,5	—	0 15 39,01	—	90 7 0,59	
July 1	22 43 24,8	—	5 21 35,89	—	68 9 0,70	
5	22 57 55,4	—	5 51 45,11	—	67 2 14,60	
7	23 6 35,1	—	6 8 19,47	—	66 38 11,01	
10	23 21 3,9	—	6 34 40,39	—	66 18 9,58	

Apparent Right Ascension and North Polar Distance of VENUS.

Date.	Mean Time.	Point Observed.	A.R.	Point Observed.	N.P.D.	REMARKS.
1831	<i>h. m. s.</i>		<i>h. m. s.</i>		<i>° m. s.</i>	
February 19	1 10 25,2	Centre.	23 4 28,98	Centre.	97 28 52,29	
20	1 11 6,3	—	23 9 6,82	—	96 59 11,14	
21	1 11 46,7	—	23 13 43,68	—	96 29 25,88	
22	1 12 25,6	—	23 18 19,25	—	95 59 21,93	
23	1 13 3,3	—	23 22 53,55	—	95 29 11,31	
24	1 13 41,4	—	23 27 28,21	—	94 58 52,48	
25	1 14 20,0	—	23 32 3,56	—	94 28 20,32	
26	1 14 55,1	—	23 36 35,49	—	93 57 42,04	
March 2		—		—	91 54 28,77	
3	1 17 52,1	—	23 59 15,62	—	91 23 30,09	
4		—		—	90 52 25,80	
5	1 19 0,7	—	0 8 17,48	—	90 21 25,37	
6	1 19 35,5	—	0 12 48,93	—	89 50 20,38	
7	1 20 8,7	—	0 17 18,80	—	89 19 13,19	
8	1 20 42,9	—	0 21 49,57	—	88 48 1,00	
10	1 21 50,8	—	0 30 50,91	—	87 45 55,60	
11	1 22 24,5	—	0 35 21,22	—	87 14 56,40	
12	1 22 58,3	—	0 39 51,63	—	86 43 57,59	
13	1 23 32,8	—	0 44 22,61	—	86 13 5,36	
14	1 23 6,9	—	0 47 53,45	—	85 42 23,91	
17		—		—	84 10 24,93	
19	1 27 2,8	—	1 11 32,63	—	83 9 38,93	
20	1 27 40,3	—	1 16 6,77	—	82 39 37,21	
21		—		—	82 9 46,58	
22		—		—	81 40 3,65	
27		—		—	79 14 18,65	
28	1 32 46,2	—	1 52 46,18	—	78 45 52,84	
29	1 33 27,6	—	1 57 24,20	—	78 17 44,96	
April 2	1 36 21,0	—	2 16 4,20	—	76 28 1,51	
10	1 42 51,1	—	2 54 7,66	—	73 4 50,97	
May 22	2 31 37,4	—	6 28 37,43	—	64 45 14,87	
23	2 32 52,7	—	6 33 49,29	—		
26	2 36 32,4	—	6 49 19,34	—	65 0 32,04	
27	2 37 44,8	—	6 54 28,50	—	65 4 6,67	
30	2 41 15,4	—	7 9 49,30	—		
June 1	2 43 30,1	—	7 19 57,52	—	65 40 48,55	
2	2 44 36,0	—	7 25 0,24	—	65 50 1,63	
3	2 45 41,3	—	7 30 2,31	—	65 59 54,77	
5	2 47 46,1	—	7 40 0,53	—	66 21 33,59	
29	3 4 41,1	—	9 31 35,72	—	73 30 59,93	
July 2	3 5 45,1	—	9 44 20,40	—	74 42 31,35	
August 12	2 50 31,7	—	12 11 2,38	—		
13	2 49 37,6	—	12 13 54,59	—	93 50 37,52	
20	2 41 48,5	—	12 33 41,18	—	96 53 56,27	
Dec. 10	20 46 48,7	2 Limb.	14 3 12,38	—	99 37 56,36	

Apparent Right Ascension and North Polar Distance of MARS.

Date.	Mean Time.	Point Observed.	A.R.	Point Observed.	N.P.D.	REMARKS.
1831	<i>h. m. s.</i>		<i>h. m. s.</i>		<i>m. s.</i>	
February 20	4 51 56,9	Centre.	2 50 33,88	Centre.	72 28 16,79	
21	4 50 26,9	—	2 52 59,84	—	72 17 13,89	
22	4 48 56,0	—	2 55 25,14	—	72 6 13,17	
23	4 47 27,1	—	2 57 52,63	—	71 55 25,01	
25		—		—	71 33 59,55	
28	4 40 5,4	—	3 10 12,47	—	71 2 50,60	
March 4	4 34 19,0	—	3 20 11,32	—	70 22 58,21	
13	4 21 44,8	—	3 43 3,94	—	69 0 31,21	
May 22		—		—	65 44 10,32	

Apparent Right Ascension and North Polar Distance of JUPITER.

Date.	Mean Time.	Point Observed.	A.R.	Point Observed.	N.P.D.	REMARKS.
1831	<i>h. m. s.</i>		<i>h. m. s.</i>		<i>m. s.</i>	
March 1	22 9 23,4	Centre.	20 46 19,34	Centre.	108 26 22,51	
3	22 3 31,1	—	20 48 19,32	—	108 19 39,93	
5	21 57 38,7	—	20 50 18,98	—	108 11 1,98	
August 23	11 9 2,2	—	21 14 6,83	—	107 7 48,31	
26	10 55 47,8	—	21 12 39,90	—	107 14 26,23	
28	10 47 0,6	—	21 11 44,25	—	107 18 37,22	
29	10 42 36,9	—	21 11 16,51	—		
30	10 38 13,9	—	21 10 49,25	—	107 22 43,96	
Sept. 1	10 29 9,2	—	21 9 56,21	—	107 26 35,25	
2	10 25 7,8	—	21 9 30,78	—	107 28 29,80	
4	10 16 25,8	—	21 8 40,28	—	107 31 11,22	
9	9 54 49,9	—	21 6 43,85	—	107 40 34,25	
11	9 46 15,8	—	21 6 1,13	—	107 43 30,75	
13	9 37 43,8	—	21 5 21,22	—	107 46 21,60	
15	9 29 14,5	—	21 4 43,49	—	107 49 2,97	
17	9 20 48,2	—	21 4 8,93	—	107 51 28,24	
20	9 8 13,6	—	21 3 21,89	—	107 54 34,17	
22	8 59 55,0	—	21 2 54,06	—	107 56 29,57	
24	8 51 37,3	—	21 2 29,12	—	107 58 3,27	
29	8 31 9,6	—	21 1 40,79	—		
30	8 27 4,0	—	21 1 31,12	—	108 1 32,63	
October 2	8 19 1,3	—	21 1 20,26	—	108 2 13,30	
3	8 15 0,5	—	21 1 15,29	—	108 2 30,18	
6		—		—	108 3 10,90	
8	7 55 7,6	—	21 1 1,98	—		
14	7 31 43,0	—	21 1 12,75	—		
16	7 24 2,8	—	21 1 24,52	—	108 0 48,00	
20	7 8 44,5	—	21 1 49,69	—		
21	7 4 59,6	—	21 2 0,79	—	107 57 36,71	
22	7 1 13,8	—	21 2 10,87	—	107 56 44,64	
23	6 57 29,4	—	21 2 22,43	—	107 55 52,67	
25	6 50 0,7	—	21 2 45,63	—	107 53 58,85	
30	6 31 33,1	—	21 3 57,91	—	107 48 14,36	
31	6 27 55,2	—	21 4 15,95	—	107 46 57,40	
Nov. 1	6 24 16,3	—	21 4 33,13	—	107 45 36,85	
3	6 17 1,7	—	21 5 10,14	—	107 42 44,09	

Apparent Right Ascension and North Polar Distance of JUPITER, continued.

Date.	Mean Time.	Point Observed.	A.R.	Point Observed.	N.P.D.	REMARKS..
1831	<i>h. m. s.</i>		<i>h. m. s.</i>		<i>m. s.</i>	
Nov. 6	6 6 15,7	Centre.	21 6 12,06	Centre.	107 38 9,67	
9	5 55 34,8	—	21 7 19,14	—	107 32 48,83	
Dec. 11	4 7 21,4	—	21 24 57,55	—	106 10 12,28	

It is proper I should here state, that the Right Ascension of the centre of Jupiter was obtained by taking the time of transit of the *first limb* at the first and second wires, of the *centre* by estimation, at the third wire, and of the *second limb* at the fourth and fifth wires.

Apparent Right Ascension and North Polar Distance of SATURN.

Date.	Mean Time.	Point Observed.	A.R.	Point Observed.	N.P.D.	REMARKS.
1831	<i>h. m. s.</i>		<i>h. m. s.</i>		<i>m. s.</i>	
March 5	11 8 38,4	Centre.	9 59 32,05	Centre.	75 56 45,81	
6	11 4 26,1	—	9 59 15,68	—	75 55 10,39	
8	10 55 59,4	—	9 58 40,58	—	75 52 7,53	
9	10 51 46,7	—	9 58 23,84	—	75 50 36,80	
10	10 47 34,0	—	9 58 7,04	—	75 49 9,29	
11	10 43 21,9	—	9 57 50,78	—	Cloudy.
12	10 39 9,4	—	9 57 33,02	—	75 46 12,36	
13	10 34 57,9	—	9 57 18,42	—	75 44 51,08	
16	10 22 25,6	—	9 56 33,83	—	75 40 44,50	
18	10 14 2,7	—	9 56 2,72	—	75 38 14,59	
19	10 9 52,9	—	9 55 48,63	—	75 37 1,04	
20	10 5 42,2	—	9 55 33,82	—	75 35 41,60	
21		—		—	75 31 14,01	
23	9 53 13,4	—	9 54 52,70	—		
25	9 44 56,2	—	9 54 27,25	—		
28		—		—	75 26 0,06	
29	9 28 25,0	—	9 53 39,81	—	75 25 2,78	
30	9 25 17,4	—	9 54 27,93	—	75 24 9,49	
April 1	9 16 5,2	—	9 53 7,66	—	75 22 26,95	
2	9 11 59,1	—	9 52 57,19	—	75 21 38,47	
3	9 7 53,8	—	9 52 47,77	—	75 21 49,66	
5	8 59 43,3	—	9 52 29,07	—	75 19 22,26	
6	8 55 39,2	—	9 52 20,79	—	75 18 41,20	
9	8 43 28,8	—	9 51 57,77	—	75 16 49,80	
10		—		—	75 16 20,30	
11	8 35 22,8	—	9 51 43,73	—	75 15 50,18	
13	8 27 17,6	—	9 51 30,56	—		
14	8 23 16,9	—	9 51 25,87	—		
15	8 19 15,8	—	9 51 20,59	—		
16	8 15 15,6	—	9 51 16,03	—		
20	7 59 17,5	—	9 51 1,56	—	75 12 59,62	
21	7 55 18,4	—	9 50 58,41	—	75 12 48,22	
22		—	9 50 56,40	—	75 12 41,72	

Apparent Right Ascension and North Polar Distance of SATURN, continued.

Date.	Mean Time.	Point Observed.	A.R.	Point Observed.	N.P.D.	REMARKS.
1831	<i>h. m. s.</i>		<i>h. m. s.</i>		<i>° m. s.</i>	
April 23	7 47 21,7	Centre.	9 50 53,47	Centre.	75 12 38,77	
25	7 39 28,3	—	9 50 51,76	—	75 12 37,91	
26	7 35 31,9	—	9 50 51,28	—	75 12 40,21	
27	7 31 36,1	—	9 50 51,41	—		
28	7 27 40,6	—	9 50 51,85	—	75 12 54,17	
29	7 23 45,1	—	9 50 52,23	—	75 13 4,37	
30	7 19 50,4	—	9 50 53,46	—	75 13 17,57	
May 1	7 15 55,2	—	9 50 54,11	—	75 13 32,64	
2	7 12 1,2	—	9 50 55,99	—	75 13 46,13	
3	7 8 8,0	—	9 50 58,72	—	75 14 2,53	
4	7 4 14,6	—	9 51 1,36	—	75 14 24,13	
6	6 56 29,0	—	9 51 7,56	—	75 16 10,95	
9		—		—	75 17 39,86	
10	6 41 1,9	—	9 51 26,31	—	75 18 6,71	
13	6 29 34,4	—	9 51 44,40	—	75 19 57,47	
14	6 25 45,2	—	9 51 51,23	—	75 20 39,68	
16	6 18 7,1	—	9 52 4,79	—	75 22 7,17	
17	6 14 18,9	—	9 52 12,54	—	75 22 54,39	

The above observations of Saturn were made with reference to the centre of the body by estimation; six cases however occur with the Transit observations, in which the transit of the first and second edges of the ring, was observed, they are as follows.

March 29	Transit of 2d edge —	Transit of 1st edge.....	3",30*
April 22	do.	do.	2",99*
27	do.	do.	2",60
28	do.	do.	2",66
29	do.	do.	2",80
30	do.	do.	2",62

Apparent Right Ascension and North Polar Distance of GEORGIAN SIDUS.

Date.	Mean Time.	Point Observed.	A.R.	Point Observed.	N.P.D.	REMARKS.
1831	<i>h. m. s.</i>		<i>h. m. s.</i>		<i>° m. s.</i>	
August 10	11 46 6,9	Centre.	21 0 2,43	Centre.		
23	10 53 59,6	—	20 58 1,56	—	107 55 30,69	
30	10 24 27,4	—	20 57 0,49	—	107 59 38,01	
Sept. 1	10 16 13,7	—	20 56 43,60	—	108 0 48,58	
2	10 12 14,5	—	20 56 35,29	—	108 1 20,81	
4	10 4 6,8	—	20 56 19,24	—	108 2 25,25	
7	9 51 55,4	—	20 55 55,68	—	108 3 57,61	
11	9 35 42,8	—	20 55 26,38	—	108 5 55,49	

* These were made by an inexperienced Assistant.

*Apparent Right Ascension and North Polar Distance of GEORGIAN SIDUS,
continued.*

Date.	Mean Time.	Point Observed.	A.R.	Point Observed.	N.P.D.	REMARKS.
1831	<i>h. m. s.</i>		<i>h. m. s.</i>		<i>° m. s.</i>	
Sept. 15	9 19 24,4	Centre.	20 54 51,81	Centre.	108 8 30,66	
October 2	8 13 13,5	—	20 53 31,56	—	108 13 9,89	
3	8 7 14,2	—	20 53 27,67	—	108 13 23,18	
6	7 55 16,4	—	20 53 17,63	—	108 14 0,89	
7	7 52 18,1	—	20 53 15,18	—	108 14 18,50	
8	7 47 19,2	—	20 53 12,30	—		
14	7 23 31,8	—	20 53 0,24	—		
16	7 15 40,9	—	20 53 1,00	—	108 15 0,86	
21	6 55 5,9	—	20 52 55,47	—		
22	6 52 0,2	—	20 52 55,80	—	108 14 58,96	
23	6 48 5,1	—	20 52 56,61	—	108 14 56,38	
25	6 40 14,2	—	20 52 57,57	—	108 14 46,79	

The above are I regret to state all the Planetary Observations which were made in the Year 1831; the four smaller Planets were altogether neglected by reason of my not having received (during my stay at Madras) the Supplement to the Nautical Almanac, in which their places are given, and from a variety of occupations, consequent to an observatory when first coming into action, I could not find time to compute an Ephemeris.

We now come to the Observations of the MOON, which we will compare with the interpolated place from the Nautical Almanac.

Comparison of the observed Right Ascension and North Polar Distance of the Moon, with the interpolated place from the Nautical Almanac.

Date.	Madras Mean Time.	Limb Observed.	Observed A.R. of D's Centre.	A.R. from Nautical Almanac.	Error of Tables.	Limb Observed.	Observed N.P.D. of D's Centre.	N.P.D. from Nautical Almanac.	Error of Tables.
1831	<i>h. m. s.</i>		<i>° m. s.</i>	<i>m. s.</i>	<i>s.</i>		<i>° ' "</i>	<i>' "</i>	<i>"</i>
Feb. 20	6 50 42,8	1	72 41 42,9	41 39,2	— 3,7	S	72 41 45,4	41 38,0	— 7,4
21	7 47 40,2	1	87 57 39,0	57 43,5	+ 4,5	S	71 15 46,5	15 46,2	— 0,3
22	8 44 47,2	1	103 15 41,5	15 47,9	+ 6,4	N	71 4 30,6	4 30,9	+ 0,3
23	9 40 58,3	1	118 19 41,5	19 56,3	+ 14,8	S	72 6 54,7	6 52,3	— 2,4
25	11 27 16,8	1	146 56 18,0	56 26,4	+ 8,4	S	77 16 53,3	16 49,5	— 3,8
26	12 17 48,7	Centre.	160 19 27,4	19 27,0	— 0,4	N	80 56 45,6	56 39,8	— 5,8
27	13 6 2,6	2	173 8 43,2	8 28,4	— 14,8	N	84 59 33,8	59 24,5	— 9,3
28						S	89 11 7,6	10 55,0	— 12,6
Mar. 1	14 35 38,1	2	197 34 52,0	34 38,6	— 13,4	S	93 19 17,1	19 18,5	+ 1,4
2	15 19 18,7	2	209 31 1,8	30 59,7	— 2,1	S	97 14 41,5	14 30,4	— 11,1
3	16 3 6,0	2	221 28 41,6	28 32,7	— 8,9	S	100 47 50,0	47 55,4	+ 5,4
4						S	103 51 37,4	51 36,5	— 0,9
5	17 32 58,4	2	245 58 23,5	58 26,2	+ 2,7	S	106 18 36,5	18 40,4	+ 3,9
6	18 19 49,1	2	258 41 52,5	41 45,8	— 6,7	S	108 1 57,7	2 5,6	+ 7,9

Date.	Madras Mean Time.	Limb Observed.	Observed A.R. of D's Centre.	A.R. from Nautical Almanac.	Error of Tables.	Limb Observed.	Observed N.P.D. of D's Centre.	N.P.D. from Nautical Almanac.	Error of Tables.
1831	<i>h. m. s.</i>		<i>° m. s.</i>	<i>m. s.</i>	<i>s.</i>		<i>° ' "</i>	<i>' "</i>	<i>"</i>
March	21	6 40 22,8	1	98 41 11,1	41 5,5	-	5,6	N	
	22	7 36 22,2	1	113 42 17,5	42 16,8	-	0,7	N	71 36 25,5 36 22,7 - 2,8
	23	8 30 25,5	1	128 14 11,3	14 15,4	+	4,1	N	73 23 45,6 23 46,7 + 1,1
	24							N	76 6 54,1
	25	10 11 23,7	1	155 29 27,0	30 22,8	-	4,2	N	79 31 46,1 31 38,6 - 7,5
	26	10 58 31,1	1	169 18 5,9	18 1,6	-	4,3	N	83 25 21,6 25 23,9 + 2,3
	27	11 43 57,9	1	180 40 33,2	40 27,3	-	5,9	N	87 32 48,0 32 47,0 - 1,0
	28	12 30 21,2	2	192 47 15,1	47 5,5	-	9,6	N	91 43 46,9 43 55,3 + 8,4
	29	13 14 6,8	2	204 44 38,9	44 34,0	-	4,0	S	95 46 34,0 46 33,9 - 0,1
	30	13 57 53,5	2	216 42 10,2	42 1,0	-	9,2	S	99 31 22,0 31 17,4 - 4,6
	31		2					S	102 49 9,6
April	2	16 13 22,7	2	253 36 59,0	36 52,0	-	7,0	S	107 34 58,7 34 46,3 - 12,4
	3	17 0 41,5	2	266 27 36,8	27 34,2	-	2,6	S	108 48 42,8 48 39,2 - 3,6
	19	6 27 2,5	1	123 55 16,1	55 11,1	-	5,0	N	72 34 12,2 34 7,3 - 4,9
	20	7 19 42,2	1	138 6 4,5	5 53,1	-	11,4	N	75 3 14,7 3 7,5 - 7,2
	21	8 9 27,2	1	151 32 50,6	32 47,7	-	2,9	N	78 17 28,5 17 28,3 - 0,2
	22	8 56 41,2	1	164 22 23,4	22 14,1	-	9,3	N	82 3 9,9 2 57,5 - 12,4
	23	9 41 58,1	1	176 42 18,9	42 15,1	-	3,8	N	86 6 43,1 6 44,0 + 0,9
	25	11 9 23,5	1	200 35 19,0	35 10,9	-	8,1	N	94 23 56,0 23 51,0 - 5,0
	26	11 53 47,9	1,2	212 27 25,5	27 25,0	-	0,5	N	98 17 5,4 17 7,7 + 2,3
	27	12 38 43,2	2	224 27 11,3	27 0,2	-	11,1	N	101 47 48,3 47 39,9 - 8,4
	29	14 9 15,8	2	249 6 59,6	6 49,5	-	10,1	N	107 6 42,2 6 33,4 - 8,8
	30	4 56 8,9	2	261 51 11,2	51 1,0	-	10,2	N	108 39 40,0 39 31,3 - 8,7
May	1	15 43 59,3	2	274 49 42,7	49 27,6	-	15,1	N	109 20 57,5 21 1,8 + 4,3
	2	16 32 29,9	2	287 58 38,6	58 23,2	-	15,4	N	109 6 58,1 7 6,3 + 8,2
	3	17 21 26,4	2	301 13 36,6	13 33,3	-	3,3	N	107 55 56,0 56 5,5 + 9,5
	20	7 41 6,7	1	173 1 22,1	1 12,0	-	10,1	N	84 39 7,1 39 0,8 - 6,3
	21	8 25 22,0	1	185 5 53,5	5 42,0	-	11,5	N	88 50 8,3 50 16,1 + 7,8
	22	9 8 36,3	1	196 55 17,9	55 10,2	-	7,7	N	93 0 19,8 0 17,8 - 2,0
	23	9 51 35,6	1	208 41 0,9	40 49,6	-	11,4	N	96 59 50,1 59 49,8 - 0,3
	24	10 34 56,5	1	220 32 16,0	32 8,0	-	8,0	N	100 40 7,7 40 8,6 + 0,9
	25	11 19 9,0	1	232 36 30,6	36 21,9	-	8,7	N	103 52 31,0 52 36,4 + 5,4
	26	12 5 34,1	1,2	244 58 38,7	58 30,0	-	8,7	N	106 28 53,9 28 55,8 + 1,9
	27	12 53 14,5	2	257 39 17,8	39 19,5	+	1,7	N	108 21 4,8 21 10,3 + 5,5
	29	14 29 20,0	2	283 42 39,0	42 36,5	-	2,5	N	109 29 13,5 29 15,6 + 2,1
	30	15 18 3,6	2	296 54 39,6	54 38,5	-	1,1	N	108 38 37,6 38 48,2 + 10,6
	31	16 6 44,0	2	310 5 58,4	5 58,8	+	0,4	N	106 52 10,4 52 17,0 + 6,6
June	1	16 55 11,3	2	323 13 53,3	13 52,6	-	0,7	N	104 12 58,9 13 11,6 + 12,7
	20	8 33 31,1	1	216 42 40,7	42 34,8	-	15,9	N	99 22 31,8 22 35,8 + 4,0
	21	9 17 17,3	1	228 40 13,2	39 39,2	-	33,3	N	102 46 49,4 46 56,3 + 6,9
	28	14 53 7,7	2	319 14 54,7	14 58,5	+	3,8	N	105 17 25,2 17 30,3 + 5,1
	29	15 41 19,1	2	332 18 48,6	18 49,1	+	0,5	N	102 6 45,2 7 7,1 + 21,9
	30	16 29 8,1	2	345 17 12,0	17 15,7	+	3,7	N	98 16 20,4 16 34,9 + 14,5
July	2	18 5 42,2	2	11 27 53,0	11 51,3	-	4,7	N	89 15 55,7 16 5,7 + 10,0
	17	6 30 29,9	1	212 29 4,7	28 58,2	-	6,5	N	97 52 38,7 52 44,8 + 6,1
	18	7 14 11,3	1	224 25 27,3	25 21,4	-	5,9	N	101 29 7,9 29 12,0 + 4,1
	29	16 3 38,9	2	7 28 48,5	28 34,5	-	14,0	N	90 48 41,0 48 38,8 - 2,2
Aug.	17	7 25 8,9	1	256 44 56,1	44 39,0	-	17,1	N	108 6 38,5 6 41,6 + 3,1
	22	11 30 42,0	1	323 14 15,0	14 1,6	-	13,4	N	104 28 10,5 28 10,6 + 0,1
Sept.	15	6 53 13,1	1	277 19 54,6	19 42,4	-	12,2	N	109 37 8,5 37 8,0 - 0,5
	16	7 42 2,8	1	290 33 33,2	33 21,1	-	12,1	S	109 18 40,3 18 38,5 - 1,8
	17	8 31 24,9	1	303 55 16,7	55 3,7	-	13,0	S	108 1 26,0 1 30,9 + 4,9
	20	11 0 22,6	1	344 13 14,1	13 9,0	-	5,1	S	98 43 11,4 43 3,8 - 7,6
	21	11 51 30,6	1,2	357 43 4,6	43 8,9	+	4,3	S	94 13 8,6 13 6,2 - 2,4
	22	12 42 52,5	2	11 21 13,9	21 13,3	-	0,6	S	89 22 58,7 22 57,0 - 1,7

Date	Madras Mean Time.	Limb Observed.	Observed A.R. of D's Centre.	A.R. from Nautical Almanac.	Error of Tables.	Limb Observed.	Observed N.P.D. of D's Centre.	N.P.D. from Nautical Almanac.	Error of Tables.
1831	<i>h. m. s.</i>		<i>m. s.</i>	<i>m. s.</i>	<i>s.</i>		<i>° ' "</i>	<i>' "</i>	<i>"</i>
Oct.	14 6 22 17,7	1	298 9 49,7	9 36,3	- 13,4	S	108 53 10,1	53 7,9	- 2,2
	17 8 48 41,1	1	337 49 6,4	49 0,2	- 6,2	S	100 47 18,9	47 22,2	+ 3,3
	20 11 19 38,8	1	18 37 29,1	37 35,5	+ 6,4	S	86 46 43,1	46 38,9	- 4,2
	21					S	81 50 44,9		
	23 14 8 45,1	2	63 25 0,5	24 59,7	- 0,8				
	25 16 8 13,1	2	95 20 3,0	20 1,3	- 1,7	S	70 7 29,7	7 21,5	- 8,2
	29 19 49 19,6	2	154 43 47,5	43 47,5	- 0,0	S	78 13 16,6	13 23,9	+ 7,3
Nov.	19					S	75 23 25,0		
	21 13 53 33,6	2	88 11 6,0	10 59,9	- 6,1	N	70 15 48,3	15 33,2	- 15,1
	26 18 36 15,4	2	164 0 43,4	0 46,6	+ 3,2	S	80 45 37,5	45 43,9	+ 6,4
Dec.	11 5 20 46,9	1	339 54 15,9	54 11,1	- 4,8	S	100 41 7,4	41 8,0	+ 0,6
	12 6 7 17,2	1	352 32 11,8	32 4,9	- 6,9	S	96 33 36,7	33 30,5	- 6,2
	13 6 54 23,4	1	5 20 42,8	20 48,0	+ 5,2	S	91 58 50,3	58 34,7	- 15,6
	14 7 43 5,1	1	18 32 35,7	32 40,2	+ 4,5	S	87 7 51,1	7 51,5	+ 0,4
	15 8 34 15,4	1	32 21 48,9	21 44,7	- 4,2	S	82 15 59,4	15 49,8	- 9,6
	16 9 28 40,3	1	47 0 10,4	0 10,3	- 0,1	S	77 41 55,4	41 55,7	+ 0,3
	21 14 39 10,9	2	129 11 47,0	11 29,1	- 17,9	S	71 59 49,8	59 46,4	- 3,4
	24 17 19 34,2	2	172 23 9,9	23 7,9	- 2,0	S	83 18 14,4	18 4,0	- 10,4

In computing the above, the semi-diameter and Parallax have been taken from the Nautical Almanac; for reducing the latter the ratio of the Polar and Equatoreal Axes has been assumed 299 : 300 whence we have the *angle of the vertical* = 5*m.* Os. and the radius of the Earth = ,999825.

In the case of the Transit observations where the first or second limb of the Moon is observed, it is obvious that after applying the semi-diameter in A.R. we obtain the Right Ascension of the Moon's centre at the instant of the *first* or *second limb* transiting the Meridian; but with the Mural Circle we get the N.P.D. due to the moment of the Moon's *centre* transiting the Meridian to avoid therefore the inconvenience which would result from having the Right Ascension and North Polar Distance at two different instants of time, I have applied to the observed N.P.D. the change of the Moon's declination in the interval occupied by the Moon's semi-diameter to pass the Meridian. In a few cases however where the Moon was not observed with the Transit Instrument as well as when both limbs were observed, the N.P.D. set down is that for the moment of the Moon's centre transiting the Meridian.

Observation of the Eclipse of the Moon on the 26th February 1831.

	Sidereal time.	Mean time.	Observed by
	<i>h. m. s.</i>	<i>h. m. s.</i>	
Beginning of the Eclipse.....	7 11 45	8 48 50,3	T.
Shadow covers Gremaldus.....	7 21 20	8 58 23,7	T.

	Sidereal time.			Mean time.			Observed by
	<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>h.</i>	<i>m.</i>	<i>s.</i>	
Shadow touches Tycho.....	7	28	49	9	5	51,5	R.
	7	29	6	9	6	8,5	T.
Shadow covers Tycho.....	7	30	35	9	7	37,2	T.
	7	30	36	9	7	38,2	R.
Shadow touches Copernicus.....	7	47	31	9	24	30,4	R.
	7	47	35	9	24	34,4	T.
Shadow covers Copernicus.....	7	52	33	9	29	31,6	R.
	7	52	40	9	29	38,6	T.
Shadow touches Aristarcus.....	7	55	53	9	32	51,1	R.
	7	56	1	9	32	59,1	T.
Shadow covers Aristarcus.....	7	59	20	9	36	17,2	T.
Shadow touches Mare Vaporium.....	8	1	59	9	38	55,7	T.
Shadow touches Mare Christium.....	8	39	35	10	16	25,9	R.
	8	40	0	10	16	50,8	T.
Shadow leaves Gremaldus.....	9	4	55	10	41	11,9	T.
Shadow leaves Tycho.....	9	19	13	10	55	57,4	T.
	9	19	35	10	56	19,3	R.
Shadow leaves Mare Christium.....	9	24	45	11	1	28,5	T.
	9	25	35	11	2	13,3	R.
End of Eclipse.....	10	8	25	11	45	1,4	R.
	10	8	55	11	45	31,3	T.

The above observations by T were made with the 5 feet Achromatic, with a power of about 40, those by R with a 46 Inch Achromatic power about 60 or 70.

Eclipses of Jupiter's Satellites observed in 1831.

March 18—Immersion of Jupiter's first Satellite was lost in consequence of my having removed my eye from the Telescope to explain to my Assistant the position of the Satellite, which was very indistinct in consequence of a thin haze, and Jupiter's proximity to the horizon; I saw the Satellite at 16h. 51m. 54s. it had disappeared at 16h. 52m. 34s. I presume I am not 20s. in error in assuming it at 16h. 52m. 24s. Sidereal time or 17h. 9m. 15,7s. Mean time.

May 13—Immersion of Jupiter's first Satellite with 5 feet Achromatic power 150 at 17h. 12m. 57s. Sidereal time or 10h. 49m. 34,3s. Mean time.

Observations indifferent, in consequence of haze.

May 15—Emersion of Jupiter's third Satellite with 5 feet Achromatic

power 120 at 17*h.* 58*m.* 40*s.* Sidereal time or 14*h.* 17*m.* 19,8*s.*
Mean time.

Indifferent, in consequence of haze.

Two Native Assistants did not see the emersion till 1 minute after the above time which I presume arises from their inexperience in this species of observation.

May 23—Immersion of Jupiter's second Satellite with 5 feet Achromatic power 120 at 18*h.* 37*m.* 9*s.* Sidereal time or 14*h.* 34*m.* 14,3*s.*
Mean time.

Do. —Do. with 42 Inches Achromatic power 75 at 18*h.* 37*m.* 11*s.*
Sidereal time or 14*h.* 34*m.* 16,3*s.* Mean time.

June 4—Immersion of Jupiter's first Satellite with 5 feet Achromatic power 120 at 18*h.* 47*m.* 59*s.* Sidereal time or 13*h.* 57*m.* 50,6*s.*
Mean time.

Do. —Do. with 42 Inches Achromatic power 75 at 18*h.* 47*m.* 57*s.* Sidereal time or 13*h.* 57*m.* 48,6*s.* Mean time.

Clear, observation good.

Aug. 30—Emersion of Jupiter's first Satellite with 5 feet Achromatic power 120 at 20*h.* 8*m.* 54*s.* Sidereal time or 9*h.* 36*m.* 28,6*s.*
Mean time.

Aug. 30—Emersion of Jupiter's first Satellite with 42 Inches Achromatic power 75 at 20*h.* 8*m.* 48*s.* Sidereal time or 9*h.* 36*m.* 28,6*s.*
Mean time.

Clear, observation good.

Sept. 14—Emersion of Jupiter's third Satellite with 42 Inches Achromatic power 75 at 22*h.* 4*m.* 58*s.* Sidereal time or 14*h.* 33*m.* 15,0*s.*
Mean time.

Observation, satisfactory.

Do. —Emersion of Jupiter's second Satellite with 42 Inches Achromatic power 75 at 22*h.* 15*m.* 50*s.* Sidereal time or 10*h.* 44*m.* 5,2*s.*
Mean time.

Sept 21.—Emersion of Jupiter's second Satellite with 42 Inches Achromatic power 75 at 0h. 19m. 2s. Sidereal time or 13h. 19m. 25,2s. Mean time.

Thin haze, but observation satisfactory.

Sept. 22—Emersion of Jupiter's first Satellite with 5 feet Achromatic power 150 at 21h. 54m. 36s. Sidereal time or 9h. 51m. 26,9s. Mean time.

Clear Moon light, observation good.

Sept. 29—Immersion of Jupiter's first Satellite with 42 Inches Achromatic power 75 at 0h 18m. 20s. Sidereal time or 11h. 47m. 26,0s. Mean time.

Oct. 20—Emersion of Jupiter's third Satellite with 5 feet Achromatic power 300 at 20h. 34m. 28s. Sidereal time or 6h. 41m. 26,8s. Mean time.

Do. —Do. with 42 Inches Achromatic power 75 at 20h. 34m. 49s. Sidereal time or 6h. 41m. 47,8s. Mean time.

Nov. 10—Emersion of Jupiter's second Satellite with 5 feet Achromatic power 150 at 22h. 46m. 10s. Sidereal time or 7h. 30m. 22,8s. Mean time.

Do. —Do. with 42 Inches Achromatic power 75 at 22h. 46m. 10s. Sidereal time or 7h. 30m. 22,8s. Mean time.

Dec. 2—Emersion of Jupiter's third Satellite with 5 feet Achromatic power 150 at 23h. 36m. 45s. Sidereal time or 6h. 54m. 20,1s. Mean time.

Do. —Do. with 42 Inches Achromatic power 120 at 23h. 37m. 24s. Sidereal time or 6h. 54m. 59,1s. Mean time.

In the absence of the Greenwich observations we will now compare the above times with the Mean times given in the Nautical Almanac for the purpose of finding the Longitude ; accordingly we have the Longitude of Madras, as follows.

1831		1st Satellite.	2d Satellite.	3d Satellite.
		<i>h. m. s.</i>	<i>h. m. s.</i>	<i>h. m. s.</i>
May	13	5 21 44,3	_____	_____
	15	_____	_____	5 22 20,8
	23	_____	5 21 44,3	_____
June	4	20 40,6	_____	_____
July	13	21 6,7	_____	_____
August	30	20 59,6	_____	_____
September	14	_____	20 53,2	_____
	14	_____	_____	21 6,0
	21	_____	20 54,2	_____
	22	20 37,9	_____	_____
	29	20 57,0	_____	_____
October	20	_____	_____	21 24,3
November	10	_____	21 15,8	_____
December	2	_____	_____	23 49,6
Mean		5 21 1,0	5 21 11,9	5 22 10,2

Taking the Mean of the observations of the first and second Satellites, we have the Longitude of the Madras Observatory, *5h. 21m. 5.4s.* differing somewhat less than a mile from the value assigned by Mr. Goldingham. Whilst upon the subject of Longitude, it will be as well here to give the observed transits of the Moon, and of Stars culminating near thereto; the comparison of observations of this nature with the corresponding observations under other Meridians appears to be the best method yet employed for determining the difference of Longitude where the difference of Meridians is small; but in the present case, from the comparison of observations at Madras with those made in Europe results of second rate accuracy only must be expected.

		<i>h. m. s.</i>			<i>h. m. s.</i>
Feb.	20—20	Orionis..... 4 47 6,96	March	3—λ	Virginis.....14 11 6,93
	▷ 1	L..... 4 49 53,28		▷ 2	L.....14 48 2,74
	1 <i>y</i>	Orionis..... 4 55 9,47		<i>f</i> 1	Libræ.....15 26 5,56
	2 <i>y</i>	Orionis..... 5 0 16,29	<hr/>		
	21—▷ 1	L..... 5 50 52,90		5—β	Scorpii.... 15 56 46,54
	ν	Orionis..... 5 58 5,97		ν	Scorpii.... 16 3 19,94
	ξ	Orionis..... 6 2 30,44		▷ 2	L.....16 26 3,92
	η	Geminorum..... 6 4 51,25	<hr/>		
	22—γ	Geminorum..... 6 27 42,62		21—ν	Geminorum..... 6 18 24,19
	▷ 1	L..... 6 51 40,71		▷ 1	L..... 6 33 5,54
	23—▷ 1	L..... 7 51 39,56		ξ	Geminorum..... 6 53 33,85
	γ	Cancri..... 8 32 58,37	<hr/>		
	27—σ	Leonis.....11 11 26,88		22—K	Geminorum..... 7 23 26,70
	▷ 2	L.....11 32 36,99		▷ 1	L..... 7 33 11,11
	β	Virginis.....11 40 55,48	<hr/>		
	π	Virginis.....11 51 14,60		23—25	Cancri..... 8 15 44,87
				θ	Cancri..... 8 21 26,67
				▷ 1	L..... 8 31 20,10
				54	Cancri..... 8 41 5,91
				π ^a	Cancri..... 9 5 23,25

		<i>h.</i>	<i>m.</i>	<i>s.</i>
March	25— <i>b</i> ¹ Leonis.....	10	15	51,77
	<i>D</i> 1 L.....	10	20	29,04
	<i>ρ</i> Leonis.....	10	23	25,20
	<i>k</i> Leonis.....	10	36	48,40
<hr/>				
	26— <i>c</i> Leonis.....	10	51	32,05
	<i>D</i> 1 Limb.....	11	11	41,47
<hr/>				
	27— <i>ν</i> Virginis.....	11	36	41,86
	<i>π</i> Virginis.....	11	51	44,30
	<i>D</i> 1 Limb.....	12	1	12,11
	35 Virginis.....	12	38	46,98
<hr/>				
	29— <i>a</i> Virginis.....	13	15	50,37
	74 Virginis.....	13	22	43,54
	82 Virginis.....	13	32	17,19
	<i>D</i> 2 Limb.....	13	39	29,19
	<i>γ</i> Virginis.....	13	59	33,66
<hr/>				
	30— <i>λ</i> Virginis.....	14	9	30,13
	106 Virginis.....	14	19	18,96
	<i>D</i> 2 Limb.....	14	27	18,77
	<i>ρ</i> Libræ.....	14	36	11,06
	<i>μ</i> Libræ.....	14	39	35,22
	<i>β</i> Libræ.....	15	7	26,64
<hr/>				
April	2— <i>m</i> Scorpii.....	16	31	20,25
	<i>D</i> 2 Limb.....	16	55	0,54
	<i>η</i> Ophiuchi.....	17	0	13,21
<hr/>				
	20— <i>D</i> 1 Limb.....	9	11	1,72
	<i>ξ</i> Leonis.....	9	22	32,18
	<i>ο</i> Leonis.....	9	31	49,80
<hr/>				
	21— <i>D</i> 1 Limb.....	10	4	39,38
	<i>i</i> Leonis.....	10	22	40,81
	50 Leonis.....	10	29	21,90
	<i>ο</i> ¹ Sextants.....	10	36	48,84
<hr/>				
	22—48 Leonis.....	10	25	31,30
	34 Sextants.....	10	33	26,43
	<i>l</i> Leonis.....	10	39	53,98
	<i>D</i> 1 Limb.....	10	55	59,43
	<i>c</i> Leonis.....	11	14	39,40
<hr/>				
	23— <i>ξ</i> ¹ Virginis.....	11	36	7,82
	<i>D</i> 1 Limb.....	11	45	21,06
	<i>r</i> Virginis.....	12	0	34,89
	<i>n</i> Virginis.....	12	9	34,20
<hr/>				
	25— <i>h</i> ¹ Virginis.....	12	50	32,72
	<i>D</i> 1 Limb.....	13	20	54,92
	<i>m</i> Virginis.....	13	32	20,25
	<i>p</i> Virginis.....	13	45	37,15
<hr/>				
	26— <i>D</i> Center.....	14	9	23,83
	<i>μ</i> Virginis.....	14	33	45,58
<hr/>				
	27— <i>D</i> 2 Limb.....	14	58	23,57
	<i>β</i> Libræ.....	15	7	31,74
	<i>ζ</i> ¹ Libræ.....	15	18	20,69

		<i>h.</i>	<i>m.</i>	<i>s.</i>
April	29— <i>D</i> 2 Limb.....	16	37	5,52
	<i>ξ</i> Serpentis.....	17	27	32,42
<hr/>				
May	1— <i>μ</i> ¹ Sagittarii.....	18	3	18,33
	<i>D</i> 2 Limb.....	18	19	58,86
	28 Sagittarii.....	18	35	47,73
<hr/>				
	3— <i>D</i> 2 Limb.....	20	5	37,23
	<i>π</i> Capricorni.....	20	17	18,21
<hr/>				
	20— <i>D</i> 1 Limb.....	11	30	56,54
	<i>β</i> Virginis.....	11	41	46,89
	<i>A</i> Virginis.....	11	46	16,17
	<i>ο</i> Virginis.....	11	56	29,29
<hr/>				
	21— <i>r</i> Virginis.....	12	0	56,17
	<i>n</i> Virginis.....	12	9	55,10
	<i>D</i> 1 Limb.....	12	19	16,62
	35 Virginis.....	12	39	9,64
<hr/>				
	22— <i>k</i> ¹ Virginis.....	12	50	52,24
	<i>D</i> 1 Limb.....	13	6	34,91
	65 Virginis.....	13	14	28,94
	<i>m</i> Virginis.....	13	32	40,16
	<i>p</i> Virginis.....	13	45	57,08
<hr/>				
	23—88 Virginis.....	13	39	23,41
	<i>D</i> 1 Limb.....	13	53	37,98
	<i>λ</i> Virginis.....	14	9	54,29
<hr/>				
	24— <i>D</i> 1 Limb.....	14	41	2,50
	<i>δ</i> Libræ.....	14	51	52,64
	<i>ν</i> ¹ Libræ.....	14	57	8,50
	<i>f</i> ¹ Libræ.....	15	24	52,57
<hr/>				
	25— <i>γ</i> Libræ.....	15	26	0,46
	<i>D</i> 1 Limb.....	15	29	18,91
	<i>ξ</i> Libræ.....	15	55	0,84
	<i>ν</i> Scorpii.....	16	2	7,13
<hr/>				
	27— <i>η</i> Ophiuchi.....	17	0	37,99
	<i>D</i> 2 Limb.....	17	11	33,50
	<i>e</i> ¹ Ophiuchi.....	17	21	3,04
	<i>ξ</i> Serpentis.....	17	27	50,93
<hr/>				
	29— <i>D</i> 2 Limb.....	18	55	49,14
	<i>d</i> Sagittarii.....	19	7	41,87
	<i>e</i> ² Sagittarii.....	19	32	47,94
<hr/>				
	31— <i>a</i> ² Capricorni.....	20	8	37,85
	<i>π</i> Capricorni.....	20	17	35,20
	<i>ο</i> Capricorni.....	20	30	22,86
	<i>D</i> 2 Limb.....	20	41	22,95
	<i>η</i> Capricorni.....	20	54	43,86
	<i>ν</i> Aquarii.....	21	0	20,32
<hr/>				
June	1— <i>D</i> 2 Limb.....	21	33	54,65
	<i>μ</i> Capricorni.....	21	44	1,59
<hr/>				
	20— <i>D</i> 1 Limb.....	14	26	1,75
	<i>μ</i> Virginis.....	14	34	22,80

RESULTS FROM OBSERVATIONS, 1831.

		h. m. s.	
June	20—ξ ^a Libræ.....	14 47	49,85
	23—D 2 Limb.....	21 18	19,82
	e Capricorni.....	21 27	55,68
	γ Capricorni.....	21 31	2,16
	δ Capricorni.....	21 38	1,53
	29—D 2 Limb.....	22 10	36,40
	τ ¹ Aquarii.....	22 38	3,97
	λ Aquarii.....	22 43	7,52
July	17—D 1 L.....	14 9	26,94
	ξ ^a Libræ.....	14 48	9,27
	δ Libræ.....	14 52	29,84
	29—D 2 Limb.....	0 31	35,95
	m Ceti.....	0 45	2,89
	e Piscium.....	1 0	20,75
Sept.	15—D 1 Limb.....	18 29	28,26
	ν ¹ Sagittarii.....	18 45	11,78
	ξ ¹ Sagittarii.....	18 48	32,04
	17—D 1 Limb.....	20 15	51,82
	19 Capricorni.....	20 46	31,40
	21 Capricorni.....	20 52	37,64
	20—λ Aquarii.....	22 44	7,94
	δ ¹ Aquarii.....	22 53	56,69
	D 1 Limb.....	22 57	6,81
	21—n Piscium.....	23 40	36,60
	24 Piscium.....	23 45	36,08
	D Center.....	23 52	10,85
	22—33 Piscium.....	0 10	29,52
	t Piscium.....	0 18	7,02
	D 2 Limb.....	0 47	48,73
October	14—f Sagittarii.....	19 38	10,05

		h. m. s.	
October	14—57 Saigtarii.....	19 44	2,64
	D 1 Limb.....	19 53	14,24
	ρ Capricorni.....	20 20	53,12
	17—D 1 Limb.....	22 31	56,91
	λ Aquarii.....	22 45	34,12
	ψ ³ Aquarii.....	23 11	56,40
	23—γ Tauri.....	4 12	5,11
	D 2 Limb.....	4 16	39,81
	Aldebaran.....	4 28	8,30
	25—μ Geminorum.....	6 14	40,25
	ν Geminorum.....	6 20	51,52
	D 2 Limb.....	6 24	23,08
Nov.	21—ζ Tauri.....	5 30	3,99
	B Tauri.....	5 41	10,02
	χ ¹ Orionis.....	5 46	53,63
	D 2 Limb.....	5 56	22,50
Dec.	12—D 1 Limb.....	23 32	0,63
	249 Piscium.....	23 53	56,81
	13—t Piscium.....	0 19	42,56
	D 1 Limb.....	0 23	15,67
	m Ceti.....	0 47	20,84
	14—189 Piscium... ..	0 42	31,36
	27 Ceti.....	1 0	37,18
	D 1 Limb.....	1 16	3,72
	15—ξ ¹ Ceti.....	2 7	5,75
	D 1 Limb.....	2 11	20,99
	21—D 2 Limb.....	11 33	51,26
	ξ ¹ Virginis.....	11 39	52,12
	β Virginis.....	11 45	11,53
	π Virginis.....	11 54	30,48

Selecting from the above those of which corresponding observations have been made at the Greenwich Royal Observatory, we have

Greenwich Observations.				Madras Observations.			
1831		h. m. s.	t m. s.	h. m. s.	τ m. s.	Δ m. s.	
March	23 Moon 1 L.....	8 45 15,92		8 31 20,10			
	π ^a Cancræ.....	9 6 40,57	— 21 24,65	9 5 23,25	— 34 3,15	— 12 38,50	
May	25 γ Libræ.....	15 27 24,72	+ 14 12,78	15 26 0,46	+ 3 18,45	10 54,33	
	Moon 1 L.....	15 41 37,50		15 29 18,91			
June	20 Moon 1 L.....	14 36 51,54		14 26 1,75			
	ξ ^a Libræ.....	14 48 5,04	— 11 13,50	14 47 49,85	— 21 48,10	10 34,60	
October	23 Aldebaran.....	4 27 3,40	+ 2 35,58	4 28 8,30	— 11 28,49	14 4,07	
	Moon 2 L.....	4 29 38,98		4 16 39,81			
Dec.	12 Moon 1 L.....	23 41 15,68		23 32 0,63			
	+ 1,1 249 Piscium.....	23 51 52,68	— 10 37,00	23 53 56,81	— 21 56,18	11 19,18	

In a similar manner we have:

Cambridge Observations.				Madra: Observations.				
1831		<i>h. m. s.</i>	<i>t</i> <i>m. s.</i>		<i>s.</i>	<i>h. m. s.</i>	<i>τ</i> <i>m. s.</i>	<i>t - τ</i> <i>m. s.</i>
Mar. 23	Moon 1 Limb.	8 42 56,32				8 31 20,10		
	π Cancri.....	9 4 22,02	- 21 25,69	+ 0,10		9 5 23,25	- 34 3,15	- 12 37,46
May 25	γ Libræ.....	15 24 0,67	+ 14 11,86	+ 0,13		15 26 0,46	- 3 18,45	10 53,41
	Moon 1 Limb.	15 38 12,54				15 29 18,91		
June 20	Moon 1 Limb.	14 34 3,57				14 26 1,75		
	ξ^a Libræ....	14 45 17,92	- 11 14,35	+ 0,73		14 47 49,85	- 21 48,10	10 33,75
Sept. 22	β Piscium...	0 6 6,55	+ 49 34,83			0 10 29,52	- 37 19,18	12 15,65
	t Piscium....	0 13 41,17	+ 41 57,21	+ 0,10		0 18 7,02	- 29 41,69	12 15,52
	Moon 2 Limb.	0 55 41,38				0 47 48,73		
Oct. 17	Moon 1 Limb.	22 38 53,30				22 31 56,91		
	χ^3 Aquarii...	+23 7 2,29	- 28 8,99	+ 0,98		23 11 56,40	- 39 59,46	11 50,47
Dec. 14	189 Piscium..	0 36 26,95	+ 45 35,38	+ 1,64		0 42 31,36	- 33 32,32	12 3,06
	Moon 1 Limb.	1 22 2,33				1 16 3,72		

The above observations at Greenwich and Cambridge are extracted from the Monthly reports of the proceedings of the Royal Astronomical Society; the former are deficient in not having the rate of the Clock annexed, but from the custom at Greenwich of keeping the daily rate of the Clock within one second, its omission altogether will not introduce an error exceeding 0,6s. of Longitude. In computing the Longitude from these observations I have assumed $x = - 5h. 21m.$ and computed the values of e^* in preference to computing the horary motion for the middle of the times of passage, hence we find the Longitude of the Madras Observatory, from

1831	Greenwich Observations.		Cambridge Observations.	
	Δ 1 L. <i>h. m. s.</i>	Δ 2 L. <i>h. m. s.</i>	Δ 1 L. <i>h. m. s.</i>	Δ 2 L. <i>h. m. s.</i>
March.....23	5 21 14,7	—————	5 20 48,4	—————
May.....25	20 54,2	—————	20 26,2	—————
June.....20	20 42,7	—————	20 16,1	—————
September.....22	—————	—————	—————	5 20 52,5
October.....23	—————	5 21 14,6	—————	—————
December.....12	20 35,7	—————	—————	—————
December.....14	—————	—————	20 22,4	—————

† This is not χ^3 Aquarii probably ψ^3 Aquarii.

* See Mr. Baily's very excellent paper on this subject, inserted in the 2d Volume of the Memoirs of the Astronomical Society.

I have remarked at Page 69, that an *irradiation* is found to exist with every observer, as well as with the Telescope with which he observes; to obviate the error which would thus be introduced into the determination of Longitude by Lunar Observations, it is necessary to employ an equal number of observations of *each* limb, or which is the same thing, we must take the mean of the results from observations of the *first* and *second* limbs as being two independent results, without reference to the number of observations made of each, thus we obtain the Longitude of Madras, from

Greenwich.	Cambridge.
<i>h. m. s.</i>	<i>h. m. s.</i>
5 21 3,2	5 20 40,8

Mr. Airy states the Longitude of the Cambridge Observatory, to be 23,54s. East of Greenwich, hence we have the Longitude of the Madras Observatory.

5 21 3,2	East of Greenwich	} by Greenwich Observations.
5 21 4,34	—————	

LATITUDE OF THE MADRAS OBSERVATORY.

To determine the Latitude, observations have been made with the Mural Circle, of the North Polar Distance of the image of Stars, as reflected from a trough of quicksilver; the Stars selected for this purpose are those which are situated within 30° of the Zenith, these being reduced to the mean places at the beginning of the year, and compared with the mean places, similarly deduced from the Observations by direct vision, give the following.

NAMES.	No. of Observations.	N.P.D. by Direct Vision.			Extreme Difference.	No. of Observations.	N.P.D. by Reflection.			Extreme Difference.	Latitude.		
		°	'	"			°	'	"		°	'	"
β Geminorum.....	33	61	34	21,66	4,24	7	272	17	20,12	5,06	13	4	9,11
γ Cancrī.....	6	67	55	42,71	4,58	8	265	55	56,20	3,60	13	4	10,54
ρ Lyncis.....	5	52	29	15,74	1,69	8	281	22	25,29	3,70	13	4	9,48
κ Leonis.....	—	not observed				5	270	45	59,35	2,27			
λ Leonis.....	2	66	17	29,26	0,92	8	267	34	11,67	4,04	13	4	9,53
b Leonis Minor.....	5	52	51	23,35	2,03	8	281	0	14,19	4,05	13	4	11,23
e Leonis.....	9	65	27	8,06	1,99	10	268	24	32,57	3,21	13	4	9,68
μ Leonis.....	5	63	12	2,65	0,60	5	270	39	39,84	5,86	13	4	8,75
d Leonis Minor.....	3	53	56	7,26	1,22	4	279	55	38,63	1,42	13	4	7,05

NAMES.	No. of Observations.	N.P.D. by Direct Vision.			Extreme Difference.	No. of Observations.	N.P.D. by Reflection.			Extreme Difference.	Latitude.		
		°	'	"			°	'	"		°	'	"
ζ Leonis.....	6	65	44	38,30	2,11	5	268	7	3,86	2,37	13	4	8,92
μ Ursæ Major.....	5	47	39	13,35	3,60	3	286	12	26,75	3,84	13	4	9,95
f Leonis Minor.....	4	55	20	46,10	5,38	5	278	30	53,08	2,77	13	4	10,41
g Leonis Minor.....	5	52	25	47,77	3,15	3	281	25	53,35	4,30	13	4	9,44
l Leonis Minor.....	5	57	8	56,13	2,52	7	276	42	46,47	4,47	13	4	8,70
n Leonis Minor.....	6	58	25	43,96	4,13	5	275	25	56,36	2,92	13	4	9,84
o Leonis Minor.....	5	54	52	40,56	5,10	7	278	59	5,63	2,37	13	4	6,91
54 Leonis Minor.....	6	54	21	4,50	2,00	5	269	30	39,25	2,37	13	4	8,12
ξ Ursæ Major.....	4	57	31	13,81	4,05	8	276	20	27,39	4,83	13	4	9,40
ν Ursæ Major.....	1	55	59	6,74	0,00	7	277	52	32,98	3,19	13	4	10,14
h Comæ Ber.....	—	not observed				5	268	44	48,27	3,64			
e Comæ Ber.....	2	62	12	53,77	2,92	5	270	38	48,75	2,49	13	4	8,74
f Comæ Ber.....	5	62	57	44,41	2,28	5	270	53	52,78	0,89	13	4	11,20
c Comæ Ber.....	—	not observed				5	273	4	10,63	2,31			
a Comæ Ber.....	5	62	14	13,40	2,51	2	271	37	27,14	3,97	13	4	9,73
d Canum Ven.....	—	not observed				6	286	8	19,11	1,06			
k Comæ Ber.....	1	66	26	17,43	0,00	5	267	25	23,83	3,87	13	4	9,37
q Comæ Ber.....	2	67	50	0,43	3,81	5	266	1	40,90	3,21	13	4	9,33
a Canum Ven.....	9	50	46	3,20	3,99	7	283	5	36,23	4,76	13	4	10,28
h Canum Ven.....	2	48	31	46,48	1,14	7	285	19	36,82	3,09	13	4	8,85
l Bootis.....	—	not observed				3	272	10	50,45	3,42			
γ Bootis.....	4	50	56	56,70	1,52	6	282	54	44,31	3,57	13	4	9,45
σ Bootis.....	5	59	31	3,40	4,80	5	274	19	40,92	2,81	13	4	7,84
34 Bootis.....	5	62	44	58,41	6,23	5	271	6	44,95	2,14	13	4	8,32
e Bootis.....	—	not observed				3	271	39	10,07	3,67			
β Bootis.....	5	48	56	19,00	3,93	5	284	55	27,33	2,01	13	4	6,83
δ Bootis.....	—	not observed				5	277	48	40,44	3,36			

Taking the Mean; from 160 Observations by direct vision combined with 171 by reflection we have the Latitude of the Madras Observatory, $13^{\circ} 4' 9", 21$.

OBSERVATIONS OF THE COMET OF JANUARY 1831.

In the foregoing statements I have endeavoured to represent as nearly as the case would permit the degree of accuracy attained in each particular species of Observation, but in the present case, the Observations of an ill defined object, made with a Telescope supported upon a wooden stand, and that too in the open air; subject to flexure from its own weight, and to tremor from every breath of air which may happen to blow, render it desirable that the whole of the particulars of each observation should be stated, accordingly the following is copied from the book "Miscellaneous Observations."

1831, 7th January at 4h. 50m. A. M. Saw a Comet towards the East about 20 degrees high but approaching twilight prevented observation.

8th January, 5h. A. M. Adjusted the five feet Achromatic by Dollond as an Equatoreal, saw the Comet with a power of 60 but it was too faint to allow the field being illuminated, the following observations were made, at the time of its occupying the centre of the field of view.

	Sidereal Time. h. m. s.	Horary Circle. h. m. s.	Declination Circle. ° ' "	REMARKS.
Comet.....	12 52 0	4 36 0	11 50 S.	Very faint, tail 4° long observations not to be depended upon to 5m.
	13 2 0	4 25 30	11 43 —	
Antares.....	12 58 0	3 19 0	—————	
	13 5 40	3 12 0	25 15 S.	

9th January at 5h. A. M.

	Sidereal Time. h. m. s.	Horary Circle. h. m. s.	Declination Circle. ° ' "	REMARKS.
Comet.....	12 22 41	6 58 45	10 52 10 S.	The Comet appeared very distinct notwithstanding its being situated within 30° of the Moon.
	12 24 54	7 0 50	10 52 30 —	
	12 25 59	7 1 58		
	12 26 48	7 2 50		
γ Serpentis.....	12 31 53	7 32 0	14 33 30 S.	The wires did not require illumination.
	12 34 21	7 34 20		
	12 35 50	7 35 55	14 33 20 —	

1831, January 10.—In consequence of the difficulty attending the adjustment of the Instrument as an Equatoreal I have availed myself of a suspension spirit level which belongs to the Telescope, to adjust it as an Altitude and Azimuth Instrument; the error of adjustment of the vertical Axis cannot I imagine exceed 20 or 30 seconds.

	Sidereal Time. h. m. s.	Altitude. ° ' "	Azimuth from North Meridian. h. m. s.	Azimuth from South. ° ' "
1831				
10th January.....	12 11 14	9 25 30	6 44 10	78 57 30
	12 14 30	10 9 0	6 45 5	78 43 45
	12 17 27	10 53 0	6 45 50	78 30 00
Comet.....	12 19 44	11 24 30	6 46 35	78 21 15
	12 21 18	11 46 10	6 46 55	78 16 15
	12 23 52	12 24 0	6 47 45	78 3 45

RESULTS FROM OBSERVATIONS, 1831.

	Sidereal Time.	Altitude.	Azimuth from North Meridian.	Azimuth from South.
1831	<i>h. m. s.</i>	<i>° ' "</i>	<i>h. m. s.</i>	<i>° ' "</i>
10th January.....	12 33 18	11 38 40	6 52 6	76 58 30
γ Ophiuchi.....	12 41 54	14 47 00	7 3 30	74 7 30
	12 45 21	17 25 0	6 54 5	76 28 45
Comet.....	12 52 47	19 11 0	6 56 35	75 51 15
	12 54 46	19 36 0	6 57 5	75 43 45
	12 56 25	19 59 0	6 57 45	75 33 45
	12 59 56	23 56 0	7 22 45	69 18 45
γ Ophiuchi.....	13 1 54	24 22 30	7 23 30	69 7 30
	13 3 37	24 45 0	7 24 0	69 0 00
	13 6 24	25 23 30	7 25 20	68 40 00
11th January... Clouds prevented any Observation.				
12th January.....	12 22 7	14 6 0	7 8 20	72 55 00
Comet.....	12 28 55	15 44 0	7 10 25	72 23 45
	12 43 47	19 9 0	7 15 5	71 13 45
	12 45 32	20 44 30	7 38 12	65 27 0
γ Ophiuchi.....	12 59 23	23 50 30	7 43 45	64 3 45
	13 1 23	24 16 30	7 44 15	63 56 15
	13 13 12	26 55 0	7 49 35	62 36 15
13th January.....	12 56 41	23 11 30	7 13 55	71 31 15
	13 0 51	24 10 30	7 15 30	71 7 30
	13 3 56	24 53 30	7 16 25	70 53 45
Comet.....	13 6 13	25 24 40	7 17 20	70 40 00
	13 10 0	26 16 20	7 18 50	70 32 30
13th January.....	13 15 20	27 24 0	7 44 25	63 53 45
	13 17 16	27 48 0	7 45 25	63 38 45
γ Ophiuchi.....	13 18 29	28 5 0	7 45 55	63 31 15
	13 20 1	28 25 0	7 46 45	63 18 45
	13 21 43	28 47 0	7 47 35	63 6 15
14th January.....	12 44 1	21 17 30	7 14 10	71 27 30
	12 47 19	22 3 30	7 15 20	71 10 00
	12 49 11	22 30 0	7 16 20	70 55 00
Comet.....	12 52 24	23 15 30	7 17 0	70 45 00
	12 55 20	23 56 30	7 18 10	70 27 30
	12 58 26	23 39 0	7 42 0	64 30 0
Ophiuchi.....	13 1 0	24 14 30	7 43 0	64 15 0
Flying Clouds but observations good.				

	Sidereal Time.	Altitude.	Azimuth from North Meridian.	Azimuth from South.
1831	<i>h. m. s.</i>	<i>° ' "</i>	<i>h. m. s.</i>	<i>° ' "</i>
15th January.....	12 45 48	22 45 0	7 19 30	70 7 30
	12 47 59	23 9 0	7 20 0	70 0 00
	12 48 30	23 22 40	7 20 15	69 56 15
Comet.....	12 49 33	23 37 0	7 20 50	69 47 30
	12 50 42	23 54 0	7 21 5	69 43 45
	12 53 9	22 30 0	7 44 10	63 57 30
	12 54 0	22 42 0	7 44 35	63 51 15
γ Ophiuchi.....	12 55 6	22 56 0	7 44 55	63 46 15
	12 56 8	23 10 0	7 45 30	63 37 30
	12 57 12	23 23 30	7 45 55	63 31 15
16th January.....	12 28 17	19 43 30	7 12 50	71 47 30
	12 29 40	19 43 1	7 13 15	71 41 15
Comet.....	12 30 55	20 20 0	7 13 45	71 33 45
	12 32 14	20 38 0	7 14 10	71 27 30
	12 33 53	21 3 0	7 14 50	71 17 30
	12 37 16	18 58 0	7 37 35	65 36 15
	12 38 10	19 9 0	7 37 53	65 31 15
γ Ophiuchi.....	12 39 10	19 23 0	7 38 20	65 25 00
	12 40 0	19 36 0	7 38 40	65 20 00
	12 41 23	19 53 30	7 39 10	65 12 30
19th January.....	12 39 4	25 29 0	6 50 20	77 25 00
	12 41 39	26 6 0	6 51 10	77 12 30
	12 44 28	26 47 0	6 52 10	76 57 30
Comet.....	12 48 17	27 38 0	6 53 20	76 40 00
	12 50 39	28 16 0	6 54 0	76 30 00
	12 53 32	32 6 0	7 8 30	72 52 30
	12 55 27	32 29 30	7 9 20	72 40 00
ζ Ophiuchi.....	12 57 9	32 53 0	7 10 10	72 27 30
	12 58 41	33 17 0	7 11 0	72 15 00
	13 0 28	33 35 0	7 11 40	72 5 00
20th January.....	12 55 11	30 24 0	7 7 40	73 5 00
	12 57 37	30 57 30	7 8 35	72 51 15
Comet.....	12 59 28	31 23 0	7 9 20	72 40 00
	13 1 3	31 44 30	7 10 10	72 27 30
	13 3 47	32 22 0	7 11 20	72 10 00
	13 6 24	34 57 0	7 25 35	68 36 15
ζ Ophiuchi.....	13 7 24	35 12 0	7 26 0	68 30 00
	13 8 24	35 24 30	7 26 30	68 22 30

	Sidereal Time.	Altitude.	Azimuth from North Meridian.	Azimuth from South.
1831	<i>h. m. s.</i>	<i>° ' "</i>	<i>h. m. s.</i>	<i>° ' "</i>
20th January.....	13 9 28	35 38 30.	7 27 0	68 15 00
	13 10 32	35 54. 0	7 27 40	68 5 00.
23d January....	13 16 0	38.43.30.	7 31 5	67 13 45
	13 18 10	39 12 0.	7 32 30	66 55 00
	13 19 43	39 33 30.	7 33 0	66 45 00
Comet.....	13 21 2	39.52 0	7 33 50	66 32 30
	13 22 39.	40 14 30.	7 34 45	66 18 45
ζ Ophiuchi.....	13 26 33	39 28 30	7 48 50	62 47 30
	13 28 14	39 53.30	7 49.35	62 36 15
	13 29 50	40 12. 0	7 50 40	62 20 00
	13 31 27.	40 34.30	7 51 45	62 3 45
	13 33 6	40 55.30.	7 52 25	61 53 45

It is necessary I should here remark that the Instrument was removed every day after the observations were made, to the inside of the Observatory and brought out again early in the morning for adjustment previously to the above observations being made, in performing the adjustment, no pains was taken to adjust the Azimuth Circle, which will account for the changes which are found from day to day in the Index Error. The Sidereal time set down is the *true* Sidereal time, found every morning by the Transit of Spica Virginis over the wires of the Mural Circle, the Transit Instrument not having been erected at this time. Employing the Sidereal time in conjunction with the Apparent Places of η and ζ Ophiuchi computed from the Astronomical Society's Tables we obtain the *true* Altitude and Azimuth, which being compared with the *observed* (the Altitude being corrected for refraction) gives the Index Error in Altitude and Azimuth which we can now apply to the observed Altitude and Azimuth of the Comet as follows.

1831	Sidereal Time.	Observed Altitude.	Refraction.	Index Error.	True Altitude.	Observed Azimuth.	Index Error.	True Azimuth.
	<i>h. m. s.</i>	<i>° ' "</i>	<i>' "</i>	<i>' "</i>	<i>° ' "</i>	<i>° ' "</i>	<i>' "</i>	<i>° ' "</i>
January	10 12 17 59	11 0 22	- 4 42	+ 3 56	10 59 36	78 28 46	- 3 8 16	75 20 30
	12 12 52 20	19 2. 45	- 2. 42	+ 3 56	19 3 59	75 54. 21	- 3 8 16	72 46 5
	12 12 25 31	14 55 00	- 3 30	+ 2 43	14 54 13	72 39 23	+ 2 10 30	74 49 53
	12 12 43 57	19 9 00	- 2 42	+ 2 43	19 9 1	71 13 45	+ 2 10 30	73 24 15
	13 13 3 32	24 47 18	- 2 2	+ 0 44	24 46 0	70 57. 0	+ 0 43 12	71 40 12
	14 12 49 39	22 36 36	- 2 17	+ 0 43	22 35 2	70 57 0	+ 1 50 48	72 47 48
	15 12 48 24	23 21 32	- 2 9	- 0 53	23 18 30	69 55 0	+ 2 55 42	72 50 42
	16 12 31 0	20 17 30	- 2 32	- 1 55	20 13 3	71 33 30	+ 2 43 52	74 17 22
	19 12 44 49	26 51 12	- 1 51	+ 1 20	26 50 41	76 57 0	- 3 58 57	72 58 03
	21 12 59 25	31 22 12	- 1 32	- 5 11	31 15 29	72 38 45	- 1 10 0	71 28 45
	23 13 19 31	39 31 6	- 1 9	- 7 53	39 22. 4	66 45 0	+ 1 59. 47	68 44 47

From the true Altitudes and Azimuths we obtain as follows.

1831	Mean Time	A.R.	N.P.D.	REMARKS.
	<i>h. m. s.</i>	<i>h. m. s.</i>	<i>• ′ ″</i>	
January..... 8	17 45 47,9	17 29 27,0	102 34 10	
9	17 10 2,8	17 24 44,0	101 49 15	
10	16 59 1,7	17 20 49,0	101 28 12	
10	17 33 17,0	17 20 41,0	101 28 15	
12	16 58 40,7	17 12 30,0	100 50 41	
12	17 17 3,6	17 12 37,5	100 52 25	
13	17 32 39,5	17 8 35,0	100 34 12	
14	17 14 52,8	17 4 28,0	100 19 17	
15	17 9 42,3	17 0 29,5	100 2 36	
16	16 48 25,3	16 56 43,3	99 45 4	
19	16 50 24,0	16 43 43,5	98 46 1	
20	17 1 1,9	16 39 33,0	98 27 47	
23	17 9 16,8	16 25 57,0	97 27 39	
February.....19	14 52 48,0	12 49 7,0	80 7 53,7	Very faint.
20	14 38 36,5	12 38 49,9	79 23 52,6	do. do.

The prevalence of haze and the presence of the Moon, added to the diminished brightness of the Comet, prevented observation after the 23d of January till the 19th February, on the latter day as well as on the 20th I was fortunate enough to obtain meridional observations with the Transit Instrument and Mural Circle, but these being made without illuminating the wires, in consequence of the extreme faintness of the Comet, cannot be depended upon to 1 or 2 minutes of space.

PLACES OF THE FIXED STARS.

I had originally intended to complete the Catalogue of 2881 Stars, before submitting to the notice of Astronomers any determination of the places of the fixed Stars, but finding on comparison with the Greenwich and Astronomical Society's Catalogue, discordances greater than can be reasonably accounted for from errors of observation, I have thought it advisable at once to offer the places of that portion of the Catalogue which is determined from the observations of 1831, together with the comparisons above named; I have added a column "Extreme difference" which with the number of observations will assist in forming an estimate of the probable error to which each result is liable; the annual variations, (those which have been employed) are copied from the Astronomical Society's Catalogue.

Right Ascension of Stars from the Madras Catalogue compared with the Greenwich, and Astronomical Society's Catalogue.

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.		Difference of Ex-tremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
				h. m. s.	s.				Greenwich...	A.S.	
1	2,3	β Cassiopeæ....	6	0 0 12,43	0,18	0 12,59	0 11,60	-0,16	+ 0,83	+ 3,124	
2	4	ε Phœnicis....	6	0 0 48,75	0,73		0 48,44		+ 0,31	3,063	
3	5	β Andromedæ..	5	0 1 33,96	0,32		1 33,91		+ 0,05	3,077	
4	5,6	ω ² App. Sculp...	6	0 2 58,71	0,78		2 58,59		+ 0,12	3,059	
5	2,3	γ Pegasi.....	12	0 4 32,59	0,46	4 32,63	4 32,43	-0,04	+ 0,16	3,075	
6	5	θ Andromedæ..	6	0 8 17,16	0,29		8 16,37		+ 0,79	3,105	
7	6,7	33 Piscium.....	6	0 9 7,24	0,35		9 6,82		+ 0,42	3,069	
8	4	ι Ceti.....	6	0 10 49,08	0,16	10 49,09	10 48,70	-0,01	+ 0,38	3,057	
9	5	ζ Toucanæ....	6	0 11 12,88	0,93		11 9,90		+ 2,98	2,923	
10	5,6	δ Piscium.....	4	0 11 54,39	0,13	11 54,52	11 54,75	-0,13	- 0,36	3,077	
11	6	9 Ceti.....	6	0 14 12,05	0,30		14 12,17		- 0,12	3,049	
12	6	ε Piscium.....	7	0 16 44,56	0,47		16 44,12		+ 0,44	3,070	
13	5	κ Phœnicis....	5	0 17 52,64	0,24		17 51,33		+ 1,31	2,966	
14	2	α Phœnicis....	6	0 17 54,89	1,08		17 54,24		+ 0,65	2,970	
15	6	48 Piscium.....	6	0 19 26,73	0,53		19 25,86		+ 0,87	3,099	
16	5	λ Cassiopeæ....	6	0 22 29,61	0,34		22 28,75		+ 0,86	3,245	
17	5	λ ¹ Phœnicis....	6	0 23 14,73	0,23		23 13,77		+ 0,06	2,909	
18	4	κ Cassiopeæ....	5	0 23 27,57	0,50	23 27,47	23 26,47	+ 0,10	+ 1,10	3,324	
19	4	β ¹ Toucanæ....	6	0 23 45,57	0,77		23 45,63		- 0,06	2,786	
20	4	ζ Cassiopeæ....	7	0 27 35,99	0,25	27 35,90	27 35,58	+ 0,09	+ 0,41	3,280	
21	4,5	π Andromedæ..	6	0 27 52,52	0,44	27 52,54	27 52,60	-0,02	- 0,08	3,172	
22	6	53 Piscium.....	5	0 27 59,75	0,18		27 59,05		+ 0,70	3,109	
23	4	ε Andromedæ..	6	0 29 38,57	0,14	29 38,65	29 38,08	-0,08	+ 0,49	3,161	
24	3	δ Andromedæ..	6	0 30 18,57	0,44	30 18,64	30 18,15	-0,07	+ 0,42	3,169	
25	3	α Cassiopeæ....	12	0 30 58,09	0,54	30 57,93	30 57,43	+ 0,16	+ 0,66	2,330	
26	6,7	* Ceti.....	5	0 32 6,33	0,42		32 5,84		+ 0,49	3,051	
27	5	μ Phœnicis....	6	0 33 19,44	0,23		33 19,42		+ 0,02	2,861	
28	5	π Cassiopeæ....	6	0 34 9,11	0,33		34 8,30		+ 0,81	3,274	
29	2,3	β Ceti.....	7	0 35 6,18	0,35	35 6,27	35 6,23	-0,09	+ 0,05	2,998	
30	5	φ ¹ Ceti.....	5	0 35 39,79	0,49		35 39,61		+ 0,18	3,026	
31	5	η Phœnicis....	4	0 35 44,13	0,44		35 43,07		+ 1,06	2,731	
32	4	ζ Andromedæ..	6	0 38 23,78	0,31	38 23,85	38 23,84	-0,07	- 0,06	3,164	
33	4	η Cassiopeæ....	4	0 38 55,79	0,10	38 55,63	38 54,73	+ 0,16	+ 1,06	3,533	
34	5	δ Piscium.....	6	0 39 55,12	0,36		39 55,20		- 0,08	2,095	
35	4	ν Andromedæ..	5	0 40 31,52	0,17	40 31,57	40 30,99	-0,05	+ 0,53	3,266	
36	6	ι Piscium.....	6	0 40 49,63	0,53		40 48,96		+ 0,67	3,187	
37	5	m Ceti.....	7	0 44 22,58	0,38	44 22,62	44 22,66	-0,04	- 0,08	3,059	
38	6	36 Andromedæ..	6	0 45 56,06	0,43		45 55,76		+ 0,30	3,179	
39	3	γ Cassiopeæ....	5	0 46 34,32	0,25	46 34,10	46 33,26	+ 0,22	+ 1,06	3,531	
40	4	μ Andromedæ..	5	0 47 24,06	0,31	47 24,11	47 23,81	-0,05	+ 0,25	3,359	
41	5	η Andromedæ..	6	0 48 12,02	0,19		48 11,16		+ 0,86	3,183	
42	5	α App. Sculp...	6	0 50 27,41	0,30		50 27,46		- 0,05	2,898	
43	7	π Piscium.....	3	0 51 4,62	0,12		51 3,94		+ 0,68	3,097	

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
				h.	m.	s.				Greenwich.	A.S.	
44	6,7	Piscium.....	5	0 53 42,06	0,22		53 42,10		+ 0,56	+ 3,111		
45	4	^e Piscium.....	6	0 54 10,85	0,20	54 10,85	54 10,90	- 0,00	- 0,05	3,106		
46	6	27 Ceti.....	4	0 57 9,06	0,63		57 8,76		+ 0,30	3,005		
47	6	28 Ceti.....	8	0 57 36,56	0,77		57 36,48		+ 0,08	3,005		
48	3,4	^β Phœnicis....	6	0 58 31,96	0,37		58 31,80		+ 0,16	2,698		
49	6,7	88 Piscium.....	6	1 5 55,46	0,45		5 54,87		+ 0,59	3,108		
50	6	38 Ceti.....	7	1 6 11,86	0,23		6 11,65		+ 0,21	3,056		
51	5,6	^v Piscium.....	6	1 10 11,62	0,37		10 11,55		+ 0,07	3,268		
52	5	^ξ Andromedæ..	6	1 12 25,62	0,39		12 25,06		+ 0,56	3,478		
53	4,5	^ψ Cassiopeæ....	6	1 14 6,22	0,43	14 6,37	14 5,35	- 0,15	+ 0,87	4,079		
54	5	Phœnicis....	6	1 17 11,68	0,30		17 11,62		+ 0,06	2,665		
55	5	^c Ceti.....	6	1 17 18,68	0,19		17 18,46		+ 0,22	2,946		
56	3	^γ Phœnicis....	6	1 21 1,24	0,37		21 0,58		+ 0,66	2,619		
57	5	^μ Piscium.....	5	1 21 20,41	0,51	21 20,21	21 19,98	+ 0,20	+ 0,43	3,111		
58	4	^η Piscium.....	6	1 22 27,27	0,15	22 27,16	22 27,08	+ 0,11	+ 0,19	3,189		
59	4	^δ Phœnicis....	5	1 24 12,50	0,14		24 12,20		+ 0,30	2,497		
60	5	^v Andromedæ..	6	1 26 54,68	0,27		26 54,01		+ 0,67	3,491		
61	3,4	^R Andromedæ..	6	1 27 39,73	0,47	27 39,66	27 39,32	+ 0,07	+ 0,41	3,617		
62	6	^τ Piscium.....	4	1 28 9,02	0,12	28 9,01	28 8,73	+ 0,01	+ 0,29	3,168		
63	1	^α Eridani.....	5	1 31 24,96	0,22		31 24,27		+ 0,69	2,235		
64	5	^v Piscium.....	6	1 32 38,46	0,21	32 38,75	32 38,49	- 0,29	- 0,03	3,111		
65	5	54 Andromedæ..	5	1 33 6,84	0,21		33 6,66		+ 0,18	3,693		
66	5,6	107 Piscium....	4	1 33 20,16	0,30		33 19,94		+ 0,22	3,255		
67	3,4	^τ Ceti.....	5	1 36 13,20	0,30	36 13,23	36 12,86	- 0,03	+ 0,34	2,779		
68	5	^o Piscium.....	6	1 36 28,70	0,40	36 28,67	36 28,55	+ 0,03	+ 0,15	3,148		
69	5	^e App. Sculp....	6	1 37 43,98	0,33		37 43,21		+ 0,77	2,800		
70	5	^χ Ceti.....	5	1 41 17,35	0,61		41 16,45		+ 0,90	2,952		
71	3,4	^e Cassiopeæ....	6	1 42 19,92	0,33	49 19,30	42 19,28	+ 0,62	+ 0,64	4,191		
72	3	^ξ Ceti.....	6	1 43 7,30	0,20	43 7,40	43 6,86	- 0,10	+ 0,44	2,953		
73	3,4	^α Trianguli....	6	1 43 28,20	0,31	43 28,13	43 27,84	+ 0,07	+ 0,36	3,388		
74	4,5	^γ Arietis.....	5	1 44 16,30	0,36	44 16,32	44 16,24	- 0,02	+ 0,06	3,264		
75	5	^φ Phœnicis....	6	1 47 21,28	0,36		47 21,27		+ 0,01	2,499		
76	5	48 Cassiopeæ....	4	1 48 13,41	0,28		48 13,42		- 0,01	4,744		
77	4,5	50 Cassiopeæ....	4	1 49 10,30	0,16	49 10,55	49 9,32	- 0,25	+ 0,98	4,908		
78	4	^χ Eridani.....	6	1 49 22,85	0,55		49 21,59		+ 1,26	2,270		
79	4,5	^v Ceti.....	6	1 52 2,68	0,33	52 2,54	52 2,18	+ 0,14	+ 0,50	2,816		
80	5	^α Piscium.....	6	1 53 18,59	0,26	53 18,56	53 18,64	+ 0,03	- 0,05	3,090		
81	5	^χ Phœnicis....	6	1 54 55,57	0,96		54 55,92		- 0,35	2,414		
82	3	^α Arietis.....	10	1 57 39,79	0,45	57 39,83	57 39,46	- 0,04	+ 0,33	3,342		
83	4	^β Trianguli....	6	1 59 30,83	0,52	59 30,82	59 30,97	+ 0,01	- 0,14	3,520		
84	5,6	14 Arietis.....	6	1 59 49,32	0,27		59 49,09		+ 0,23	3,381		
85	6,7	62 Ceti.....	3	2 0 52,12	0,66		0 51,96		+ 0,16	3,108		
86	6	^η Arietis.....	3	2 3 21,39	0,16		3 20,78		+ 0,61	3,323		
87	7	19 Arietis.....	7	2 3 51,06	0,20	3 50,99	3 50,37	+ 0,07	+ 0,69	3,245		
88	5	^ξ Ceti.....	7	2 4 3,24	0,36	4 3,30	4 2,67	- 0,06	+ 0,57	3,165		
89	6	Ceti.....	6	2 9 15,02	0,60		9 14,33		+ 0,69	3,080		

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.			Differ-ence of Ex-tremes.	Green-wich Catalogue	A.S. Cata-logue.	Difference from		Annual Variation.
				<i>h.</i>	<i>m.</i>	<i>s.</i>				<i>s.</i>	<i>s.</i>	
90	4	ϕ Eridani.....	6	2	10	28,41	0,40		10	27,98		+ 0,43 + 2,136
91	5	ι Persei.....	6	2	10	38,13	0,44		10	37,82		+ 0,31 4,100
92	4,5	δ Cassiopeæ....	4	2	15	15,71	1,07	15 15,43	15	15,69	+ 0,28	+ 0,02 4,788
93	5	ρ Ceti.....	6	2	17	47,44	0,26		17	47,23		+ 0,21 2,893
94	4,5	κ Eridani.....	4	2	20	47,43	0,54		20	45,40		+ 2,03 2,199
95	5	σ Ceti.....	6	2	24	4,97	0,48		24	4,69		+ 0,28 2,843
96	4,5	ν Ceti.....	6	2	27	0,86	0,35	27 0,88	27	0,40	- 0,02	+ 0,46 3,136
97	6,7	ζ Ceti.....	6	2	27	38,51	0,27		27	38,36		+ 0,15 3,166
98	4	δ Ceti.....	6	2	30	49,75	0,31	30 49,75	30	49,71	0,00	+ 0,04 3,062
99	4,5	ϵ Ceti.....	6	2	31	23,59	0,29	31 23,70	31	23,52	- 0,11	+ 0,07 2,885
100	3	γ Ceti.....	6	2	34	33,00	0,25	34 33,15	34	32,87	- 0,15	+ 0,13 3,105
101	4	μ Ceti.....	6	2	35	49,09	0,19	35 49,02	35	48,46	+ 0,07	+ 0,63 3,207
102	4	π Ceti.....	6	2	36	5,01	0,32	36 4,94	36	4,61	+ 0,07	+ 0,40 2,849
103	3	ζ Arietis.....	3	2	40	3,27	0,15		40	3,14		+ 0,13 3,497
104	5	ν Fornacis....	5	2	41	53,12	0,48		41	53,35		- 0,23 2,388
105	5	τ Persei.....	6	2	42	19,50	0,58	42 19,36	42	18,81	+ 0,23	+ 0,78 4,182
106	3	η Eridani.....	6	2	48	10,53	0,25	48 10,65	48	10,70	- 0,12	- 0,17 2,917
107	5	ϵ Arietis....	6	2	49	33,89	0,08	49 33,86	49	33,57	+ 0,03	+ 0,32 3,408
108	4	γ Persei.....	6	2	52	36,52	0,83	52 36,33	52	36,00	+ 0,19	+ 0,52 4,273
109	2,3	α Ceti.....	9	2	53	27,23	0,14	53 27,20	53	27,02	+ 0,03	+ 0,21 3,123
110	4	ρ Persei.....	5	2	54	22,44	0,28	54 22,48	54	21,98	+ 0,04	+ 0,46 3,792
111	5	ρ^2 Eridani.....	3	2	54	24,80	0,51		54	24,66		+ 0,14 2,933
112	2,3	β Persei.....	5	2	57	12,17	0,42	57 12,11	57	11,57	+ 0,06	+ 0,60 3,859
113	5	κ Persei.....	6	2	58	8,01	0,76		58	7,55		+ 0,46 3,979
114	4	δ Arietis.....	6	3	1	58,76	0,23	1 58,69	1	58,63	+ 0,07	+ 0,13 3,398
115	3,4	λ Eridani.....	6	3	4	53,64	0,28	4 53,85	4	54,03	- 0,21	- 0,39 2,561
116	4	ζ Eridani.....	5	3	7	37,76	0,33	7 37,76	7	37,58	- 0,00	+ 0,18 2,906
117	2,3	α Persei.....	8	3	12	17,83	0,41	12 17,92	12	17,55	- 0,09	+ 0,28 4,221
118	4	ϵ Eridani.....	6	3	13	11,08	0,47		13	2,43		+ 8,65 2,114
119	4	\ast Camelopard ..	6	3	15	27,21	0,36	15 26,73	15	26,76	+ 0,48	+ 0,45 4,765
120	5	σ Persei.....	6	3	18	41,81	0,37		18	42,42		- 0,61 4,178
121	4,5	λ Eridani.....	6	3	22	14,26	0,26	22 14,34	22	14,22	- 0,08	+ 0,04 2,966
122	5	ζ Eridani.....	3	3	24	12,27	0,18		24	12,28		- 0,01 2,134
123	5	ν Persei.....	6	3	24	31,14	0,24		24	30,38		+ 0,76 4,208
124	4	ι^2 Eridani.....	6	3	26	19,52	0,37		26	19,33		+ 0,19 2,641
125	5	ϵ Tauri.....	4	3	28	15,45	0,53		28	15,35		+ 0,10 3,065
126	3,4	δ Persei.....	6	3	30	55,65	0,34	30 55,55	30	55,55	+ 0,10	+ 0,10 4,217
127	5	γ Eridani.....	5	3	31	1,99	0,41		31	2,06		- 0,07 2,149
128	4,5	ν Persei.....	5	3	33	44,46	0,24	33 44,39	33	43,45	+ 0,07	+ 1,01 4,035
129	4,5	b Pleiadum....	3	3	34	51,47	0,22	34 51,21	34	51,05	+ 0,26	+ 0,42 3,538
130	5	e Pleiadum....	6	3	35	9,87	0,15	35 9,83	35	9,33	+ 0,04	+ 0,54 3,546
131	3	η Tauri.....	3	3	37	27,22	0,19	37 27,11	37	26,73	+ 0,11	+ 0,49 3,542
132	5	f Pleiadum....	3	3	39	7,44	0,47		39	7,17		+ 0,27 3,543
133	5	m^1 Eridani....	6	3	39	34,85	0,33		39	34,73		+ 0,12 2,587
134	5	g Eridani.....	5	3	43	7,70	0,37		43	8,25		- 0,55 2,244
135	3,4	ζ Persei.....	6	3	43	31,62	0,17	43 31,61	43	31,33	+ 0,01	+ 0,29 3,742

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.		Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
				<i>h. m. s.</i>	<i>s.</i>				<i>s.</i>	A.S.	
136	5	ϵ Eridani.....	5	3 47 12,54	0,31		47 12,75		- 0,21	+ 2,278	
137	5	ζ Persei.....	6	3 48 1,29	0,33		48 0,51		+ 0,78	3,861	
138	4	λ Tauri.....	6	3 51 19,66	0,58	51 19,57	51 19,05	+ 0,09	+ 0,61	3,309.	
139	5	k Eridani.....	4	3 52 43,53	0,12		52 43,13		+ 0,40	2,551	
140	5	A^1 Tauri.....	6	3 54 42,90	0,93	54 42,96	54 42,60	- 0,06	+ 0,30	3,520.	
141	5	c Persei.....	4	3 56 25,49	0,66		56 24,87		+ 0,62	4,308.	
142	5	γ Reticuli.....	5	3 58 28,57	0,25		58 26,48		+ 2,09	0,841	
143	4,5	μ Persei.....	5	4 2 31,12	0,44	2 31,09	2 30,98	+ 0,03	+ 0,14	4,360.	
144	4,5	o Eridani.....	6	4 3 37,24	0,60	3 37,27	3 36,68	- 0,03	+ 0,56	2,919	
145	5	b Persei.....	4	4 5 34,01	0,28		5 33,56		+ 0,45	4,459	
146	5	a Horologii.....	5	4 8 24,34	0,72		8 24,00		+ 0,34	1,978	
147	3,4	γ Tauri.....	6	4 10 11,22	0,21	10 11,03	10 10,84	+ 0,19	+ 0,38	3,390.	
148	3,4	X Eridani.....	5	4 11 30,12	0,42	11 30,16	11 29,37	- 0,04	+ 0,75	2,259.	
149	4	δ^1 Tauri.....	5	4 13 11,86	0,15	13 11,87	13 11,45	- 0,01	+ 0,41	3,436.	
150	5	δ^2 Tauri.....	6	4 15 43,47	0,27	15 43,33	15 42,65	+ 0,14	+ 0,82	3,447	
151	5	π Tauri.....	6	4 17 3,97	0,25		17 3,81		+ 0,16	3,375	
152	4	ϵ Tauri.....	3	4 18 45,45	0,10	18 45,45	18 45,15	+ 0,00	+ 0,30	3,479.	
153	5	θ^1 Tauri.....	6	4 18 55,74	0,15	18 55,72	18 55,27	+ 0,02	+ 0,47	3,405	
154	5	ρ Tauri.....	6	4 24 16,26	0,61		24 15,66		+ 0,60	3,383	
155	5	δ Cæli Scalp...	6	4 25 39,56	0,21		25 39,50		+ 0,06	1,830	
156	5	47 Eridani.....	6	4 26 3,49	0,31		26 3,35		+ 0,14	2,883	
157	1	a Tauri.....	12	4 26 13,93	0,72	26 13,93	26 13,51	+ 0,00	+ 0,42	3,423	
158	5	d Tauri.....	4	4 26 22,48	0,10		26 22,19		+ 0,29	3,280	
159	4	54 Eridani.....	6	4 33 3,20	0,65	33 3,24	33 2,99	- 0,04	+ 0,21	2,616	
160	4,5	a Cæli Scalp...	6	4 35 7,15	0,36		35 7,60		- 0,45	1,939	
161	5	β Cæli Scalp...	5	4 36 5,26	0,21		36 4,99		+ 0,28	2,111	
162	5	μ Eridani.....	4	4 37 3,67	0,72		37 3,06		+ 0,61	2,990.	
163	4	q Orionis.....	6	4 40 40,31	0,40	40 40,41	40 40,54	- 0,10	- 0,23	3,251	
164	5	π^1 Orionis.....	6	4 41 24,49	0,27	41 24,47	41 24,33	+ 0,02	+ 0,16	3,258.	
165	4	r Orionis.....	5	4 42 12,49	0,40		42 12,93		- 0,44	3,185	
166	5	o^1 Orionis.....	5	4 42 58,96	0,56		42 58,24		+ 0,72	3,382	
167	4	i Aurigæ.....	4	4 45 59,92	0,21	45 59,99	45 56,73	- 0,07	+ 3,19	3,887	
168	5	o^2 Orionis.....	6	4 46 52,68	0,62		46 52,10		+ 0,58	3,367	
169	5	w Aurigæ.....	6	4 47 47,68	0,51		47 47,59		+ 0,09	4,047	
170	4	e Aurigæ.....	4	4 49 51,39	0,30	49 51,48	49 51,04	- 0,09	+ 0,35	4,280.	
171	4,5	i Tauri.....	6	4 53 0,20	0,24	53 0,19	52 59,73	+ 0,01	+ 0,47	3,568	
172	5	ψ Eridani.....	3	4 53 14,91	0,04		53 14,71		+ 0,20	2,901	
173	4	η Aurigæ.....	5	4 54 40,53	0,31	54 40,57	54 40,29	- 0,04	+ 0,24	4,182	
174	5	γ^1 Orionis.....	3	4 54 55,02	0,57		54 54,75		+ 0,27	3,416	
175	4	e Leporis.....	5	4 58 18,61	0,19	58 18,74	58 17,87	- 0,13	+ 0,74	2,532	
176	3	β Eridani.....	6	4 59 32,74	0,27	59 33,06	59 32,63	- 0,32	+ 0,11	2,948	
177	1	a Aurigæ.....	17	5 4 13,06	0,25	4 13,02	4 12,71	+ 0,04	+ 0,35	4,402	
178	5	ρ^1 Orionis.....	4	5 4 27,66	0,13		4 27,33		+ 0,33	3,128	
179	5	μ Leporis.....	6	5 5 20,58	0,14		5 20,26		+ 0,32	2,686	
180	1	β Orionis.....	10	5 6 25,20	0,37	6 25,17	6 25,01	+ 0,03	+ 0,19	2,876	
181	4	r Orionis.....	7	5 9 24,19	0,33	9 24,21	9 24,31	- 0,02	- 0,12	2,907	

RESULTS FROM OBSERVATIONS, 1831.

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.		Differ-ence of Ex-tremes.	Green-wich Catalogue	A.S. Cata-logue.	Difference from		Annual Variation.
				<i>h. m. s.</i>	<i>s.</i>				<i>m. s.</i>	<i>m. s.</i>	
182	5	o Columbæ....	6	5 11 23,53	0,15		11 23,17		+ 0,36	+ 2,151	
183	4,5	λ Leporis.....	5	5 11 47,40	0,37	11 47,58	11 47,14	- 0,18	+ 0,26	2,758	
184	5	m Orionis.....	5	5 13 57,19	0,60		13 57,22		- 0,03	3,145	
185	2	β Tauri.....	12	5 16 36,91	0,23	15 36,92	15 36,75	- 0,01	+ 0,16	3,779	
186	4,5	γ Orionis.....	6	5 15 58,99	0,23	15 59,02	15 58,85	- 0,03	+ 0,14	3,009	
187	5	o Tauri.....	3	5 17 29,56	0,03	17 29,37	17 29,18	+ 0,19	+ 0,38	3,593	
188	5	ψ ² Orionis.....	6	5 17 59,23	0,19	17 59,20	17 58,95	+ 0,03	+ 0,28	3,136	
189	4	β Leporis.....	6	5 21 0,40	0,30	21 0,43	21 0,39	- 0,03	+ 0,01	2,565	
190	5	ι Orionis.....	4	5 21 9,22	0,73		21 8,39		+ 0,83	3,040	
191	5	A Orionis.....	6	5 21 45,54	0,25		21 44,31		+ 1,23	3,202	
192	2	δ Orionis.....	6	5 23 22,48	0,67	23 22,63	23 22,49	- 0,15	- 0,01	3,058	
193	5	c ¹ Orionis.....	5	5 27 3,15	0,65		27 2,57		+ 0,58	2,953	
194	3,4	ζ Tauri.....	6	5 27 32,98	0,25	27 32,93	27 32,24	+ 0,05	+ 0,74	3,577	
195	4	σ Orionis.....	11	5 30 15,86	0,41	30 15,94	30 15,56	- 0,08	+ 0,30	3,005	
196	3	ζ Orionis.....	6	5 32 13,90	0,37	32 14,08	32 14,10	- 0,18	- 0,20	3,021	
197	2	α Columbæ....	7	5 33 31,97	0,77	33 32,00	33 31,65	- 0,03	+ 0,32	2,167	
198	4	γ Leporis.....	15	5 37 25,20	0,85	37 25,17	37 24,81	+ 0,03	+ 0,39	2,517	
199	5	B Tauri.....	5	5 38 38,88	0,83	38 38,03	38 38,19	- 0,15	+ 0,69	3,674	
200	4,5	ζ Leporis.....	5	5 39 17,98	0,27	39 18,13	39 17,52	- 0,15	+ 0,46	2,714	
201	4,5	C Tauri.....	13	5 42 42,53	0,49	42 42,56	42 42,24	- 0,03	+ 0,29	3,763	
202	5	δ Leporis.....	4	5 44 3,33	0,31		44 3,11		+ 0,22	2,559	
203	5	χ ¹ Orionis.....	5	5 44 22,78	0,34	44 22,71	44 22,26	+ 0,07	+ 0,52	3,559	
204	3	β Columbæ....	3	5 45 0,50	0,39	45 0,21	45 0,22	+ 0,29	+ 0,28	2,105	
205	1	α Orionis.....	7	5 46 1,43	0,32	46 1,54	46 1,42	- 0,11	+ 0,01	3,241	
206	2	β Aurigæ.....	5	5 47 7,96	0,24	47 8,01	47 7,85	- 0,05	+ 0,11	4,398	
207	4	θ Aurigæ.....	14	5 48 11,87	0,38	48 11,84	48 11,55	+ 0,03	+ 0,32	4,081	
208	4	γ Leporis.....	6	5 48 42,83	0,61	48 42,67	48 42,33	+ 0,16	+ 0,50	2,730	
209	4	η Columbæ....	9	5 51 32,74	0,68	51 32,74	51 32,68	+ 0,00	+ 0,06	2,122	
210	5	μ Orionis.....	6	5 53 5,26	0,41	53 5,29	53 4,55	- 0,03	+ 0,71	3,295	
211	5	H Geminor.....	6	5 53 51,05	0,52	53 50,99	53 50,68	+ 0,06	+ 0,37	3,642	
212	5	χ ² Orionis.....	3	5 53 52,99	0,27		53 53,11		- 0,12	3,558	
213	4,5	ν Orionis.....	19	5 57 55,50	0,61	57 55,43	57 55,03	+ 0,07	+ 0,47	3,421	
214	5	θ Columbæ....	5	6 1 44,19	0,18		1 43,62		+ 0,57	2,053	
215	5	α Lyncis.....	4	6 2 19,32	0,39		2 19,15		+ 0,17	5,535	
216	5	ξ Orionis.....	8	6 2 19,87	0,45		2 19,73		+ 0,14	3,407	
217	4	κ Aurigæ.....	5	6 4 36,64	0,55	4 36,61	4 35,83	+ 0,03	+ 0,81	3,825	
218	4,5	α Monocer.....	18	6 6 36,82	0,58	6 36,87	6 36,57	+ 0,05	+ 0,25	2,922	
219	4,5	κ Columbæ....	16	6 10 32,43	0,63	10 32,61	10 32,43	- 0,18	+ 0,00	2,130	
220	5	ι Aurigæ.....	3	6 11 52,69	0,68		11 52,37		+ 0,32	4,623	
221	3	μ Geminor.....	8	6 12 44,04	0,47	12 44,21	12 43,81	- 0,17	+ 0,23	3,623	
222	3	ζ Can. Maj.....	6	6 13 49,64	0,48	13 49,71	13 49,65	- 0,07	- 0,01	2,298	
223	2,3	β Can. Maj.....	7	6 15 15,46	0,33	15 15,57	15 15,30	- 0,11	+ 0,16	2,638	
224	5	ν Geminor....	18	6 18 55,70	0,58	18 55,61	18 55,20	+ 0,09	+ 0,50	3,561	
225	1	α Argus.....	12	6 20 11,89	1,02		20 11,91		- 0,02	1,327	
226	5	D Can. Maj....	5	6 21 54,76	0,73		21 54,45		+ 0,31	2,221	
227	5	f Monocer.....	6	6 23 45,94	0,58		23 45,80		+ 0,14	3,242	

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.		Difference of Ex-tremes.	Green-wich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
				<i>h. m. s.</i>	<i>s.</i>				Green-wich.	A.S.	
228	3	γ Geminor.	3	6 27 56,60	0,71	27 56,83	27 56,47	- 0,23	+ 0,13	+ 3,462	
229	5	ξ^a Can. Maj.	4	6 27 58,75	0,76		27 58,23		+ 0,52	2,510	
230	5	ν^a Can. Maj.	4	6 29 18,78	0,54		29 18,71		+ 0,07	2,609	
231	5	55 Aurigæ.	2	6 30 46,62	0,36		30 46,49		+ 0,13	4,377	
232	3	ν Argus.	4	6 32 35,41	0,50		32 35,49		- 0,08	1,832	
233	5	42 Camelopardi..	5	6 33 17,09	0,84		33 16,82		+ 0,27	6,299	
234	3	c Geminor.	5	6 33 32,00	0,73	33 31,94	33 31,58	+ 0,06	+ 0,42	3,693	
235	4	ξ^a Geminor.	6	6 35 48,38	0,57	35 48,15	35 48,19	+ 0,23	+ 0,19	3,375	
236	1	a Can. Maj.	21	6 37 41,98	1,10	37 41,91	37 41,91	+ 0,07	+ 0,07	2,643	
237	5	i Monocer.	4	6 38 9,21	0,44		38 8,77		+ 0,44	3,258	
238	5	α Agr. in Pup..	7	6 41 34,35	1,07		41 34,38		- 0,03	2,051	
239	5	θ Geminor.	3	6 41 38,66	0,75		41 38,18		+ 0,48	3,960	
240	5	e Lyncis.	4	6 42 37,08	0,58		42 36,56		+ 0,52	5,222	
241	4	κ_2 Can. Maj.	5	6 43 31,82	0,36	43 31,89	43 31,67	- 0,07	+ 0,15	2,238	
242	5	Can. Maj.	7	6 44 43,51	0,48		44 43,52		- 0,01	2,178	
243	4	σ^1 Can. Maj.	5	6 47 7,36	0,17	47 7,43	47 7,17	- 0,07	+ 0,19	2,486	
244	4,5	ι Can. Maj.	10	6 48 36,14	0,67	48 36,09	48 35,77	+ 0,05	+ 0,37	2,673	
245	2,3	e Can. Maj.	11	6 51 59,30	0,87	51 59,24	51 58,93	+ 0,06	+ 0,37	2,354	
246	4	ζ Geminor.	7	6 54 4,91	0,70	54 4,86	54 4,77	+ 0,05	+ 0,14	3,562	
247	3,4	σ Can. Maj.	6	6 54 59,34	0,10	54 59,42	54 59,15	- 0,08	+ 0,19	2,387	
248	4	σ^2 Can. Maj.	4	6 55 58,14	0,32	55 58,29	55 57,88	- 0,15	+ 0,26	2,502	
249	4	γ Can. Maj.	7	6 56 6,79	1,02	56 6,75	56 6,50	- 0,04	+ 0,29	2,711	
250	5	63 Aurigæ.	6	7 0 1,17	0,43		0 1,44		- 0,27	4,135	
251	5	τ Geminorum ..	5	7 0 22,38	0,80		0 22,19		+ 0,19	3,829	
252	3,4	δ Can. Maj.	5	7 1 31,07	0,60		1 31,09		- 0,02	2,436	
253	4,5	m Monocerotis ..	7	7 3 13,94	0,71	3 14,02	3 14,05	- 0,08	- 0,11	3,063	
254	5	51 Geminorum ..	5	7 3 39,81	0,60	3 39,84	3 39,71	- 0,03	+ 0,10	3,447	
255	5	64 Aurigæ.	5	7 6 16,36	0,51		6 15,97		+ 0,39	4,188	
256	4,5	e^1 Can. Maj.	9	7 7 22,04	0,56	7 22,07	7 21,88	- 0,03	+ 0,16	2,443	
257	5	1 Arg. in Pup ..	5	7 7 44,61	0,51		7 44,64		- 0,03	1,722	
258	4,5	λ Geminorum ..	4	7 8 22,75	0,34	8 22,52	8 22,67	+ 0,23	+ 0,08	3,455	
259	3,4	δ Geminorum ..	6	7 10 1,46	0,47	10 1,52	10 1,23	- 0,06	+ 0,23	3,590	
260	5	65 Aurigæ.	5	7 10 44,35	0,31		10 43,63		+ 0,72	4,030	
261	3,4	π Argus.	2	7 11 10,54	0,38		11 9,70		+ 0,84	2,116	
262	4	60 Geminorum ..	13	7 15 13,38	0,44	15 13,30	15 13,19	+ 0,08	+ 0,19	3,744	
263	5	δ Piscis Vol.	5	7 16 53,32	0,43		16 53,50		- 0,18	0,000	
264	3	η Can. Maj.	6	7 17 24,78	0,33	17 24,85	17 24,00	- 0,07	+ 0,78	2,370	
265	3	β Can. Minoris ..	6	7 17 59,14	0,46	17 58,95	17 58,47	+ 0,19	+ 0,67	3,259	
266	5	ρ Geminorum ..	4	7 18 14,01	0,37		18 13,35		+ 0,66	3,858	
267	3	α Geminorum ..	30	7 23 48,10	0,85	23 48,33	23 48,15	- 0,23	- 0,05	3,856	
268	4	σ Argus.	5	7 23 52,30	0,28		23 51,79		+ 0,51	1,906	
269	5	68 Geminorum ..	5	7 23 57,70	0,42		23 57,37		+ 0,33	3,430	
270	1,2	a Canis Minoris.	41	7 30 27,13	1,10	30 27,20	30 26,99	- 0,07	+ 0,14	3,143	
271	4,5	n Monocerotis ..	5	7 33 10,24	0,54	33 10,43	33 10,15	- 0,19	+ 0,09	2,870	
272	4	κ Geminorum ..	5	7 34 14,38	0,44	34 14,17	34 13,41	+ 0,21	+ 0,97	3,634	
273	2	β Geminorum ..	24	7 34 57,94	0,82	34 57,91	34 57,46	+ 0,03	+ 0,48	3,682	

No.	May.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.		Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
				h. m. s.	s.				m. s.	m. s.	
274	5	3 Argus.....	4	7 37 1,51	0,40		37 1,26		+ 0,25	+ 2,405	
275	4	c Argus in Pup.	6	7 39 14,19	1,10		39 14,29		- 0,10	2,135	
276	4	5 Argus.....	7	7 42 11,47	0,95	42 11,38	42 11,21	+ 0,09	+ 0,26	2,520	
277	5	6 Geminorum..	7	7 43 8,66	0,79	43 8,68	43 8,20	- 0,02	+ 0,46	3,686	
278	5	9 Argus.....	5	7 43 56,72	0,93		43 56,43		+ 0,29	2,781	
279	4,5	p Argus in Pup.	4	7 44 5,49	0,62		44 5,54		- 0,05	1,827	
280	5	b Argus in Pup.	6	7 46 39,92	0,66		46 39,53		+ 0,39	2,121	
281	5	R Argus in Pup.	6	7 48 20,19	0,69		48 20,32		- 0,13	1,762	
282	3	X Argus.....	5	7 52 28,58	1,26		52 28,81		- 0,23	1,530	
283	5	13 Argus.....	8	7 53 28,31	0,90		53 28,18		+ 0,13	3,125	
284	5	27 Lyncis — k.	7	7 55 42,71	0,84		55 42,42		+ 0,29	4,564	
285	5	55 Camelopardi..	5	7 55 53,01	0,74		55 52,09		+ 0,92	6,107	
286	3	z Argus.....	8	7 57 38,81	0,91		57 38,63		+ 0,18	2,108	
287	3,4	15 Argus.....	6	8 0 20,76	0,80	0 20,99	0 20,38	- 0,23	+ 0,38	2,558	
288	5	γ Argus.....	14	8 4 17,02	1,01		4 17,05		- 0,03	1,847	
289	2	γ Argus.....	16	8 4 19,34	0,89		4 19,65		- 0,31	1,848	
290	5	20 Argus.....	5	8 5 34,19	0,53		5 33,68		+ 0,51	2,756	
291	5	* Argus.....	5	8 5 43,81	0,80		5 43,03		+ 0,78	2,024	
292	5	r Argus in Pup.	6	8 7 6,88	0,45		7 6,78		+ 0,10	2,261	
293	4	β Cancri.....	7	8 7 20,78	0,44	7 20,74	7 20,30	+ 0,04	+ 0,48	3,262	
294	5	m Lyncis or 31..	9	8 11 14,34	0,67		11 13,85		+ 0,49	4,143	
295	4,5	q Argus in Pup.	5	8 12 14,18	0,45		12 14,31		- 0,13	2,250	
296	4,5	o Ursæ Maj. ..	5	8 16 9,21	0,55	16 9,29	16 8,38	- 0,08	+ 0,83	5,089	
297	2	c Argus.....	12	8 17 2,41	0,87		17 2,38		+ 0,03	1,243	
298	5	η Piscis. Vol. . .	7	8 23 31,32	0,74		23 30,72		+ 0,60	- 0,441	
299	5	β Piscis. Vol. . .	6	8 23 52,78	0,68		23 44,25		+ 8,53	+ 0,686	
300	5	π Ursæ Maj. . .	7	8 25 20,72	0,84		25 20,71		+ 0,01	5,368	
301	4	δ Hydra.....	11	8 28 42,14	0,77	28 42,29	28 41,81	- 0,15	+ 0,33	3,185	
302	5	σ Hydra.....	6	8 29 55,52	0,83		29 54,92		+ 0,60	3,141	
303	5	ε Arg. in Vel. . .	8	8 31 42,21	0,43		31 42,27		- 0,06	2,106	
304	5	γ Cancri.....	3	8 33 29,59	0,64	33 29,71	33 29,33	- 0,12	+ 0,26	3,493	
305	5	β Pixed Naut. . .	8	8 33 29,44	0,89		33 28,80		+ 0,64	2,342	
306	5	η Hydra.....	8	8 34 23,39	0,82		34 23,02		+ 0,37	3,141	
307	5	δ Arg. in Vel. . .	6	8 35 1,33	0,64		35 1,30		+ 0,03	1,987	
308	4,5	δ Cancri.....	5	8 35 4,33	0,30	35 4,36	35 3,93	- 0,03	+ 0,40	3,422	
309	4	o Argus.....	4	8 35 27,17	0,55		35 27,44		- 0,27	1,721	
310	4,5	a Pixed Naut. . .	6	8 36 48,44	0,87	36 48,51	36 48,25	- 0,07	+ 0,19	2,406	
311	4	ε Hydra.....	4	8 37 49,37	0,42	37 49,36	37 49,05	- 0,01	+ 0,32	3,195	
312	5	ρ Hydra.....	8	8 39 28,44	0,92		39 28,31		+ 0,13	3,184	
313	3	δ Argus.....	5	8 40 2,11	1,03		40 2,25		- 0,14	1,655	
314	5	α Arg. in Vel. . .	8	8 40 17,98	0,55		40 18,03		- 0,05	2,030	
315	4	z Hydra.....	18	8 43 27,62	0,90	43 27,37	43 27,79	+ 0,25	- 0,17	3,183	
316	3,4	ι Ursæ Maj. . . .	7	8 47 35,92	0,78		47 35,42		+ 0,50	4,131	
317	5	α Cancri.....	7	8 49 14,27	0,86	49 14,21	49 13,96	+ 0,06	+ 0,31	3,287	
318	4,5	η Ursæ Maj. or κ	6	8 52 3,08	0,53		52 3,15		- 0,07	4,147	
319	5	δ Arg. in Car. . .	9	8 52 50,36	1,02		52 50,19		+ 0,17	1,474	

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.		Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
				<i>h. m. s.</i>	<i>s.</i>				Greenwich.	A.S.	
320	5	<i>b</i> ¹ Arg. in Car...	7	8 55 15,13	1,06		55 15,60		— 0,47	+ 1,498	
321	5	Lyncis.....	7	8 55 45,37	0,67		55 45,03		+ 0,34	3,851	
322	5	<i>c</i> Arg. in Vel...	14	8 58 19,96	0,95		58 20,19		— 0,23	2,068	
323	5	<i>a</i> Piscis. Vol...	7	8 59 45,74	0,87		59 45,29		+ 0,45	0,966	
324	3,4	λ Argus....	10	9 1 47,51	0,93		1 47,27		+ 0,24	2,201	
325	5	<i>c</i> Ursæ Maj....	8	9 3 58,44	0,64		3 58,50		— 0,06	4,380	
326	4,5	<i>o</i> Hydræ.....	10	9 5 34,24	0,66	5 34,07	5 33,92	+ 0,17	+ 0,32	3,117	
327	5	<i>a</i> Arg. in Car...	5	9 6 31,47	1,10		6 30,96		+ 0,51	1,584	
328	4	<i>p</i> Lyncis.....	8	9 8 18,23	0,34	8 18,30	8 17,69	— 0,07	+ 0,54	3,767	
329	5	<i>l</i> Arg. in Vel...	6	9 8 57,34	0,62		8 57,27		+ 0,07	2,363	
330	4,5	<i>r</i> Lyncis.....	7	9 10 44,29	0,67	10 44,43	10 44,19	— 0,14	+ 0,10	3,701	
331	2-8	β Argus.....	4	9 11 18,72	0,78		11 19,73		— 1,01	0,729	
332	2	<i>c</i> Argus.....	5	9 12 33,93	0,75		12 32,27		+ 1,66	1,609	
333	5	θ Pixed Naut...	10	9 14 1,10	0,77		14 0,69		+ 0,41	2,650	
334	5	κ Leonis.....	9	9 14 47,88	0,49		14 47,74		+ 0,14	3,516	
335	3	κ Argus.....	6	9 16 53,19	1,20		16 53,15		+ 0,04	1,854	
336	4	<i>h</i> Ursæ Maj....	8	9 18 6,74	0,49	18 6,91	18 5,97	— 0,17	+ 0,77	4,831	
337	2	<i>a</i> Hydræ.....	15	9 19 16,96	0,78	19 17,02	19 16,64	— 0,06	+ 0,32	2,948	
338	5	<i>d</i> Ursæ Maj....	5	9 19 22,99	0,93		19 23,86		— 0,87	5,512	
339	3	θ Ursæ Maj....	5	9 21 30,38	1,01	21 30,52	21 29,85	— 0,14	+ 0,53	4,057	
340	4,5	λ Leonis.....	5	9 22 4,18	0,54	22 4,01	22 3,85	+ 0,17	+ 0,33	3,441	
341	5	<i>n</i> Arg. in Car...	4	9 23 3,89	0,79		23 7,04		— 3,15	1,320	
342	4,5	ψ Argus.....	4	9 24 3,62	0,87		24 3,34		+ 0,28	2,369	
343	5	<i>N</i> Arg. in Vel...	8	9 26 5,60	0,74		26 4,04		+ 1,56	1,822	
344	5	<i>h</i> Arg. in Car...	10	9 29 32,87	0,56		29 32,88		— 0,01	1,738	
345	5	<i>t</i> Hydræ.....	10	9 31 13,65	0,85		31 12,96		+ 0,69	3,063	
346	4	<i>o</i> Leonis.....	8	9 32 7,44	0,44	32 7,51	32 6,96	— 0,07	+ 0,48	3,219	
347	5	κ Hydræ.....	5	9 32 12,47	0,70		32 12,33		+ 0,14	2,874	
348	6	ψ Leonis.....	3	9 34 31,22	0,25		34 31,01		+ 0,21	3,277	
349	3	<i>e</i> Leonis.....	14	9 36 14,63	0,79	36 14,78	36 14,31	— 0,15	+ 0,32	3,426	
350	4,5	<i>v</i> Ursæ Maj....	10	9 38 54,10	0,87	38 54,10	38 54,71	+ 0,00	— 0,61	4,356	
351	5	ϕ Ursæ Maj....	7	9 40 32,98	0,67		40 33,63		— 0,65	4,153	
352	5	<i>l</i> Arg. in Car...	5	9 40 36,02	0,32		40 34,75		+ 1,27	1,648	
353	3,4	<i>v</i> Argus.....	5	9 42 52,50	0,70		42 52,56		— 0,06	1,505	
354	3	μ Leonis.....	6	9 43 8,14	0,72	43 8,32	43 7,57	— 0,18	+ 0,57	3,448	
355	5	<i>v</i> ¹ Hydræ.....	5	9 43 21,16	0,17		43 21,21		— 0,05	2,880	
356	4	ϕ Argus.....	5	9 50 56,69	0,58		50 56,49		+ 0,20	2,095	
357	4,5	π Leonis.....	7	9 51 16,59	0,37	51 16,72	51 16,50	— 0,13	+ 0,09	3,179	
358	5	<i>d</i> Leonis Min...	10	9 57 26,56	0,64		57 25,99		+ 0,57	3,564	
359	3,4	η Leonis.....	8	9 58 6,80	0,62	58 6,65	58 6,23	+ 0,15	+ 0,57	3,283	
360	5	Λ Leonis.....	5	9 58 56,00	0,49	58 55,79	58 55,55	+ 0,21	+ 0,45	3,197	
361	1	<i>a</i> Leonis.....	7	9 59 21,97	0,58	59 21,93	59 21,51	+ 0,04	+ 0,46	3,221	
362	4,5	λ ² Hydræ.....	11	10 2 21,13	0,58	2 21,20	2 20,66	— 0,07	+ 0,47	2,934	
363	6	21 Sextantis...	5	10 5 43,19	0,69		5 43,13		+ 0,06	2,988	
364	3,4	λ Ursæ Maj....	9	10 6 52,30	0,65	6 52,44	6 52,23	— 0,14	+ 0,07	3,675	
365	4,5	ζ Leonis.....	6	10 7 16,65	0,32	7 16,67	7 16,31	— 0,02	+ 0,34	3,353	

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.		Difference of Ex-tremes.	Green-wich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
				<i>h. m. s.</i>	<i>s.</i>				<i>m. s.</i>	<i>m. s.</i>	
366	4	<i>γ</i> Arg. in Vel...	9	10 7 39,58	0,48		7 39,60		- 0,02	+ 2,516	
367	4,5	<i>ω</i> Argus.....	6	10 9 42,90	0,74		9 42,28		+ 0,62	1,440	
368	2	<i>γ</i> Leonis.....	6	10 10 38,69	0,53	10 38,66	10 38,31	+ 0,03	+ 0,38	3,300	
369	5	<i>q</i> Arg. in Car...	4	10 11 27,24	0,49		11 25,63		+ 1,61	1,991	
370	3	<i>μ</i> Ursæ Maj....	7	10 12 13,86	0,37	12 13,91	12 13,81	- 0,05	+ 0,05	3,620	
371	6	42 Leonis.....	3	10 12 44,54	0,52		12 44,28		+ 0,26	3,239	
372	5	<i>Γ</i> Arg. in Vel...	8	10 14 38,46	1,24		14 36,67		+ 1,79	2,215	
373	4,5	<i>r</i> Arg. in Vel...	6	10 15 5,58	0,53		15 5,61		- 0,03	2,558	
374	4,5	<i>f</i> Leonis Min...	6	10 16 12,42	0,47	16 12,44	16 11,42	- 0,02	+ 1,00	3,473	
375	4	<i>μ</i> Hydræ.....	7	10 17 55,46	0,56	17 55,41	17 54,75	+ 0,05	+ 0,71	2,903	
376	4,5	<i>g</i> Leonis.....	5	10 18 5,25	0,20	18 5,31	18 3,82	- 0,06	+ 1,43	3,511	
377	4,5	<i>a</i> Antl. Penum..	4	10 19 25,80	0,27	19 25,74	19 25,88	+ 0,06	- 0,08	2,737	
378	5	36 Ursæ Maj. . .	6	10 19 45,17	0,53	19 45,13	19 46,22	+ 0,04	- 1,05	3,935	
379	4	<i>p</i> Leonis.....	14	10 23 54,47	0,39	23 54,44	23 54,31	+ 0,03	+ 0,16	3,166	
380	5	<i>κ</i> Ursæ Maj....	5	10 24 12,79	0,85	24 13,02	24 12,77	- 0,23	+ 0,02	3,935	
381	5,6	48 Leonis.....	5	10 25 58,88	0,38	25 58,80	25 58,42	+ 0,08	+ 0,46	3,141	
382	4	<i>p</i> Arg. in Car...	7	10 26 2,13	0,69		26 2,67		- 0,54	2,114	
383	4	<i>l</i> Leonis Min...	9	10 29 11,34	0,82	29 11,46	29 11,18	- 0,12	+ 0,16	3,401	
384	5	<i>p</i> Arg. in Vel...		10 30 13,78	0,96		30 13,27		+ 0,51	2,514	
385	5	<i>φ</i> ³ Hydræ & Crat.	7	10 30 21,23	0,47		30 21,35		- 0,12	2,922	
386	5,6	40 Leonis Min..	5	10 33 44,28			33 43,69		+ 0,59	3,321	
387	5	<i>θ</i> ¹ Argus.....	5	10 36 16,02	0,76		36 13,87		+ 2,15	2,106	
388	4,5	<i>n</i> Leo. Min. or 42	8	10 36 26,94	0,41	36 27,01	36 26,28	- 0,07	+ 0,66	3,361	
389	2,3	<i>θ</i> ² Argus.....	15	10 36 56,84	0,87		36 55,04		+ 1,80	2,117	
390	2	<i>q</i> Argus.....	6	10 38 31,81	0,48		38 32,30		- 0,49	2,300	
391	3	<i>u</i> Argus.....	7	10 39 31,22	0,98		39 30,95		+ 0,27	2,548	
392	4	<i>v</i> Hydræ & Crat.	2	10 41 17,56	0,03	41 17,54	41 17,06	+ 0,02	+ 0,50	2,945	
393	4,5	<i>o</i> Leonis Min...	8	10 43 50,39	0,64	43 50,22	43 50,09	+ 0,17	+ 0,30	3,375	
394	5	<i>ω</i> Ursæ Maj....	10	10 44 13,44	0,40		44 12,89		+ 0,55	3,488	
395	4,5	54 Leonis.....	8	10 46 27,02	0,54	46 27,09	46 26,89	- 0,07	+ 0,13	3,271	
396	5	<i>z</i> Arg. in Car...	6	10 46 39,66	0,62		46 40,74		- 1,08	2,396	
397	5	Antl. Penum..	7	10 48 51,46	0,84		48 51,64		- 0,18	2,769	
398	4	<i>a</i> Hydræ & Crat.	10	10 51 32,75	0,27	51 32,92	51 32,32	- 0,17	+ 0,43	2,905	
399	2	<i>β</i> Ursæ Maj....	5	10 51 35,36	0,24	51 35,35	51 35,48	+ 0,01	- 0,12	3,680	
400	5	<i>d</i> Leonis.....	6	10 51 49,86	0,23	51 49,79	51 49,58	+ 0,07	+ 0,28	3,099	
401	1,2	<i>a</i> Ursæ Maj....	6	10 53 13,68	0,55	53 13,66	53 13,57	+ 0,02	+ 0,11	3,811	
402	5	<i>b</i> Leonis.....	7	10 53 17,85	0,55		53 17,41		+ 0,44	3,216	
403	4,5	<i>χ</i> Leonis.....	7	10 56 18,03	0,81	56 17,86	56 17,34	+ 0,17	+ 0,69	3,086	
404	5	<i>χ</i> ¹ Hydræ & Crat.	8	10 57 12,20	0,51		57 12,01		+ 0,19	2,889	
405	5,6	<i>p</i> ² Leonis.....	6	10 58 16,86	0,41		58 16,60		+ 0,26	3,086	
406	3,4	<i>ψ</i> Ursæ Maj....	8	11 0 7,75	0,66	0 7,98	0 8,17	- 0,23	- 0,42	3,419	
407	5	10 Hydræ & Crat.	6	11 0 34,32	0,59		0 34,17		+ 0,15	2,892	
408	4	<i>β</i> Hydræ & Crat.	6	11 3 21,56	0,81	3 21,46	3 21,55	+ 0,10	+ 0,01	2,937	
409	5,6	<i>p</i> ⁴ Leonis.....	9	11 5 6,55	0,80		5 6,36		+ 0,19	3,073	
410	3	<i>δ</i> Leonis.....	4	11 5 6,92	0,50	5 6,79	5 6,30	+ 0,13	+ 0,62	3,193	
411	3	<i>θ</i> Leonis.....	7	11 5 21,85	0,52	5 21,98	5 21,45	- 0,13	+ 0,40	3,161	

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.		Difference from		Annual Variation.	
				<i>h.</i>	<i>m.</i>	<i>s.</i>			<i>s.</i>	<i>m.</i>	<i>s.</i>	Greenwich.		A.S.
412	5	φ Leonis.....	7	11	8	4,42	0,53		8	3,67	+ 0,75	+ 3,054		
413	4	ξ Ursæ Maj....	6	11	9	9,16	0,61	9	9,21	9	8,45	+ 0,71	3,221	
414	4	v Ursæ Maj....	3	11	9	19,93	0,23	9	19,95	9	19,82	- 0,11	3,266	
415	5	p Ursæ Maj....	3	11	9	53,89	0,56		9	53,53	+ 0,36	3,304		
416	4	σ Leonis.....	4	11	12	25,54	0,32	12	25,26	12	24,84	+ 0,28	+ 0,70	3,071
417	4	π Centauri....	4	11	13	19,79	0,40		13	19,10	+ 0,69	2,702		
418	7	* Leonis.....	7	11	14	38,84	0,59		14	38,63	+ 0,21	3,073		
419	4	ι Leonis.....	9	11	15	6,85	0,55	15	6,00	15	6,26	+ 0,25	+ 0,59	3,121
420	5	e Hydræ & Crat.	7	11	16	4,85	0,54		16	4,77	+ 0,08	3,023		
421	4	γ Hydræ & Crat.	3	11	16	26,89	0,10	16	26,78	16	26,27	+ 0,11	+ 0,62	2,992
422	4	τ Leonis.....	9	11	19	14,96	0,66	19	14,84	19	14,39	+ 0,12	+ 0,57	3,084
423	4,5	e Leonis.....	7	11	21	40,97	0,41	21	40,97	21	40,91	0,00	+ 0,06	3,060
424	7	* Hydræ & Crat.	5	11	23	20,89	0,25		23	20,82	+ 0,07	3,047		
425	4	ξ Hydræ & Crat.	7	11	24	42,64	0,38	24	42,46	24	41,71	+ 0,18	+ 0,93	2,945
426	6	* Ursæ Min....	5	11	27	23,34	0,52		27	23,31	+ 0,03	3,172		
427	4	λ Centauri....	8	11	28	2,16	0,69		28	0,18	+ 1,98	2,717		
428	4	θ Hydræ & Crat.	8	11	28	6,85	0,58	28	6,96	28	6,45	- 0,11	+ 0,40	3,039
429	4,5	v Leonis.....	6	11	28	18,00	0,37	28	18,05	28	17,58	- 0,05	+ 0,42	3,068
430	5,6	92 Leonis.....	3	11	31	59,23	0,01		31	59,03	+ 0,20	3,135		
431	6,7	* Virginis.....	7	11	35	17,46	0,66		35	17,24	+ 0,22	3,054		
432	4	ζ Hydræ & Crat.	8	11	36	12,50	0,51	36	12,46	36	11,95	+ 0,04	+ 0,55	3,025
433	5	ξ Virginis.....	9	11	36	34,27	0,28	36	34,10	36	33,38	+ 0,17	+ 0,89	3,090
434	4	χ Ursæ Maj....	5	11	37	5,77	0,20	37	5,78	37	4,99	- 0,01	+ 0,78	3,220
435	4,5	v Virginis.....	5	11	37	10,20	0,25	37	10,34	37	10,12	- 0,14	+ 0,08	3,086
436	2,3	β Leonis.....	3	11	40	26,01	0,21	40	26,11	40	25,83	+ 0,10	+ 0,18	3,064
437	3,4	β Virginis.....	8	11	41	53,67	0,51	41	53,78	41	53,30	- 0,11	+ 0,37	3,124
438	6	B Virginis.....	6	11	42	24,12	0,48		42	23,65	+ 0,47	3,060		
439	4	β Hydræ & Crat.	6	11	44	23,64	0,39	44	23,41	44	23,50	+ 0,23	+ 0,14	3,009
440	2	γ Ursæ Maj....	7	11	44	54,23	0,50	44	54,21	44	54,02	+ 0,02	+ 0,21	3,192
441	6	A Virginis.....	6	11	46	22,77	0,35		46	22,52	+ 0,25	3,081		
442	5	π Virginis.....	13	11	52	12,77	0,47		52	12,68	+ 0,09	3,074		
443	6	l Comæ Ber....	5	11	53	4,50	1,08		53	3,62	+ 0,88	3,085		
444	7	* Virginis.....	6	11	55	6,85	0,51		55	7,04	- 0,19	3,071		
445	4,5	o Virginis.....	15	11	56	36,05	0,44	56	35,95	56	35,78	+ 0,10	+ 0,27	3,071
446	4,5	η Crucis.....	6	11	58	7,88	0,52		58	8,29	- 0,11	3,046		
447	3	δ Centauri....	6	11	59	38,26	0,82		59	38,62	- 0,36	3,065		
448	4,5	a Corvi.....	3	11	59	42,76	0,09	59	42,72	59	42,59	+ 0,04	+ 0,17	3,067
449	6	r Virginis.....	3	12	1	1,66	0,70		1	1,46	+ 0,20	3,068		
450	4	e Corvi.....	6	12	1	27,00	0,50	1	26,82	1	26,57	+ 0,18	+ 0,43	3,071
451	4	ρ Centauri....	6	12	2	51,48	0,50		2	50,33	+ 1,15	3,088		
452	3	δ Crucis.....	7	12	6	13,70	0,42		6	11,64	+ 2,06	3,125		
453	3	δ Ursæ Maj....	8	12	7	1,40	0,85	7	1,38	7	0,58	+ 0,02	+ 0,82	3,003
454	3	γ Corvi.....	6	12	7	7,61	0,25	7	7,61	7	7,29	0,00	+ 0,32	3,080
455	5	h Comæ Ber....	5	12	7	47,23	0,33		7	46,84	+ 0,39	3,047		
456	6	n Virginis.....	6	12	10	0,70	0,46	10	0,71	10	0,43	- 0,01	+ 0,27	3,068
457	3,4	γ Virginis.....	5	12	11	15,66	0,20	11	15,62	11	15,70	+ 0,04	- 0,04	3,068

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.			Differ-ence of Ex-tremes.	Green-wich Catalogue	A.S. Cata-logue.	Difference from		Annual Variation.		
				<i>h.</i>	<i>m.</i>	<i>s.</i>				<i>s.</i>	<i>s.</i>		<i>s.</i>	
458	5	<i>s</i> Comæ Ber...	5	12	12	10,44	0,65		12	9,81	+ 0,63	+ 3,044		
459	5	<i>e</i> Comæ Ber...	9	12	14	0,15	0,56		13	59,92	+ 0,23	3,027		
460	5	<i>f</i> Comæ Ber...	6	12	15	49,40	0,69		15	48,60	+ 0,80	3,021		
461	1	<i>a</i> * Crucis.....	5	12	17	15,72	0,70		17	14,36	+ 1,36	3,258		
462	4,5	<i>a</i> Comæ Ber...	4,5	12	18	31,85	0,54	18	31,81	18	31,12	+ 0,04	+ 0,73	3,011
463	5	<i>σ</i> Centauri.....	5	12	18	56,48	0,60		18	57,04	- 0,56	3,196		
464	2,3	<i>γ</i> Crucis.....	5	12	21	50,79	0,31		21	51,26	- 0,47	3,257		
465	4,5	<i>η</i> Corvi.....	6	12	23	22,52	0,50	23	22,34	23	22,36	+ 0,18	+ 0,16	3,105
466	2,3	<i>β</i> Corvi.....	7	12	25	31,60	0,77	25	31,53	25	30,92	+ 0,07	+ 0,68	3,129
467	4,5	<i>d</i> Canum. Ven.	5	12	25	42,04	0,55	25	42,01	25	41,97	0,00	+ 0,07	2,864
468	3,4	<i>κ</i> Draconis.....	3	12	26	13,35	0,38	26	13,24	26	12,32	+ 0,11	+ 1,03	2,600
469	5	<i>τ</i> Centauri.....	5	12	28	30,02	0,51		28	30,05	- 0,03	3,249		
470	5	<i>ι</i> Centauri.....	10	12	30	45,53	0,68		30	45,46	+ 0,07	3,213		
471	3	<i>γ</i> Centauri.....	7	12	32	14,30	0,62		32	15,00	- 0,70	3,276		
472	4	<i>γ</i> ¹ Virginis.....	3	12	33	6,11	0,30	33	6,11	33	5,81	0,00	+ 0,30	3,022
473	4	<i>β</i> Muscæ.....	4	12	36	1,16	0,73		36	1,92	- 0,76	3,564		
474	2	<i>β</i> Crucis.....	8	12	37	54,59	0,61		37	55,41	- 0,82	3,430		
475	6	<i>n</i> Comæ Ber...	7	12	38	12,17	0,38		38	11,89	+ 0,28	2,998		
476	6,7	* Virginis....	3	12	38	50,19	0,67		38	49,64	+ 0,55	3,089		
477	6	35 Virginis....	3	12	39	15,18	0,17		39	14,93	+ 0,25	3,050		
478	5	<i>n</i> Centauri.....	10	12	44	6,55	0,93		44	5,27	+ 1,28	3,277		
479	5	<i>o</i> Centauri.....	7	12	44	43,31	0,83		44	43,55	- 0,24	3,455		
480	5	<i>q</i> Comæ Ber...	5	12	44	58,18	0,42		44	58,21	- 0,03	2,962		
481	3	<i>e</i> Ursæ Maj....	4	12	46	34,29	0,20	46	34,26	46	33,58	+ 0,03	+ 0,71	2,655
482	3,4	<i>δ</i> Virginis.....	5	12	47	5,64	0,42	47	5,59	47	5,37	+ 0,05	+ 0,27	3,004
483	2,3	<i>a</i> Canum. Ven..	4	12	48	6,76	0,21	48	6,74	48	6,27	+ 0,02	+ 0,49	2,841
484	4,5	<i>γ</i> Comæ Ber...	6	12	50	33,70	0,68	50	33,80	50	33,34	- 0,10	+ 0,36	2,971
485	6	<i>k</i> ¹ Virginis....	3	12	50	57,60	0,41		50	57,32	+ 0,28	3,083		
486	5	37 Comæ Ber...	9	12	52	10,77	0,75		52	11,26	- 0,49	2,832		
487	3,4	<i>e</i> Virginis.....	7	12	53	46,06	0,56	53	46,01	53	45,77	+ 0,05	+ 0,29	3,003
488	6	<i>k</i> ⁴ Virginis....	6	12	55	12,27	0,34		55	11,96	+ 0,31	3,083		
489	5	53 Virginis.....	7	13	3	4,83	0,53		3	4,24	+ 0,59	3,167		
490	6	* Virginis.....	4	13	4	8,45	0,27		4	8,29	+ 0,16	2,987		
491	4,5	61 Virginis....	12	13	9	34,87	0,44	9	35,04	9	34,27	- 0,17	+ 0,60	3,166
492	4,5	<i>γ</i> Hydræ Con...	6	13	9	45,17	0,57	9	45,22	9	44,91	- 0,05	+ 0,26	3,232
493	5	<i>h</i> Canum. Ven..	5	13	9	57,29	0,48		9	56,68	+ 0,61	2,713		
494	5	21 Canum. Ven..	4	13	11	2,32	0,20		11	2,11	+ 0,21	2,573		
495	6	65 Virginis....	4	13	14	33,97	0,21		14	34,15	- 0,18	3,098		
496	1	<i>a</i> Virginis....	15	13	16	18,14	0,72	16	18,00	16	17,86	+ 0,14	+ 0,28	3,147
497	3	<i>ζ</i> Ursæ Maj....	9	13	17	6,37	0,50	17	6,21	17	4,91	+ 0,16	+ 1,46	2,419
498	5	<i>i</i> Virginis.....	6	13	17	48,26	0,46	17	48,20	17	47,61	+ 0,06	+ 0,65	3,161
499	5,6	<i>p</i> Virginis.....	3	13	18	27,18	0,32		18	26,76	+ 0,42	3,189		
500	4	<i>d</i> Centauri.....	8	13	21	16,68	0,75		21	16,61	+ 0,07	3,437		
501	6	<i>l</i> ² Virginis....	4	13	23	11,28	0,30	23	11,57	23	10,94	- 0,29	+ 0,34	3,113
502	6	75 Virginis....	4	13	23	50,66	0,40		23	50,19	+ 0,47	3,191		
503	4	<i>ζ</i> Virginis.....	9	13	26	5,37	0,48	26	5,39	26	5,33	- 0,02	+ 0,04	3,066

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
				<i>h.</i>	<i>m.</i>	<i>s.</i>				<i>s.</i>	<i>s.</i>	
504	6	l ³ Virginis.	6	13	26	44,29	0,27	26	44,06		+ 0,23	+ 3,107
505	6	t Centauri.	4	13	29	13,76	0,14	29	13,93		- 0,17	3,345
506	3	e Centauri.		13	29	14,24	0,51	29	14,91		- 0,67	3,731
507	6	l Bootis.	7	13	32	36,12	0,54	32	35,48		+ 0,64	2,868
508	5,6	m Virginis.	5	13	32	45,15	0,10	32	44,95	- 0,08	+ 0,20	3,140
509	5	i Centauri.	9	13	36	6,70	0,76	36	7,44		- 0,74	3,410
510	5	τ Bootis.	6	13	39	13,82	0,46	39	13,41		+ 0,41	2,883
511	4	μ Centauri.	5	13	39	28,52	0,35	39	28,55		- 0,03	3,567
512	5	g Centauri.	3	13	39	40,98	0,19	39	40,88		+ 0,10	3,442
513	2,3	η Ursæ Maj.	2	13	40	52,33	0,47	40	51,72	- 0,01	+ 0,61	2,353
514	4	v Bootis.	3	13	41	19,58	0,16	41	19,56	+ 0,02	+ 0,46	2,897
515	5	h Centauri.	5	13	43	30,58	0,33	43	30,46		+ 0,12	3,419
516	3	ζ Centauri.	6	13	45	2,65	0,60	45	2,84		- 0,19	3,690
517	4,5	i Draconis.	5	13	46	29,66	0,18	46	29,61	+ 0,05	+ 0,18	1,751
518	3	γ Bootis.	6	13	46	38,39	0,31	46	38,49	- 0,10	+ 0,35	2,859
519	5	φ Centauri.	5	13	48	2,15	0,49	48	2,13		+ 0,02	3,600
520	5	v ^a Centauri.	6	13	51	13,63	0,68	51	13,64		- 0,01	3,686
521	1	β Centauri.	5	13	51	58,57	1,07	51	59,57		- 1,00	4,134
522	4,5	τ Virginis.	6	13	53	3,25	0,51	53	3,14	+ 0,11	+ 0,22	3,042
523	5	χ Centauri.	6	13	55	45,93	1,03	55	45,52		+ 0,41	3,617
524	4,5	π Hydræ Con.	5	13	56	46,17	0,61	56	46,17	0,00	+ 0,15	3,384
525	2	θ Centauri.	4	13	56	46,02	0,43	56	45,98	+ 0,04	- 0,56	3,491
526	3,4	α Draconis.	4	13	59	49,02		59	49,10	- 0,08	+ 0,83	1,625
527	6	* Virginis.	3	14	1	37,47	0,57	1	36,96		+ 0,51	3,255
528	5	6 Hydræ Con.	8	14	3	6,51	0,51	3	6,44		+ 0,07	3,408
529	4	κ Virginis.	8	14	3	53,58	0,22	3	53,61	- 0,03	+ 0,04	3,183
530	4	ι Virginis.	3	14	7	9,91	0,10	7	9,85	+ 0,06	+ 0,21	3,132
531	1	α Bootis.	9	14	7	57,30	0,40	7	57,37	- 0,07	+ 0,28	2,731
532	4,5	ι Lupi.	6	14	8	37,68	0,26	8	37,16		+ 0,52	3,786
533	4	λ Virginis.	5	14	9	58,84	0,39	9	59,00	- 0,16	+ 0,30	3,228
534	5	τ ¹ Lupi.	6	14	15	19,89	0,18	15	19,63		+ 0,26	3,797
535	5	τ ² Lupi.	5	14	15	21,46	0,18	15	20,97		+ 0,49	3,802
536	4	θ Bootis.	6	14	19	26,63	0,56	19	26,50	+ 0,13	+ 0,91	2,015
537	5	φ Virginis.	8	14	19	30,26	0,26	19	29,97		+ 0,29	3,088
538	5	σ Lupi.	5	14	21	17,19	0,31	21	17,70		- 0,51	3,979
539	4	ρ Bootis.	6	14	24	32,76	0,15	24	32,75	+ 0,01	+ 0,06	2,592
540	3	η Centauri.	4	14	24	48,73	0,49	24	48,76		- 0,03	3,764
541	5	ρ Lupi.	6	14	26	34,16	0,82	26	35,19		- 1,03	3,975
542	5	σ Bootis.	4	14	27	19,21	0,54	27	18,77		+ 0,44	2,597
543	1	α ² Centauri.	3	14	28	11,78	0,76	28	18,13		- 6,35	4,470
544	3	α Lupi.	6	14	30	44,17	0,85	30	44,71		- 0,54	3,933
545	3,4	π Bootis.	6	14	32	47,08	0,58	32	47,13	- 0,05	+ 0,66	2,813
546	3,4	ζ Bootis.	5	14	33	4,90	0,26	33	4,32	+ 0,02	+ 0,58	2,855
547	5	C ¹ Centauri.	5	14	33	20,93	0,43	33	21,25		- 0,32	3,636
548	3	e Bootis.	6	14	37	36,38	0,23	37	36,36	+ 0,02	+ 0,48	2,621
549	5	6 Libræ.	5	14	40	23,13	0,48	40	22,90		+ 0,23	3,511

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.		Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
				h. m. s.	s.				Greenwich.	A.S.	
550	5	o Lupi.....	6	14 40 39,01	0,63		40 38,52		+ 0,49	+ 3,868	
551	3	a ² Libræ.....	4	14 41 32,56	0,22	41 32,63	41 32,60	- 0,07	- 0,04	3,305	
552	3,4	ξ Bootis.....	6	14 43 35,76	0,34	43 35,84	43 35,46	- 0,08	+ 0,30	2,753	
553	3,4	β Lupi.....	6	14 47 30,09	0,44		47 30,11		- 0,02	3,883	
554	5	ξ ² Libræ.....	6	14 47 36,77	0,18	47 36,72	47 35,85	+ 0,05	+ 0,92	3,237	
555	3	χ Centauri.....	5	14 48 12,31	0,51		48 12,27		+ 0,04	3,857	
556	4,5	δ Libræ.....	6	14 51 57,35	0,20	51 57,28	51 57,01	+ 0,07	+ 0,34	3,198	
557	5	π Lupi.....	4	14 53 39,40	0,53		53 39,29		+ 0,11	4,031	
558	3,4	γ Libræ.....	5	14 54 12,00	0,14	54 11,99	54 11,64	+ 0,01	+ 0,36	3,490	
559	5	110 Virginis....	2	14 54 22,06	0,08		54 21,95		+ 0,11	3,024	
560	5	ψ Bootis.....	5	14 57 12,32	0,43		57 11,67		+ 0,65	2,580	
561	5	c Bootis.....	7	14 59 52,86	0,35		59 52,10		+ 0,76	2,617	
562	4	ζ Lupi.....	4	15 0 11,35	0,59		0 12,59		- 1,24	4,254	
563	5	β Circini.....	6	15 4 21,40	0,39		4 22,58		- 1,18	4,618	
564	5	μ Lupi.....	4	15 6 49,49	0,52		6 49,94		- 0,45	4,119	
565	5	χ Bootis.....	6	15 7 25,39	0,40		7 24,99		+ 0,40	2,510	
566	2,3	β Libræ.....	2	15 7 55,50	0,09	7 55,49	7 55,21	+ 0,01	+ 0,29	3,218	
567	5	φ ² Lupi.....	6	15 12 23,32	0,41		12 23,37		- 0,05	3,797	
568	4	μ Bootis.....	8	15 18 6,38	0,23	18 6,50	18 5,93	- 0,12	+ 0,45	2,275	
569	4	β Cor. Bor.....	7	15 20 51,85	0,25	20 51,86	20 51,74	- 0,01	+ 0,11	2,483	
570	4	γ Lupi.....	3	15 23 54,77	0,22		23 54,72		+ 0,05	3,957	
571	4	f ¹ Libræ.....	7	15 24 57,19	0,16	24 57,13	24 56,78	+ 0,06	+ 0,41	3,242	
572	4,5	γ Libræ.....	3	15 26 4,91	0,62	26 5,13	26 5,24	- 0,22	- 0,33	3,333	
573	4,5	40 Libræ.....	6	15 28 17,90	0,39	28 18,02	28 17,55	- 0,12	+ 0,35	3,657	
574	5	κ Libræ.....	7	15 32 13,44	0,31	32 13,54	32 13,31	- 0,10	+ 0,13	3,524	
575	5	ζ Cor. Bor.....	4	15 33 1,06	0,65		33 0,52		+ 0,54	2,256	
576	5	i Serpentis.....	6	15 34 1,26	0,47		34 0,46		+ 0,80	2,672	
577	4,5	λ Serpentis.....	6	15 38 14,94	0,46	38 14,94	38 14,70	0,00	+ 0,24	2,917	
578	3,4	β Serpentis.....	6	15 38 23,46	0,31	38 23,29	38 23,12	+ 0,17	+ 0,34	2,757	
579	4,5	χ Lupi.....	4	15 40 14,50	0,14	40 14,44	40 14,43	+ 0,06	+ 0,07	3,782	
580	4,5	θ Libræ.....	5	15 44 13,00	0,10	44 13,02	44 13,20	- 0,02	- 0,20	3,390	
581	4	ρ Scorpil.....	2	15 46 28,21	0,02	46 28,20	46 27,98	+ 0,01	+ 0,23	3,679	
582	3,4	π Scorpil.....		15 48 38,70	0,47	48 38,85	48 38,64	- 0,15	+ 0,06	3,606	
583	3	δ Scorpil.....	5	15 50 21,26	0,29	50 21,28	50 21,33	- 0,02	- 0,07	3,527	
584	4,5	e Cor. Bor.....	4	15 50 35,63	0,27		50 35,59		+ 0,04	2,484	
585	4,5	ξ Libræ.....	5	15 55 5,25	0,10	55 5,04	55 4,93	+ 0,21	+ 0,32	3,288	
586	2	β Scorpil.....	8	15 55 37,48	0,26		55 37,46		+ 0,02	3,469	
587	4,5	w ¹ Scorpil.....	5	15 56 56,19	0,10	56 56,19	56 56,25	0,00	- 0,06	3,496	
588	5	δ Trian. Aus....	4	16 0 7,94	0,63		0 9,22		- 1,28	5,363	
589	4	v Scorpil.....	6	16 2 11,22	0,51	2 11,19	2 11,27	+ 0,03	- 0,05	3,469	
590	5	z Scorpil.....	6	16 6 26,69	0,26		6 26,20		+ 0,49	3,231	
591	5	γ ² Normæ.....	5	16 7 13,98	0,12		7 13,62		+ 0,36	4,158	
592	3	e Ophiuchi....	5	16 9 23,22	0,16	9 23,16	9 23,01	+ 0,08	+ 0,21	3,156	
593	5	σ Serpentis....	4	16 13 31,09	0,44		13 31,51		- 0,42	3,098	
594	5	ψ Ophiuchi....	6	16 14 13,63	0,56	14 13,55	14 13,50	+ 0,08	+ 0,13	3,495	
595	3,4	γ Herculis.....	5	16 14 28,99	0,25	14 28,07	14 27,84	+ 0,02	+ 0,25	2,643	

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.			Differ-ence of Ex-tremes.	Green-wich Catalogue	A.S. Cata-logue.	Difference from		Annual Variation.
				<i>h. m. s.</i>	<i>s.</i>	<i>m. s.</i>				<i>m. s.</i>	<i>s.</i>	
596	5	χ Ophiuchi	3	16 17 14,36	0,13		17 13,62		+ 0,74	+ 3,461		
597	5	α Normæ	5	16 20 21,48	0,30		20 21,56		- 0,08	3,895		
598	4,5	ϕ Ophiuchi	6	16 21 28,67	0,15	21 28,52	21 28,62	+ 0,15	+ 0,05	3,422		
599	3	η Draconis	3	16 22 43,00	0,36	21 42,90	21 42,27	+ 0,10	+ 0,73	0,792		
600	3,4	τ Scorpii	6	16 25 22,65	0,20	25 22,59	25 22,59	+ 0,06	+ 0,06	3,715		
601	3,4	ζ Ophiuchi	6	16 27 51,74	0,06	27 51,73	27 51,58	+ 0,01	+ 0,16	3,290		
602	5	ν Scorpii	5	16 31 48,55	0,39	31 48,59	31 48,42	- 0,04	+ 0,13	3,456		
603	3	ζ Herculis	6	16 34 55,08	0,21	34 55,08	34 54,62	0,00	+ 0,46	2,246		
604	4	η Aræ	5	16 35 14,41	0,42		35 14,94		- 0,53	5,119		
605	3	η Herculis	5	16 37 6,21	0,31	37 6,35	37 5,56	- 0,14	+ 0,65	2,047		
606	5	ν Ophiuchi	7	16 40 29,59	0,54		40 29,27		+ 0,32	3,330		
607	5	k Herculis	5	16 42 7,43	0,54		42 7,21		+ 0,22	2,901		
608	5	s Herculis	5	16 44 3,45	0,35		44 2,82		+ 0,63	2,336		
609	5	q Ophiuchi	3	16 45 34,13	0,16		45 33,95		+ 0,18	3,198		
610	5	53 Herculis	7	16 46 33,68	0,33		46 33,11		+ 0,57	2,276		
611	4	κ Ophiuchi	5	16 49 40,55	0,30	49 40,51	49 40,19	+ 0,04	+ 0,36	2,852		
612	5	k Scorpii	5	16 53 43,20	0,21		53 43,20		+ 0,00	3,928		
613	3	e Herculis	5	16 53 49,61	0,09	53 49,62	53 49,11	- 0,01	+ 0,50	2,293		
614	5	60 Herculis	4	16 57 32,53	0,56		57 32,58		- 0,05	2,771		
615	4	η Scorpii	5	17 0 3,26	0,67		0 3,70		- 0,44	4,272		
616	2,3	η Ophiuchi	6	17 0 41,78	0,38	0 41,72	0 41,58	+ 0,06	+ 0,20	3,426		
617	4	μ Draconis	3	17 1 50,37	0,11	1 50,34	1 50,05	+ 0,03	+ 0,32	1,242		
618	5	37 Ophiuchi	6	17 4 30,04	0,29		4 29,89		+ 0,15	2,821		
619	4,5	A Ophiuchi	3	17 4 58,00	0,24		4 58,29		- 0,29	3,671		
620	3,4	a Herculis	8	17 6 56,73	0,34	6 56,71	6 56,29	+ 0,02	+ 0,44	2,729		
621	4,5	o Ophiuchi	3	17 7 56,05	0,04	7 55,92	7 56,57	+ 0,13	- 0,52	3,074		
622	3,4	π Herculis	5	17 9 9,91	0,26	9 9,97	9 9,66	- 0,06	+ 0,25	2,086		
623	4,5	ρ Ophiuchi	5	17 10 52,86	0,33	10 52,82	10 52,49	+ 0,04	+ 0,37	3,567		
624	3,4	θ Ophiuchi	5	17 11 38,46	0,14	11 38,41	11 38,21	+ 0,05	+ 0,25	3,672		
625	5	d Ophiuchi	4	17 16 34,26	0,66		16 34,28		- 0,02	3,817		
626	4	ρ Herculis	3	17 17 51,38	0,43	17 51,43	17 50,97	- 0,05	+ 0,41	2,067		
627	4,5	σ Ophiuchi	4,5	17 18 7,96	0,54	18 8,13	18 7,92	- 0,17	+ 0,04	2,969		
628	3	λ Scorpii	3	17 22 8,73	0,42	22 8,59	22 8,71	+ 0,14	+ 0,02	4,060		
629	4,5	λ Herculis	6	17 23 54,69	0,23	23 54,70	23 54,52	- 0,01	+ 0,17	2,417		
630	5	* Scorpii	4	17 24 55,45	0,48		24 55,05		+ 0,40	4,119		
631	2	a Ophiuchi	4	17 27 5,50	0,23	27 5,62	27 5,20	- 0,12	+ 0,30	2,770		
632	5	ξ Serpentis	4	17 27 54,99	0,41		27 54,68		+ 0,31	3,430		
633	5	μ Ophiuchi	6	17 28 39,78	0,53	28 39,77	28 39,33	+ 0,01	+ 0,45	3,254		
634	3	κ Scorpii	3	17 30 48,55	0,18		30 48,18		+ 0,37	4,139		
635	4,5	o Serpentis	2	17 31 55,39	0,06	31 55,27	31 55,00	+ 0,12	+ 0,39	3,369		
636	4	ι Herculis	5	17 34 41,89	0,22	34 41,90	34 41,11	- 0,01	+ 0,78	1,688		
637	3	β Ophiuchi	5	17 35 7,63	0,19	35 7,70	35 7,38	- 0,07	+ 0,25	2,960		
638	4,5	ι Scorpii	3	17 35 46,56	0,17		35 45,89		+ 0,67	4,185		
639	5	P Sagittarii	5	17 36 55,65	0,60	36 55,65	36 55,27	0,00	+ 0,38	3,768		
640	4	ν Ophiuchi	3	17 39 25,33	0,12	39 25,36	39 25,00	- 0,03	+ 0,33	3,003		
641	4	μ Herculis	6	17 39 50,92	0,15	39 50,97	39 50,20	- 0,05	+ 0,72	2,366		

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from		Annual Variation.
				<i>h.</i>	<i>m.</i>	<i>s.</i>		<i>s.</i>	<i>m.</i>	<i>s.</i>	<i>m.</i>	<i>s.</i>	Greenwich.	
642	5	* Sagittarii	5	17	48	14,43	0,31			48	14,19		+ 0,24	+ 3,845
643	5	b Sagittarii....	5	17	49	28,60	0,73			49	28,62		- 0,02	3,656
644	4	v Ophiuchi....	5	17	49	43,54	0,23	49	43,55	49	43,27	- 0,01	+ 0,27	3,297
645	4	o Hercules.....	3	17	50	27,63	0,06	50	27,67	50	26,94	- 0,04	- 0,31	2,052
646	2	γ Draconis.....	2	17	52	41,03	0,42	52	41,17	52	40,78	- 0,14	+ 0,25	1,388
647	5	τ Ophiuchi....	5	17	53	53,05	0,22			53	52,49		+ 0,56	3,260
648	4	γ* Sagittarii....	5	17	54	57,32	0,20	54	57,42	54	57,54	- 0,10	- 0,22	3,852
649	5	Q Hercules....	4	17	55	9,83	1,07			55	9,13		+ 0,70	2,560
650	4,5	p Ophiuchi....	5	17	56	55,06	0,24	56	55,01	56	54,03	+ 0,05	+ 0,13	3,009
651	4	S* Ophiuchi...	5	17	59	20,16	0,38			59	20,00		+ 0,16	2,843
652	4	o Hercules....	6	18	0	57,24	0,30	0	57,24	0	56,90	0,00	+ 0,34	2,335
653	3,4	a* Sagittarii...	5	18	3	39,68	0,33	3	39,64	3	39,20	+ 0,04	+ 0,48	3,583
654	5	A Hercules....	6	18	5	32,70	0,86			5	32,44		+ 0,26	2,254
655	3,4	δ Sagittarii....	5	18	10	10,60	0,19	10	10,59	10	10,26	+ 0,01	+ 0,34	3,835
656	5	G Hercules....	5	18	12	13,49	0,57			12	12,95		+ 0,54	2,463
657	3	a Sagittarii....	5	18	12	57,45	0,30	12	57,37	12	57,30	- 0,12	- 0,05	3,983
658	4	λ Sagittarii....	6	18	17	32,52	0,30	17	32,47	17	32,42	+ 0,05	+ 0,10	3,704
659	5	κ Cygni Sob....	6	18	18	33,86	0,58							3,416
660	3	δ Ursæ Min...	18	18	26	48,31	2,81	26	49,37	26	41,38	- 1,06	+ 6,93	- 19,168
661	1	a Lyrae.....	15	18	31	13,20	0,50	31	13,11	31	12,78	+ 0,09	+ 0,42	+ 2,010
662	5	o Aquilæ.....	6	18	33	1,32	0,12			33	1,13		+ 0,19	3,282
663	1,5	β Sagittarii....	3	18	35	5,87	0,04	35	5,80	35	6,14	+ 0,07	+ 0,27	3,745
664	5	K Hercules....	6	18	38	23,30	0,59			38	22,88		+ 0,42	2,578
665	5	a Lyrae.....	2	18	38	44,60	0,10			38	43,91		+ 0,69	1,982
666	3	β Lyrae.....	9	18	43	50,63	0,29	43	50,58	43	50,00	+ 0,05	+ 0,63	2,211
667	5	v* Sagittarii...	7	18	43	57,95	0,25	43	57,89	43	57,64	+ 0,06	+ 0,31	3,623
668	3	σ Sagittarii...	5	18	44	47,02	0,64	44	46,99	44	47,01	+ 0,03	+ 0,01	3,722
669	5	v* Sagittarii...	3	18	44	53,95	0,39	44	54,18	44	54,07	- 0,23	- 0,12	3,621
670	5	O Hercules....	5	18	47	37,28	0,52			47	36,91		+ 0,37	2,528
671	3,4	ζ Sagittarii...	6	18	51	51,38	0,30	51	51,44	51	51,19	- 0,06	+ 0,19	3,823
772	3	γ Lyrae.....	5	18	52	37,31	0,69	52	37,36	52	36,78	- 0,05	+ 0,53	2,240
673	4,5	o Sagittarii....	4	18	54	32,85	0,38	54	32,59	54	33,09	+ 0,26	- 0,24	3,592
671	3	ζ Aquilæ.....	14	18	57	38,67	0,46	57	38,61	57	38,11	+ 0,06	+ 0,56	2,754
675	5	B Aquilæ.....	5	19	3	30,83	0,40			3	30,27		+ 0,56	3,254
676	5	d Sagittarii....	8	19	7	44,73	0,66	7	44,69	7	44,36	+ 0,04	+ 0,37	3,514
677	5	η Lyrae.....	3	19	8	0,23	0,65			7	59,96		+ 0,27	2,038
678	5	n Draconis....	2	19	8	28,32	0,12			8	27,78		+ 0,54	1,133
679	5	l Vulpeculæ...	6	19	8	57,24	0,31			8	57,11		+ 0,13	2,576
680	3	δ Draconis....	10	19	12	29,63	0,36	12	29,69	12	29,12	- 0,06	+ 0,51	0,023
681	4	κ Cygni.....	5	19	13	11,59	0,38	13	11,72	13	11,18	- 0,13	+ 0,41	1,381
682	5	b Aquilæ.....	5	19	16	54,78	0,17			16	54,32		+ 0,46	2,871
683	3,4	δ Aquilæ....	6	19	16	58,66	0,19	16	58,58	16	58,33	+ 0,08	+ 0,33	3,007
684	4	τ Draconis....	4	19	19	46,80	0,47	19	47,27	19	46,74	- 0,47	+ 0,06	0,326
685	4	b Vulpeculæ...	5	19	21	40,53	0,36	21	40,55	21	39,80	- 0,02	+ 0,73	2,502
686	3	β* Cygni.....	5	19	23	54,47	0,38	23	54,42	23	54,06	+ 0,05	+ 0,41	2,416
687	5	K Aquilæ.....	5	19	25	48,59	0,57			25	48,14		+ 0,45	3,308

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.			Difference of Extremes:	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.		
				h.	m.	s.				Greenwich.	A.S.			
688	5	ι Aquilæ.....	4	19	27	58,52	0,43	27	58,62	—	0,10	+ 3,104		
689	5	σ Aquilæ.....	6	19	30	51,29	0,31	30	51,57	—	0,28	2,960		
690	4	θ Cygni.....	3	19	31	54,32	0,20	31	54,54	+ 0,22	+ 0,29	1,611		
691	4	α Sagittæ.....	3	19	32	32,64	0,33	32	32,71	— 0,07	+ 0,41	2,678		
692	5	β Sagittæ.....	2	19	33	27,60	0,08	33	27,33	+	0,27	2,691		
693	5	15 Cygni.....	6	19	38	10,99	0,36	38	10,69	+	0,30	2,154		
694	3	γ Aquilæ.....	24	19	38	13,55	0,47	38	13,44	0,00	+ 0,11	2,849		
695	3,4	δ Cygni.....	6	19	39	41,35	0,67	39	41,68	— 0,33	+ 0,03	1,868		
696	4	δ Sagittæ.....	4	19	39	51,23	0,09	39	51,27	— 0,04	+ 0,28	2,672		
697	5	χ Cygni.....	4	19	40	0,97	0,45	40	0,36	+	0,61	2,271		
698	1,2	α Aquilæ.....	29	19	42	32,30	0,36	42	32,29	42	32,11	+ 0,01	+ 0,19	2,924
699	4,5	E Sagittarij....	6	19	43	35,71	0,90	43	34,96	+	0,75	4,162		
700	5	ξ Aquilæ.....	3	19	46	3,55	0,23	46	3,06	+	0,49	2,899		
701	5	13 Vulpeculæ...	4	19	46	16,84	0,28	46	16,44	+	0,40	2,545		
702	3,4	β Aquilæ.....	23	19	47	0,75	0,26	47	0,79	— 0,04	+ 0,15	2,943		
703	5	22 Cygni.....	3	19	49	49,38	0,23	49	49,08	+	0,30	2,140		
704	4,5	γ Sagittæ.....	6	19	51	14,52	0,13	51	14,59	— 0,07	+ 0,14	2,660		
705	5	f Vulpeculæ...	7	19	51	55,57	0,43	51	54,96	+	0,61	2,576		
706	5	L ¹ Sagittarij....	2	19	53	36,18	0,13	53	35,03	+	1,15	3,818		
707	5	g Vulpeculæ...	4	19	54	8,71	0,43	54	8,09	+	0,62	2,462		
708	5	ρ Draconis.....	6	20	2	1,57	0,41	2	1,99	— 0,42	+ 0,94	0,304		
709	3,4	θ Aquilæ.....	6	20	2	35,05	0,58	2	35,02	2	34,88	+ 0,03	+ 0,17	3,095
710	5	66 Draconis....	4	20	2	50,75	0,48	2	50,02	+	0,73	0,952		
711	5	δ^2 Cygni.....	3	20	3	9,35	0,47	3	8,71	+	0,64	2,223		
712	4	α^1 Capricorni...	6	20	8	16,61	0,56	8	16,50	8	16,35	+ 0,11	+ 0,26	3,330
713	4	α^2 Cygni.....	3	20	8	18,72	0,29	8	18,70	8	18,34	+ 0,02	+ 0,38	1,886
714	3	α^2 Capricorni...	16	20	8	40,39	0,84	8	40,34	8	40,31	+ 0,05	+ 0,08	3,331
715	4,5	n Vulpeculæ...	5	20	8	46,07	0,29	8	46,12	8	45,76	— 0,05	+ 0,31	2,484
716	4,5	κ Cephei.....	5	20	14	25,59	0,49	14	29,49	14	26,03	— 3,90	— 0,44	1,882
717	3	γ Cygni.....	6	20	16	9,69	0,20	16	10,04	16	9,45	— 0,35	+ 0,24	2,148
718	5	h Cygni.....	6	20	17	6,73	0,45	17	6,63	+	0,10	2,387		
719	5	ρ Capricorni...	3	20	19	12,80	0,11	19	12,84	19	12,33	— 0,04	+ 0,47	3,432
720	5	G Aquilæ.....	6	20	20	48,91		20	48,56	+	0,35	3,134		
721	4,5	i Cygni.....	6	20	22	29,48	0,29	22	29,62	22	29,39	— 0,14	+ 0,09	2,446
722	5	w ² Cygni.....	6	20	24	49,48	0,32	24	48,90	+	0,58	1,854		
723	4	c Delphini.....	7	20	25	8,42	0,40	25	8,35	25	8,01	+ 0,07	+ 0,41	2,864
724	5	θ Cephei.....	3	20	26	44,06	0,36	26	43,04	+	1,02	1,016		
725	5	ζ Delphini....	6	20	27	24,52	0,34	27	23,98	+	0,54	2,800		
726	5	l Aquilæ.....	4	20	29	36,49	0,78	29	36,20	+	0,29	3,099		
727	4,5	θ Delphini....	6	20	30	45,50	0,26	30	45,43	30	44,93	+ 0,07	+ 0,57	2,829
728	3,4	a Delphini.....	5	20	31	47,38	0,16	31	47,65	31	47,27	— 0,27	+ 0,11	2,779
729	5	δ Delphini....	4	20	35	34,36	0,53	35	34,11	+	0,25	2,800		
730	1	a Cygni.....	31	20	35	40,46	0,63	35	40,42	35	39,93	+ 0,04	+ 0,53	2,040
731	4,5	ψ Capricorni...	5	20	36	4,76	0,59	36	4,72	36	5,01	+ 0,04	+ 0,25	3,572
732	4,5	e Aquarii.....	6	20	38	31,61	0,29	38	31,44	38	31,26	+ 0,17	+ 0,35	3,252
733	4	k Aquarii.....	3	20	38	48,87	0,19	38	48,99	38	48,53	— 0,12	+ 0,34	3,170

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.		Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
				h. m. s.	s.				Greenwich.	A.S.	
734	4	γ Delphini.....	6	20 38 49,25	0,33	38 49,16	38 49,93	+ 0,09	- 0,68	+ 2,783	
735	5	λ Cygni.....	4	20 40 49,96	0,17		40 49,08		+ 0,88	2,330	
736	4,5	μ Aquarii.....	6	20 43 32,23	0,34	43 32,05	43 31,83	+ 0,18	+ 0,40	3,230	
737	5	57 Cygni.....	5	20 47 16,19	0,33		47 15,90		+ 0,29	2,115	
738	4,5	η Vulpeculæ...	7	20 47 21,69	0,34		47 21,19		+ 0,50	2,552	
739	4	ν Cygni.....	6	20 50 52,36	0,27	50 52,65	50 52,13	- 0,29	+ 0,23	2,229	
740	5	η Capricorni...	3	20 54 46,67	0,09	54 46,72	54 46,64	- 0,05	+ 0,03	3,430	
741	5	γ Equulei.....	4	21 2 7,46	0,34		2 7,19		+ 0,27	2,912	
742	3	ζ Cygni.....	3	21 5 44,85	0,09	5 44,87	5 44,27	- 0,02	+ 0,58	2,546	
743	5	4 Piscis. Aust..	5	21 7 40,62	0,55		7 39,77		+ 0,85	3,658	
744	4,5	σ Cygni.....	7	21 10 47,00	0,24	10 47,10	10 45,91	- 0,10	+ 1,09	2,348	
745	5	ι Capricorni. . .	4	21 12 49,71	0,48	12 49,66	12 49,78	+ 0,05	- 0,07	3,350	
746	5	γ Indi.....	3	21 14 8,96	0,38		14 9,21		- 0,25	4,350	
747	3	α Cephei.....	3	21 14 32,70	0,46	14 32,44	14 31,66	+ 0,26	+ 1,04	1,416	
748	5	6 Cephei.....	7	21 15 51,23	0,74		15 50,55		+ 0,68	1,257	
749	4	ζ Capricorni...	6	21 17 0,44	0,56	17 0,36	16 59,96	+ 0,08	+ 0,48	3,441	
750	3	β Aquarii.....	6	21 22 39,66	0,34	22 39,39	22 39,31	+ 0,27	+ 0,35	3,162	
751	5	η Cygni.....	5	21 23 13,12	0,32		23 12,19		+ 0,93	2,200	
752	3	β Cephei.....	2	21 26 26,82	0,22	26 26,03	26 25,88	- 0,11	+ 0,94	0,811	
753	5	ϵ Capricorni...	6	21 27 36,64	0,41	27 36,62	27 36,41	+ 0,02	+ 0,23	3,372	
754	5	ρ Cygni.....	5	21 27 37,78	0,17	27 37,84	27 37,50	- 0,06	+ 0,28	2,248	
755	5	ξ Aquarii.....	5	21 28 44,97	0,17	28 44,93	28 45,02	+ 0,04	- 0,05	3,192	
756	4	γ Capricorni....	5	21 30 43,40	0,52	30 43,21	30 42,94	+ 0,19	+ 0,46	3,322	
757	5	41 Capricorni...	5	21 32 22,67	0,40		32 22,78		- 0,11	3,426	
758	5	κ Capricorni...	4	21 33 12,61	0,29	33 12,89	33 12,31	- 0,28	+ 0,30	3,353	
759	4,5	ι Piscis. Aust..	6	21 34 51,79	0,44	34 51,56	34 51,31	+ 0,23	+ 0,48	3,598	
760	2,3	α Pegasi.....	6	21 35 53,20	0,52	35 53,21	35 53,04	- 0,01	+ 0,16	2,942	
761	4,5	π^1 Cygni.....	3	21 36 6,25	0,27	36 6,19	36 5,61	+ 0,06	+ 0,64	2,118	
762	3,4	δ Capricorni...	6	21 37 42,40	0,14	37 42,28	37 42,11	+ 0,12	+ 0,29	3,304	
763	5	0 Piscis. Aust..	5	21 37 48,05	0,08		37 48,17		- 0,12	3,548	
764	5	π^2 Cygni.....	7	21 40 33,37	0,57	40 33,72	40 33,08	- 0,35	+ 0,29	2,204	
765	5	14 Pegasi.....	6	21 42 22,28	0,64		42 22,36		- 0,08	2,643	
766	4	γ Gruis.....	4	21 43 40,40	0,26		43 39,60		+ 0,80	3,657	
767	5	μ Capricorni...	6	21 44 4,47	0,38	44 4,64	44 4,00	- 0,17	+ 0,47	3,259	
768	5	δ Indi.....	4	21 46 22,00	0,37		46 20,91		+ 1,09	4,151	
769	5	ν Aquarii.....	6	21 54 34,31	0,27	54 34,27	54 33,83	+ 0,04	+ 0,48	3,104	
770	3	α Aquarii.....	9	21 57 6,12	0,50	57 6,18	57 5,96	- 0,06	+ 0,16	3,082	
771	5	ν Pegasi.....	5	21 57 9,37	0,25		57 9,23		+ 0,14	3,018	
772	4,5	ι Aquarii.....	8	21 57 18,22	0,39	57 18,19	57 18,24	+ 0,03	- 0,02	3,247	
773	2	α Gruis.....	4	21 57 32,74	0,24		57 32,84		- 0,10	3,818	
774	5	ξ Cephei.....	3	21 58 54,50	0,21		58 52,84		+ 1,66	1,699	
775	4	ι Pegasi.....	3	21 59 8,99	0,15	59 9,03	59 8,93	- 0,04	+ 0,06	2,761	
776	4	0 Pegasi.....	6	22 1 40,50	0,36	1 40,50	1 40,63	0,00	- 0,13	3,006	
777	5	π^1 Pegasi.....	2	22 1 44,68	0,11		1 45,06		- 0,38	2,650	
778	4	π^2 Pegasi.....	7	22 2 29,34	0,76	2 29,33	2 29,61	+ 0,01	- 0,27	2,653	
779	5	μ^1 Gruis.....	4	22 5 24,41	0,35		5 23,42		+ 0,99	3,649	

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.			Differ-ence of Ex-tremes.	Green-wich Catalogue		A.S. Cata-logue.		Difference from		Annual Variation.
				h.	m.	s.		m.	s.	m.	s.	Green-wich.	A.S.	
780	5	μ^2 Gruis.....	6	22	6	14,99	0,42			6	14,76		+ 0,23	+ 3,651
781	4,5	θ Aquarii.....	6	22	7	54,79	0,39	7	54,61	7	54,44	+ 0,18	+ 0,35	3,163
782	5	a Lacertæ.....	5	22	8	36,87	0,52			8	35,95		+ 0,92	2,599
783	4,5	e Cephei.....	2	22	8	49,53	0,26	8	49,40	8	48,91	+ 0,13	+ 0,62	2,137
784	5	β Pegasi.....	6	22	11	57,41	0,22			11	57,04		+ 0,37	3,016
785	4,5	d Pegasi.....	7	22	13	12,22	0,37	13	12,20	13	12,09	+ 0,02	+ 0,13	2,947
786	4,5	b Lacertæ.....	4	22	14	3,46	0,50			14	2,86		+ 0,60	2,458
787	7	* Aquarii.....	4	22	14	39,93	0,28			14	39,64		+ 0,29	3,152
788	5	π Aquarii.....	7	22	16	38,81	0,28			16	38,48		+ 0,33	3,063
789	5	d Lacertæ.....	4	22	17	40,67	0,66			17	39,58		+ 1,09	2,413
790	4	δ^1 Gruis.....	3	22	19	8,20	0,61			19	8,01		+ 0,19	3,625
791	4	ζ Aquarii.....	5	22	20	7,73	0,35	20	7,74	20	7,46	- 0,01	+ 0,27	3,077
792	4	β Piscis. Aust..	4	22	21	52,78	0,12	21	52,87	21	52,81	- 0,09	- 0,03	3,431
793	4,5	δ Cephei.....	6	22	22	54,54	0,41	22	54,68	22	53,65	- 0,14	+ 0,89	2,204
794	4	g Lacertæ.....	3	22	24	20,65	0,04	24	20,73	24	20,45	- 0,08	+ 0,20	2,436
795	5	v Aquarii.....	5	22	25	26,32	0,13	25	26,36	25	26,23	- 0,04	+ 0,09	3,280
796	4	η Aquarii.....	6	22	26	40,38	0,18	26	40,34	26	40,06	+ 0,04	+ 0,32	3,077
797	4	18 Piscis. Aust..	6	22	31	17,68	0,53	31	17,64	31	17,55	+ 0,04	+ 0,13	3,336
798	5	31 Cephei.....	4	22	31	35,16	0,43			31	35,30		- 0,14	1,447
799	3	β Gruis.....	6	22	32	32,26	0,38			32	32,24		+ 0,02	3,617
800	5	30 Cephei.....	4	22	32	40,72	0,85			32	40,34		+ 0,38	2,105
801	3	ζ Pegasi.....	6	22	33	2,18	0,42	33	2,25	33	1,92	- 0,07	+ 0,26	2,981
802	5	ξ Pegasi.....	6	22	38	15,36	0,52			38	14,98		+ 0,38	2,975
803	4	e Gruis.....	3	22	38	18,37	0,11			38	17,35		+ 1,02	3,679
804	4,5	λ Pegasi.....	6	22	38	23,94	0,15	38	23,96	38	24,35	- 0,02	- 0,41	2,873
805	4	μ Pegasi.....	5	22	41	51,31	0,20	41	51,29	41	50,53	+ 0,02	+ 0,78	2,872
806	5	22 Piscis. Aust..	5	22	43	6,86	0,29			43	6,75		+ 0,11	3,362
807	4	λ Aquarii.....	7	22	43	47,74	0,23	43	47,65	43	47,26	+ 0,09	+ 0,48	3,133
808	3	δ Aquarii.....	6	22	45	40,53	0,36	45	40,55	45	40,34	- 0,02	+ 0,19	3,196
809	5,6	ρ Pegasi.....	2	22	46	43,53	0,16			46	42,58		+ 0,95	3,010
810	1	a Piscis. Aust..	23	22	48	17,69	0,50	48	17,65	48	17,65	+ 0,04	+ 0,04	3,311
811	5	ζ Gruis.....	5	22	50	51,73	0,57			50	51,17		+ 0,56	3,608
812	6	χ^2 Piscium....	4	22	51	57,98	0,15			51	57,86		+ 0,12	3,073
813	4	o Andromedæ..	6	22	54	9,75	0,43	54	9,74	54	9,04	+ 0,01	+ 0,71	2,734
814	5	β Piscium.....	6	22	55	16,79	0,40	55	16,66	55	16,30	+ 0,13	+ 0,49	3,049
815	2	β Pegasi.....	2	22	55	35,39	0,07	55	35,48	55	35,25	- 0,09	+ 0,14	2,878
816	2	a Pegasi.....	17	22	56	20,91	0,39	56	20,96	56	20,68	- 0,05	+ 0,23	2,975
817	5	θ Gruis.....	4	22	57	20,05	0,21			57	19,95		+ 0,10	3,422
818	4,5	h Pegasi.....	6	22	58	53,37	0,37	58	53,48	58	53,34	- 0,11	+ 0,03	2,907
819	4,5	c^2 Aquarii.....	6	23	0	25,71	0,38	0	25,57	0	25,57	+ 0,14	+ 0,14	3,208
820	5	i Gruis.....	3	23	0	45,82	0,69			0	42,20		+ 3,62	3,424
821	5	π Cephei.....	2	23	2	32,45	0,77			2	32,23		+ 0,22	1,875
822	5	w Andromedæ..	6	23	4	49,68	0,20			4	49,16		+ 0,52	2,708
823	5	ϕ Aquarii.....	6	23	5	34,09	0,17	5	34,18	5	33,84	- 0,09	+ 0,25	3,106
824	4	γ Toucanæ....	4	23	7	30,93	0,53			7	30,06		+ 0,87	3,577
825	4,5	γ Piscium.....	5	23	8	24,26	0,52	8	24,35	8	24,15	- 0,09	+ 0,11	3,108

No.	Mag.	NAMES.	No. of Observations.	Observed A.R. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.	
				h.	m.	s.				Greenwich.	A.S.		
826	5	γ App. Sculp...	6	23	9	41,14	0,48		9	40,92	+ 0,22	+ 3,261	
827	5	δ Andromedæ...	3	23	9	55,85	0,28		9	55,23	+ 0,62	2,745	
828	5	ψ^3 Aquarii....	5	23	10	10,07	0,48	10 10,10	10	10,81	- 0,03	3,122	
829	5	τ Pegasi.....	5	23	12	17,04	0,33		12	16,50	+ 0,54	2,952	
830	5	b^1 Aquarii....	6	23	14	5,28	0,22		14	4,69	+ 0,59	3,170	
831	5	v Pegasi.....	6	23	16	57,40	0,54		16	56,89	+ 0,51	2,965	
832	5	b^2 Aquarii....	5	23	17	9,63	0,68		17	9,04	+ 0,59	3,166	
833	5	θ Piscium.....	6	23	19	23,94	0,11		19	23,31	+ 0,63	3,046	
834	5	s Cephei.....	5	23	20	10,54	0,70					2,452	
835	5	* Cassiopeæ...	4	23	22	15,58	0,27		22	14,99	+ 0,59	2,722	
836	7	ω^3 Piscium....	3	23	23	17,39	0,58		23	16,76	+ 0,63	3,076	
837	5	b^4 Aquarii....	6	23	24	25,57	0,34		24	25,46	+ 0,11	3,151	
838	5	y Pegasi.....	5	23	25	1,02	0,34		25	0,66	+ 0,36	2,988	
839	5	t Phœnicis....	5	23	25	57,44	0,49		25	57,63	- 0,19	3,256	
840	4,5	λ Andromedæ..	4	23	29	19,11	0,24	29 19,08	29	18,40	+ 0,03	2,886	
841	5	θ Phœnicis....	7	23	30	21,03	0,53		30	21,32	+ 0,61	3,257	
842	5	ω^4 Aquarii....	6	23	31	0,96	0,20		31	0,61	- 0,08	3,114	
843	4,5	i Piscium.....	6	23	31	15,60	0,33	31 15,68	31	15,61	- 0,01	3,054	
844	5	κ Andromedæ..	3	23	32	6,41	0,68		32	6,05	+ 0,36	2,914	
845	5	A^3 Aquarii....	6	23	35	25,95	0,31		35	25,65	+ 0,30	3,118	
846	5	i Pegasi.....	6	23	35	30,31	0,56		35	30,49	- 0,18	2,991	
847	7	* Piscium.....	4	23	36	11,78	0,72		36	11,18	+ 0,60	3,053	
848	5	ψ Andromedæ..	6	23	37	41,05	0,57		37	41,89	- 0,84	2,936	
849	5	τ Cassiopeæ....	4	23	38	50,10	0,30		38	49,36	+ 0,74	2,873	
850	5	* Draconis.....	3	23	39	53,07	0,24		39	52,79	+ 0,28	2,793	
851	5	δ App. Sculp...	6	23	40	6,88	0,20		40	6,42	+ 0,46	3,133	
852	6	ϵ^1 Piscium.....	7	23	40	48,50	0,65		40	48,35	+ 0,15	3,068	
853	6,7	* Piscium.....	4	23	46	7,89	0,33		46	7,81	+ 0,08	3,069	
854	6	ϵ^2 Piscium.....	6	23	46	29,38	0,51		46	29,07	+ 0,31	3,060	
855	5	η Toucanæ....	6	23	48	39,54	0,38		48	38,43	+ 1,11	3,212	
856	5	p Piscium.....	5	23	50	1,33	0,35	50 1,23	50	0,63	+ 0,10	+ 0,70	3,073
857	4,5	ω Piscium.....	5	23	50	38,34	0,20	50 38,23	50	38,28	+ 0,11	+ 0,06	3,062
858	5	e Toucanæ....	3	23	51	3,69	0,67		51	3,89	- 0,20	3,189	
859	5	q Piscium.....	4	23	53	9,91	0,41	53 9,95	53	9,04	- 0,04	+ 0,87	3,071
860	4,5	r Piscium.....	6	23	53	17,66	0,27	53 17,67	53	17,19	- 0,01	+ 0,47	3,073
861	4	g Ceti.....	6	23	55	4,70	0,56	55 4,69	55	4,64	+ 0,01	+ 0,06	3,078
862	5	s Piscium.....	7	23	56	41,13	0,24	56 44,16	56	40,82	- 0,03	+ 0,31	3,070
863	7	5 Ceti.....	4	23	59	32,78	0,59		59	32,70		+ 0,08	3,068
864	1	α Andromedæ..	29	23	59	40,17	0,54	59 40,12	59	39,66	+ 0,05	+ 0,51	3,067

In the above 423 comparisons between the Madras and Greenwich Catalogues there are

150	Cases in which the difference is within 0",05
122	— — — — — 0",10
60	— — — — — 0",15
45	— — — — — 0",20

i. e. Out of the 423 comparisons, there are 376 cases in which the difference does not amount to two-tenths of a second of time; of the remaining 46 there are 34 which are within three-tenths of a second; these I have carefully re-examined, and find them affected with a much less *probable* error than this amount; the disagreement no doubt arises from the discrepancies in either Catalogue acting with contrary signs, the 12 cases which exceed three-tenths of a second, and seem to merit some further attention are as follows.

No.		A.R.	N.P.D.	Difference.
		<i>h. m.</i>	<i>° ′</i>	<i>s.</i>
1	... ϵ Cassiopeæ.....	1 42	27 20	+ 0,62
2	...* Camelopardali.....	3 15	30 39	+ 0,48
3	... β Eridani.....	4 59	95 18	- 0,32
4	... π Virginis... ..	11 52	82 26	+ 0,69
5	... δ Sagittarii	17 49	113 47	+ 0,69
6	... δ Ursæ Minoris.....	18 26	3 24	- 1,06
7	... π Draconis	19 19	24 36	- 0,47
8	... δ Cygni...†.....	19 39	45 16	- 0,33
9	... ρ Draconis.....	20 2	22 36	- 0,42
10	... κ Cephei.....	20 14	12 48	- 3,90
11	... γ Cygni... ..	20 16	50 17	- 0,35
12	... π ² Cygni.....	21 40	41 28	- 0,35

No. 1.—Comparing the Greenwich places of ψ and δ Cassiopeæ (Stars situated in the neighbourhood of ϵ) with the Astronomical Society's Catalogue; it appears that the places in the former Catalogue are in excess 1,02s. and 0,34s. respectively, whereas the place of ϵ is only 0,02s. in excess; or in other words the Astronomical Society's Catalogue agrees with the Madras Catalogue in assigning to ϵ Cassiopeæ a place above half a second less than that given in the Greenwich Catalogue.

No. 2.—From a comparison similar to the above, the testimony of * Camelopardali (which follows the Stars under consideration only 1m. 2s. and is situated within 1°,5 of the same in Polar Distance) is in favor of the Madras Catalogue.

No. 3, 8, 11, 12, require to be re-examined before any thing decisive can be said.

No. 4, 6, 10, the Astronomical Society's Catalogue so strongly confirms the Madras result that we may very safely conclude the error to rest with the Greenwich Catalogue.

No. 5.—The proximity of this Star to the Pole fully accounts for the difference, which is somewhat less than $^{\circ}07$ of a great Circle.

On inspecting the column "difference from A.S." the numbers though not so small as those in the preceding column, are nevertheless much within the limits which might be expected from Catalogues brought forward from the observations of 1755 and 1800; out of 863 comparisons which this column affords, there are 615 which do not exceed half a second of time, and among the remaining 248 there are only 51 which exceed one second of time; the latter with one or two exceptions only have been brought forward from the observations of La Caille, these having been made with Instruments capable of but limited accuracy, can throw very little, if any doubt upon the Madras results; not willing however to think too favorably of the Madras Catalogue, I have from the observations of the present year examined a few of the most discordant results, and propose with the observations of next year to furnish the remainder.

The cases most deserving notice are as follows:

No.	Disagrees to the amount of the Annual Variation.				
94	—	—	—	—	—
118	—	—	—	8",65	Madras result confirmed by the observations of 1832.
142	—	—	—	2",09	Do. Do.
167	—	—	—	3",19	Madras result confirmed by the Greenwich Catalogue.
299	—	—	—	8",53	
331	—	—	—	— 1",01	} Madras result confirmed by the observations of 1832.
332	—	—	—	+ 1",66	
338	—	much from <i>h</i> and <i>o</i> of the same constellation.			
341	—	to the amount of		3",15	Madras result confirmed by the observations in 1832.
343	—	—	—	1",56	Do. Do.
352	—	—	—	1",37	
369	—	—	—	1",61	
372	—	—	—	1",79	
387	—	—	—	2",15	} Madras result confirmed by the observations of 1832.
389	—	—	—	1",80	
396	—	—	—	1",08	
427	—	—	—	1",98	
451	—	—	—	1",15	
452	—	—	—	2",06	
461	—	—	—	1",36	
478	—	—	—	1",28	
543	—	—	—	6",35	Madras result confirmed by the observations of 1832.
562	—	—	—	1",24	
563	—	—	—	1",18	
588	—	—	—	1",28	

No. 660. Differs to the amount of $6''{,}93$ This disagreement is but small considering the proximity of the Star to the Pole.

774 — — — $1''{,}66$
 822 — — — $3''{,}62$ or the Annual Variation.

MEMORANDUM.—It must be recollected that Right Ascension in the Madras Catalogue is reckoned from the Equinoctial point as assumed in the construction of the Greenwich Catalogue, which is the point assumed by Dr. Maskelyne $+ 0,20s$. Now Dr. Maskelyne having assumed the place of α Aquilæ from Bradley and determined the position of the other Stars with reference to this Star, it follows that the places of Stars in the Astronomical Society's Catalogue are not reckoned from the same point as that assumed in the construction of the Madras Catalogue. If Piazzi's Catalogue is reckoned from the Equinoctial point as assumed by Dr. Maskelyne, (in order to render the two Catalogues strictly comparable) it is necessary to subtract $0,20s$. from the Madras Catalogue or add $0,20s$. to the Astronomical Society's Catalogue.

North Polar Distance of Stars from the Madras Catalogue, compared with the Greenwich, and Astronomical Society's Catalogue.

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Differ-ence of Ex-tremes.	Green-wich Catalogue	A.S. Cata-logue.	Difference from		Annual Variation.	
					a	l	u				s.	m. s.		m. s.
1	2,3	β Cassiopeæ....	0 0	5	31	46	57,10	2,08	46 58,91	46 59,51	-1,81	-	2,41	20,043
2	4	ε Phœnicis....	0 1	5	136	40	44,34	1,45		40 45,66	-	-	1,32	20,042
3	5	B Andromedæ..	0 1	5	44	52	10,06	3,64		52 6,50	+	+	3,56	20,042
4	6	f Ceti.....	0 3	4	106	23	43,46	2,35		23 48,00	-	-	4,54	20,041
5	2,3	γ Pegasi.....	0 4	22	75	45	23,71	4,70	45 22,92	45 21,85	+0,79	+	1,86	20,039
6	6	χ Pegasi.....	0 6	5	70	44	0,56	1,57		43 55,14	+	+	5,42	20,036
7	5	θ Andromedæ..	0 8	4	52	15	25,62	3,49		15 25,26	+	+	0,36	20,030
8	6,7	36 Piscium....	0 8	3	82	41	53,74	2,30		41 52,34	+	+	1,40	20,031
9	6,7	33 Piscium....	0 9	5	89	14	56,38	2,34		15 4,54	-	-	8,16	20,027
10	4	i Ceti.....	0 11	5	99	45	43,46	3,65	45 40,55	45 37,10	+2,91	+	6,36	20,021
11	5	ζ Tucanæ....	0 13	5	155	52	5,56	3,94		54 7,80	-	-	122,24	20,019
12	6	40 Piscium....	0 11	5	74	41	17,22	2,89		41 12,60	+	+	4,62	20,018
13	5,6	d Piscium....	0 12	6	82	44	53,40	3,53	44 56,50	44 53,94	-3,10	-	0,54	20,016
14	6	9 Ceti.....	0 14	3	103	8	56,53	1,30		8 58,54	-	-	2,01	20,004
15	6,7	* Ceti.....	0 16	5	93	9	12,53	2,38		9 16,47	-	-	3,94	19,995
16	5	κ Phœnicis....	0 18	5	134	37	6,72	2,81		37 0,36	+	+	6,36	19,982
17	2	α Phœnicis....	0 18	8	133	13	25,01	4,88		13 16,17	+	+	8,84	19,982
18	6	10 Ceti.....	0 18	5	90	59	10,27	1,35		59 7,35	+	+	2,92	19,981
19	6	47 Piscium....	0 19	8	73	2	32,93	2,45		2 34,05	-	-	1,12	19,972
20	5	λ Cassiopeæ....	0 22	5	36	24	41,47	3,84		24 31,80	+	+	9,67	19,947
21	5	λ ¹ Phœnicis...	0 23	5	139	44	21,54	1,27		43 26,36	+	+	55,13	19,940
22	4	κ Cassiopeæ....	0 23	5	28	0	4,64	2,13	0 8,34	0 9,70	-3,70	-	5,06	19,938
23	6,7	51 Piscium....	0 24	4	83	58	40,60	1,17		58 39,84	+	+	0,76	19,936
24	5	β ³ Tucanæ....	0 23	4	153	57	42,79	2,91		57 47,50	-	-	4,71	19,924
25	6,7	* Piscium....	0 27	3	77	33	29,33	1,55		33 31,43	-	-	2,10	19,913
26	4	ζ Cassiopeæ....	0 28	5	37	2	4,93	3,36	2 3,47	2 2,84	+1,46	+	2,09	19,898
27	4,5	π Andromedæ..	0 28	6	57	12	44,43	1,84	12 43,34	12 41,70	+1,09	+	2,73	19,895
28	6	53 Piscium....	0 28	5	76	41	57,45	1,62		41 54,96	+	+	2,49	19,894
29	4	ε Andromedæ..	0 30	4	61	36	25,09	0,63	36 24,00	36 24,99	+1,09	+	0,10	19,876
30	3	δ Andromedæ..	0 30	5	60	3	57,42	2,08	3 52,59	3 51,41	+4,83	+	6,01	19,868
31	3	α Cassiopeæ....	0 31	16	34	23	28,60	4,62	23 27,34	23 27,35	+1,26	+	1,25	19,861
32	6	55 Piscium....	0 31	5	69	29	23,49	1,40		29 21,94	+	+	1,55	19,860
33	6,7	* Ceti.....	0 32	4	95	16	51,57	1,27		16 47,42	+	+	4,15	19,847
34	5	μ Phœnicis....	0 33	5	137	0	48,63	3,37		0 38,13	+	+	10,50	19,832
35	6	* Ceti.....	0 33	4	102	43	50,45	0,82		43 53,23	-	-	2,78	19,827
36	5	π Cassiopeæ....	0 34	5	43	54	3,85	1,88		54 5,85	-	-	2,00	19,821
37	2,3	β Ceti.....	0 35	5	108	54	53,04	2,60	54 54,73	54 55,02	-1,69	-	1,98	19,809
38	5	φ ¹ Ceti.....	0 35	4	101	31	50,22	1,55		31 53,37	-	-	3,15	19,801
39	5	η Phœnicis....	0 35	5	148	23	23,22	1,57		23 23,34	-	-	0,11	19,800
40	6	* Ceti.....	0 36	3	112	56	6,09	1,82		56 11,62	-	-	5,53	19,791
41	4	ζ Andromedæ..	0 38	5	66	39	10,61	2,14	39 12,56	39 12,60	-1,95	-	1,99	19,763
42	6	60 Piscium....	0 38	4	84	10	58,08	3,09		10 53,44	+	+	4,64	19,759
43	4	η Cassiopeæ....	0 39	5	33	4	58,74	1,05	5 1,23	4 59,73	-2,49	-	0,99	19,035

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
					h. m. s.	s.	m. s.				m. s.	Greenwich.	
44	5	δ Piscium.....	0 40	5	83 20 7,65	3,43		20 6,35		+ 1,30		19,740	
45	4	ν Andromedæ..	0 40	5	49 50 39,31	1,63	50 34,75	50 37,02	+ 4,56	+ 2,29		19,731	
46	6	i Piscium.....	0 41	5	63 12 39,61	3,18		12 42,39		— 2,78		19,726	
47	5	m Ceti.....	0 44	5	92 3 48,60	1,31	3 48,83	3 48,53	— 0,23	+ 0,07		19,669	
48	6	66 Piscium....	0 45	5	71 43 44,76	2,40		43 47,00		— 2,24		19,647	
49	6	36 Andromedæ..	0 46	5	67 17 16,18	0,84		17 17,20		— 1,02		19,642	
50	3	γ Cassiopeæ....	0 46	9	30 12 5,23	4,39	12 2,08	11 57,11	+ 3,15	+ 8,12		19,632	
51	5	* Cephei.....	0 47	5	4 39 17,97	1,14		39 19,21		— 1,24		19,625	
52	4	μ Andromedæ..	0 47	5	52 25 7,59	1,95	25 8,87	25 8,73	— 1,28	— 1,14		19,617	
53	5	η Andromedæ..	0 48	5	67 29 43,51	1,03		29 45,27		— 1,76		19,602	
54	6	h Piscium.....	0 49	4	61 55 22,17	3,43		55 23,50		— 1,33		19,593	
55	5	a App. Sculp...	0 50	5	120 16 22,26	2,33		16 19,69		+ 2,57		19,560	
56	7	* Piscium.....	0 54	5	82 5 20,94	2,90		5 18,11		+ 2,83		19,496	
57	4	e Piscium.....	0 54	7	83 1 15,37	3,06	1 17,31	1 17,31	— 1,94	— 1,94		19,486	
58	5,6	ψ * Piscium....	0 57	5	69 26 0,67	2,19		26 0,66		+ 0,01		19,435	
59	3,4	β Phœnicis	0 58	5	137 37 31,34	1,52		37 22,21		+ 9,13		19,394	
60	2,3	a Ursæ Minoris.	0 59	10	1 35 31,54		35 32,00	35 31,58	— 0,46	— 0,04		19,375	
61	5	e Piscium.....	1 0	5	85 14 50,11	1,66	14 47,76	14 47,38	+ 2,35	+ 2,73		19,368	
62	5	φ Andromedæ..	1 0	5	43 39 38,59	3,37		39 41,39		— 2,80		19,367	
63	3,4	η Ceti.....	1 0	5	101 4 46,76	3,35	4 48,70	4 44,44	— 1,94	+ 2,32		19,359	
64	2	β Andromedæ..	1 0	5	55 16 42,29	3,24	16 38,81	16 36,97	+ 3,48	+ 5,32		19,355	
65	4,5	θ Cassiopeæ....	1 1	4	35 45 6,35	1,80	45 6,05	45 7,75	+ 0,30	— 1,40		19,341	
66	5	χ Piscium.....	1 2	6	69 52 1,29	3,53		51 57,48		+ 3,81		19,306	
67	6,7	34 Ceti.....	1 3	3	93 8 57,60	1,29		9 1,55		— 3,95		19,288	
68	6	b Ceti.....	1 6	5	98 49 55,18	3,26		49 55,49		— 0,31		19,221	
69	6	39 Ceti.....	1 8	4	93 23 27,64	1,44		23 27,57		+ 0,07		19,167	
70	6	f Piscium.....	1 9	4	87 16 38,69	2,13		16 33,54		+ 5,15		19,140	
71	6	l Piscium.....	1 12	3	62 8 49,40	1,20		8 50,23		— 0,83		19,069	
72	5	ξ Andromedæ..	1 12	7	45 21 37,50	4,19		21 30,29		+ 7,21		19,052	
73	7	* Ceti.....	1 14	5	89 9 27,76	2,10		9 28,40		— 0,64		19,011	
74	4,5	ψ Cassiopeæ....	1 14	6	22 45 17,74	3,32	45 20,67	45 21,29	— 2,93	— 3,55		19,006	
75	3	δ Cassiopeæ....	1 15	6	30 38 48,51	3,98	38 51,08	38 44,77	— 2,57	+ 3,74		18,986	
76	3	θ * Ceti.....	1 16	4	99 3 25 56	2,59	3 27,80	3 26,22	— 2,24	— 0,66		18,964	
77	5	* Phœnicis	1 17	5	132 22 24,69	1,65		22 27,93		— 3,24		18,918	
78	5	c Ceti.....	1 17	5	105 28 51,54	2,59		28 45,53		+ 6,01		18,915	
79	6,7	* Ceti.....	1 18	3	91 16 44,14	1,65		17				18,900	
80	6,7	97 Piscium....	1 21	5	72 31 10,63	1,89		31 13,05		— 2,42		18,813	
81	3	γ Phœnicis	1 21	5	134 11 8,67	1,79		11 4,72		+ 3,95		18,805	
82	5	μ Piscium.....	1 21	4	84 43 50,84	1,68	43 42,54	43 52,68	+ 8,30	— 1,84		18,795	
83	4	η Piscium....	1 22	6	75 31 42,02	3,38	31 40,41	31 38,84	+ 1,61	+ 3,18		18,761	
84	4	δ Phœnicis	1 24	5	139 57 14,94	2,52		56 58,97		+ 15,97		18,706	
85	7	* Piscium.....	1 24	5	82 39 36,35	3,70		39 37,66		— 1,31		18,699	
86	6	* Piscium.....	1 25	5	72 24 16,35	1,74		24 12,23		+ 4,12		18,660	
87	5	ν Andromedæ..	1 27	5	49 26 36,83	2,33		26 34,51		+ 2,32		18,621	
88	3,4	R * Andromedæ.	1 28	5	42 13 53,95	2,17	13 53,21	13 52,94	+ 0,74	+ 1,01		18,596	
89	6	π Piscium.....	1 28	5	78 43 27,91	2,71	43 33,58	43 29,93	— 5,67	— 2,02		18,580	

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.		Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
					h.	m.				m.	s.	
90	1	α Eridani.....	1 31	7	148	5 49,08	1,13		5 53,05		- 4,57	- 18,471
91	7	* Piscium.....	1 32	3	82	5 56,35	2,34		15 56,64		- 0,29	18,462
92	5	ν Piscium.....	1 32	5	85	22 17,90	1,78	22 14,17	22 13,72	+ 3,73	+ 4,18	18,429
93	5	54 Andromedæ..	1 33	5	40	10 2,88	4,38		9 58,65		+ 4,03	18,413
94	5,6	107 Piscium....	1 33	4	70	33 22,29	2,63		33 16,71		+ 5,58	17,835
95	5,7	109 Piscium....	1 36	3	70	45 50,24	0,67		45 47,98		+ 2,26	18,322
96	3,4	γ Ceti.....	1 36	5	106	40 46,43	4,57	49 48,85	49 43,92	- 2,42	+ 2,51	19,144
97	5	σ Piscium.....	1 36	5	81	41 44,59	2,73	41 44,97	41 43,96	- 0,38	+ 0,63	18,295
98	5	ϵ App. Sculp...	1 38	5	175	53 58,71	2,02		53 59,46		- 0,75	18,250
99	6,7	δ Arietis.....	1 39	4	73	53 17,83	2,15		53 18,51		- 0,68	18,203
100	5	χ Ceti.....	1 41	5	101	31 31,65	2,90		31 31,98		- 0,33	18,119
101	5	54 Ceti.....	1 42	5	79	47 51,24	1,85		47 49,01		+ 2,23	18,096
102	3,4	α Cassiopeæ...	1 42	6	27	10 0,87	3,90	10 2,03	10 2,33	- 1,16	- 1,46	18,081
103	3	ζ Ceti.....	1 43	5	101	10 22,12	3,30	10 24,50	10 27,54	- 2,38	- 5,42	18,050
104	3,4	α Trianguli...	1 43	5	61	14 53,18	0,91	14 51,80	14 54,85	+ 1,18	- 1,67	18,037
105	4,5	γ Arietis.....	1 44	3	71	32 16,68	0,07	32 18,83	32 13,90	- 2,15	+ 3,48	18,006
106	5,0	ϵ Piscium.....	1 45	3	87	38 56,30	1,54		39 2,48		- 6,18	17,985
107	3	β Arietis.....	1 45	5	70	1 20,81	4,00	1 17,91	1 14,77	+ 2,90	+ 6,04	17,966
108	5	ϕ Phœnicis...	1 47	5	133	19 39,64	5,56		19 43,79		- 4,15	17,885
109	5	48 Cassiopeæ...	1 48	4	19	55 9,04	2,60		55 5,72		+ 3,32	17,852
110	5,6	λ Arietis.....	1 48	3	67	13 54,94	4,02	13 55,47	13 51,11	- 0,53	+ 3,83	17,839
111	4,5	50 Cassiopeæ...	1 49	5	18	24 6,15	2,94	24 7,21	24 8,32	- 1,06	- 2,17	17,815
112	4	χ Eridani.....	1 49	5	142	27 5,62	1,53		27 16,28		- 10,66	17,805
113	4,5	η Hydri.....	1 51	5	158	28 46,43	2,34		28 52,69		- 6,26	17,753
114	4,5	ν Ceti.....	1 52	4	111	53 58,79	1,78	54 9,68	53 55,78	- 10,89	+ 3,01	17,697
115	5	α Piscium.....	1 53	5	88	3 18,55	3,00	3 21,92	3 18,08	- 3,37	+ 0,47	17,645
116	3	α Hydri.....	1 53	4	152	23 37,59	2,10		23 27,46		+ 10,13	17,638
117	3,4	γ Andromedæ..	1 53	5	48	29 4,86	3,68	29 7,53	29 8,38	- 2,67	- 3,52	17,635
118	5	χ Phœnicis.....	1 55	4	135	31 47,01	3,58		31 57,53		- 10,52	17,576
119	5	* Arietis.....	1 57	3	64	58 50,83	4,17		59			17,483
120	3	α Arietis.....	1 58	30	67	20 23,11	3,74	20 26,17	20 24,08	- 1,06	+ 1,03	17,461
121	4	β Trianguli....	1 59	6	55	48 58,87	1,43	48 59,33	48 54,52	- 0,46	+ 4,35	17,381
122	6,7	64 Ceti.....	2 2	4	82	13 30,27	2,15		13 30,03		+ 0,24	17,252
123	5	ϵ Ceti.....	2 4	5	81	56 59,96	1,05	56 58,77	56 58,18	+ 1,19	+ 1,78	17,180
124	6	ϵ Ceti.....	2 9	5	97	12 17,19	3,46		12 16,22		+ 0,97	16,974
125	6	θ Arietis.....	2 9	4	70	53 4,15	2,64	53 4,23	53 3,33	- 0,08	+ 0,82	16,966
126	4	ϕ Eridani.....	2 10	5	142	17 50,12	2,85		17 50,72		- 0,60	16,883
127	5	i Persi.....	2 11	4	34	56 5,83	0,71		56 4,66		+ 1,19	16,877
128	6	69 Ceti.....	2 13	5	90	22 52,24	3,90		22 49,05		+ 3,19	16,750
129	6	70 Ceti.....	2 13	5	91	39 27,62	4,45		39 28,43		- 0,81	16,735
130	4,5	* Cassiopeæ 35 H.	2 15	5	23	21 49,61	2,42	21 50,78	21 53,38	- 1,17	- 3,77	16,656
131	6	71 Ceti.....	2 16	6	93	32 53,89	3,02		32 54,56		- 0,67	16,597
132	6	* Arietis.....	2 18	4	80	12 1,31	2,17		11 57,60		+ 4,21	16,535
133	5	ρ Ceti.....	2 18	5	103	3 20,40	1,67		3 19,33		+ 1,16	16,531
134	4	δ Hydri.....	2 19	5	159	25 51,93	4,09		25 52,58		- 0,65	16,481
135	5	ϵ Ceti.....	2 19	6	82	18 5,75	2,03	18 5,18	18 7,00	+ 0,57	- 1,25	16,461

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from		Annual Variation.		
					h.	m.	s.		m.	s.	m.	s.	Greenwich.	A.S.C.			
136	4.5	κ Eridani.....	2 20	5	138	27	53,16	1,30		28	0,22		-	7,06	-	16,382	
137	6,7	26 Arietis.....	2 21	5	70	53	58,05	2,58		53	53,75		+	4,30		16,362	
138	6	ν Arietis.....	2 21	5	73	2	50,48	3,94		2	48,67		+	1,81		16,343	
139	5	σ Ceti.....	2 24	5	105	59	25,45	2,27		59	18,03		+	7,42		16,213	
140	6	ε Ceti.....	2 26	5	98	36	1,53	3,88		36	1,49		+	0,04		16,094	
141	6	* Fornacis.....	2 26	5	118	58	40,18	1,21		58	41,14		-	0,96		16,090	
142	4,5	ν Ceti.....	2 27	5	85	9	0,02	2,50	8	54,46	8	52,30	+	5,56	+	7,72	16,062
143	4	δ Ceti.....	2 31	5	90	24	20,72	2,39	24	17,90	24	19,38	+	2,82	+	1,34	15,859
144	7	* Ceti.....	2 31	3	84	37	12,98	0,91		37	12,23		+	0,75		15,831	
145	4,5	ε Ceti.....	2 31	5	102	35	35,91	2,54	35	27,53	35	35,26	+	8,38	+	0,65	15,829
146	4	θ Persei.....	2 32	4	41	29	31,39	1,41	29	32,51	29	31,85	-	1,12	-	0,46	15,760
147	5	* Eridani.....	2 33	5	133	37	9,55	1,69		37	11,65		-	2,10		15,723	
148	4	α Arietis.....	2 33	4	63	0	57,85	0,48	1	2,14	1	1,04	-	4,29		3,19	15,713
149	4,5	κ Eridani.....	2 34	4	130	34	56,63	1,81		34	54,16		+	2,47		15,688	
150	3	γ Ceti.....	2 34	5	87	28	50,29	2,10	28	51,49	28	52,04	-	1,20	-	1,75	15,658
151	5,6	38 Arietis.....	2 36	3	78	16	8,10	1,90		16	9,30		-	1,20		15,593	
152	4	μ Ceti.....	2 36	5	80	36	17,67	2,50	36	14,43	36	9,61	+	3,24	+	8,06	15,590
153	4	π Ceti.....	2 36	5	104	34	45,87	2,87	34	44,28	34	32,92	+	1,59	+	12,95	15,574
154	4	b Arietis.....	2 38	5	61	27	38,11	3,83	27	36,89	27	35,73	+	1,22	+	2,38	15,476
155	5	* Persei.....	2 38	6	34	48	48,22	1,90		48	46,35		+	1,87		15,446	
156	5	π Arietis.....	2 40	5	73	14	33,52	2,86		14	34,06		-	0,54		15,364	
157	5	β Fornacis.....	2 42	3	123	7	7,41	2,39		7	17,40		-	9,99		15,242	
158	6	σ Arietis.....	2 42	3	75	37	9,26	0,57		37	7,13		+	2,13		15,234	
159	5	τ Persei.....	2 42	5	37	56	7,54	1,41	56	10,08	56	13,00	-	2,54	-	5,46	15,227
160	6	γ Fornacis.....	2 42	5	115	15	34,87	3,42		15	28,64		+	6,23		15,222	
161	5	ζ Hydri.....	2 43	5	158	19	41,43	3,20		19	30,67		+	10,76		15,187	
162	4,5	τ Eridani.....	2 44	5	111	42	17,93	3,10	42	16,63	42	11,79	+	1,30	+	6,14	15,165
163	7	* Arietis.....	2 44	5	74	12	39,97	4,06		12	37,66		+	2,31		15,142	
164	6	ρ Arietis.....	2 46	3	72	21	29,03	1,66		21	24,11		+	4,92		14,996	
165	6	ρ Arietis.....	2 47	5	72	39	21,01	3,13	39	21,35	39	20,74	-	0,34	+	0,27	14,961
166	6,7	* Arietis.....	2 47	5	92	18	10,50	1,17		18	9,46		+	1,04		14,944	
167	3	γ Eridani.....	2 48	4	99	34	34,42	1,64	34	31,06	34	25,56	+	3,36	+	8,86	14,887
168	5	ε Arietis.....	2 49	5	69	20	26,79	2,00	20	27,89	20	23,82	-	1,10	+	2,97	14,806
169	5,6	κ Eridani.....	2 51	5	114	17	25,92	3,22		17	21,83		+	4,09		14,745	
170	4,5	θ Eridani.....	2 51	5	130	59	7,80	2,38		59	8,71		-	0,91		14,669	
171	4	γ Persei.....	2 53	3	37	9	46,93	1,10	9	45,67	9	46,26	+	1,26	+	0,67	14,627
172	5,6	ρ Eridani.....	2 53	2	98	20	0,68	0,28		20	58,70		+	1,98		14,610	
173	5	* Persei.....	2 53	5	33	57	54,49	1,59		57	58,31		-	3,82		14,609	
174	2,3	α Ceti.....	2 53	6	86	34	40,35	2,81	34	42,30	34	40,38	-	1,95	-	0,03	14,575
175	4	ρ Persei.....	2 54	4	51	49	16,71	3,95	49	14,67	49	14,12	+	2,04	+	2,59	14,520
176	5	ρ Eridani.....	2 54	3	98	21	16,06	1,37		21	19,09		-	3,03		14,517	
177	4	* Persei.....	2 57	5	41	2	21,88	3,24	2	22,61	2	22,60	-	0,73	-	0,72	14,370
178	2,3	β Persei.....	2 57	12	49	42	10,33	2,31	42	7,23	42	5,19	+	3,10	+	5,14	14,349
179	5	κ Persei.....	2 58	5	45	47	22,80	2,85		47	24,05		-	1,25		14,292	
180	6	σ Fornacis.....	3 0	4	118	28	56,34	2,49		28	3,05		-	6,71		14,136	
181	4	δ Arietis.....	3 2	4	70	55	6,39	1,15	55	5,43	55	3,99	+	0,96	+	2,40	14,053

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.		Difference from		Annual Variation.	
					°	'	"			m.	s.	m.	s.		Greenwich.
182	5,6	k ¹ Ceti.....	3 4	5	91	50	1,96	3,22		50	0,39	+	1,57	13,916	
183	3,4	12 Eridani.....	3 5	5	119	39	30,42	1,95	39	27,50	39	22,93	+ 2,92	+ 7,44	14,689
184	5	ζ Arietis.....	3 5	5	69	35	16,46	3,94	35	15,60	35	12,96	+ 0,86	+ 3,50	13,851
185	4	ζ Eridani.....	3 8	5	99	27	11,75	2,95	27	9,31	27	6,86	+ 2,44	+ 4,89	13,696
186	6	14 Eridani.....	3 8	5	99	47	8,78	3,59		47	5,54	+	3,24	13,646	
187	5,6	k ² Ceti.....	3 10	5	91	33	2,95	3,45		33	5,25	-	2,30	13,561	
188	3,4	16 Eridani.....	3 12	5	112	22	37,48	3,34	22	40,75	22	41,00	- 3,27	- 3,52	13,414
189	2,3	α Persei.....	3 12	29	40	44	54,69	4,71	44	52,31	44	49,72	+ 2,38	+ 4,97	13,397
190	6	κ ² Ceti.....	3 12	5	86	56	18,60	3,99		56	16,92	+	1,68	13,396	
191	4	e Eridani.....	3 13	4	133	44	10,57	3,28		43	39,22	+	31,35	13,346	
192	6	65 Arietis.....	3 14	3	69	48	9,93	1,36		48	3,37	+	6,56	13,239	
193	4	* Camelopardi..	3 15	5	30	39	29,46	4,88	39	27,78	39	28,71	+ 1,68	+ 0,75	13,191
194	4,5	* Camelopardi..	3 16	5	31	42	57,98	2,76	42	58,98	42	58,84	- 1,00	- 0,86	13,123
195	4	ξ Tauri.....	3 18	5	80	51	44,85	2,84	51	42,75	51	41,49	+ 2,10	+ 3,36	13,019
196	6,7	66 Arietis.....	3 19	5	67	47	1,66	2,39		47	1,83	-	0,17	12,982	
197	5	σ Persei.....	3 19	5	42	35	42,86	3,17		35	46,90	-	4,04	12,974	
198	6	s Tauri.....	3 21	3	79	14	59,20	0,56		14	51,88	+	7,32	12,808	
199	4,5	17 Eridani.....	3 22	4	95	39	38,67	4,32	39	36,56	39	35,71	+ 2,21	+ 2,96	12,736
200	6,7	t Tauri.....	3 23	5	81	12	12,58	3,68		12	11,08	+	1,50	12,654	
201	5	z Eridani.....	3 24	4	131	56	38,43	2,58		56	29,53	+	8,90	12,602	
202	6	7 Tauri.....	3 24	5	66	6	30,74	2,94		6	31,05	-	0,31	12,588	
203	5	ψ Persei.....	3 24	5	42	22	37,81	1,23		22	43,11	-	5,30	12,584	
204	4	e Eridani.....	3 25	5	100	2	11,48	3,87	2	7,94	2	8,10	+ 3,54	+ 3,33	12,550
205	4	t ² Eridani.....	3 26	5	112	12	17,53	5,68	12	16,24	12	15,52	+ 1,29	+ 2,01	12,458
206	6	9 Tauri.....	3 27	6	67	21	15,13	3,15		21	11,89	+	3,24	12,409	
207	5	E Tauri.....	3 28	8	90	8	28,23	2,80		8	20,51	+	7,72	12,325	
208	3,4	δ Persei.....	3 31	5	42	45	38,34	1,01	45	39,13	45	41,53	- 0,79	- 3,19	12,142
209	5	y Eridani.....	3 31	5	130	50	0,99	1,93		49	59,77	+	1,22	12,132	
210	4,5	v Persei.....	3 34	5	47	57	43,36	0,53	57	47,12	57	45,62	- 3,76	- 2,26	11,946
211	4	* Persei.....	3 34	5	58	15	12,73	2,14		15	14,44	-	1,71	11,944	
212	4,5	b Pliedum.....	3 35	4	66	25	27,52	0,42	25	28,35	25	23,12	- 0,83	+ 4,40	11,866
213	5	e Pliedum.....	3 35	4	66	4	8,79	0,99	4	11,95	4	6,30	- 3,16	+ 2,49	11,845
214	3,4	δ Eridani.....	3 35	5	100	20	28,40	2,89	20	30,63	20	24,44	- 2,23	+ 3,96	11,243
215	5	δ Fornacis.....	3 35	5	122	23	56,43	4,06		23	55,90	+	0,53	11,817	
216	3	η Tauri.....	3 37	5	66	25	24,42	3,56	25	26,95	25	26,18	- 2,53	- 1,76	11,682
217	5	π Eridani.....	3 38	4	102	38	11,41	1,44		38	12,52	-	1,11	11,631	
218	5	f Pliedum.....	3 39	4	66	28	10,75	3,85		28	11,59	-	0,84	11,563	
219	5	m ¹ Eridani.....	3 39	5	113	45	10,75	2,82		45	14,46	-	3,71	10,989	
220	5	m ² Eridani.....	3 40	5	114	24	12,68	5,12		24	8,61	+	4,07	11,470	
221	5	g Eridani.....	3 43	5	126	42	57,11	1,51		42	56,30	+	0,81	11,273	
222	3,4	ζ Persei.....	3 43	5	58	37	29,05	1,46	37	30,56	37	34,02	- 1,51	- 4,97	11,247
223	5	32 Eridani.....	3 46	5	93	27	34,71	2,77		27	35,45	-	0,74	11,080	
224	3,4	e Persei.....	3 46	5	50	29	13,49	1,76	29	11,65	29	10,56	+ 1,84	+ 2,93	11,028
225	5	i Eridani.....	3 47	5	125	14	15,01	1,83		14	10,45	+	4,56	10,976	
226	5	ξ Persei.....	3 48	5	54	42	13,57	2,50		42	9,36	+	4,21	10,920	
227	2,3	γ Eridani.....	3 50	7	103	59	39,01	3,88	59	41,33	59	39,14	- 2,32	- 0,13	10,762

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.
					°	'	"				s.	s.	
228	6,7	* Tauri.....	3 51	5	72	17	20,04	1,64		17 19,59		+ 0,45	10,693
229	4	λ Tauri.....	3 51	5	77	59	32,36	2,21	59 36,56	59 30,48	- 4,20	+ 1,88	10,676
230	5	k Eridani.....	3 53	4	114	30	1,65	1,20		29 57,87		+ 3,78	10,571
231	5	35 Eridani.....	3 53	5	92	1	46,59	2,39		1 45,76		+ 0,83	10,552
232	5	ν Tauri.....	3 54	5	84	29	8,14	0,83		29 5,32		+ 2,82	10,463
233	5	A Tauri.....	3 55	5	68	23	8,57	1,72	23 13,08	23 14,47	- 4,51	- 5,90	10,424
234	6,7	A Tauri.....	3 55	5	68	27	10,58	0,68		27 11,83		- 1,25	10,376
235	5	δ Reticuli.....	3 56	5	151	52	45,10	1,23		52 28,70		+ 16,40	10,318
236	6	41 Tauri.....	3 56	5	62	51	43,57	2,86		51 44,80		- 1,23	10,308
237	5	c Persei.....	3 56	5	42	44	49,64	1,66		44 53,89		- 4,25	10,297
238	5	γ Reticuli.....	3 58	5	152	37	57,36	2,16		37 53,96		+ 3,40	10,140
239	6,7	* Tauri.....	4 0	3	77	3	20,61	0,69		3 19,74		+ 0,87	10,056
240	6,7	P Tauri.....	4 0	5	63	58	0,85	1,83		58 5,57		- 4,72	9,984
241	5,6	37 Eridani.....	4 2	5	97	22	17,70	1,72		22 16,14		+ 1,56	9,863
242	4,5	μ Persei.....	4 2	5	42	1	46,27	4,26	1 44,62	1 48,22	+ 1,65	- 1,95	9,836
243	4,5	o Eridani.....	4 4	5	97	17	3,75	4,15	17 4,32	17 2,57	- 0,57	+ 1,18	9,750
244	5	b Persei.....	4 5	5	40	7	47,01	1,49		7 45,73		+ 1,28	9,603
245	5	μ Tauri.....	4 6	4	81	32	15,31	1,09		32 13,68		+ 1,63	9,540
246	5	A Eridani.....	4 6	5	100	40	56,14	2,98		40 51,59		+ 4,55	9,540
247	5,6	ω Tauri.....	4 7	5	69	50	42,99	1,81		50 36,69		+ 6,30	9,463
248	5	d Eridani.....	4 7	5	97	55	13,33	2,62		55 12,55		+ 0,78	5,852
249	7	51 Tauri.....	4 8	5	68	50	25,43	4,03		50 25,97		- 0,54	9,384
250	5	α Horologii.....	4 8	5	132	42	50,59	1,62		42 47,52		+ 3,07	9,381
251	3,4	γ Tauri.....	4 10	5	74	47	17,04	1,93	47 13,98	47 14,17	+ 3,06	+ 2,87	9,245
252	6,7	* Tauri.....	4 11	3	76	32	45,86	1,37		32 43,58		+ 2,28	9,152
253	3,4	X Eridani.....	4 11	5	124	12	56,07	1,82	12 57,00	12 55,78	- 0,93	+ 0,29	9,142
254	4	γ Doradus.....	4 12	4	141	54	59,08	2,04		55 18,97		- 19,89	9,133
255	3,4	α Reticuli.....	4 12	4	152	53	52,83	2,55		53 37,38		+ 15,45	9,080
256	4	δ Tauri.....	4 13	5	72	51	37,25	2,49	51 39,36	51 33,04	- 2,11	+ 4,21	9,011
257	5	ε Reticuli.....	4 14	5	149	42	39,64	3,88		42 39,08		+ 0,56	8,977
258	4,5	δ Tauri.....	4 14	5	72	57	13,13	1,38	57 16,86	57 12,62	- 3,73	+ 0,61	8,919
259	5	δ Tauri.....	4 16	4	72	27	55,73	2,37	27 57,44	27 54,95	- 1,71	+ 0,78	8,813
260	5	θ Reticuli.....	4 16	5	153	39	56,81	6,22		39 56,34		+ 0,47	8,805
261	5	ν Tauri.....	4 16	5	67	34	34,42	3,32		34 37,12		- 2,70	8,775
262	5	π Tauri.....	4 17	5	75	40	33,97	4,84		40 29,07		+ 4,90	8,707
263	4,5	43 Eridani.....	4 18	5	124	24	51,50	3,33	24 49,10	24 50,81	+ 2,40	+ 0,69	8,656
264	4	ε Tauri.....	4 19	5	71	12	5,28	2,93	12 6,02	12 4,28	- 0,74	+ 1,00	8,574
265	5	θ Tauri.....	4 19	5	74	25	12,56	1,48	25 12,68	25 7,13	- 0,12	+ 5,43	8,560
266	6	b Tauri.....	4 19	5	77	20	2,14	3,26		20 2,06		+ 0,08	8,525
267	5	η Reticuli.....	4 20	5	153	47	15,68	2,93		47 17,72		- 2,04	8,465
268	6	85 Tauri.....	4 20	6	74	31	9,28	3,05		31 0,70		+ 8,58	8,299
269	5	ρ Tauri.....	4 24	5	75	31	6,94	2,20		30 57,72		+ 9,22	8,136
270	6	46 Eridani.....	4 25	5	97	5	59,27	2,83		5 58,44		+ 0,83	8,022
271	5	δ Cæli Scalp...	4 26	5	135	19	15,82	4,64		19 18,21		- 2,39	8,022
272	5	47 Eridani.....	4 26	5	98	35	23,66	3,63		35 26,46		- 2,80	7,991
273	1	α Tauri.....	4 26	44	73	50	13,39	4,60	50 15,77	50 11,71	- 2,38	+ 1,68	7,979

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.		Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from		Annual Variation.
					h. m.	° ' "		s.	m. s.	m. s.	s.	s.		
274	5	<i>t</i> Tauri.....	4 26	5	80 11 33,14	1,17		11 36,09		—	2,88	—	7,967	
275	6	<i>k</i> Eridani.....	4 28	5	89 21 1,89	2,51		20 56,12		+	5,77	+	7,794	
276	5	<i>c</i> Tauri.....	4 29	5	77 50 6,52	2,02		50 6,57		—	0,05	—	7,778	
277	3	<i>v</i> Eridani.....	4 29	7	120 54 48,79	5,53	54 51,20	54 49,14	—	2,41	—	0,35	7,755	
278	3	<i>a</i> Doradas.....	4 30	5	145 23 54,23	2,78		23 41,40		+	12,83	+	7,644	
279	4	<i>53</i> Eridani.....	4 30	5	104 38 25,82	2,70	38 24,26	38 19,84	+	1,56	+	5,98	7,638	
280	5	<i>c</i> Tauri.....	4 31	5	78 8 23,17	3,52		8 28,36		—	5,19	—	7,622	
281	6	<i>k</i> Eridani.....	4 31	4	104 41 36,57	2,39		41 29,69		+	6,88	+	7,548	
282	5	<i>r</i> Tauri.....	4 32	5	67 22 27,60	2,06	22 28,76	22 27,64	—	1,16	—	0,04	7,504	
283	4	<i>51</i> Eridani.....	4 33	4	110 0 9,52	1,07	0 6,37	0 1,91	+	3,15	+	7,61	7,426	
284	6	<i>2</i> Eridani.....	4 33	5	114 49 9,78	2,54		49 3,05		+	6,73	+	7,423	
285	4,5	<i>u</i> Caeli Sculp...	4 35	5	132 11 26,84	3,32		11 23,85		+	2,99	+	7,257	
286	5	<i>β</i> Caeli Sculp...	4 36	5	127 28 41,79	1,21		28 50,65		—	8,86	—	7,179	
287	5	<i>μ</i> Eridani.....	4 37	5	93 34 11,14	1,66		34 11,56		—	0,42	—	7,101	
288	4,5	<i>γ</i> Camelopard...	4 37	5	23 57 25,17	4,75	57 29,19	57 27,78	—	4,02	—	2,01	7,084	
289	6	<i>1</i> Eridani.....	4 40	5	118 23 52,06	3,84		23 51,60		+	0,46	+	6,884	
290	4	<i>q</i> Orionis.....	4 41	5	83 20 24,42	4,72	20 27,61	20 28,19	—	3,19	—	3,77	6,804	
291	5	<i>π</i> Orionis.....	4 41	5	81 23 40,22	0,97	23 49,23	23 48,15	—	0,01	+	1,07	6,744	
292	4	<i>ρ</i> Orionis.....	4 42	5	84 41 30,28	0,90	41 27,13	41 24,24	+	3,15	+	6,04	6,677	
293	5	<i>σ</i> Orionis.....	4 43	5	76 2 18,53	3,92		2 17,06		+	1,47	+	6,615	
294	5	<i>β</i> Camelopard...	4 43	5	36 31 45,61	2,38		31 46,77		—	1,16	—	6,554	
295	5	<i>ω</i> Eridani.....	4 45	4	95 44 33,87	4,32		44 26,30		+	7,57	+	6,480	
296	4,5	<i>z</i> Orionis.....	4 45	5	87 50 32,80	2,18	50 35,72	50 31,48	—	2,92	+	1,32	6,410	
297	4	<i>ε</i> Aurigæ.....	4 46	5	57 6 36,28	2,34		6 33,14		+	3,14	+	6,369	
298	5	<i>ν</i> Orionis.....	4 47	5	76 45 34,94	3,88		45 31,62		+	3,32	+	6,292	
299	5	<i>ω</i> Aurigæ.....	4 48	5	52 22 26,57	2,18		22 24,91		+	1,66	+	6,216	
300	4,5	<i>d</i> Camelopard...	4 48	5	29 49 3,42	3,74	49 2,56	49 0,33	+	0,86	+	3,09	6,166	
301	4	<i>ν</i> Aurigæ.....	4 50	5	46 26 10,86	1,37	26 11,50	26 10,40	—	0,64	+	0,46	6,045	
302	4	<i>ξ</i> Aurigæ.....	4 51	5	49 10 49,36	1,73	10 50,13	10 49,42	—	0,77	—	0,06	5,976	
303	5	<i>63</i> Eridani.....	4 52	5	100 31 1,15	5,79		31 0,23		+	0,92	+	5,877	
304	4,5	<i>δ</i> Tauri.....	4 53	5	68 39 30,03	3,25	39 33,38	39 31,72	—	3,35	—	1,69	5,782	
305	5	<i>ψ</i> Eridani.....	4 53	5	97 25 39,99	3,34		25 39,28		+	0,71	+	5,760	
306	4	<i>η</i> Aurigæ.....	4 55	3	49 0 12,43	2,49	0 13,43	0 13,18	—	1,00	—	0,75	5,642	
307	5	<i>η</i> Orionis.....	4 55	5	74 50 22,80	1,81		50 17,10		+	5,70	+	5,621	
308	5,6	<i>*</i> Leporis.....	4 55	3	116 31 10,64	2,80		31 8,24		+	2,40	+	5,588	
309	5	<i>m</i> Tauri.....	4 57	5	71 35 21,81	4,81	35 23,33	35 19,35	—	0,52	+	2,45	5,405	
310	4	<i>ε</i> Leporis.....	4 58	5	112 36 11,26	5,59	36 15,35	36 7,67	—	4,09	+	3,59	5,335	
311	5	<i>γ</i> Caeli Sculp...	4 58	5	125 43 10,29	4,16		43 6,56		+	3,73	+	5,331	
312	3	<i>β</i> Eridani.....	4 59	5	95 18 43,92	2,01	18 41,62	18 39,19	+	2,30	+	4,73	5,230	
313	5	<i>η</i> Orionis.....	4 59	5	74 37 36,56	2,74		37 27,80		+	8,76	+	5,190	
314	6	<i>68</i> Eridani.....	5 0	3	94 40 53,40	2,19		40 53,20		+	6,26	+	5,162	
315	4	<i>λ</i> Eridani.....	5 1	5	98 58 37,39	1,60	58 37,20	58 36,95	+	0,19	+	0,44	5,102	
316	5	<i>α</i> Aurigæ.....	5 2	5	51 43 31,18	1,13		43 35,04		—	3,83	—	5,035	
317	1	<i>α</i> Aurigæ.....	5 4	40	44 11 2,23	6,60	11 0,94	11 2,86	+	1,29	—	0,63	4,837	
318	4,5	<i>ε</i> Leporis.....	5 4	4	102 4 43,09	1,82	4 43,53	4 38,39	—	0,44	+	4,70	4,818	
319	5	<i>μ</i> Leporis.....	5 5	5	106 24 40,48	2,98		24 39,24		+	1,24	+	4,739	

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from		Annual Variation.
					°	'	"		s.	m.	s.	m.	s.	Greenwich.	
320	4	* Orionis.....	h. m. 5 5	5	98	21	10,20	2,13	21	12,38	21		-2,18		-4,732
321	1	β Orionis.....	5 6	28	98	24	9,71	4,08	24	12,38	24	10,04	-2,67	-0,33	4,647
322	5	λ Aurigæ.....	5 7	5	50	3	44,06	4,67			3	36,78		+7,28	4,578
323	4	τ Orionis.....	5 9	5	97	2	0,04	5,17	2	2,02	2	0,51	-1,98	-0,47	4,393
324	5	ο Columbæ.....	5 11	5	125	3	59,48	2,17			3	40,71		+18,77	4,223
325	4,5	λ Leporis.....	5 11	5	103	21	25,98	3,43	21	27,38	21	25,24	-1,40	+0,74	4,189
326	5	ζ Orionis.....	5 14	5	86	37	31,38	3,22			37	30,69		+0,69	4,004
327	2	β Tauri.....	5 16	32	61	32	36,45	4,61	32	37,19	32	38,18	-0,74	-1,73	3,863
328	4,5	η Orionis.....	5 16	5	92	33	32,31	6,17	33	34,64	33	33,78	-2,33	-1,47	3,830
329	2	γ Orionis.....	5 16	6	83	48	33,62	4,30	48	38,01	48	39,08	-4,39	-5,46	3,823
330	5	φ Aurigæ.....	5 16	5	55	40	40,47	1,68			40	41,58		-1,11	3,790
331	5	ο Tauri.....	5 17	5	65	12	53,98	2,84	12	55,08	12	51,58	-1,10	+2,40	3,702
332	5	ψ* Orionis.....	5 18	5	87	3	27,10	3,41	3	29,10	3	27,05	-2,00	+0,05	3,658
333	4	β Leporis.....	5 21	5	110	54	1,85	2,31	54	0,20	53	57,42	+1,65	+4,43	3,397
334	5	ε Orionis.....	5 21	5	91	13	56,82	0,93			13	54,31		+2,51	3,387
335	5	χ Aurigæ.....	5 22	5	57	56	27,89	4,57			56	32,30		-4,41	3,337
336	5	Α Orionis.....	5 22	5	84	11	17,00	4,33			11	13,21		+3,79	3,335
337	2	δ Orionis.....	5 23	5	90	25	55,99	2,19	25	53,30	25	53,15	+2,69	+2,84	3,194
338	5	ν Orionis.....	5 24	2	97	25	56,61	0,01			25	58,68		-2,07	3,160
339	4	ε Columbæ.....	5 25	5	125	35	59,06	3,26			35	53,58		+5,48	3,034
340	3,4	α Leporis.....	5 25	6	107	56	54,64	3,96	56	57,00	56	57,22	-2,36	-2,58	3,029
341	4,5	φ ¹ Orionis.....	5 25	5	80	37	59,67	3,36			37	56,02		+3,65	3,007
342	4	λ Orionis.....	5 26	5	80	11	15,97	1,33	11	11,42	11	7,03	+4,55	+8,04	2,982
343	5	c ¹ Orionis.....	5 27	5	94	57	27,33	2,40			57	19,36		+7,97	2,877
344	3,4	ε Orionis.....	5 27	4	96	1	37,13	1,73	1	39,83	1	32,72	-2,75	+4,41	2,865
345	3,4	ζ Tauri.....	5 27	6	68	58	3,75	1,51	58	6,49	58	4,68	-2,74	-0,93	2,835
346	5	φ ² Orionis.....	5 28	3	80	48	30,14	4,10			48	30,50		-0,36	2,827
347	2,3	ε Orionis.....	5 28	10	91	19	0,29	4,75	19	0,90	18	59,85	-0,61	+0,44	2,825
348	5	ι Aurigæ.....	5 28	5	59	37	3,67	2,81			36	52,54		+11,13	2,814
349	4	σ Orionis.....	5 30	5	92	42	13,74	2,55	42	16,36	42	11,70	-2,62	+2,04	2,598
350	5	d Orionis.....	5 31	5	97	18	49,67	2,77			18	46,83		+2,84	2,558
351	4	β Doradus.....	5 32	4	152	36	4,95	3,80			36	2,93		+2,02	2,429
352	3	ζ Orionis.....	5 32	5	92	2	17,39	5,07	2	20,34	2	19,06	-2,95	-1,67	2,427
353	2	α Columbæ.....	5 33	40	124	10	7,42	7,02	10	9,70	10	7,88	-2,28	-0,46	2,313
354	4	γ Leporis.....	5 37	5	112	30	31,26	3,25	30	33,05	30	32,56	-1,79	-1,30	1,976
355	5	B Tauri.....	5 39	4	65	29	51,91	3,24	29	52,65	29	48,79	-0,74	+3,12	1,871
356	4,5	ζ Leporis.....	5 39	5	104	53	31,16	1,98	53	28,10	53	25,26	+3,06	+5,90	1,812
357	5	μ Columbæ.....	5 40	5	122	22	32,16	2,05			22	31,03		+1,13	1,775
358	3	κ Orionis.....	5 40	5	99	44	13,01	1,72	44	10,41	44	6,06	+2,60	+6,95	1,774
359	5	ν Aurigæ.....	5 40	5	50	54	39,28	2,39			54	39,67		-0,39	1,772
360	5	31 Camelopard...	5 40	5	30	9	49,08	0,92			9	44,16		+4,92	1,770
361	5	ξ Aurigæ.....	5 41	4	34	20	42,28	5,30			20	37,31		+4,97	1,694
362	4,5	136 Tauri....C.	5 43	5	62	26	8,64	5,53	26	10,10	26	12,63	-1,46	-3,99	1,517
363	5	δ Leporis.....	5 44	5	110	53	57,44	1,80			53	56,86		+0,58	0,777
364	5	χ ¹ Orionis.....	5 44	5	69	45	49,41	2,50	45	47,07	45	46,61	+2,34	+2,80	1,371
365	5	δ Doradus.....	5 44	6	155	47	58,77	3,86			47	55,48		+3,29	1,357

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Differ-ence of Ex-tremes.	Green-wich Catalogue		A.S. Cata-logue.		Difference from		Annual Variation.
					°	'	"		s.	m.	s.	m.	s.	s.	
366	3	β Columbae.....	5 45	5	125	50	13,48	5,16		50	21,90		8,42	-1,314	
367	3,4	δ Aurigae.....	5 46	5	35	44	25,98	2,00	44	22,76	44	24,69	+ 3,22	+ 1,29	1,265
368	1	α Orionis.....	5 46	43	82	37	53,39	4,71	37	54,90	37	54,64	- 1,51	- 1,25	1,226
369	2	β Aurigae.....	5 47	6	45	4	51,66	4,53	4	46,75	4	47,51	+ 4,91	+ 4,15	1,131
370	5	τ Aurigae.....	5 47	5	44	5	17,72	3,37		5	18,72		- 1,00	1,109	
371	4	θ Aurigae.....	5 48	5	52	48	23,87	3,56	48	30,37	48	27,98	- 1,50	+ 0,89	1,036
372	4	η Leporis.....	5 49	5	104	12	14,49	1,64	12	18,10	12	11,22	- 3,61	+ 3,27	0 991
373	5	δ Doradus.....	5 50	5	156	56	38,62	5,17		56	52,91		- 14,29	0,870	
374	4	γ Columbae.....	5 52	5	125	18	25,29	3,47		18	22,60		+ 2,69	0,742	
375	5	μ Orionis.....	5 53	5	80	21	41,15	4,8	21	35,93	21	37,93	+ 5,22	+ 3,22	0,610
376	5	Π Geminorum...	5 54	5	66	44	8,14	3,26	44	9,27	44	7,18	- 1,13	+ 0,96	0,543
377	5	χ^3 Orionis.....	5 54	5	69	52	0,99	5,28		51	56,59		+ 4,40	0,540	
378	5,6	β Monocerotis...	5 54	4	100	36	25,15	2,43		36	23,86		+ 1,29	0,539	
379	4,5	ν Orionis.....	5 58	11	75	13	12,44	3,37	13	7,41	13	3,01	+ 5,03	+ 9,43	0,187
380	4,5	θ Leporis.....	5 58	7	104	55	37,47	1,72	55	35,58	55	39,72	+ 1,89	- 2,25	0,134
381	5	κ Camelopard...	6 0	5	20	38	12,01	4,98		38	6,43		+ 5,58	+ 0,009	
382	5	λ Camelopard...	6 0	4	29	58	6,99	3,24		58	1,03		+ 5,96	0,034	
383	5	θ Columbae.....	6 2	4	127	14	1,61	3,27		14	2,75		- 1,14	0,148	
384	5	α Lynceis.....	6 2	4	28	26	32,38	2,14		26	38,53		- 6,15	0,195	
385	5	ξ Orionis.....	6 2	4	75	45	43,53	4,19		45	40,54		+ 2,99	0,199	
386	4	κ Aurigae.....	6 4	5	60	26	56,45	2,74	26	53,18	26	57,24	+ 3,27	- 0,79	0,396
387	4,5	η Geminorum...	6 5	10	67	27	5,25	2,25	27	9,07	27	7,49	- 3,82	- 2,24	0,403
388	4,5	β Lynceis.....	6 5	7	30	56	26,79	3,60	56	26,28	56	25,74	+ 0,51	+ 1,05	0,403
389	4,5	α Monocerotis...	6 6	5	96	13	48,18	3,54	13	47,78	13	42,87	+ 0,40	+ 5,31	0,574
390	5,6	κ^* Orionis.....	6 7	5	77	41	17,01	6,56		41	15,27		+ 1,74	0,603	
391	4,5	κ Columbae.....	6 11	5	125	5	25,11	3,30		5	21,08		+ 4,03	0,918	
392	5	δ Aurigae.....	6 12	5	40	38	14,32	1,96		38	15,07		- 0,75	1,008	
393	3	μ Geminorum...	6 13	19	67	24	26,75	5,14	24	28,57	24	26,36	- 1,82	+ 0,39	1,107
394	3	ζ Canis Maj....	6 14	5	119	59	41,62	1,94	59	37,60	59	42,97	+ 4,02	- 1,35	1,205
395	6	ν^* Monocerotis..	6 14	3	86	9	26,77	2,71		9	25,30		+ 1,47	1,254	
396	2,3	β Canis Maj....	6 15	27	107	52	38,79	3,59	52	37,60	52	47,03	+ 1,19	- 8,24	1,329
397	4	λ Canis Maj....	6 15	5	123	21	19,16	1,70	21	22,50	21	24,63	- 3,34	- 5,47	1,389
398	5	ν Geminorum...	6 19	5	69	41	22,55	2,22	41	19,79	41	17,02	+ 2,76	+ 5,53	1,648
399	1	α Argus.....	6 20	59	142	36	20,43	6,80		36	29,26		- 8,83	1,762	
400	5	δ Canis Maj....	6 22	6	122	28	42,93	2,85		28	45,26		- 2,33	1,910	
401	5	f Monocerotis..	6 24	5	82	33	2,01	3,18		33	4,41		- 2,40	2,070	
402	3	δ Ursae Minor.SP.	6 27	31	3	24	50,99	6,53	24	50,90	24	51,94	+ 0,09	- 0,95	2,356
403	3	γ Geminorum...	6 28	9	73	27	48,47	4,94	27	51,26	27	48,92	- 2,79	- 0,45	2,432
404	5	ξ^* Canis Maj....	6 28	5	112	50	8,64	4,70		50	5,40		+ 3,24	2,436	
405	5	ν^* Canis Maj....	6 29	5	109	7	1,69	2,89		7	1,16		+ 0,53	2,553	
406	5	55 Aurigae.....	6 31	5	45	19	24,69	3,61		19	21,26		+ 3,02	2,677	
407	6	h Monocerotis..	6 32	5	79	57	24,45	2,44		57	19,50		+ 4,95	2,756	
408	3	ν Argus.....	6 33	5	133	3	5,18	5,57		3	7,80		- 2,62	2,838	
409	5	42 Camelopard..	6 33	5	22	15	23,45	4,26		15	23,67		- 0,22	2,891	
410	3	c Geminorum...	6 33	16	64	42	37,38	3,25	42	36,14	42	33,66	+ 1,24	+ 3,72	2,916
411	5	κ^* Camelopard...	6 35	5	12	49	36,40	3,48		49	37,64		- 1,24	3,061	

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.				Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from Greenwich.		Annual Variation.
					°	'	"	s.		m.	s.	m.	s.	s.	s.	
412	5	q Camelopard...	6 35	5	20	55	48,40	2,99			55	46,58	+	1,82	+	3,080
413	4	ξ ² Geminorum...	6 36	5	76	55	45,90	2,98	55	46,14	55	42,12	- 0,24	+	3,78	3,113
414	1	a Canis Maj....	6 37	52	106	29	25,98	5,62	29	26,68	29	23,16	- 0,70	+	2,82	4,418
415	5	i Monocerotis...	6 38	5	81	47	15,10	4,92			47	14,54		+	0,56	3,316
416	5	k Monocerotis...	6 39	5	87	24	33,85	0,81			24	27,94		+	5,91	3,394
417	5	x Arg. in Pup...	6 42	5	127	44	50,35	2,79			44	52,68		-	2,33	3,613
418	5	θ Geminorum...	6 42	5	55	50	38,70	2,89			50	39,81		-	1,11	3,615
419	5	e Lynceis.....	6 43	5	31	22	5,60	2,76			22	7,81		-	2,21	3,697
420	4	κ ² Canis Maj...	6 43	5	122	19	5,57	4,03	19	5,40	19	4,42	+ 0,17	+	1,15	3,731
421	5	* Canis Maj....	6 45	5	124	10	20,34	1,38			10	23,99		-	3,65	3,883
422	5,6	e ¹ Geminorum...	6 45	5	76	36	51,90	3,58			36	51,22		+	0,68	3,914
423	4	τ Argus.....	6 46	5	140	25	0,47	1,49			24	55,73		+	4,74	3,971
424	5,6	π ¹ Canis Maj...	6 46	5	110	1	19,21	0,19			1	16,10		+	3,11	4,013
425	5	θ Canis Maj....	6 46	5	111	49	33,77	4,77			49	59,70		-	25,93	4,021
426	4	α Equulei Pict..	6 46	5	151	45	40,34	2,88			45	43,21		-	2,87	4,034
427	4	ο ¹ Canis Maj...	6 47	5	113	58	39,34	2,97	58	42,43	58	39,42	- 3,09	-	0,08	4,088
428	4,5	ε Canis Maj....	6 49	6	106	50	27,17	3,95	50	28,30	50	23,89	- 1,13	+	3,98	4,214
429	2,3	ε Canis Maj....	6 52	35	118	44	45,98	4,72	44	49,87	44	47,29	- 3,89	-	1,31	4,504
430	4	ζ Geminorum...	6 54	5	69	11	21,36	4,22	11	22,69	11	23,92	- 1,33	-	2,56	4,681
431	4,5	* Camelopard...	6 55	5	7	17	28,33	3,80	17	27,50	17	26,75	+ 0,83	-	1,58	4,747
432	3,4	σ Canis Maj....	6 55	6	117	41	51,42	3,62	41	52,00	41	54,90	- 0,58	-	3,48	4,759
433	4	ο ² Canis Maj...	6 56	5	113	35	26,62	3,07	35	33,04	35	31,28	- 6,42	-	4,66	4,842
434	4	γ Canis Maj....	6 56	5	105	23	24,75	2,84	23	23,73	23	18,92	+ 1,02	+	5,83	4,854
435	5	β ³ Aurigæ.....	7 0	5	50	24	49,84	1,63			24	43,24		+	6,60	5,184
436	5	τ Geminorum...	7 0	4	59	29	13,97	2,80			29	7,12		+	6,35	5,213
437	6	47 Geminorum..	7 1	4	62	52	22,51	0,55			52	25,26		+	2,75	5,257
438	3,4	δ Canis Maj....	7 1	5	116	7	48,15	3,21	7	48,92	7	47,28	- 0,77	+	0,87	5,312
439	5,6	20 Monocerotis..	7 2	5	93	58	44,72	3,31			58	42,10		+	2,62	5,338
440	6	m Geminorum...	7 2	5	65	35	47,91	1,81			35	46,88		+	1,03	5,364
441	3,5	m Monocerotis..	7 3	5	90	13	13,87	1,99	13	11,05	13	9,63	- 2,82	+	4,24	5,456
442	5	51 Geminorum...	7 4	5	73	33	38,96	1,80	33	40,56	33	39,30	- 1,60	-	0,34	5,491
443	6	26 Canis Maj...	7 5	3	115	39	55,62	1,88			39	47,42		+	8,20	5,630
444	5	β ⁴ Aurigæ.....	7 6	5	48	49	28,30	2,15			49	32,71		-	4,41	5,709
445	4,5	e ¹ Canis Maj...	7 7	5	116	3	58,36	4,44	3	59,53	3	55,87	- 1,17	+	2,49	5,803
446	5	l Arg. in Pup..	7 8	5	136	28	49,59	2,88			28	52,76		-	3,17	5,836
447	6	ω Canis Maj....	7 8	5	116	29	3,29	1,95			29	1,23		+	2,06	5,852
448	5	L ¹ Arg. in Pup..	7 8	5	134	53	38,52	3,16			53	27,88		+	10,64	5,871
449	4,5	λ Geminorum...	7 8	5	73	9	38,80	4,43	9	43,66	9	41,58	- 4,86	-	2,78	5,887
450	3,4	δ Geminorum...	7 10	5	67	42	48,84	3,49	42	51,28	42	50,29	- 2,44	-	1,45	6,024
451	5	γ Piscis Vol....	7 10	3	160	13	32,48	4,33			13	25,28		+	7,20	6,040
452	5	β ⁵ Aurigæ.....	7 11	5	52	55	47,12	2,97			55	44,32		+	2,80	6,082
453	3,4	π Argus.....	7 11	5	126	47	56,36	1,82			47	55,14		+	1,22	6,121
454	6	29 Canis Maj....	7 12	5	114	15	19,21	2,45			15	19,71		-	0,50	6,159
455	6	d Canis Maj....	7 12	5	114	39	6,62	1,26			39	4,34		+	2,28	6,165
456	4	ε Geminorum...	7 15	4	61	52	28,47	2,26	52	26,41	52	22,20	+ 2,06	+	6,27	6,456
457	5	ο Piscis Vol....	7 17	4	157	38	50,29	3,28			38	45,20		+	5,09	6,592

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from		Annual Variation.
					h.	m.	s.		m.	s.	m.	s.	Greenwich.	A.S.C.	
458	3	γ Canis Maj....	7 17	13	118 58 40,12	3,26	58 42,78	58 42,21	- 2,66	- 2,09	+ 6,638				
459	3	β Canis Min....	7 18	8	81 22 37,57	2,12	22 37,54	22 33,34	+ 0,03	+ 4,23	6,684				
460	5	ρ Geminorum...	7 18	5	57 53 14,65	1,92		53 13,88		+ 0,77	6,703				
461	5,6	b ¹ Geminorum..	7 19	7	61 32 28,67	3,46		32 28,87		- 0,20	6,751				
462	3	a Geminorum...	7 24	33	57 44 56,75	6,71	44 57,06	44 56,29	- 0,31	+ 0,46	7,161				
463	4	σ Argus.....	7 24	5	132 57 46,33	2,52		57 51,59		- 5,26	7,169				
464	5	k Geminorum...	7 24	5	73 48 58,81	2,12		48 55,58		+ 3,23	7,174				
465	5,6	δ ² Canis Min...	7 24	6	86 21 22,07	3,36		21 20,23		+ 1,84	7,205				
466	6	δ ³ Canis Min...	7 25	6	86 16 6,18	2,73		16 3,01		+ 3,17	7,291				
467	5	v Geminorum...	7 25	5	62 44 8,73	3,11	44 10,22	44 6,06	- 1,49	+ 2,67	7,300				
468	1,2	a Canis Min....	7 30	60	84 20 55,10	5,40	20 54,24	20 48,79	+ 0,86	+ 6,31	8,682				
469	4,5	n Monocretis...	7 33	4	99 9 43,90	2,23	9 46,00	9 43,18	- 3,00	+ 0,72	7,921				
470	4	κ Geminorum...	7 34	5	65 12 17,06	1,40	12 14,51	12 14,69	+ 2,52	+ 2,37	8,005				
471	2	β Geminorum...	7 35	33	61 34 21,66	4,24	34 22,20	34 22,79	- 0,54	- 1,13	8,064				
472	5	3 Argus.....	7 37	5	118 33 16,63	3,82		33 18,72		- 2,09	8,230				
473	4	c Arg. in Pup..	7 39	5	127 33 43,95	3,33		33 46,37		- 2,42	8,407				
474	5,6	o Arg. in Pup..	7 41	9	115 31 23,65	4,25		31 19,21		+ 4,44	8,551				
475	5,6	6 Argus.....	7 42	5	106 48 11,44	4,05		48 9,17		+ 2,27	8,630				
476	4	ξ Argus.....	7 42	5	114 26 28,38	1,17	26 27,23	26 24,83	+ 1,15	+ 3,55	8,640				
477	5	φ Geminorum...	7 43	5	62 48 19,44	3,42	48 15,22	48 16,20	+ 4,22	+ 3,24	8,713				
478	5	9 Argus.....	7 44	5	103 27 15,83	4,13		27 19,66		- 3,83	8,778				
479	4,5	p Arg. in Pup..	7 44	5	135 57 4,04	4,23		57 3,46		+ 0,58	8,791				
480	6	* Canis Min....	7 46	6	80 41 43,88	1,50		41 39,42		+ 4,46	8,965				
481	5	b Arg. in Pup..	7 47	5	128 25 44,33	4,09		25 43,52		+ 0,81	8,992				
482	6	1 Cancr.....	7 47	7	73 45 51,58	4,58		45 49,63		+ 1,95	9,047				
483	5	R Arg. in Pup..	7 48	5	137 39 52,13	1,51		39 58,78		- 6,65	9,123				
484	6	3 Cancr.....	7 51	4	72 14 5,65	2,78	14 4,34	14 3,79	+ 1,31	+ 1,86	9,335				
485	3	χ Argus.....	7 52	15	142 31 54,43	5,83		32 0,08		- 5,65	9,445				
486	5	13 Argus.....	7 53	5	87 12 26,91	3,37		12 21,96		+ 4,95	9,519				
487	5	k Lyncis.....	7 56	5	38 0 52,77	1,27		1 0,69		- 7,92	9,689				
488	5	56 Camelopard..	7 56	5	21 2 24,64	3,70		3			9,699				
489	3	ζ Argus.....	7 58	12	129 31 54,58	1,97		31 50,65		+ 3,93	9,840				
490	5,6	29 Monocrotis..	8 0	3	92 29 49,05	1,77		29 49,23		- 0,18	10,025				
491	3,4	15 Argus.....	8 0	5	113 49 16,22	1,97	49 20,34	49 14,72	- 4,12	+ 1,50	10,044				
492	5,6	16 Argus.....	8 1	5	108 45 15,91	2,52		45 20,29		- 4,38	10,128				
493	5	γ ¹ Argus.....	8 4	5	136 51 0,47	3,97		51 2,32		- 1,85	10,342				
494	2	γ ² Argus.....	8 4	9	136 50 20,32	3,28		50 33,32		- 13,00	10,345				
495	5	20 Argus.....	8 6	5	105 17 7,29	2,02		16 58,29		+ 9,00	10,436				
496	5	* Argus.....	8 6	5	132 29 12,11	1,73		29 11,80		+ 0,31	10,440				
497	5	r Arg. in Pup..	8 7	5	125 23 40,38	1,22		23 33,88		+ 6,50	10,552				
498	4	β Cancr.....	8 7	8	80 18 2,84	4,27	17 58,75	17 56,50	+ 4,09	+ 6,34	10,568				
499	5	m Lyncis.....	8 11	5	46 16 38,54	2,27		16 37,14		+ 1,40	10,854				
500	4,5	q Arg. in Pup..	8 12	5	126 8 24,46	4,87		8 23,14		+ 1,32	10,931				
501	6	d ¹ Cancr.....	8 14	5	71 7 53,61	4,94		7 53,42		+ 0,19	11,035				
502	4,5	o Ursæ Maj....	8 16	5	28 43 31,89	2,77	43 33,51	43 36,05	- 1,65	+ 4,16	11,212				
503	2	e Argus.....	8 19	20	148 58 3,32			58 11,36		- 8,04	11,426				

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Differ-ence of Ex-tremes.	Green-wich Catalogue		A.S. Cata-logue.		Difference from		Annual Variation.
					h.	m.	s.		m.	s.	m.	s.	Green-wich.	A.S.C.	
504	5,6	θ Cancrī.....	8 22	3	71	20	23,00	0,27	20	25,33	20	23,31	- 2,33	- 0,31	+ 11,630
505	5	γ Piscis Vol....	8 24	5	162	51	30,21	5,24			51	20,25		+ 9,96	11,747
506	5	β Piscis Vol....	8 24	6	155	34	28,46	6,78			34	16,38		+ 12,08	11,761
507	5	π ² Ursæ Maj...	8 25	3	25	5	33,32	1,46			5	30,14		+ 3,18	11,869
508	4	δ Hydræ.....	8 29	5	83	42	43,50	2,15	42	44,92	42	40,20	- 1,42	+ 3,30	12,107
509	5	σ Hydræ.....	8 30	5	86	4	13,83	2,18			4	12,94		+ 0,89	12,192
510	5	e ¹ Arg. in Vel..	8 33	5	132	24	4,96	5,85			24	4,26		+ 0,70	12,317
511	5	γ Cancrī.....	8 33	6	67	55	42,71	4,68	55	45,68	55	42,78	- 2,97	- 0,07	12,438
512	5	β Pix Naut....	8 33	5	124	42	50,05	5,40			42	45,42		+ 4,63	12,439
513	5	η Hydræ.....	8 34	5	86	0	1,01	2,13			59	59,86		+ 1,15	12,499
514	5	b Arg. in Vel..	8 35	6	136	3	3,93	1,37			3	1,08		+ 2,85	12,544
515	4,5	δ Cancrī.....	8 35	5	71	13	47,51	3,31	13	48,23	13	47,46	- 0,72	+ 0,05	12,546
516	4	o Argus.....	8 35	3	142	19	26,64	2,47			19	29,93		- 3,29	12,574
517	4,5	a Pix Naut....	8 37	5	122	34	52,16	2,62	34	52,13	34	53,62	+ 0,03	- 1,46	12,665
518	6	12 Hydræ.....	8 38	3	102	56	0,32	0,15			56	4,56		- 4,24	12,771
519	5	ρ Hydræ.....	8 39	6	83	32	34,25	1,90			32	34,08		+ 0,17	12,844
520	3	δ Argus.....	8 40	6	144	5	31,61	1,68			5	39,10		- 7,49	12,884
521	5	a ¹ Arg. in Vel..	8 40	4	135	25	38,61	5,22			25	35,74		+ 2,87	12,901
522	6	γ Pix Naut....	8 43	5	117	5	14,66	2,00			5	13,69		+ 0,97	13,104
523	4	ζ Hydræ.....	8 46	5	83	24	56,80	2,89	24	59,31	24	56,51	- 2,51	+ 0,29	13,307
524	3,4	ι Ursæ Maj....	8 47	6	41	18	4,85	2,58	18	3,01	18	4,12	+ 1,84	+ 0,73	13,380
525	6	o ² Cancrī.....	8 48	5	73	46	28,66	1,18			46	33,56		- 4,90	13,416
526	6	δ Pix Naut....	8 48	5	117	2	7,22	2,81			2	3,98		+ 3,24	13,426
527	5	a ² Cancrī.....	8 49	5	77	29	34,68	5,00	29	35,84	29	36,21	- 1,16	- 1,53	13,487
528	7	* Cancrī.....	8 50	4	71	12	44,18	2,97			12	41,26		+ 2,92	13,512
529	4,5	κ Ursæ Maj....	8 52	5	42	10	56,18	2,01	10	54,62	10	57,81	+ 1,56	- 1,63	13,668
530	6	ν Cancrī.....	8 53	5	64	53	20,14	4,42			53	13,08		+ 7,06	13,719
531	5	δ ¹ Arg. in Car..	8 53	5	148	34	44,12	4,30			34	45,97		- 1,85	13,721
532	5	δ ² Arg. in Car..	8 55	4	148	26	18,17	5,06			26	24,03		- 5,86	13,875
533	5	* Lyncis.....	8 56	5	50	52	42,86	2,39			52	39,00		+ 3,86	13,903
534	6	w Hydræ.....	8 57	5	84	14	14,75	3,04			14	14,59		+ 0,16	13,987
535	5	c Arg. in Vel..	8 58	5	136	25	41,79	1,66			25	46,37		- 4,58	14,067
536	5,6	ξ Cancrī.....	8 59	5	67	16	32,46	6,04	16	34,37	16	30,54	- 1,91	+ 1,92	14,146
537	5	a Piscis Vol....	9 0	5	155	43	25,40	3,14			43	24,06		+ 1,34	14,156
538	3,4	λ Argus.....	9 2	5	132	45	13,08	1,79			45	9,70		+ 3,38	14,280
539	6	e Pix Naut....	9 3	5	119	40	49,17	5,96			40	45,43		+ 3,74	14,341
540	5	e Ursæ Maj....	9 4	5	35	17	16,85	5,39			17	12,76		+ 4,09	14,412
541	5	G Arg. in Car..	9 5	5	161	55	35,88	2,30			55	27,16		+ 8,72	14,456
542	4,5	θ Hydræ.....	9 6	5	86	58	39,50	1,74	58	38,00	58	37,73	+ 1,50	+ 1,77	14,509
543	5	a Arg. in Car..	9 6	5	148	16	36,59	2,44			16	51,93		- 15,34	14,568
544	5	i Arg. in Car..	9 7	4	151	37	28,49	4,32			37	38,28		- 9,79	14,623
545	4	p Lyncis.....	9 8	5	52	29	15,74	1,69	29	15,57	29	9,33	+ 0,17	+ 6,41	14,672
546	6	K ² Hydræ.....	9 8	3	95	39	9,74	2,03			39	6,77		+ 2,97	14,673
547	5	l Arg. in Vel..	9 9	4	127	52	8,42	2,75			52	14,07		- 5,65	14,713
548	4,5	r Lyncis.....	9 11	5	54	53	58,93	1,46	53	53,51	53	50,20	+ 5,42	+ 8,73	14,817
549	2	β Argus.....	9 11	5	159	1	18,55	6,53			1	30,95		- 12,40	14,855

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.		Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.	Difference from		Annual Variation.
					<i>h.</i>	<i>m.</i>		<i>m.</i>	<i>s.</i>		<i>m.</i>	<i>s.</i>	
550	5	θ Pix Naut....	9 14	5	115 15	3,88	2,96		15 4,16		—	0,28	+ 15,009
551	3	κ Argus.....	9 17	5	144 17	30,52	3,73		17 36,19		—	5,67	15,175
552	4	h Ursæ Maj....	9 18	7	26 12	22,48	4,69	12 22,32	12 17,69	+ 0,16	+	4,79	15,242
553	2	a Hydræ.....	9 19	10	97 55	46,45	5,28	55 47,96	55 46,09	— 1,51	+	0,36	15,310
554	5	d Ursæ Maj....	9 19	5	19 26	5,93	4,36		26 3,88		+	2,05	15,314
555	3	θ Ursæ Maj....	9 21	3	37 33	25,94	3,32	33 28,37	33 30,27	— 2,43	—	4,33	16,034
556	4,5	λ Leonis.....	9 22	2	66 17	29,26	0,92	17 29,26	17 25,09	0,00	+	4,17	15,466
557	5	b Leonis Min...	9 24	5	52 51	23,35	2,03	51 25,98	51 19,91	— 2,63	+	3,44	15,564
558	4,5	ψ Argus.....	9 24	5	129 43	53,40	1,78		43 51,94		+	1,46	15,577
559	5	N Arg. in Vel...	9 26	9	146 17	29,74	6,11		17 32,77		—	3,03	15,687
560	5	h Arg. in Car...	9 30	5	148 28	40,76	1,76		28 47,09		—	6,33	15,875
561	5,6	b Sextantis....	9 30	5	84 35	32,54	2,47		35 23,18		+	4,36	15,878
562	5	e Hydræ.....	9 31	5	90 22	51,15	4,84		22 45,39		+	5,76	15,963
563	6	13 Leonis.....	9 32	3	63 19	15,99	1,86		19 21,97		—	5,98	15,998
564	4	o Leonis.....	9 32	7	79 20	36,07	4,20	20 34,14	20 32,10	+ 1,93	+	3,97	16,010
565	5	κ Hydræ.....	9 32	5	103 34	6,50	3,17		34 9,75		—	3,25	16,015
566	6	ψ Leonis.....	9 34	5	75 12	35,90	4,42		12 33,62		+	2,28	16,135
567	3	e Leonis.....	9 36	9	65 27	8,06	1,99	27 6,24	27 3,24	+ 1,82	+	4,82	16,224
568	6	θ Antle Pneum..	9 37	5	116 59	55,56	4,13		59 54,67		+	0,89	16,247
569	4,5	v Ursæ Maj....	9 39	5	30 10	23,72	2,26	10 17,66	10 21,06	+ 6,06	+	2,66	16,359
570	5	φ Ursæ Maj....	9 40	7	35 9	7,71	2,43		9 5,55		+	2,16	16,443
571	5	l Arg. in Car...	9 41	5	151 43	48,87	2,74		43 52,09		—	3,22	16,446
572	6	g Leonis.....	9 42	3	64 48	33,82	2,17		48 31,32		+	2,50	16,528
573	3,4	v Argus.....	9 43	6	154 17	24,60	3,87		17 21,86		+	2,74	16,560
574	3	μ Leonis.....	9 43	5	63 12	2,65	0,60	12 4,92	12 2,66	— 2,27	—	0,01	16,570
575	5	v Hydræ.....	9 43	5	104 3	23,56	5,16		3 21,27		+	2,29	16,582
576	5,6	v Leonis.....	9 49	3	76 45	9,86	2,19	45 9,85	45 4,69	+ 0,01	+	5,17	16,860
577	6	11 Sextantis....	9 49	5	80 53	0,10	2,31		52 57,22		+	2,88	16,862
578	4	φ Argus.....	9 51	5	143 45	55,79	3,35		45 59,59		—	3,80	16,946
579	4,5	π Leonis.....	9 51	5	81 8	55,80	2,45	8 54,49	8 52,65	+ 1,31	+	3,15	16,961
580	5	d Leonis Min...	9 57	3	53 56	7,26	1,22		56 4,87		+	2,39	17,241
581	5	A Leonis.....	9 59	4	79 10	37,94	1,67	10 38,37	10 35,84	— 0,53	+	2,00	17,308
582	1	a Leonis.....	9 59	10	77 12	38,58	2,28	12 35,16	12 32,50	+ 3,42	+	6,08	17,327
583	6	16 Sextantis....	10 0	5	83 0	11,44	4,75		0 9,64		+	1,80	17,372
584	4,5	λ Hydræ.....	10 2	5	101 31	18,24	5,84	31 19,50	31 16,36	— 1,26	+	1,88	17,457
585	6	21 Sextantis....	10 6	3	97 9	29,47	4,65		9 21,36		+	3,11	17,600
586	3,4	λ Ursæ Maj....	10 6	6	46 14	43,81	4,90	14 43,30	14 43,72	+ 0,51	+	0,09	17,647
587	4,5	ζ Leonis.....	10 7	6	65 44	38,30	2,11	44 41,06	44 32,20	— 2,76	+	6,10	17,664
588	4	q Arg. in Vel...	10 8	5	131 17	15,08	1,87		17 7,37		+	7,71	17,661
589	4,5	w Argus.....	10 10	2	159 12	2,94	3,37		12 10,32		—	7,38	17,765
590	2	γ Leonis.....	10 11	6	69 18	24,49	2,30	18 24,00	18 22,42	+ 0,49	+	2,07	17,802
591	5	q Arg. in Car...	10 11	4	150 29	24,15	3,87		29 29,81		—	5,66	17,834
592	3	μ Ursæ Maj....	10 12	5	47 39	13,35	3,60	39 15,12	39 11,08	— 1,77	+	2,27	17,865
593	5	T Arg. in Vel...	10 15	5	145 11	43,92	5,81		11 40,66		+	3,16	17,959
594	4,5	r Arg. in Vel...	10 15	5	130 48	8,45	5,08		48 4,71		+	3,74	17,978
595	4,5	f Leonis Min...	10 16	4	55 20	46,10	5,38	20 46,01	20 45,27	+ 0,09	+	0,33	18,019

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from		Annual Variation.
					°	'	"		s.	m.	s.	m.	s.	s.	
596	4	<i>μ</i> Hydræ.....	10 18	5	105	58	34,95	1,92	58	34,24	58	32,28	+ 0,71	+ 2,67	+ 18,085
597	4,5	<i>g</i> Leonis Min...	10 18	5	52	25	47,77	3,15	25	46,67	25	45,54	+ 1,10	+ 2,23	18,090
598	4,5	<i>α</i> Antl. Pneum...	10 19	5	120	12	38,52	4,04	12	33,57	12	33,88	+ 4,95	+ 4,64	18,142
599	5	<i>36</i> Ursæ Maj...	10 20	5	33	9	19,58	3,97	9	23,43	9	19,50	- 3,85	+ 0,08	18,154
600	5	<i>1</i> Arg. in Car...	10 21	4	163	10	43,21	4,06							18,201
601	4	<i>ρ</i> Leonis.....	10 24	5	79	49	40,01	4,52	49	34,01	49	31,94	+ 6,00	+ 8,07	18,305
602	5	<i>m</i> Ursæ Maj....	10 24	5	32	3	2,15	3,78	3	1,72	2	59,76	+ 0,43	+ 2,39	18,315
603	4	<i>p</i> Arg. in Car...	10 26	6	150	49	4,16	4,41			49	6,60		- 2,44	18,381
604	6	<i>1</i> Hyd. & Crat...	10 28	3	105	28	18,62	0,43			28	12,52		+ 6,10	18,449
605	4	<i>l</i> Leonis Min...	10 29	5	57	8	56,13	2,52	8	54,55	8	52,06	+ 1,58	+ 4,07	18,488
606	5	<i>p</i> Arg. in Vel...	10 30	4	137	20	58,22	4,41			21	3,64		- 5,42	18,523
607	5	<i>φ</i> ³ Hyd. & Crat.	10 30	5	106	0	6,36	1,12			0	0,71		+ 5,65	18,527
608	5	* Ursæ Maj....	10 31	5	20	2	40,58	1,34			2	34,74		+ 5,84	18,543
609	6	<i>m</i> Sextantis....	10 33	3	90	51	21,45	3,67			51	17,27		+ 4,18	18,608
610	5	<i>θ</i> ¹ Argus.....	10 36	5	153	34	58,39	2,24			35	6,98		- 8,59	18,718
611	4,5	<i>n</i> Leonis Min...	10 36	6	58	25	43,96	4,13	25	47,76	25	44,79	- 3,80	- 0,83	18,723
612	2,3	<i>θ</i> ² Argus.....	10 37	5	153	30	35,81	6,26			30	34,12		+ 1,69	18,739
613	2	<i>η</i> Argus.....	10 38	5	148	47	47,64	2,94			47	54,47		- 6,83	18,789
614	3	<i>u</i> Argus.....	10 39	6	138	31	40,65	4,97			31	26,68		+ 13,97	18,818
615	4	<i>ν</i> Hyd. & Crat...	10 41	5	105	18	43,08	4,88	18	40,50	18	37,33	+ 2,58	+ 5,75	18,871
616	4,5	<i>o</i> Leonis Min...	10 44	5	54	52	40,83	4,54	52	34,13	52	32,08	+ 6,70	+ 3,75	18,944
617	5	<i>w</i> Ursæ Maj....	10 44	3	45	54	44,91	3,80			54	43,64		+ 1,27	18,955
618	4,5	<i>54</i> Leonis.....	10 46	6	64	21	4,50	2,00	21	3,56	20	58,90	+ 0,94	+ 5,60	19,018
619	5	<i>u</i> Arg. in Car...	10 47	5	147	57	23,68	3,61			57	29,14		- 5,46	19,025
620	5	* Antl. Pneum...	10 49	5	126	13	52,33	2,75			13	44,57		+ 7,76	19,084
621	4	<i>α</i> Hyd. & Crat...	10 51	5	107	24	0,70	4,76	24	2,17	23	59,84	- 1,47	+ 0,86	19,154
622	5	<i>d</i> Leonis.....	10 52	5	85	28	40,03	4,08	28	36,98	28	31,55	+ 3,05	+ 8,48	19,161
623	5,6	<i>s</i> Leonis.....	10 53	4	91	34	31,69	1,91			34	32,30		- 0,61	19,196
624	1,2	<i>α</i> Ursæ Maj....	10 53	5	27	20	20,21	1,95	20	18,04	20	19,26	+ 2,17	+ 0,95	19,196
625	5	<i>β</i> Leonis.....	10 53	5	68	54	52,02	1,78			54	51,18		+ 0,84	19,198
626	4,5	<i>χ</i> Leonis.....	10 56	5	81	45	7,47	3,45	45	7,19	45	4,16	+ 0,28	+ 3,31	19,272
627	5	<i>χ</i> ¹ Hyd. & Crat.	10 57	4	116	22	58,72	4,78			22	56,86		+ 1,86	19,294
628	5,6	<i>p</i> ² Leonis.....	10 58	5	87	7	43,81	5,35			7	38,36		+ 5,45	19,319
629	3,4	<i>ψ</i> Ursæ Maj....	11 0	5	44	35	12,93	0,96	35	10,34	35	9,45	+ 2,59	+ 3,48	19,361
630	5	<i>10</i> Hyd. & Crat.	11 0	5	117	9	57,55	5,64			9	54,92		+ 2,63	19,371
631	7	* Leonis.....	11 3	5	74	41	0,42	5,08			40	55,10		+ 5,32	19,422
632	4	<i>β</i> Hyd. & Crat...	11 3	5	111	54	15,34	5,39	54	15,94	54	13,58	- 0,60	+ 1,76	19,433
633	3	<i>β</i> Leonis.....	11 5	5	68	33	3,17	1,65	33	5,38	33	2,41	- 2,21	+ 0,76	19,469
634	3	<i>θ</i> Leonis.....	11 5	5	73	38	48,30	2,22	38	52,54	38	48,67	- 4,24	- 0,37	19,475
635	5	<i>φ</i> Leonis.....	11 5	5	92	43	45,40	2,44			43	37,75		+ 7,65	19,529
636	4	<i>ξ</i> Ursæ Maj....	11 9	4	57	31	13,81	4,05	31	16,34	31	17,42	- 2,53	- 3,61	20,190
637	4	<i>σ</i> Leonis.....	11 12	5	83	2	41,40	2,53	2	45,11	2	44,05	- 3,71	- 2,65	19,611
638	4	<i>π</i> Centauri....	11 13	5	143	33	56,29	4,19			34	4,35		- 8,06	19,627
639	4	<i>ν</i> Leonis.....	11 15	4	78	32	24,58	1,50	32	27,27	32	22,57	- 2,69	+ 2,01	19,658
640	5	<i>c</i> Hyd. & Crat...	11 16	5	99	56	4,48	4,40			55	53,73		+ 10,75	19,675
641	4	<i>γ</i> Hyd. & Crat...	11 16	7	106	45	22,40	2,19	45	24,12	45	19,86	- 1,72	+ 2,54	19,681

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from		Annual Variation.
					°	'	"		s.	m.	s.	m.	s.	s.	
642	4	τ Leonis.....	h. m. 11 19	6	86	12	50,07	1,59	12	50,49	12	46,28	-0,42	+ 3,79	+ 19,726
643	6	86 Leonis.....	11 22	4	70	39	37,71	3,44			39	33,00		+ 4,71	19,762
644	4,5	e Leonis.....	11 22	5	92	4	20,29	2,82			4	17,00		+ 3,29	19,762
645	7	* Hyd. & Crat..	11 23	5	95	32	3,65	4,86			32	0,41		+ 3,24	19,786
646	7	* Hyd. & Crat..	11 24	5	96	53	41,99	2,94			53	39,47		+ 2,52	19,798
647	4	ξ Hyd. & Crat..	11 25	5	120	55	24,18	3,89	55	21,84	55	23,80	+ 2,34	+ 0,38	19,805
648	4	λ Centauri.....	11 28	5	152	5	9,57	2,43			5	9,89		- 0,32	19,847
649	4	θ Hyd. & Crat..	11 28	6	98	52	1,66	2,88	52	5,89	51	59,94	- 4,23	+ 1,72	19,848
650	4,5	v Leonis.....	11 28	6	89	53	31,00	3,68	53	29,43	53	26,76	+ 1,57	+ 4,24	19,850
651	6,7	ω Virginis.....	11 30	4	30	55	51,65	0,66			55	50,73		+ 0,92	19,868
652	7	* Virginis.....	11 30	3	91	30	4,25	2,43			30	2,68		+ 1,57	19,868
653	5,6	92 Leonis.....	11 32	5	67	42	27,08	3,09			42	27,43		+ 0,35	19,892
654	4	ζ Hyd. & Crat..	11 36	5	107	24	41,02	4,19	24	41,97	24	39,58	- 0,95	+ 1,44	19,934
655	5	ξ' Virginis.....	11 36	5	80	48	12,83	2,53	48	10,06	48	2,36	+ 2,77	+ 10,47	19,937
656	4	χ Ursæ Major..	11 37	5	41	16	59,33	2,43	17	1,81	17	0,98	- 2,48	- 1,66	19,942
657	4,5	v Virginis.....	11 37	5	82	31	22,67	5,15	31	25,01	31	24,03	- 2,34	- 1,26	19,943
658	4	E Leonis.....	11 39	5	68	50	28,49	2,36	50	32,09	50	29,36	- 3,60	- 0,87	19,960
659	2,3	β Leonis.....	11 40	7	74	28	59,54	4,32	29	0,38	28	55,67	- 0,84	+ 3,87	19,969
660	3,4	β Virginis.....	11 42	8	87	16	59,26	2,44	16	59,68	16	57,62	- 0,42	+ 1,64	19,980
661	6	B Virginis.....	11 42	4	94	23	37,87	2,13			23	33,76		+ 4,11	19,983
662	4	β Hyd. & Crat..	11 44	6	122	58	0,90	4,86			58	1,96		+ 4,06	19,996
663	2	γ Ursæ Major..	11 45	7	35	21	58,72	3,16	21	55,55	21	58,27	+ 3,17	+ 0,45	19,999
664	6	A Virginis.....	11 46	3	80	37	0,55	3,43			36	55,19		+ 5,26	20,007
665	6	29 Hyd. & Crat..	11 47	5	117	32	2,34	4,32			32	3,98		- 1,64	20,011
666	5,6	δ Virginis.....	11 51	5	85	24	14,40	2,57			24	10,54		+ 3,86	20,028
667	5	π Virginis.....	11 52	9	82	26	35,05	3,85	26	30,30	26	33,76	+ 4,74	+ 1,29	20,031
668	4,5	o Virginis.....	11 57	12	80	19	44,47	5,42	19	40,44	19	40,61	+ 4,03	+ 3,96	20,040
669	7	* Virginis.....	11 57	3	92	11	23,56	2,48			11	19,29		+ 4,27	20,041
670	4,5	γ Crucis.....	11 58	5	153	40	10,22	3,02			40	17,28		- 7,06	20,042
671	3	δ Centauri.....	12 0	9	139	46	55,63	4,31			46	56,28		- 1,25	20,043
672	4,5	α Corvi.....	12 0	5	113	47	5,66	5,46	17	3,44	47	7,55	- 2,78	- 1,89	20,043
673	4	ρ Centauri.....	12 3	5	141	25	36,79	1,61			25	47,28		- 10,49	20,041
674	3	δ Crucis.....	12 6	6	147	48	26,34	1,65			48	35,27		- 8,46	20,035
675	3	δ Ursæ Major..	12 7	6	32	1	42,67	2,50	1	40,22	1	44,37	+ 2,45	- 1,70	20,033
676	3,4	γ Virginis.....	12 11	7	89	43	39,24	4,64	43	36,93	43	33,67	+ 2,31	+ 5,57	20,019
677	5	σ Comæ Ber....	12 12	4	71	16	21,09	1,71			16	19,38		+ 1,71	20,015
678	4	e Crucis.....	12 12	5	149	28	1,06	3,04			28	4,13		+ 3,07	20,014
679	5	f Comæ Ber....	12 16	5	62	57	44,41	2,28			57	47,27		- 2,86	19,995
680	1	a* Crucis.....	12 17	5	152	9	41,89	2,70			9	47,99		- 6,10	19,986
681	4,5	α Comæ Ber....	12 18	5	62	14	13,40	2,51	14	14,62	14	11,75	- 1,22	+ 1,65	19,978
682	5	σ Centauri.....	12 19	4	139	17	34,69	2,76			17	21,85		+ 12,84	19,975
683	3	δ Corvi.....	12 21	3	105	34	27,66	5,65	34	25,23	34	21,23	+ 2,43	+ 6,43	19,958
684	2,3	γ Crucis.....	12 22	5	146	9	56,18	2,60			9	44,71		+ 11,47	19,952
685	4,5	η Corvi.....	12 23	4	105	15	36,17	0,37	15	32,43	15	27,79	+ 3,74	+ 8,38	19,939
686	2,3	β Corvi.....	12 25	3	112	27	39,91	2,98	27	38,17	27	39,48	+ 1,74	+ 0,43	19,919
687	3,4	κ Draconis.....	12 26	6	19	16	45,22	2,47	16	44,51	16	46,72	+ 0,71	+ 1,50	19,912

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from Greenwich. A.S.C.		Annual Variation.
					h.	m.	s.		m.	s.	m.	s.	s.	s.	
688	4	<i>a</i> Muscæ.....	12 27	5	158	12	8,07	3,94		12	6,36	+	1,71	+ 19,902	
689	5	<i>r</i> Centauri.....	12 28	5	137	36	32,43	4,81		36	29,23	+	3,20	19,888	
690	5	<i>l</i> Centauri.....	12 31	3	129	3	19,26	5,20		3	17,07	+	2,19	19,863	
691	3	<i>γ</i> Centauri.....	12 32	5	138	1	42,31	4,63		1	40,95	+	1,36	19,845	
692	4	<i>β</i> Muscæ.....	12 36	6	157	10	53,64	4,33		10	50,60	+	3,04	19,796	
693	6	<i>33</i> Virginis.....	12 38	4	79	30	52,76	3,40		30	47,38	+	5,38	19,771	
694	2	<i>β</i> Crucis.....	12 38	6	148	45	42,30	3,69		45	42,39	-	0,09	19,770	
695	5	<i>n</i> Centauri.....	12 44	4	129	15	29,02	3,79		15	23,33	+	5,69	19,674	
696	5	<i>o</i> Centauri.....	12 45	5	146	15	27,16	1,30		15	24,68	+	2,48	19,663	
697	3	<i>e</i> Ursæ Major..	12 46	5	33	7	16,47	4,23	7	16,93	7	19,84	- 0,46	- 3,37	19,631
698	2,3	<i>a</i> Canum Ven..	12 48	9	50	46	3,20	3,99	46	2,84	46	1,60	+ 0,36	+ 1,60	19,604
699	4,5	<i>r</i> Comæ Ber...	12 50	6	71	40	38,19	5,23	40	39,54	40	31,00	- 1,35	+ 7,19	19,558
700	4	<i>δ</i> Muscæ.....	12 51	5	160	38	12,23	2,74		38	1,91			+ 10,32	19,554
701	5	<i>37</i> Comæ Ber...	12 52	5	58	19	1,39	5,36		18	1,07			+ 0,32	19,526
702	3,4	<i>c</i> Virginis.....	12 54	4	78	7	47,66	1,97	7	49,46	7	46,27	- 1,80	+ 1,39	19,495
703	5	<i>z</i> Centauri.....	13 2	5	132	27	53,33	5,65		27	53,57			- 0,24	19,320
704	5	<i>53</i> Virginis.....	13 3	3	105	17	8,31	3,01		17	0,32			+ 7,99	19,290
705	4,5	<i>61</i> Virginis.....	13 9	5	107	22	5,47	4,59	22	7,18	22	5,20	- 1,71	+ 0,27	20,208
706	4,5	<i>γ</i> Hydræ Con...	13 10	6	112	16	37,14	3,51	16	37,86	16	33,15	- 0,72	+ 3,99	19,123
707	5	<i>21</i> Canum Ven..	13 11	4	39	25	40,52	5,72		25	40,34			+ 0,18	19,089
708	1	<i>a</i> Virginis.....	13 16	21	100	16	39,27	5,05	16	35,06	16	32,07	+ 4,21	+ 7,20	18,944
709	3	<i>ζ</i> Ursæ Maj....	13 17	7	34	11	24,38	4,67	11	23,44	11	22,08	+ 0,94	+ 2,30	18,921
710	5	<i>i</i> Virginis.....	13 18	5	101	49	32,65	4,44	49	31,42	49	31,85	+ 1,23	+ 0,80	18,901
711	7	* Virginis.....	13 21	3	90	27	9,30	4,03							18,818
712	4	<i>d</i> Centauri.....	13 21	5	128	31	51,65	2,68		31	49,03			+ 2,62	18,797
713	6	<i>73</i> Virginis.....	13 23	4	107	51	14,69	4,58		51	15,21			- 0,52	18,746
714	4	<i>ζ</i> Virginis.....	13 26	6	89	43	46,33	2,82	43	44,13	43	39,40	+ 2,20	+ 6,93	18,647
715	6	<i>ε</i> Centauri.....	13 29	4	118	41	34,49	3,78		41	35,74			- 1,25	18,544
716	3	<i>e</i> Centauri.....	13 29	7	142	36	9,18	5,76		36	4,84			+ 4,34	18,544
717	6	<i>l</i> Bootis.....	13 32	5	69	11	12,56	3,79		11	11,33			+ 1,23	18,431
718	5	<i>i</i> Centauri.....	13 36	5	122	11	7,75	5,69		11	8,37			- 0,62	18,307
719	5	<i>r</i> Bootis.....	13 39	5	71	41	50,11	3,90		41	48,33			+ 1,78	18,195
720	4	<i>μ</i> Centauri.....	13 39	5	131	37	39,64	2,61		37	36,30			+ 3,34	18,186
721	5	<i>g</i> Centauri.....	13 40	3	123	36	11,08	4,23		36	13,44			- 2,36	18,179
722	2,3	<i>η</i> Ursæ Maj....	13 41	14	39	50	29,37	3,35	50	25,90	50	23,74	+ 3,47	+ 5,63	18,134
723	4	<i>v</i> Bootis.....	13 41	5	73	21	36,45	3,32	21	37,67	21	33,10	- 1,92	+ 3,35	18,118
724	4,5	<i>k</i> Centauri.....	13 42	4	122	9	9,65	3,41	9	4,70	9	5,78	+ 4,95	+ 3,87	18,088
725	4,5	<i>i</i> Draconis.....	13 46	3	24	26	27,51	3,76		26	26,77			+ 0,74	17,919
726	5	<i>η</i> Bootis.....	13 47	5	70	45	8,45	2,99	45	5,84	45	4,02	+ 2,61	+ 4,43	17,914
727	3	<i>φ</i> Centauri.....	13 48	4	131	16	15,52	2,18		16	10,04			+ 5,48	17,859
728	5	<i>v</i> Centauri.....	13 48	4	133	58	27,43	2,61		58	18,45			+ 8,98	17,849
729	5	<i>v</i> Centauri.....	13 51	5	134	46	52,34	4,63		46	46,31			+ 6,03	17,731
730	1	<i>β</i> Centauri.....	13 52	7	149	33	9,09	3,39		33	2,96			+ 6,13	17,700
731	5	<i>χ</i> Centauri.....	13 56	5	130	21	57,14	6,09		21	49,00			+ 8,14	17,542
732	4,5	<i>π</i> Hydræ Con...	13 57	5	115	51	51,26	6,32		51	48,83			+ 2,43	17,499
733	2	<i>θ</i> Centauri.....	13 57	5	125	32	6,63	3,20		32	4,44			+ 2,19	17,499

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Differ-ence of Ex-tremes.	Green-wich Catalogue		A.S. Cata-logue.		Difference from		Annual Variation.
					°	'	"		s.	m.	s.	m.	s.	s.	
734	3,4	α Draconis.....	13 59	5	24	48	57,92	2,85	48	52,00	48	51,55	+ 5,92	+ 6,37	+ 17,367
735	5 6	Hydræ Con...	14 3	4	116	27	41,95	1,19			27	36,50		+ 5,45	17,323
736	4	κ Virginis.....	14 4	4	99	29	1,03	5,40	28	58,51	29	3,52	+ 2,52	- 2,49	17,187
737	4	ι Virginis.....	14 7	5	95	11	29,40	4,36	11	23,62	11	20,17	+ 5,78	+ 9,23	17,038
738	1	α Bootis.....	14 8	14	69	56	5,63	4,49	56	2,62	55	59,77	+ 3,01	+ 5,86	18,962
739	4,5	ι Lupi.....	14 9	4	135	16	26,11	2,87			16	17,46		+ 8,65	16,971
740	4	λ Bootis.....	14 10	4	43	7	57,37	1,69	7	57,83	7	56,43	- 0,46	+ 0,94	16,908
741	5	τ^1 Lupi.....	14 15	5	134	27	5,80	1,72			26	59,33		+ 6,47	16,652
742	5	τ^2 Lupi.....	14 15	4	134	36	38,52	2,01			36	32,60		+ 5,92	16,651
743	4	θ Bootis.....	14 19	5	37	21	56,11	4,46	21	55,17	21	56,41	+ 0,94	- 0,30	16,988
744	5	σ Lupi.....	14 21	4	139	42	8,21	3,70			42	0,58		+ 7,63	16,356
745	4	ρ Bootis.....	14 25	5	58	52	58,56	5,26	52	58,37	52	59,30	+ 0,19	- 0,74	16,189
746	3,4	γ Bootis.....	14 25	4	50	56	56,70	1,52	56	56,66	56	53,79	+ 0,04	+ 2,91	16,151
747	5	σ Bootis.....	14 27	5	59	31	3,40	4,80			30	59,68		+ 3,72	16,045
748	1	α^2 Centauri.....	14 28	3	150	7	53,43	2,37			7	47,11		+ 6,32	15,995
749	3,4	ζ Bootis.....	14 33	5	75	32	33,89	2,39	32	32,51	32	27,07	+ 1,38	+ 6,82	15,738
750	5	ϵ^1 Centauri.....	14 33	5	124	26	27,37	2,35			26	15,44		+ 11,93	15,724
751	4,5	β^4 Bootis.....	14 36	4	62	44	59,33	4,11	44	58,81	44	56,73	+ 0,52	+ 2,60	15,579
752	5	δ Libræ.....	14 40	3	117	15	2,06	3,09			15	0,11		+ 1,95	15,335
753	5	σ Lupi.....	14 41	4	132	52	10,26	5,29			52	5,11		+ 5,15	15,321
754	3	α^2 Libræ.....	14 41	8	105	20	5,11	5,09	20	2,93	19	59,00	+ 2,18	+ 6,11	15,270
755	3,4	β Lupi.....	14 47	7	132	26	46,79	4,20			26	42,48		+ 4,31	14,927
756	5	ξ^2 Libræ.....	14 47	5	100	43	21,96	3,23	43	21,08	43	16,93	+ 0,88	+ 5,03	14,921
757	3	κ Centauri.....	14 48	5	131	25	10,84	2,88			25	10,20		+ 0,64	14,886
758	5	π Lupi.....	14 54	5	136	22	59,20	1,37			22	53,30		+ 5,90	14,563
759	3,4	γ Libræ.....	14 54	5	114	36	44,43	4,46	36	41,48	36	37,31	+ 2,95	+ 7,12	14,530
760	3	β Bootis.....	14 55	5	48	56	19,00	3,93	56	20,63	56	19,45	- 1,63	- 0,45	14,446
761	5	ϵ Bootis.....	14 59	5	64	28	0,01	2,71			28	5,44		+ 0,57	14,183
762	4	ζ Lupi.....	15 0	5	141	26	56,45	4,02			26	54,30		+ 2,15	14,164
763	5	κ Lupi.....	15 0	3	138	5	12,97	2,46			5	15,30		- 2,33	14,162
764	5	μ Lupi.....	15 7	3	137	14	45,24	4,05			14	34,43		+ 10,81	13,748
765	5	χ Bootis.....	15 7	4	60	12	18,31	0,51			12	5,62		+ 12,69	13,709
766	2,3	β Libræ.....	15 8	5	98	45	9,79	5,32	45	13,10	45	9,36	- 3,31	- 0,43	13,678
767	5	ϕ^2 Lupi.....	15 12	5	126	14	43,58	3,49			14	41,47		+ 2,11	13,390
768	4	μ Bootis.....	15 18	5	52	1	36,14	4,08	1	33,87	1	28,50	+ 2,27	+ 7,64	13,013
769	4	β Cor. Bor.....	15 21	5	60	18	27,24	4,54	18	25,48	18	22,59	+ 1,76	+ 4,65	12,828
770	3,4	γ^2 Ursæ Min...	15 21	5	17	33	51,65	2,11	33	52,40	33	50,59	- 0,75	+ 1,06	12,813
771	3	ι Draconis.....	15 21	5	30	26	24,91	3,26	26	22,78	26	23,08	+ 2,13	+ 1,83	12,806
772	4,5	γ Libræ.....	15 26	3	104	13	9,38	0,91	13	9,46	13	2 80	- 0,08	+ 6,58	12,475
773	2	α Cor. Bor.....	15 27	8	62	42	40,16	3,45	42	41,82	42	37,10	- 1,66	+ 3,06	12,375
774	4,5	δ Libræ.....	15 28	4	119	12	52,73	2,46	12	51,47	12	49,39	+ 1,26	+ 3,34	12,324
775	5	ι Serpentic.....	15 34	5	69	46	51,19	2,13			46	46,08		+ 5,11	11,924
776	4,5	γ Libræ.....	15 35	4	105	7	43,18	1,65			7	35,03		+ 8,15	11,885
777	2,3	α Serpentic.....	15 36	7	83	2	11,49	4,34	2	12,21	2	10,76	- 0,72	+ 0,73	11,788
778	4,5	λ Serpentic.....	15 38	5	82	6	43,21	5,44	6	42,38	6	41,63	+ 0,83	+ 1,58	11,625
779	3,4	μ Serpentic.....	15 41	3	92	54	22,27	1,66			54	18,71		+ 3,56	11,442

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from		Annual Variation.	
					°	'	"		s.	m.	s.	m.	s.	s.		+
780	4,5	δ Cor. Bor.	h. m. 15 42	5	63	24	31,16	4,13	24	32,72	24	27,95	-1,56	+	3,21	+ 11,319
781	3,4	π Scorpil.	15 49	5	115	37	13,97	3,62	37	10,45	37	10,44	+ 3,52	+	3,53	10,873
782	3	γ Serpenti.	15 49	6	73	46	49,11	1,69	46	52,27	46	48,94	- 3,16	+	0,17	12,182
783	3	δ Scorpil.	15 50	6	112	8	2,04	5,84			7	53,11		+	8,93	10,747
784	5	δ Norma.	15 54	4	134	42	23,00	2,78			42	16,82		+	6,18	10,434
785	4,5	π Serpenti.	15 55	5	66	43	14,45	2,27	43	16,57	43	11,34	- 2,12	+	3,11	10,400
786	5	v Herculis.	15 57	4	43	29	23,01	3,55	29	27,13	29	24,82	- 4,12	-	1,81	10,211
787	3,4	θ Draconis.	15 59	6	30	58	53,88	4,27	58	54,11	58	51,89	- 0,23	+	1,99	10,118
788	5	δ Triang. Aus.	16 0	4	153	14	29,25	5,86			14	26,32		+	2,93	10,016
789	3	δ Ophiuchi	16 5	6	93	15	5,87	3,62	15	8,57	15	3,91	- 2,70	+	1,96	9,607
790	5	n Scorpil.	16 6	5	97	54	53,06	3,26			54	50,64		+	2,42	10,064
791	3	e Ophiuchi	16 9	5	94	16	24,55	3,86	16	26,20	16	20,47	- 1,65	+	4,08	9,306
792	4	σ Scorpil.	16 11	5	115	10	48,82	2,54	10	45,15	10	39,89	+ 3,67	+	8,93	9,187
793	5	σ Serpenti.	16 13	4	88	34	0,65	1,04			33	59,88		+	0,77	8,984
794	5	ξ Cor. Bor.	16 15	5	58	42	36,18	2,85			42	36,33		-	0,15	8,827
795	5	v Ophiuchi	16 19	5	97	59	6,87	3,60			59	8,93		-	2,06	8,580
796	1	α Scorpil.	16 19	5	116	2	56,03	6,13	2	54,43	2	52,70	+ 1,60	+	3,33	8,550
797	5	25 Herculis.	16 19	4	52	13	0,36	1,63			12	55,34		+	5,02	8,522
798	2,3	β Herculis.	16 23	6	68	8	8,95	3,06	8	11,98	8	10,03	- 3,03	-	1,08	8,239
799	3,4	ζ Ophiuchi	16 23	6	100	13	7,87	5,86	13	3,74	12	58,81	+ 4,13	+	9,03	7,847
800	4	σ Herculis.	16 29	5	47	12	37,26	2,27	12	37,40	12	34,47	- 0,14	+	2,79	7,781
801	5	m Scorpil.	16 32	6	107	24	25,62	1,69			24	22,65		+	2,97	7,528
802	3	ζ Herculis.	16 35	4	58	5	8,74	1,17	5	11,86	5	4,69	- 3,12	+	4,05	7,275
803	3	η Herculis.	16 37	7	50	45	8,58	4,47	45	8,24	45	3,96	+ 0,34	-	4,62	7,096
804	3	e Scorpil.	16 39	5	123	58	33,79	0,60			58	39,37		-	0,58	6,923
805	5	r Ophiuchi	16 40	4	100	28	36,66	1,85			28	29,26		+	7,40	6,819
806	5	K Herculis.	16 42	5	82	27	14,51	4,16			27	9,92		+	4,59	6,684
807	5	s Herculis.	16 44	4	59	54	3,74	2,10			53	55,49		+	8,25	6,525
808	3,4	ζ Ara.	16 45	5	145	42	44,82	3,54			42	38,60		+	6,22	6,476
809	5	53 Herculis.	16 47	5	58	0	48,09	1,19			0	49,49		-	1,40	6,317
810	4	κ Ophiuchi	16 50	7	80	21	25,41	2,29	21	25,22	21	15,07	+ 0,19	+	10,34	6,058
811	5	k Scorpil.	16 54	5	123	52	32,37	2,08			52	32,84		-	0,47	5,721
812	3	e Herculis.	16 54	5	58	49	8,67	3,33	49	10,77	49	7,22	- 2,10	+	1,48	5,711
813	5	h Draconis.	16 55	5	24	36	26,35	2,72			36	19,89		+	6,46	5,601
814	5	60 Herculis.	16 57	5	77	1	15,29	2,77			1	9,26		+	6,03	5,399
815	4	γ Scorpil.	17 0	4	133	0	8,89	1,81			0	10,55		-	1,66	5,188
816	2,3	γ Ophiuchi	17 0	5	105	30	29,41	6,81	30	26,95	30	21,14	+ 2,46	+	8,27	5,134
817	4	μ Draconis.	17 2	5	35	18	19,49	3,20	18	17,55	18	15,82	+ 1,94	+	3,67	5,034
818	4,5	A Ophiuchi	17 5	5	116	20	46,21	2,82	20	43,60	20	40,94	+ 2,61	+	5,27	6,021
819	3,4	a Herculis.	17 7	9	76	24	40,53	2,85	24	38,40	24	33,09	+ 2,13	+	7,44	4,603
820	5,6	o Ophiuchi 41..	17 8	5	90	14	49,81	3,51	14	52,32	14	48,85	- 2,51	+	0,96	4,518
821	4	u Herculis.	17 11	5	56	42	45,80	1,69	42	45,73	42	44,24	+ 0,07	+	1,56	4,248
822	3	γ Ara.	17 11	4	146	12	23,60	1,75			12	22,98		+	0,62	4,241
823	4,5	e Herculis.	17 12	5	52	31	33,90	1,22	31	36,58	31	27,01	- 2,68	+	6,89	4,184
824	4	δ Ara.	17 16	5	150	31	47,83	0,41			31	46,18		+	1,65	3,841
825	5	d Ophiuchi	17 17	5	119	42	21,24	2,15			42	19,79		+	1,45	3,781

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from		Annual Variation.	
					a	l	u		s.	m.	s.	m.	s.	Greenwich.		A.S.C.
826	4	ρ Herculis.....	17 18	6	52	41	37,16	2,42	41	37,88	41	30,36	-0,72	+	6,80	+ 3,668
827	4,5	σ Ophiuchi....	17 18	5	85	42	23,66	3,82	42	21,77	42	15,50	+1,89	+	8,16	3,645
828	5	e^* Ophiuchi....	17 21	5	113	49	21,21	4,13	49	24,94	49	20,02	-3,73	+	1,19	3,389
829	3	λ Scorpii.....	17 22	5	126	58	14,02	2,97			58	6,78		+	7,24	3,301
830	4,5	λ Herculis.....	17 24	5	63	45	20,06	2,34	45	22,93	45	21,86	-2,87	-	1,80	3,147
831	5	θ Scorpii.....	17 25	4	132	52	45,29	3,85			52	42,20		+	3,09	3,039
832	2	β Draconis....	17 27	4	37	34	11,58	4,01	34	13,78	34	11,80	-2,20	-	0,22	2,912
833	2	α Ophiuchi....	17 27	8	77	18	36,62	2,68	18	36,65	18	31,51	-0,03	+	5,11	2,872
834	5	ξ Serpentis....	17 28	5	105	17	7,39	1,99			16	58,06		+	9,33	2,802
835	5	μ Ophiuchi....	17 29	3	98	0	27,08	1,11	0	30,46	0	24,68	-3,38	+	2,40	2,737
836	3	κ Scorpii.....	17 31	5	128	55	57,23	1,85			55	55,82		+	1,41	2,552
837	4,5	σ Serpentis....	17 32	4	102	46	33,38	1,23	46	37,94	46	32,11	-4,56	+	1,27	2,455
838	5	D Ophiuchi....	17 33	3	111	35	31,91	0,34	35	32,42	35	25,14	-0,51	+	6,77	2,334
839	3	β Ophiuchi....	17 35	7	85	21	21,98	3,90	21	19,46	21	13,90	+2,52	+	6,08	2,176
840	4,5	ι^* Scorpii.....	17 36	5	130	3	15,09	4,02			3	1,00		+	14,09	2,121
841	4	γ Telescopii....	17 38	5	126	58	45,56	2,38			58	43,44		+	2,12	1,896
842	4	γ Ophiuchi....	17 39	4	87	13	21,50	1,26	13	20,32	13	15,44	+1,18	+	6,06	1,802
843	4	μ Herculis....	17 40	6	62	10	29,31	2,82	10	30,71	10	29,77	-1,40	-	0,46	2,604
844	5	* Sagittarii....	17 48	6	120	13	38,03	2,09			13	33,56		+	4,47	1,723
845	5	b Sagittarii....	17 49	6	113	47	27,58	3,05	47	31,14	47	25,06	-3,56	+	2,52	0,925
846	4	ν Ophiuchi....	17 50	5	99	44	45,29	2,01	44	42,31	44	33,00	+2,98	+	12,29	0,903
847	4	θ Herculis....	17 50	5	52	43	18,51	3,48	43	20,58	43	16,33	-2,07	+	2,18	0,838
848	4	ξ Herculis....	17 51	4	60	43	43,13	1,15	43	42,10	43	37,27	+1,03	+	5,86	0,773
849	2	γ Draconis....	17 53	18	38	29	14,21	4,34	29	17,80	29	17,92	-3,59	-	3,71	0,642
850	4	θ Aræ.....	17 53	5	140	5	31,93	3,81			5	28,30		+	3,63	0,576
851	5	τ Ophiuchi....	17 54	5	98	10	17,17	5,99			10	18,05		-	0,88	0,540
852	5	Q Herculis....	17 55	3	69	9	36,91	2,60			9	34,29		+	2,62	0,428
853	4,5	p Ophiuchi....	17 57	5	87	27	11,49	2,41	27	13,21	27	13,76	-1,72	-	2,27	1,444
854	5	* Sagittarii....	17 57	5	118	27	56,16	4,49			27	58,56		-	2,40	0,335
855	4	S ² Ophiuchi....	17 59	5	80	27	13,99	1,63	27	13,12	27	6,65	+0,87	+	7,34	+ 0,062
856	4	\circ Herculis....	18 0	5	61	15	18,28	3,40	15	18,60	15	15,83	-0,32	+	2,45	- 0,080
857	3,4	μ Sagittarii....	18 4	5	111	5	40,43	1,49	5	40,41	5	36,85	+0,02	+	3,58	0,314
858	5	A Herculis....	18 5	4	58	37	49,65	1,88			37	46,56		+	3,09	0,481
859	4	β Telescopii....	18 6	5	126	48	8,74	3,38			48	0,74		+	8,00	0,536
860	3,4	δ Sagittarii....	18 10	6	119	53	29,34	5,62	53	24,25	53	26,34	+5,09	+	3,00	0,884
861	5	G Herculis....	18 12	5	65	37	8,50	1,22			37	1,51		+	6,99	1,064
862	4	η Serpentis....	18 12	4	92	56	4,55	2,96	56	8,76	56	3,35	-4,21	+	1,20	0,414
863	3	e Sagittarii....	18 13	5	124	27	19,32	1,44			27	6,30		+	13,02	1,127
864	4,5	α Telescopii....	18 14	4	136	3	7,90	1,05			2	56,00		+	11,90	1,255
865	4	λ Sagittarii....	18 17	5	115	30	25,88	5,56	30	21,76	30	22,68	+4,12	+	3,20	1,527
866	5	* Clypei Sob....	18 19	6	104	39	58,07	6,28			39	55,89		+	2,18	1,707
867	5	b Draconis....	18 21	6	31	17	42,73	0,47			17	41,53		+	1,20	1,871
868	4,5	χ Draconis....	18 24	5	17	20	32,67	2,64	20	32,20	20	31,04	+0,47	+	1,63	2,104
869	3	δ Ursæ Min....	18 27	5	13	24	48,43	1,93	24	50,90	24	51,27	-2,47	-	2,84	2,357
870	5	* Pavonis.....	18 29	5	155	1	0,58	3,74			1	22,25		-	21,67	2,508
871	1	α Lyrae.....	18 31	40	51	22	8,64	4,07	22	8,55	22	6,94	+0,09	+	1,70	2,718

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from Greenwich.		Annual Variation.
					h.	m.	s.		m.	s.	m.	s.	s.	s.	
872	5	o Aquilæ.....	18 33	5	99	12	23,01	3,41		12	17,94		+ 5,07	- 2,873	
873	4,5	φ Sagittarii.....	18 35	5	117	9	20,46	2,82	9	19,22	9	17,89	+ 1,24	+ 2,57	3,052
874	5	K Herculis.....	18 38	4	69	36	36,14	1,03			36	31,24		+ 4,90	3,337
875	5	e Lyræ.....	18 38	5	50	30	8,80	1,60			30	2,37		+ 6,43	3,368
876	5	5 Lyræ.....	18 39	6	50	33	37,65	3,52			33	32,05		+ 5,60	3,371
877	3	β Lyræ.....	18 44	7	56	49	42,48	6,04	49	42,84	49	39,64	- 0,36	+ 2,84	3,807
878	5	v ¹ Sagittarii.....	18 44	5	112	56	37,49	1,75	56	38,38	56	36,77	- 0,89	+ 0,72	3,816
879	3	σ Sagittarii.....	18 45	5	116	29	50,29	1,83	29	51,58	29	50,97	- 1,29	- 0,68	3,886
880	5	O Herculis.....	18 48	5	67	33	44,33	1,86			33	44,22		+ 0,11	4,131
881	5	δ ² Lyræ.....	18 49	5	53	18	38,91	1,84			18	38,78		+ 0,13	4,214
882	2,3	ζ Sagittarii.....	18 52	5	120	6	48,99	2,59	6	43,29	6	45,07	+ 5,70	+ 3,92	4,491
883	3,4	e Aquilæ.....	18 52	3	75	9	21,21	1,34	9	16,80	9	14,13	+ 4,41	+ 7,08	4,500
884	3	γ Lyræ.....	18 53	5	57	32	12,45	2,21	32	14,05	32	10,83	- 1,60	+ 1,62	4,558
885	5	γ Cor. Aust.....	18 55	5	127	17	48,43	4,81			17	34,82		+ 13,61	4,756
886	4	τ Sagittarii.....	18 56	3	117	54	29,30	1,43	54	29,86	54	28,71	- 0,56	+ 0,59	4,876
887	3	λ Aquilæ.....	18 57	5	95	7	47,72	2,58	7	43,16	7	37,34	+ 4,56	+ 10,38	4,952
888	3	ζ Aquilæ.....	18 58	10	76	22	51,97	3,97	22	53,02	22	43,57	- 1,05	+ 8,40	4,983
889	5	α Cor. Aust.....	18 58	3	128	9	26,29	0,63			9	23,00		+ 3,29	5,009
890	4,5	π Sagittarii.....	19 00	5	111	17	1,03	2,79			16	59,13	- 2,19	+ 1,90	5,158
891	5	B Aquilæ.....	19 3	5	98	12	50,41	2,36			12	48,45		+ 1,96	5,478
892	5	d Sagittarii.....	19 8	5	109	14	43,72	1,83	14	44,23	14	41,94	- 0,51	+ 1,78	5,833
893	5	η Lyræ.....	19 8		51	8	24,30	5,83			8	22,76		+ 1,54	5,857
894	5	n Draconis.....	19 8	3	33	25	32,58	0,47			25	34,55		- 1,97	5,897
895	5	ω ¹ Aquilæ.....	19 10	5	78	42	8,27	0,89			42	3,55		+ 4,72	6,013
896	4	β ¹ Sagittarii.....	19 10	5	134	46	2,35	1,84			45	54,82		+ 7,53	6,060
897	3	δ Draconis.....	19 12	12	22	38	4,44	2,87	38	8,38	38	9,16	- 3,94	- 4,72	6,234
898	4	κ Cygni.....	19 13	5	36	56	25,83	1,43	56	25,76	56	25,64	+ 0,07	+ 0,19	6,290
899	5	b Aquilæ.....	19 17	5	78	24	36,45	2,63			24	33,22		+ 3,23	6,316
900	3,4	δ Aquilæ.....	19 17	12	87	12	56,37	3,25	12	56,01	12	50,52	+ 0,36	+ 5,85	6,601
901	4,5	τ Draconis.....	19 19	5	16	57	38,72	2,07	57	39,10	57	35,71	- 0,38	+ 3,01	6,753
902	4	κ Draconis.....	19 20	5	24	36	38,57	3,15	36	37,49	36	35,74	+ 1,08	+ 2,83	6,836
903	4	β Vulpeculæ.....	19 22	5	65	40	19,52	1,63	40	18,96	40	15,67	+ 0,56	+ 3,85	6,988
904	3	β ² Cygni.....	19 24	6	62	23	23,56	3,54	23	24,42	23	21,19	- 0,86	+ 2,37	7,171
905	5	ε Cygni.....	19 25	5	38	37	36,36	4,46	37	37,48	37	36,30	- 1,12	+ 0,06	7,298
906	4,5	μ Aquilæ.....	19 26	5	82	58	21,32	0,74	58	25,11	58	19,40	- 3,79	+ 1,92	7,328
907	4	κ Aquilæ.....	19 28	5	97	23	45,36	2,47	23	47,34	23	42,45	- 1,98	+ 2,91	7,487
908	5	ε Aquilæ.....	19 28	5	91	39	15,29	1,09			39	11,03		+ 4,26	7,502
909	4	θ Cygni.....	19 32	5	40	10	4,61	2,23	10	0,86	9	58,54	+ 3,75	+ 6,07	7,821
910	4	α Sagittæ.....	19 32	5	72	22	7,39	3,56	22	8,54	22	1,87	- 1,15	+ 5,52	7,871
911	4	φ Cygni.....	19 33	4	60	13	55,78	3,07	13	52,09	13	47,23	+ 3,69	+ 8,55	7,885
912	5	β Sagittæ.....	19 33	5	72	54	32,13	2,30			54	30,87		+ 1,26	7,945
913	5	15 Cygni.....	19 38	5	53	2	55,57	5,93			2	51,84		+ 3,73	8,323
914	3	γ Aquilæ.....	19 38	37	79	47	36,50	5,38	47	33,40	47	30,64	+ 3,10	+ 5,86	8,326
915	4	δ ¹ Sagittæ.....	19 40	5	71	52	36,65	1,67	52	38,55	52	35,53	- 1,90	+ 1,12	8,455
916	5	χ Cygni.....	19 40	5	56	39	39,59	2,55			39	37,14		+ 2,45	8,468
917	5	ζ Sagittæ.....	19 41	4	71	16	32,59	1,30			16	28,12		+ 4,47	8,583

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.				Difference of Extremes.		Greenwich Catalogue		A.S. Catalogue.		Difference from Greenwich.		A.S.C.	Annual Variation.
					°	'	"	s.	m.	s.	m.	s.	s.	s.				
918	1,2	<i>a</i> Aquilæ.....	19 42	70	81	34	16,88	5,99	34	19,35	34	14,14	-2,47	+	2,74	-	8,667	
919	4	<i>γ</i> Aquilæ.....	19 44	5	89	25	19,50	2,02	25	18,90	25	14,01	+0,60	+	5,49	+	8,771	
920	5	<i>ξ</i> Aquilæ.....	19 46	3	81	58	11,99	1,70			58	4,64		+	7,35		8,943	
921	5	<i>b</i> Sagittariii...	19 47	5	117	36	34,06	1,22	36	35,50	36	30,39	-1,44	+	3,67		8,982	
922	3,4	<i>β</i> Aquilæ.....	19 47	26	84	0	31,34	4,52	0	33,40	0	29,97	-2,06	+	1,37		8,478	
923	4,5	<i>γ</i> Sagittæ.....	19 51	5	70	57	40,89	0,74	57	40,84	57	30,08	+0,05	+	10,81		9,347	
924	5	<i>f</i> Vulpeculæ...	19 52	4	67	21	17,35	2,52			21	10,05		+	7,30		9,400	
925	4,5	<i>c</i> Sagittariii...	19 52	5	118	10	17,26	2,42	10	18,20	10	13,29	-0,94	+	3,97		9,424	
926	5	<i>L</i> Sagittariii...	19 54	5	122	30	22,39	1,53			31	19,75		+	2,64		9,527	
927	5	<i>g</i> Vulpeculæ...	19 54	4	62	42	29,09	2,35			42	26,54		+	2,55		9,571	
928	5	<i>ρ</i> Draconis....	20 2	5	22	36	26,05	1,38	36	28,58	36	28,71	-2,53	-	2,66		10,173	
929	3,4	<i>θ</i> Aquilæ.....	20 3	6	91	18	59,48	1,71	18	58,33	18	56,05	+1,15	+	3,43		10,212	
930	5	66 Draconis...	20 3	4	28	29	33,55	0,46			29	39,40		-	5,85		10,234	
931	5	<i>b</i> ² Cygni.....	20 3	4	53	39	8,93	1,93			39	8,44		+	0,49		10,256	
932	4	<i>a</i> ¹ Capricorni...	20 8	5	103	1	33,50	0,46	1	25,46	1	20,77	-2,96	+	2,73		10,637	
933	4	<i>o</i> ² Cygni.....	20 8	5	43	46	1,42	1,96	46	3,94	45	59,12	-2,52	+	2,30		10,641	
934	3	<i>a</i> ² Capricorni...	20 9	23	103	3	38,41	5,90	3	43,66	3	38,76	-5,25	-	0,35		10,667	
935	5	<i>o</i> Vulpeculæ...	20 10	5	65	50	37,90	2,27			50	36,54		+	1,36		10,733	
936	7	<i>β</i> ¹ Capricorni...	20 11	5	105	18	34,71	2,17			18	29,65		+	5,06		10,858	
937	2	<i>α</i> Pavonis.....	20 12	5	147	16	0,86	2,07			15	57,73		+	3,13		10,927	
938	4,5	<i>κ</i> Cephei.....	20 14	5	12	48	1,59	3,64	48	4,60	48	8,21	-3,01	-	6,62		11,092	
939	3	<i>γ</i> Cygni.....	20 16	5	50	16	51,90	5,99	16	48,95	16	46,64	+2,95	+	5,26		11,216	
940	5	<i>h</i> Cygni.....	20 17	5	58	21	0,41	2,26			21	1,30		-	0,89		11,285	
941	5	<i>π</i> Capricorni...	20 18	5	118	45	30,56	2,14			45	28,13		+	2,43		11,322	
942	5	<i>ρ</i> Capricorni...	20 19	4	108	21	53,84	3,67	21	57,72	21	52,87	-3,88	+	0,97		11,435	
943	5	<i>G</i> Aquilæ.....	20 21	7	93	26	26,49	4,76			26	24,29		+	2,20		11,550	
944	4,5	<i>i</i> Cygni.....	20 22	6	60	11	30,59	1,87	11	26,98	11	23,30	+3,61	+	7,29		11,671	
945	5	<i>w</i> ² Cygni.....	20 25	5	41	36	47,34	1,17			36	46,70		+	0,64		11,836	
946	4	<i>e</i> Delphini.....	20 25	5	79	15	56,74	2,08	15	57,29	15	50,73	-0,55	+	6,01		11,857	
947	5	<i>θ</i> Cephei.....	20 27	5	27	34	16,94	1,60			34	22,82		-	5,88		11,971	
948	5	<i>ζ</i> Delphini.....	20 27	5	75	54	13,40	2,05			54	4,13		+	9,27		12,017	
949	3	<i>β</i> Pavonis.....	20 30	5	156	47	57,89	6,70			48	4,05		-	6,16		12,169	
950	4,5	<i>θ</i> Delphini.....	20 31	4	77	16	20,92	2,66	16	22,54	16	18,56	-1,62	+	2,36		12,250	
951	3,4	<i>a</i> Delphini.....	20 32	5	74	40	48,00	2,84	40	44,79	40	38,26	+3,21	+	9,74		12,322	
952	5	<i>δ</i> Delphini.....	20 36	6	75	31	37,70	2,26			31	30,57		+	7,13		12,581	
953	1	<i>a</i> Cygni.....	20 36	70	45	19	14,34	6,46	19	12,76	19	9,66	+1,58	+	4,68		12,588	
954	4,5	<i>e</i> Aquarii.....	20 38	4	100	6	35,42	0,66	6	32,14	6	23,62	+3,28	+	11,80		12,780	
955	4	<i>k</i> Aquarii.....	20 39	5	95	38	28,79	2,53	38	28,17	38	21,92	+0,62	+	6,87		12,800	
956	4	<i>γ</i> Delphini.....	20 39	4	74	28	46,42	0,97	28	47,08	28	41,26	-0,66	+	5,16		12,802	
957	3	<i>e</i> Cygni.....	20 39	5	56	39	29,38	4,50	39	29,05	39	24,22	+0,33	+	5,16		12,838	
958	5	<i>λ</i> Cygni.....	20 40	5	54	7	35,95	3,21			7	27,86		+	8,09		12,935	
959	4	<i>β</i> Indi.....	20 41	4	149	4	56,72	1,61			4	48,32		+	8,40		12,981	
960	4,5	<i>μ</i> Aquarii.....	20 43	5	99	36	44,99	2,32			36	34,95		+	10,04		13,114	
961	5	57 Cygni.....	20 47	5	46	14	56,96	2,18			14	56,51		+	0,45		13,361	
962	4,5	<i>g</i> Vulpeculæ...	20 47	6	62	34	48,02	4,26	34	51,42	34	45,15	-3,40	+	2,87		13,366	
963	4	<i>ν</i> Cygni.....	20 51	5	49	28	48,46	2,43	28	48,23	28	42,22	+0,23	+	6,24		13,594	

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from		Annual Variation.
					°	'	"		s.	m.	s.	m.	s.	s.	
964	5	K Cephei.....	20 51	6	33	45	35,68	3,50			45	35,44	+	0,24	13,652
965	5	76 Draconis....	20 54	4	8	6	3,14	2,57			6	9,64	-	6,50	13,823
966	5	η Capricorni....	20 55	4	110	31	3,18	0,81	31	2,19	30	58,97	+	0,99	13,842
967	4	ξ Cygni.....	20 59	5	46	44	33,83	1,29	44	34,91	44	33,89	-	1,08	14,094
968	5	ν Aquarii.....	21 0	5	102	3	2,65	1,21	3	4,12	2	57,25	-	1,47	14,192
969	5	γ Equulei.....	21 2	4	80	32	41,17	1,86			32	38,11	+	3,06	14,300
970	3	ζ Cygni.....	21 6	5	60	27	46,03		27	44,54	27	42,24	+	1,49	14,520
971	4,5	α Equulei.....	21 7	6	85	26	50,18	2,55	26	46,35	26	42,65	+	3,83	14,618
972	5	4 Piscis Aust....	21 7	4	122	52	20,03	1,18			52	13,47	+	6,56	14,634
973	5	τ Cygni.....	21 8	3	52	40	17,49	0,51			40	13,16	+	4,33	15,158
974	4,5	σ Cygni.....	21 10	5	51	18	35,64	2,62	18	36,88	18	40,19	-	1,24	14,820
975	4,5	ν Cygni.....	21 10	5	55	48	32,29	2,27	48	32,30	48	29,52	-	0,01	14,831
976	3	γ Pavonis.....	21 12	3	156	7	20,43	1,74			7	25,81	-	5,38	14,911
977	5	γ Indi.....	21 14	5	145	23	3,75	4,28			22	49,04	+	14,71	15,016
978	4	ϵ Pegasi.....	21 14	5	70	54	52,76	1,98	54	51,23	54	48,44	+	1,53	15,024
979	5	6 Cephei.....	21 15	5	25	50	34,68	2,35			50	36,58	-	1,90	15,116
980	4	ζ Capricorni....	21 16	6	113	8	14,25	4,60	8	17,07	8	11,84	-	2,82	15,180
981	3	β Aquarii.....	21 23	5	96	18	36,23	2,46	18	37,22	18	30,62	-	0,99	15,499
982	5	ξ Cygni.....	21 23	6	44	12	4,36	2,80			12	5,84	-	1,48	15,530
983	3	β Cephei.....	21 26	6	20	10	53,13	2,96	10	48,63	10	53,82	+	4,50	15,708
984	5	ϵ Capricorni....	21 28	5	110	13	11,78	0,59	13	5,69	13	3,09	+	6,09	15,770
985	5	ρ Cygni.....	21 28	5	45	9	15,88	4,55	9	7,97	9	9,46	+	7,91	15,771
986	5	ξ Aquarii.....	21 29	5	98	36	23,68	2,16	36	28,79	36	22,25	-	5,11	15,831
987	5	T ¹ Pegasi.....	21 30	5	84	59	16,30	3,38			59	7,09	+	9,21	15,902
988	4	γ Capricorni....	21 31	5	107	25	13,99	2,05	25	16,07	25	11,79	-	2,08	15,936
989	5	41 Capricorni....	21 32	5	114	1	20,61	2,84			1	16,46	+	4,15	16,024
990	5	κ Capricorni....	21 33	6	109	37	55,95	1,73	37	54,86	37	50,65	+	1,09	16,067
991	5	9 Cephei.....	21 33	6	28	40	39,44	3,04			40	42,98	-	3,54	16,077
992	4,5	ϵ Piscis Aust....	21 35	4	123	47	30,22	0,97			47	27,23	+	2,99	16,153
993	2,3	ϵ Pegasi.....	21 36	5	80	53	45,57	0,69	53	45,85	53	41,00	-	0,28	16,206
994	4,5	π^1 Cygni.....	21 36	5	39	34	46,70	1,68	34	44,47	34	43,84	+	2,23	16,218
995	5	μ Cygni.....	21 37	4	62	1	2,78	3,38			1	1,60	+	1,18	16,242
996	3,4	δ Capricorni....	21 38	5	106	53	24,24	4,14	53	22,32	53	17,78	+	1,92	16,299
997	4,5	τ Cephei.....	21 39	6	19	27	57,13	3,26	27	57,90	28	1,09	-	0,77	16,386
998	5	π^2 Cygni.....	21 41	5	41	28	11,42	3,19	28	11,41	28	11,36	+	0,01	16,444
999	4,5	σ Cephei.....	21 41	5	29	39	27,44	1,96	39	25,86	39	28,29	-	1,58	16,444
1000	5	z Draconis....	21 41	5	18	27	10,09	2,39			27	15,27	-	5,18	16,465
1001	4	γ Gruis.....	21 44	5	128	9	17,70	1,74			9	14,02	+	3,68	16,596
1002	5	μ Capricorni....	21 44	5	104	20	35,31	2,47	20	35,14	20	29,22	+	0,17	16,617
1003	5	δ Indi.....	21 46	5	145	47	25,89	4,46			47	14,01	+	11,88	16,727
1004	5	σ Aquarii.....	21 55	5	92	58	0,53	1,51	58	4,47	57	59,95	-	3,94	17,112
1005	3	α Aquarii.....	21 57	33	91	8	12,33	6,30	8	14,74	8	11,46	-	2,41	17,227
1006	2	α Gruis.....	21 57	6	137	46	26,67	3,12			46	22,37	+	4,30	17,246
1007	5	ξ Cephei.....	21 59	5	26	11	37,35	3,93			11	39,40	-	2,05	17,307
1008	4	ϵ Pegasi.....	22 2	5	65	28	38,49	2,13	28	38,24	28	32,49	+	0,25	17,318
1009	4	θ Pegasi.....	22 2	5	84	37	53,48	2,93	37	48,46	37	43,88	+	5,02	17,428

No.	Mag.	NAME.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.	Difference of Extremes.	Greenwich Catalogue		A.S. Catalogue.		Difference from		Annual Variation.
							m.	s.	m.	s.	Greenwich.	A.S.C.	
1010	5	π^1 Pegasi.....	h. m. 22 2	5	57 38 50,44	5,96	38 53,36	39 1,87	— 2,92	— 11,43	— 17,432		
1011	4	π^2 Pegasi.....	22 2	5	57 38 50,80	3,04	38 53,36	38 49,42	— 2,56	+ 1,38	17,463		
1012	4	ζ Cephei.....	22 5	5	32 37 46,15	1,63	37 48,12	37 50,18	— 1,97	— 4,03	17,570		
1013	5	μ^1 Gruis.....	22 5	5	132 11 2,95	5,01		10 58,65		+ 4,30	17,586		
1014	5	μ^2 Gruis.....	22 6	5	132 27 48,91	3,90		27 46,34		+ 2,57	17,622		
1015	5	m Lacertæ.....	22 7	5	51 7 13,77	2,13		7 11,78		+ 1,99	16,838		
1016	3	a Toucanæ....	22 7	5	151 5 49,14	3,92		5 40,57		+ 8,57	17,647		
1017	4,5	θ Aquarii.....	22 8	5	98 37 12,82	2,81	37 17,18	37 14,62	— 4,36	— 1,80	17,691		
1018	5	a Lacertæ.....	22 9	6	53 5 24,44	1,53		5 19,95		+ 4,49	17,719		
1019	4,5	e Cephei.....	22 9	5	33 47 46,08	2,86	47 49,43	47 51,34	— 3,35	— 5,26	17,728		
1020	5	β Pegasi.....	22 12	5	85 3 28,12	1,05		3 21,05		+ 7,07	17,854		
1021	5	l Aquarii.....	22 12	5	112 24 27,75	0,75		24 25,10		+ 2,65	17,867		
1022	4	γ Aquarii.....	22 13	4	92 14 10,74	2,24	14 9,16	14 2,77	+ 1,58	+ 7,97	17,893		
1023	5	b Lacertæ.....	22 14	5	44 18 42,94	4,27		18 44,39		— 1,45	17,937		
1024	5	δ Toucanæ....	22 15	4	155 49 20,00	2,61		49 4,20		+ 15,80	17,981		
1025	5	π Aquarii.....	22 17	6	89 28 39,92	4,25	28 39,12	28 35,21	+ 0,80	+ 4,71	18,037		
1026	4	c Lacertæ.....	22 17	5	38 36 53,51	2,75	36 55,58	36 56,69	— 2,07	— 3,18	18,048		
1027	5	d Lacertæ.....	22 18	5	41 22 39,65	1,55		22 42,69		— 3,04	18,076		
1028	4	δ^1 Gruis.....	22 19	4	134 21 18,92	0,75		21 18,84		+ 0,08	18,130		
1029	5	δ^2 Gruis.....	22 19	3	134 36 40,41	2,79		36 32,87		+ 7,54	18,149		
1030	4	ζ Aquarii.....	22 21	5	90 52 55,97	1,95	52 56,84	52 53,04	— 0,87	+ 2,93	18,167		
1031	5	σ Aquarii.....	22 22	5	101 32 21,93	1,93	32 24,10	32 13,45	— 2,17	+ 8,48	18,225		
1032	4	β Piscis Aust...	22 22	5	123 12 33,13	4,69	12 30,90	12 32,41	+ 2,23	+ 0,72	18,232		
1033	4,5	δ Cephei.....	22 23	5	32 26 50,34	1,99	26 49,62	26 52,69	+ 0,72	— 2,35	18,269		
1034	4	g Lacertæ.....	22 24	5	40 35 4,78	2,88	35 3,80	35 3,64	+ 0,98	+ 1,14	18,320		
1035	5	v Aquarii.....	22 25	5	111 34 13,55	4,59	34 3,44	34 10,65	+10,11	+ 2,90	18,359		
1036	4	η Aquarii.....	22 27	5	90 59 10,84	3,48	59 10,34	59 2,42	+ 0,50	+ 8,42	18,402		
1037	4	18 Piscis Aust...	22 31	5	117 55 15,39	3,52	55 18,95	55 15,52	— 3,56	— 0,13	18,558		
1038	5	β Cephei.....	22 32	5	17 13 57,04	2,83		13 57,06		— 0,02	18,569		
1039	3	β Gruis.....	22 32	5	137 45 53,14	2,19		45 46,92		+ 6,22	18,599		
1040	5	β Cephei.....	22 33	5	27 17 31,88	2,55		17 36,14		— 4,26	18,604		
1041	3	ζ Pegasi.....	22 33	5	80 3 0,03	1,81	2 53,63	2 48,47	+ 6,40	+ 11,56	18,615		
1042	5	σ Pegasi.....	22 34	5	61 34 20,14	1,49		34 16,93		+ 3,21	18,641		
1043	3	η Pegasi.....	22 35	3	60 39 38,16	3,14	39 37,64	39 34,08	+ 0,52	+ 4,08	18,681		
1044	5	ζ Pegasi.....	22 38	5	78 41 26,85	1,63		41 21,57		+ 5,28	18,779		
1045	4	e Gruis.....	22 38	5	142 12 11,13	3,39		12 10,60		+ 0,53	18,780		
1046	4,5	λ Pegasi.....	22 38	3	67 19 15,75	0,99	19 17,48	19 8,62	— 1,73	+ 7,13	18,784		
1047	4	μ Pegasi.....	22 42	5	66 17 18,14	1,55	17 19,37	17 18,68	— 1,23	— 0,54	18,887		
1048	5	22 Piscis Aust...	22 43	3	123 46 6,00	2,37		46 2,16		+ 3,84	18,924		
1049	4	i Cephei.....	22 44	5	24 41 15,13	1,74	41 12,99	41 14,77	+ 2,14	+ 0,36	18,940		
1050	4	λ Aquarii.....	22 44	5	98 28 32,50	2,32	28 36,38	28 33,41	— 3,88	— 0,91	18,943		
1051	5,6	σ Pegasi.....	22 44	3	81 3 40,84	1,73		3 36,46		+ 4,38	18,944		
1052	5	e Cephei.....	22 44	4	29 12 0,53	3,29		12 6,86		— 6,33	18,971		
1053	3	δ Aquarii.....	22 46	7	106 43 2,17	5,95	43 1,12	42 56,39	+ 1,05	+ 5,78	18,996		
1054	1	α Piscis Aust...	22 48	48	120 30 57,25	6,07	30 55,95	30 54,81	+ 1,30	+ 2,44	19,068		
1055	5	ζ Gruis.....	22 51	5	143 39 24,67	1,01		39 26,08		— 1,41	19,136		

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Differ-ence of Ex-tremes.	Green-wich Catalogue		A.S. Cata-logue.		Difference from		Annual Variation.
					h.	m.	s.		m.	s.	m.	s.	Green-wich.	A.S.C.	
1056	6	x [*] Piscium	22 52	4	90	43	17,36	3,24		43	6,79		+ 10,57	- 19,165	
1057	4	o Andromedæ . . .	22 54	5	48	34	47,38	1,78	34	49,83	34	50,75	- 2,45	- 3,37	19,220
1058	5	β Piscium	22 55	5	87	5	18,33	2,47	5	17,62	5	11,66	+ 0,71	+ 6,67	19,247
1059	2	β Pegasi	22 56	5	62	49	54,79	1,15	49	56,72	49	51,64	+ 1,93	+ 3,15	19,255
1060	2	a Pegasi	22 56	37	75	42	9,41	5,12	42	7,63	42	4,07	+ 1,78	+ 5,34	19,273
1061	5	θ Gruis	22 57	4	134	25	50,09	1,54		25	47,77		+ 2,32	19,297	
1062	5	l Pegasi	22 58	5	81	30	6,82	2,83	30	7,24	29	59,37	- 0,42	+ 7,45	19,324
1063	4,5	c [*] Aquarii	23 0	5	112	5	15,24	2,09	5	16,25	5	9,64	- 1,01	+ 5,60	19,368
1064	5	i Gruis	23 0	5	136	9	34,51	3,81		9	36,41		- 1,90	19,374	
1065	5	c [*] Aquarii	23 1	4	113	22	15,53	3,05		22	13,45		+ 2,08	19,378	
1066	5	π Cephei	23 3	5	15	31	33,10	5,94		31	34,77		- 1,67	19,415	
1067	5,6	p Pegasi	23 3	4	82	11	46,44	3,99		11	40,99		+ 5,45	19,429	
1068	5	u Andromedæ . . .	23 5	4	41	30	55,71	3,83		30	57,58		+ 1,87	19,464	
1069	5	φ Aquarii	23 6	5	96	57	30,31	2,31	57	31,15	57	27,87	- 0,84	+ 2,44	19,479
1070	4	γ Toucanæ	23 7	5	149	9	37,87	5,82		9	43,42		- 5,55	19,518	
1071	4,5	γ Piscium	23 8	5	87	38	21,39	2,17	38	22,71	38	21,16	- 1,32	+ 0,23	19,536
1072	5	ψ [*] Aquarii	23 9	4	100	6	16,55	3,97	6	14,24	6	8,01	+ 2,31	+ 8,54	19,550
1073	5	γ App. Sculp . . .	23 10	4	123	27	2,25	0,89		27	2,86		- 0,61	19,560	
1074	5	8 Andromedæ . . .	23 10	3	41	54	25,00	2,19		54	27,31		- 2,31	19,565	
1075	6	b Piscium	23 12	3	85	32	24,16	2,79		32	22,36		+ 1,80	19,599	
1076	5	τ Pegasi	23 12	5	67	10	59,12	1,82		10	56,04		+ 3,08	19,609	
1077	5	b [*] Aquarii	23 14	5	111	1	18,79	4,80		1	17,52		+ 1,27	19,641	
1078	6	66 Pegasi	23 15	3	78	36	36,97	1,92		36	38,22		- 1,25	19,649	
1079	5	ν Pegasi	23 17	5	67	31	27,44	5,18		31	28,62		- 1,18	19,689	
1080	5	b [*] Aquarii	23 17	5	111	34	0,21	1,13		34	2,18		- 1,97	19,692	
1081	5	d Cassiopeæ	23 17	5	28	38	34,96	2,35	38	39,85	38	41,87	- 4,89	- 6,91	19,696
1082	5	o Piscium	23 19	5	84	32	38,16	3,47		32	55,00		- 16,84	19,728	
1083	5	s Cephei	23 20	5	20	34	11,72	2,16		34	10,28		+ 1,44	19,740	
1084	5	q Pegasi	23 21	5	78	10	13,52	3,43		10	8,81		+ 4,71	19,746	
1085	6,7	ω [*] Piscium	23 21	3	92	43	11,73	0,85		43	10,01		+ 1,72	19,749	
1086	5	* Cassiopeæ	23 22	5	32	22	54,31	4,74		22	56,56		- 2,25	19,771	
1087	5	b [*] Aquarii	23 24	5	111	50	50,87	3,38		50	47,98		+ 2,89	19,801	
1088	5	y Pegasi	23 25	6	68	25	54,83	2,73		25	47,89		+ 6,94	19,809	
1089	6,7	ω [*] Piscium	23 25	3	92	10	48,07	0,65		10	42,52		+ 5,55	19,815	
1090	5	i Phœnicis	23 26	5	133	32	50,41	4,22		32	47,82		+ 2,59	19,821	
1091	4,5	λ Andromedæ . . .	23 29	6	44	27	26,79	4,08	27	23,66	27	20,00	+ 3,13	+ 6,79	19,863
1092	5	θ Phœnicis	23 30	5	137	34	27,06	2,66		34	27,91		- 0,85	19,875	
1093	5	ω [*] Aquarii	23 31	5	105	9	24,96	6,21		9	14,32		+ 10,64	19,882	
1094	4,5	ι Piscium	23 31	5	85	17	23,36	2,88	17	20,38	17	19,03	+ 2,98	+ 4,33	19,335
1095	5	κ Andromedæ . . .	23 32	5	46	36	2,17	3,84		36	4,62		- 2,45	19,894	
1096	3	γ Cephæ	23 32	4	13	18	35,74	3,07	18	37,25	18	40,10	- 1,51	- 4,36	19,898
1097	5	A [*] Aquarii	23 33	4	108	57	36,40	4,96		57	30,99		+ 5,41	19,901	
1098	5	λ Piscium	23 33	3	89	8	56,64	1,11	8	58,44	8	57,82	- 1,80	- 1,18	19,907
1099	5	A [*] Aquarii	23 35	5	109	12	51,54	2,10		12	46,77		+ 4,77	19,927	
1100	5	i Pegasi	23 35	5	61	34	23,04	2,76		34	25,95		- 2,91	19,928	
1101	7	* Piscium	23 36	5	83	44	39,38	2,62		44	41,92		- 2,54	19,934	

No.	Mag.	NAMES.	A.R.	No. of Observations.	Observed N.P.D. reduced to Jan. 1, 1831.			Difference of Extremes.	Greenwich Catalogue	A.S. Catalogue.	Difference from		Annual Variation.	
					°	'	"				s.	m.		s.
1102	5	ψ Andromedæ..	23 38	6	44	31	5,76	2,51		31	2,96	+ 2,80	- 19,947	
1103	5	γ Cassiopeæ....	23 39	5	32	17	20,48	3,08		17	21,58	- 1,10	19,957	
1104	5	* Draconis.....	23 40	5	23	7	54,03	2,43		7	57,95	- 3,92	19,965	
1105	5	δ App. Sculp...	23 40	6	119	3	49,37	5,90		3	48,17	+ 1,20	19,967	
1106	6	21 Piscium.....	23 41	5	89	51	46,31	4,82		51	41,82	+ 4,49	19,972	
1107	6	22 Piscium.....	23 43	3	88	0	28,97	1,76		0	30,54	- 1,57	19,989	
1108	5,6	ψ Pegasi.....	23 49	3	65	47	50,78	1,48		47	47,34	+ 3,44	20,020	
1109	5	p Piscium.....	23 50	5	94	29	36,81	2,81	29	36,55	29 32,33	+ 0,26	+ 4,48	20,023
1110	4,5	ω Piscium.....	23 51	5	84	4	17,57	3,84	4	19,82	4 18,79	- 2,25	- 1,22	20,026
1111	5	* Cassiopeæ....	23 53	5	29	43	7,32	2,16		43	5,52	+ 1,80	20,033	
1112	5	q Piscium.....	23 53	5	93	58	3,43	5,80	58	5,89	58 0,73	- 2,45	+ 2,70	20,034
1113	4,5	r Piscium.....	23 53	5	96	57	11,40	4,35	57	12,03	57 8,85	- 0,63	+ 2,55	20,034
1114	6	85 Pegasi.....	23 53	4	63	48	41,13	2,31		48	53,56	- 12,43	18,884	
1115	4	g Ceti.....	23 53	5	108	16	35,60	2,17	16	35,52	16 30,01	+ 0,08	+ 5,59	20,038
1116	5	s Piscium.....	23 57	5	96	39	8,38	2,13	39	11,70	39 10,09	- 3,32	- 1,71	20,040
1117	7	5 Ceti.....	23 59	5	93	23	15,20	4,83		23	12,19	+ 3,01	20,043	
1118	1	α Andromedæ..	23 59	50	61	50	33,16	6,41	50	34,37	50 33,31	- 1,21	- 0,15	20,043

In constructing the Madras Catalogue (as I have before stated) the tables of refraction employed are those by Atkinson, published in the 2d Volume of the Astronomical Society's Memoirs; whereas the Greenwich Catalogue with which I have compared the Madras results, was computed from Bradley's table of refractions. Now to render the two Catalogues strictly comparable, it is necessary that each should be reduced by the same table; I have consequently applied to the Greenwich Catalogue the corrections which have already been employed at Pages 61 and 62.

Continuation of table at Page 62.

N.P.D.


93	Add to Greenwich Catalogue.....	+ 1,57
98	do. do. do.	+ 1,76
103	do. do. do.	+ 1,96
108	do. do. do.	+ 2,30
113	do. do. do.	+ 2,48
116	do. do. do.	+ 2,33
118	do. do. do.	+ 2,14
119	do. do. do.	+ 1,78
120	do. do. do.	+ 1,19
121	do. do. do.	+ 0,13
122	do. do. do.	- 1,45
123	do. do. do.	- 4,06

On examining the column "difference from Greenwich" out of 489 comparisons there are

197	in which the difference is less than...	1",5		
122	do.	greater than.1",5 and less than 2",5		
115	do.	do.	.2",5	do. 4",0

and 55 cases in which the difference exceeds four seconds I forbear at present entering into a detailed investigation of these discordances and will only remark, that in about 40 cases out of the 55 the Madras results are confirmed by further observation in the present year (1832.)

The column "difference from A. S. C." furnishes 1114 comparisons between the Madras and Astronomical Society's Catalogue, of these there are 693 which are less than 4"; 315 between 4" and 8", and 105 which exceed 8"; among this last class 53 of the larger numbers are marked in the Astronomical Society's Catalogue with an Asterisk, signifying that the place of the Star has been brought forward from the observations of La Caille, and is less to be depended upon than the other portions of the Catalogue; by way of concluding this work I may remark that two of the larger errors, viz. ζ Toucanæ and λ^1 Phœnicis are confirmed by further observation in 1832, and with the observations of this year when completed, the remaining discordancies will I imagine all be confirmed.



Errata in Result of Observations for 1831.

Page 32	Correction for Azimuth	line 3	for 13,310	read	13,267
—	— for Collimation	— 3	— 0,411	—	4,10
—	Transit over Meridian	— 3	— 11,41	—	7,68
—	Correction for Level	— 37	— ,175	—	1,75
—	— for Azimuth	— 37	— 2,831	—	28,31
—	— for Collimation	— 37	— ,881	—	8,81
—	Transit over Meridian	— 37	— 59,75	—	50,25
—	Name	— 52	— χ	—	κ
43	Correction for Level	— 21	— ,326	—	,295
—	— for Azimuth	— 21	— ,585	—	+ ,194
—	— for Collimation	— 21	— ,664	—	,627
—	Transit over Meridian	— 21	— 2,84	—	3,62
—	Name	— 21	— ν Aurigæ	—	A Orionis.
45	Name	— 29	— χ Aurigæ	—	A Orionis.
—	App. A. R.	— 29	— 2,84	—	3,62
—	Aberration, &c.	— 29	— 4,48	—	3,70
—	Mean A. R.	— 29	— 58,36	—	59,92
—	Mean Place	— 29	— 43,08	—	45,54
52	Clock Rate by Stars	— 17	—	—	+
72	Observed N. P. D.	— 28	— 19'	—	18'
81	Mean Time October 16	—	— 2'',8	—	0'',3
—	A. R.	—	— 24,52	—	21,94
84	Mean Time	—	— 40,9	—	38,3
—	A. R.	—	— 53' 1'',00	—	52' 58'',43
102	Line 36	—	— ν	—	ν
105	— 17 Difference from Greenwich	—	+ 0,03	—	— 0,07
128	— 17 Observed N. P. D.	—	— 47,01	—	50,17
—	— 17 Difference	—	+ 1,28	—	+ 4,44
132	— 11 Observed N. P. D.	—	— 0,47	—	0,87
133	— 7 — —	—	— 58,81	—	58,31

PLATE I.

