

RESULT
OF
ASTRONOMICAL OBSERVATIONS

MADE AT
THE HONORABLE
THE EAST INDIA COMPANY'S OBSERVATORY
AT MADRAS

BY
THOMAS GLANVILLE TAYLOR, Esq.
ASTRONOMER TO THE HONORABLE COMPANY.

VOL. II.
FOR THE YEARS 1832 AND 1833.

PRINTED
BY ORDER OF THE
MADRAS GOVERNMENT.

MADRAS:
PRINTED AT THE MALE ASYLUM, AND FORT SAINT GEORGE GAZETTE PRESS.

M.DCCC.XXXV.

P R E F A C E.

MORE convinced than ever of the propriety of giving the *result* of Astronomical Observations, in preference to publishing the observations in an unreduced state, I have (with the permission of the Right Honorable the Governor in Council of this Presidency) given in the following pages the result of all the Astronomical Observations which have been made in this Observatory in the years 1832 and 1833, and have forwarded to England manuscript copies of the rough observations to be deposited in the Honorable Company's Library. The plan adopted upon the present occasion is as nearly as possible the same as that followed in Volume I. for 1831; since the publication of the latter work I have been favored with part 5 of the Greenwich Observations for 1831, containing the result of the observations given in parts 1—4 of that year; in one respect (that of giving the result of *each* observation instead of the *mean* result) I should much prefer to have followed the Greenwich plan, but the inconvenient delay I have experienced in printing (arising partly from want of type) would have rendered it impossible to print the additional number of pages which would be necessary without getting very considerably in arrears; to accomplish even the present work in ten months, I have been compelled to employ two separate printing establishments, and from this cause have been obliged to introduce a different size type and a second system of pageing, but the ill appearance thus introduced, will I hope be considered sufficiently atoned for, by my being enabled to publish the work 4 or 5 months earlier than I otherwise should have been able to do. The observations with the Transit Instrument have been somewhat interfered with by the unequal wear of the pivots (which has rendered it necessary to have them returned and Steel Collars applied over the present ones of Bell Metal), and by the unstability of the Meridian Marks: the result of these casualties is, that the accuracy of half a tenth of a second of time, which at commencing the Superintendence of this Observatory, I had vainly promised myself to attain, is forfeited in *some cases* to double and treble this amount; notwithstanding this, the observations will I imagine be found to possess a very useful, if not a valuable degree of accuracy:

A tolerable opinion of their relative accuracy with regard to that attained at other Observatories can be formed by the following table; exhibiting a *Comparison of the Right Ascension of several Stars which have been frequently observed at Madras with their places observed at the Observatories of Greenwich, Cambridge, and Königsberg.*

NAMES.	Right Ascension, January 1, 1832. Madras.			Difference from Madras.			REMARKS.			
				Greenwich.	Cambridge.	Königsberg.				
16 Ceti	a	0	35	9,29	—	0,02	—	0,02	
13 Arietis	a	1	57	43,22	—	0,05	—	0,02	—	0,03
92 Ceti	a	2	53	30,36	—	0,04	+	0,03	—	0,01
33 Persei	a	3	12	22,05	+	0,09	+	0,09	+	0,25
87 Tauri	a	4	26	17,31	+	0,05	+	0,04	+	0,04
13 Au. i. æ	a	5	4	17,36	+	0,07	+	0,09	+	0,14
58 Orionis	a	5	46	4,74	+	0,05	+	0,07	+	0,01
9 Can. Maj.	a	6	37	44,63	—	0,08	+	0,03		0,00
66 Geminorum	a	7	23	52,10	+	0,08	+	0,08	
10 Can. Min.	a	7	30	30,29	+	0,06	—	0,05	—	0,02
78 Geminorum	β	7	35	1,56	+	0,04	—	0,07	—	0,01
30 Hydræ	a	9	19	20,01	—	0,04	—	0,02	—	0,15
32 Leonis	a	9	59	25,11	+	0,03	—	0,05	—	0,06
50 Urs. Maj.	α	10	53	17,37	+	0,09	+	0,06	+	0,06
94 Leonis	β	11	40	29,21	—	0,03	—	0,08	—	0,10
67 Virginis	a	13	16	21,25	—	0,11	+	0,01	—	0,04
16 Bootis	a	14	8	0,04	+	0,06	+	0,08	+	0,07
36 Bootis	ε	14	37	39,00	—	0,03	+	0,05	
9 Libræ	α ²	14	41	35,97	—	0,05	—	0,01	—	0,04
5 Cor. Bor.	a	15	27	04,65		0,00	+	0,01		0,00
24 Serpentis	a	15	35	59,95	—	0,02	+	0,02	+	0,01
1 Ophiuchi	δ	16	4	33,03	—	0,08	—	0,15	
21 Scorpii	a	16	19	7,26	—	0,17	—	0,07	—	0,04
55 Ophiuchi	a	17	27	8,42	—	0,03	—	0,02	—	0,09
3 Lyræ	a	18	37	15,13	+	0,01	—	0,08	—	0,02
10 Lyræ	β	18	43	52,78	+	0,01	—	0,10	
53 Aquilæ	a	19	42	35,23	—	0,01	—	0,03	—	0,02
60 Aquilæ	β	19	47	3,72	+	0,02	—	0,07	+	0,03
6 Capricorni	α ²	20	8	43,71	—	0,03	—	0,05	+	0,04
50 Cygni	a	20	35	42,48	—	0,02	+	0,06	—	0,03
22 Aquarii	β	21	22	42,68	—	0,14	—	0,01	
34 Aquarii	a	21	56	9,24	+	0,03	—	0,01		0,00
54 Pegasi	a	22	56	23,93	+	0,01	—	0,08	—	0,02
21 Andromedæ	a	23	59	43,19	+	0,01	—	0,05	—	0,06

The observations with the Mural Circle have proceeded without any interruption, and the results *inter se*, are as accordant as any observations

N. B.—The Catalogues of Cambridge and Königsberg are derived from Vol. V. Page 17 of the Cambridge Observations, save that the former has been *diminished* 0,10s. and the latter *increased* 0,07s. to reduce them to the Equinoctial Point assumed in the Madras and Greenwich Catalogues.

of this nature will permit; in computing the Parallax of the Planet Mars where a comparison between the Madras, Cape of Good Hope, and St. Helena observations has been instituted; two singular cases of discordance had led me to suspect an error of large amount in one set of divisions of the Madras Mural Circle; but on examining the divisions by means of two Collimators, I am enabled to state, *that the error of division is confined to very allowable limits, and may possibly extend to a very great degree of accuracy.* The observations made out of the meridian are not so numerous as I could wish, this has mostly arisen from the want of a building to shelter the observing Telescope from the wind; on this account the transit of Mercury over the Sun's disc in May 1832, was but imperfectly observed, and several occultations have been lost; in short I cannot but consider the Observatory incomplete from the want of a fixed Instrument for observing objects out of the meridian; hitherto for this purpose I have employed the 5 feet Achromatic by Dollond, mounted upon Sineaton's plan; but the utmost accuracy attainable with this sort of Instrument falls far short of that accuracy which the present state of practical Astronomy demands. The observations on the meridian have with but few exceptions been made as heretofore by the Assistants, who are natives of high *caste*; and those out of the meridian for the most part by myself: of the abilities of the Native Assistants as observers I entertain the highest possible opinion, and as computers, they possess a very serviceable degree of accuracy and despatch, notwithstanding which the reduction of the observations has for the most part been performed by myself, having trusted nothing of importance to the native computers without a strict examination or a recomputation.

With regard to the methods of reduction and finding the Index Error, &c. the plans I have adopted differ in no respect from those employed at the Greenwich Royal Observatory, which were I believe devised by Mr. Pond the present Astronomer Royal, to whom at least I am indebted for them: In employing the Greenwich Catalogue as my Zero point, it may not be amiss for me here to record my opinion, that the Greenwich Transit Observations are at present excelled by none, and the observations derived from the two Mural Circles (due as much to the judicious way in which their results are combined, as to the superior excellence of the Instruments) are very superior to any yet published observations.

It now remains for me to say a word or two with regard to the arrangement of the work. Having experienced considerable delay on the part of the printer in the execution of Volume I. I commenced printing the present work on the 20th December 1833, ten days before the observations constituting the results were completed ; and about as many months before the completion of the computation ; from this cause, circumstances (which have arisen in the course of computation and printing) have compelled me to deviate from the straight forward course of arrangement I otherwise could wish to have followed ; my object however has been to render the work complete in itself, and forward in a degree however small the cause of Astronomy.

T. G. TAYLOR,
H. C. ASTRONOMER.

OBSERVATIONS MADE WITH THE TRANSIT INSTRUMENT.

THE Intervals between the wires at the beginning of the year 1832, is assumed to be the same as that determined at Page 6, Vol. I for 1831 ; there hold good up to 18th January when the whole set were broken, and a new set put in by my Assistant Anuntacharyer ; (being absent myself at the time on other duty in Calcutta) ; from 50 Observations between the 18th and the 23d the Equatoreal Intervals were found to be :

from 1st wire to centre.....	54',462
2d	27,294
4th.....	27,438
5th.....	54,985

these wires were I imagine put in with bad varnish, for on the 23d January they were again found broken; on this occasion Mr. Law (of whose skill and abilities to perform this or any other job which he may undertake I have the highest possible opinion) applied a new set ; from the mean of 70 Observations the Equatoreal Intervals were found to be :

from 1st wire to centre.....	54',400
2d	27,280
4th.....	27,302
5th.....	54,750

On the 8th August, I found the 1st and second wires bent in consequence of which I put in a new set of silk lines; when from 70 Observations of Stars situated near the Pole, the Equatoreal Intervals were found to be :

from 1st wire to centre.....	54',988
2d	27,566
4th.....	27,352
5th.....	55,021

On the 9th November the following note appears in the Transit book.
 " Found the moveable wire had been caught by the varnish which secured the fixed wires, to remedy this I filed a groove to contain the varnish and put

MADRAS OBSERVATIONS.

in a new set of wires" from 70 Observations the Equatoreal Intervals were now found to be :

from 1st wire to centre.....	54,643
2d	27,323
4th.....	28,128
5th.....	55,281

In the month of May 1833, being desirous of ascertaining if the wires remained stable; from 72 Observations of Stars situated near to the Pole I found the Equatoreal Intervals to be :

from 1st wire to centre.....	54,619
2d	27,357
4th.....	28,071
5th.....	55,121

On the 12th May A. M. the following note appears in the Transit book. " The centre wire appears to have shifted its position since the observations of last night" and on the 13th " fearing that by reason of the hot land wind the centre wire might have become loosed, which however was not apparent I applied fresh varnish (tincture of opium) to the ends of all the five wires," from the observations of several Stars situated near to the Pole the Equatoreal Intervals were found to be :

from 1st wire to centre.....	54,961
2d	27,618
4th.....	27,878
5th.....	54,924

On the 23d of August by reason of very heavy rain a few drops of water had leaked through the roof and falling upon the eye end of the Telescope, had bent two of the wires, in consequence of which I put in a new set; from 36 Observations of Stars situated near to the Pole the Equatoreal Intervals were found to be :

from 1st wire to centre.....	55,420
2d	27,896
4th.....	27,374
5th.....	54,594

Hence to reduce observations made at the five wires to the centre wire, it becomes necessary to apply the following corrections :

				s.
1832—From	January 1st to	January 28th....	+	$\frac{,061}{\sin. N.P.D.}$
	January 18th —	January 23d	—	$\frac{,133}{\sin. N.P.D.}$
	January 23d —	August 8th....	—	$\frac{,074}{\sin. N.P.D.}$
	August 8th —	November 9th....	+	$\frac{,036}{\sin. N.P.D.}$
	November 9th —	End of the year....	—	$\frac{,288}{\sin. N.P.D.}$
1832— „	November 9th —	1833 May 12th....	—	$\frac{,243}{\sin. N.P.D.}$
1833— „	May 12th —	August 23d	—	$\frac{,045}{\sin. N.P.D.}$
	August 23d —	End of the year... +		$\frac{,270}{\sin. N.P.D.}$

In the absence of any cause which can explain why the Equatoreal Intervals in November 1832 differ from those in May 1833, I have employed between the

		s.
9th of November and 31st December 1832..	—	$\frac{,288}{\sin. N.P.D.}$
1st of January — 1st March 1833..	—	$\frac{,266}{\sin. N.P.D.}$
1st of March — 12th May 1833..	—	$\frac{,243}{\sin. N.P.D.}$

We will now examine the observations for the determination of the value of the Micrometer screw which it will be recollected was found but approximately in the year 1831; for this purpose the following are the *Intervals of time which the Pole Star took to pass over 2 R. 0 D. 1 R. 50 D. &c. to the East and West of the centre wire.*

East of the Centre Wire.					West of the Centre Wire.			
	R. D. 2 0	R. D. 1 50	R. D. 1 0	R. D. 0 50	R. D. 0 50	R. D. 1 0	R. D. 1 50	R. D. 2 0
	m. s.	m. s.	m. s.	m. s.	m. s.	m. s.	m. s.	m. s.
1832								
May	16	2 5,5	1 23,5	0 44,5	0 39,5
	17	2 45,0	2 4,0	1 22,5	0 42,0	0 41,5	1 23,0	2 5,0
	26	0 41,5	1 22,5	2 5,0
	29	2 47,0	2 4,0	1 22,0	0 41,5	0 40,0	1 21,0	2 1,0
June	9	2 46,0	2 4,5	1 23,0	0 41,5	0 41,5	1 23,5	2 4,5
November	16	2 46,0	2 4,5	1 22,0	0 40,0	0 41,5	1 23,0	2 5,5
	17	2 45,5	2 4,5	1 22,0	0 42,0	0 40,5	1 23,5	2 6,0
	22	2 49,5	2 5,5	1 23,5	0 42,5	0 42,5	1 24,5	2 5,0

MADRAS OBSERVATIONS.

1832		m. s.	m. s.	m. s.	m. s.	m. s.	m. s.	m. s.
November	23	2 46,5	2 4,5	1 23,5	0 41,5	0 42,0	1 23,5	2 5,5
	25	2 46,0	2 4,5	1 23,0	0 42,0	0 41,5	1 23,0	2 6,0
	29	2 44,5	2 3,5	1 22,0	0 41,0	0 42,0	1 25,0	2 6,5
December	4	2 47,0	2 5,0	1 23,5	0 42,0	0 41,0	1 23,0	2 4,5
	5	1 21,5	1 23,5
	7	1 25,5	1 23,0
	9	1 24,5	1 20,5
	12	2 45,0	1 22,5	1 24,5	2 47,0
	13	2 44,0	1 22,0	1 22,5	2 46,5
	14	2 49,5	1 24,5	1 23,0	2 47,5
	15	2 45,5	1 25,5	1 23,5	2 45,5
	16	2 46,0	1 23,0	1 24,5	2 46,5
	16	2 45,0	1 23,0	1 23,5	2 46,0
	17	2 45,0	1 21,5	1 23,0	2 45,5
	18	2 45,0	1 21,0	1 22,0	2 46,0
	19	2 48,0	1 25,5	1 22,0	2 44,0
	20	2 46,0	1 22,0	1 25,0	2 48,0
	21	2 46,5	1 23,0	1 22,5	2 46,5
	22	2 49,5	1 25,0	1 24,0	2 46,5
	23	2 47,0	1 23,0	1 23,5	2 47,0
	24	2 45,5	1 22,5	1 24,0	2 44,5
	24	2 48,0	1 23,5	1 24,0	2 48,0
	25	2 46,5	1 22,5	1 21,5	2 43,5
	26	2 48,0	1 25,0	1 22,0	2 46,0
	26	2 48,0	1 24,0	1 23,0	2 45,5
Mean..		2 46,46	2 4,55	1 23,25	0 41,87	0 41,25	1 23,14	2 4,95

Taking the differences we find that the Pole Star passed from

R. D.	R. D.	s.
2 0 East to 1 50 East	41,89 of time.
1 50 — 1 00 —	41,30 —
1 00 — 0 50 —	41,38 —
0 50 — 0 00 —	41,87 —
0 00 — 0 50 West	41,25 —
0 50 — 1 00 —	41,89 —
1 00 — 1 50 —	41,81 —
1 50 — 2 00 —	41,34 —

Hence we may safely assume the threads of the screw to be equidistant, and assuming the N.P.D. of the Pole Star to be $1^{\circ} 34' 46''$, the value of each revolution from the Mean of 32 Observations..... = $34', 394$
 In 1831.....10 — = $34', 277$
 hence we may safely assume the value of each revolution of the screw at $34', 366$.

ERROR OF LEVEL OF THE TRANSIT AXIS

IN the Results of Observations Vol. I for 1831, it is stated that from 80 times inverting the Transit Axis, the diameter of the illuminating pivot apparently exceeded that of the other pivot $0^{\prime\prime},58$: producing an error of level to the amount $0^{\prime\prime},29$. In the reduction of the Observations in 1831, the correction due to this was included with the correction due to the error registered by the spirit level in the column for that purpose; following this plan the observations of 1832, were reduced, and those for 1833 far advanced, when, with a view to determine if this result remained constant (of which I had some doubt by reason of disagreement in the places of Polaris and δ Ursæ Minoris,) I made the following Observations.

Cross Level East and Illuminating Pivot.

1833		Inversions.		East.		West.		Illuminating Pivot.
				"		"		
November	12	Mean of 10	Obj. Glass	N. 6,51	W. Obj. Glass	S. 26,37	W. +	9,93
	13	—	6	—	S. 2,30	"	—	N. 20,55 — + 9,13
	14	—	4	—	S. 2,64	"	—	N. 24,21 — + 10,78
December	20	—	10	—	N. 11,07	E.	—	S. 13,31 — + 12,19
	20	—	10	—	N. 11,97	"	—	S. 13,94 — + 12,25
	25	—	8	—	S. 10,07	"	—	N. 14,60 — + 12,33
	25	—	4	—	N. 5,46	"	—	S. 16,53 — + 11,00

Taking the Mean, it appears that in the year 1833 or at 1833, 93 the illuminating pivot exceeded the other pivot to the amount $11^{\prime\prime},37$, whereas at 1831, 27 it exceeded it only $0^{\prime\prime},58$,

This enormous and extraordinary wear of the one pivot above the other, is, as far as concerns the construction of the pivots and Y's, altogether unaccountable; on inspecting the former which are of Bell Metal, it is quite evident that the unilluminating pivot has worn more than the other pivot, the appearance of either being such as would result from their having been turned in a lathe; the latter which are of brass, are not more worn than might be reasonably expected; in lieu of a line of contact on each face as exhibited on the erection of this Instrument in 1831; a groove of about, 14 Inches broad has been worn by the action of the pivots; to account for this change, no circumstances offer beyond the Instrument having been kept hard at work, and that too during three years of unusual heat and dust, in which, notwith-

standing the pivots were constantly kept covered by the slips of Brass* for the purpose; it was found necessary to wipe them and apply fresh oil on every second day at farthest but more frequently every day.

With a view to discover if the pivots continued of a circular figure: I attached to the Stone Pier a microscope (into the focus of which I had fitted a pair of lines at right angles to each other) and watched the motion of an exceedingly small point which I had made in a slip of ivory and cemented to the end of the pivot; when placed on the eastern or illuminating pivot and adjusted to its axis, the centre of the dot in every position of the Instrument remained perfectly bisected by the cross wires; when placed upon the Western pivot however the bisection was not so satisfactory; having failed after much loss of time to attach the dot opposite to the center of the axis of this pivot I allowed it to remain at a distance of about ,001 + from the centre and estimated the value of the rectangular co-ordinates of the centre of the dot from the intersection of the cross lines in tenths of the diameter of the dot as follows:

No.	N.P.D.	The dot was situated.			Direction of the Telescope.
1	at 347	to the North	,00	and 0,05 too high	North Horizon.
2	— 0	—	,10	— ,10	Pole.
3	— 13	—	,15	— ,25	"
4	— 30	—	,25	— ,30	"
5	— 50	—	,30	— ,35	"
6	— 76	—	,30	— ,45	Zenith.
7	— 95	—	,28	— ,52	"
8	— 120	—	,20	— ,55	"
9	— 140	—	,09	— ,58	"
10	— 160	—	,01	— ,65	South Horizon.
11	— 180	South	,03	— ,66	South Pole.
12	— 200	—	,15	— ,65	"
13	— 220	—	,20	— ,55	"
14	— 240	—	,25	— ,50	"
15	— 257	—	,30	— ,50	Nadir.
16	— 280	—	,33	— ,45	"
17	— 300	—	,30	— ,35	"
18	— 320	—	,16	— ,23	"
19	— 347	—	,00	— ,05	North Horizon.

* It is much to be regretted that no better means has yet offered for protecting this part of the Instrument from dust; in this climate, where for several days together occasionally, the air is saturated with sand, the want of a better cover is much felt.

the above which are the mean of several readings or rather estimations, can safely be depended upon to ,03 or ,04; the diameter of the dot was determined from a very neatly engraved scale of converging lines to be ,0025 Inches; assuming the above numbers and a large scale, we may now trace the curve described by any point on the axis of the pivot see fig. 1; and comparing the above numbers with the natural sines of the angles, we can determine the circle A. B. C. which agrees best with all the measures, from whence it appears that *the deviation of the pivot from a circular figure does not entail an error in any direction to the amount of one second of space beyond which limits the means at my command do not enable me to offer an opinion or proof.*

We will now enquire to what amount the Right Ascensions of the Planets and fixed Stars for the years 1831, 1832 and 1833, are effected by this unforeseen change in the pivots; In the first place we must recollect that the above excess of the illuminating pivot over the other is only an *apparent* one, for we have $2,82 (r - r') = 11'',37$ or the true difference of the radii of the pivots $r - r' = 4',03 = ,00058$ Inches; and the error of level thus produced $= (r - r') \operatorname{cosec} \frac{20^\circ}{2} = 5'',68$. Now if we diminish this amount by $0'',29$ (the error already allowed for) we obtain the error which remains to be allowed $= 5'',39$ which produces corrections to be applied to the reduced Right Ascensions as follows :

For	1 35	N.P.D.	above the Pole	+	3,30	in Time.	
	1 35	—	below the Pole	—	2,60		—
	3 25	—	above the Pole	+	1,71		—
	3 25	—	below the Pole	—	1,05		—
	10 0	—	above the Pole	+	0,81		—
	20 0	—	—	+	0,57		—
	30 0	—	—	+	0,49		—
	40 0	—	—	+	0,45		—
	50 0	—	—	+	0,42		—
	60 0	—	—	+	0,39		—
	70 0	—	—	+	0,37		—
	80 0	—	—	+	0,36		—
	90 0	—	—	+	0,34	} + 0,34	—
	100 0	—	—	+	0,33		—
	110 0	—	—	+	0,31		—
	120 0	—	—	+	0,29		—
	130 0	—	—	+	0,27		—
	140 0	—	—	+	0,25		—
	150 0	—	—	+	0,21		—
	160 0	—	—	+	0,12		—
	165 0	—	—				—

On consulting the method employed in reducing the observations of transits at Pages 31 et seq. of Vol. I, it will readily appear, that for Stars situated above 30° or 40° from the Pole, the greater part of the above corrections will be lost sight of in the determination of the Error of the Clock; thus, suppose (as actually has been the case in the reduction of the Observations for 1832 and 1833) that the Stars selected for the determination of the clock's error be situated between the limits of 65° and 115° of North Polar Distance; in this case the error of the Clock will be, instead of e , some number between $e + ,30$ and $e + ,38$; by assuming $e + ,34$ we are liable to an error $0',04$, i. e. this is the largest error we can possibly commit by such an assumption; but in 9 cases out of 10, I find the error does not exceed the half of this amount. Subtracting then $0',34$ from the above numbers we obtain the corrections proper to be applied to the reduced Right Ascensions made towards the end of the year 1833; thus in the case of α Lyræ; N.P.D. $51^\circ 22'$; whose A.R. we will suppose to have been determined towards the end of the year 1833; (the Clock error having been determined from a comparison of the observed places of Equatoreal Stars with the Greenwich Catalogue) we have to apply the correction $,42 - ,34 = ,08$; and for the Star α Cephei N.P.D. $28^\circ 7'$ we have to apply $,50 - ,34 = ,16$ &c. These corrections it must be recollected apply only to observations made towards the end of 1833, for dates antecedent to this (on the supposition that the wear of the one pivot above the other has been uniformly accelerated with the time) it will be proper to apply corresponding smaller corrections as follows:

Corrections to be applied to the Reduced A.R,							
	1831,5	1832,0	1832,5	1833,0	1833,5	1833,93	
"	"	"	"	"	"	"	"
1 35 A. P.	+ 0,15	+ 0,73	+ 1,30	+ 1,87	+ 2,45	+ 2,96	
1 35 B. P.	- 0,12	- 0,56	- 1,00	- 1,44	- 1,87	- 2,26	
3 25 A. P.	+ 0,07	+ 0,34	+ 0,60	+ 0,87	+ 1,13	+ 1,37	
3 25 B. P.	- 0,04	- 0,18	- 0,32	- 0,46	- 0,59	- 0,71	
10 0 A. P.	+ 0,03	+ 0,12	+ 0,21	+ 0,30	+ 0,39	+ 0,47	
20 0 —	+ 0,01	+ 0,06	+ 0,10	+ 0,14	+ 0,19	+ 0,23	
30 0 —	+ 0,01	+ 0,04	+ 0,07	+ 0,10	+ 0,13	+ 0,15	
40 0 —	+ 0,01	+ 0,03	+ 0,05	+ 0,07	+ 0,09	+ 0,11	
50 0 —	0,00	+ 0,02	+ 0,03	+ 0,05	+ 0,07	+ 0,08	
60 0 —	0,00	+ 0,01	+ 0,02	+ 0,03	+ 0,04	+ 0,05	
70 0 —	0,00	+ 0,01	+ 0,01	+ 0,02	+ 0,03	+ 0,03	
80 0 —	0,00	0,00	+ 0,01	+ 0,01	+ 0,02	+ 0,02	
90 0 —	0,00	0,00	0,00	0,00	0,00	0,00	
100 0 —	0,00	0,00	0,00	0,00	0,00	0,00	
110 0 —	0,00	0,00	- 0,01	- 0,02	- 0,02	- 0,03	
120 0 —	0,00	- 0,01	- 0,02	- 0,03	- 0,04	- 0,05	
130 0 —	0,00	- 0,01	- 0,03	- 0,04	- 0,06	- 0,07	
140 0 —	0,00	- 0,02	- 0,04	- 0,06	- 0,08	- 0,09	
150 0 —	- 0,01	- 0,04	- 0,06	- 0,09	- 0,11	- 0,13	
160 0 —	- 0,01	- 0,05	- 0,10	- 0,14	- 0,18	- 0,22	

If the computation of the observations for 1832 and 1833 had not been so nearly completed, the above corrections would as heretofore have been included with the ordinary correction for Level, but the case being otherwise, it will be found as I have already shewn, that an error of comparatively little importance is committed in employing the above table, where we find, that for observations in 1831 for Stars situated between 40° and 140° of N.P.D. no correction need be applied, and that for observations in 1832 and 1833 situated between these limits by employing the columns 1832,5 and 1833,5 respectively we are liable to errors which in no case exceed $.02''$. And further that up to the limits of 10° of N.P.D. it will be found sufficient to suppose the observations to have been made at that time of the year when the Star whose place we desire to correct passes the meridian at 9 o'clock at night; for Stars situated within this limit, (of which there are very few cases) the month and day must be taken into the account. In the reductions of the observations of the Sun, Moon and Planets for 1832 and 1833, I had employed the Errors of Level which now follow together with $0'',29$ for inequality of the pivots, these have consequently been since corrected by the above table, whereas the places of the fixed Stars are set down under the respective years in which they were observed *uncorrected*; the correction being taken account of in obtaining the column *mean* as will be further explained in the proper place.

ERROR OF COLLIMATION.

FROM inverting the Transit Instrument 23 times in the month of April 1831, it appeared that the South meridian mark was distant from the meridian mark to the North, reckoning towards the West; $180^\circ + 26'',97$; this number was accordingly employed in computing the error of Collimation for this year: towards the latter end of 1832, being about to compute the Errors of Collimation for the observations of that year, I set to work to verify the angle above measured as follows:

		Illuminating Pivot.				Collimation of	
		East.		West.		North	South
		No. of	North	South	North	South	
		Invers.	mark.	mark.	mark.	mark.	mark.
		"	"	"	"	"	"
1832							
Oct.	18.. 10	..39,04 W.	.. 71,97 E.	.. 47,58 W.	.. 63,65 E.	.. 4,27 W.	.. 4,16

ERROR OF COLLIMATION.

1833	"	"	"	"	"	"
Jan. 5.. 10	..38,94 W.	.. 64,10 —	E. 39,39 W.	.. 63,50 E.	.. 0,22 W.	.. 0,30
16.. 10	..37,12 —	.. 65,75 —	.. 40,28 —	.. 63,87 —	.. 1,58 —	.. 1,44
18.. 10	..35,91 —	.. 67,11 —	.. 42,17 —	.. 62,07 —	.. 3,13 —	.. 2,52

And to see if this continued unaltered.

"	"	"	"	"	"	"
Dec. 25.. 10	..32,92 W.	.. 61,40 E.	.. 36,59 W.	.. 56,94 E.	.. 1,83 W.	.. 2,23
26.. 10	..33,83 —	.. 62,78 —	.. 36,22 —	.. 59,90 —	.. 1,20 —	.. 4,44
28.. 10	..33,71 —	.. 61,93 —	.. 37,26 —	.. 57,77 —	.. 1,78 —	.. 2,08

If the Instrument were free from Error of Collimation the readings of Illuminating end East would be identical with those of Illuminating end West and would be as follows:

		Illuminating Pivot E. or W.		
		North mark.	South mark.	Angular Distance. = 180° +
		"	"	"
1832				
October	18 43,31 W. 67,81 E. 24,50
1833				
January	5 39,16 — 63,80 — 24,64
	16 38,70 — 64,31 — 25,61
	18 39,04 — 64,59 — 25,55

And for the second series.

		"	"	"
December	25 34,75 W. 59,17 E. 24,42
	26 35,02 — 61,34 — 26,32
	28 35,48 — 59,85 — 24,37

Taking the mean it appears the South meridian mark was distant from the North meridian mark, reckoning to the *West about* ;

$$\text{at the End of the year 1832} = 180^\circ + 25',07$$

$$\text{————— 1833} = \quad \quad \quad 25',04$$

For the present I will postpone the discussion as to *when* the alteration of the angular distance between the two marks from 26',97 to 25',07 took place, and proceed to state, that the numbers which now follow in computing the corrections for Collimation for the years 1832 and 1833, have been employed, using 25',07 for the angular distance together with 0',29 for diurnal aberration.

ERROR OF COLLIMATION.

1832	Azimuth of		N. + S.	N + S + 25'' 07 2	Mean.	REMARKS.
	North mark.	South mark.				
	"	"	"	"		
January	1 + 19,46	- 28,68	- 9,22	+ 7,92		
	2 18,87	32,82	13,95	5,56		
	3 18,53	34,44	15,91	4,58		
	4	32,92				
	5 19,66	32,75	13,09	5,99		
	6 18,87	32,96	14,09	5,49		
	7 21,36	33,44	12,08	6,49		
	8 20,42	30,93	10,51	7,28		
	9 20,96	30,98	10,02	7,52		
	10 21,13	33,27	12,14	6,47	Mean of 10 + 6'',53	
	11 21,65	30,83	9,18	7,94		
	12 21,31	31,24	9,93	7,57		
	13 21,72	30,76	9,04	8,01		
	14 21,13	29,99	8,86	8,10		
	15 21,60	29,24	7,64	8,71		
	16 21,45	30,25	8,80	8,14		
	17 21,48	29,41	7,93	8,57		
	18 21,96	30,35	8,39	8,34		
	19 21,48	29,38	7,90	8,58		
	24 31,86	Mean of 8 + 8'',25	Found the fourth wire broken, a new set were put in by Mr. Law.
	25 31,41	17,18				
February	3 30,42	22,78	+ 7,64	16,35	One of the moveable wires found broken, which not being re- quired was consequently not replaced.
	4 29,58	19,24	10,34	17,70		
	5 29,34	21,52	7,82	16,44		
	6 29,75	20,72	9,03	17,05		
	7 30,18	18,39	11,79	18,43		
	8 31,52	18,43	13,09	19,08		
	9 32,21	17,70	14,51	19,79		
	10 31,30	17,56	13,74	19,40		
	11 32,65	16,63	16,02	20,54		
	12 29,58	21,72	7,86	16,46		
	13 29,92	20,96	8,96	17,01		
	14 29,72	21,10	8,62	16,84		
	15 29,89	21,48	8,41	16,74		
	16 29,61	21,65	7,96	16,51		
	17 28,44	22,89	5,55	15,31		
	18 29,71	22,58	7,13	16,10	I returned to Madras from Cal- cutta, where I had been as- sisting in the measurement of a base line.
	19 27,98	22,41	5,57	15,32		
	20 29,48	21,65	7,83	16,45		
	21 30,93	20,72	10,21	17,64		
	22 31,89	21,03	10,86	17,96		
	23 28,92	22,65	6,27	15,67		
	24 29,65	21,92	7,73	16,40		
	25 31,10	20,59	10,51	17,79		
	26 30,96	21,21	9,75	17,41	Mean of 25 + 17'',26	
	27 30,73	21,41	9,32	17,20		
	28 32,13	21,55	10,58	17,82		
	29 31,96	21,48	10,48	17,78		

ERROR OF COLLIMATION.

1832	Azimuth of		N. + S.	N + S + 25",07 2	Mean.	REMARKS.
	North mark.	South mark.				
March	"	"	"	"		
1	+ 32,20	- 21,03	+ 11,17	+ 18,12		
2	32,27	20 69	11,58	18,32		
3	32,17	21,03	11,14	18,11		
4	31,83	20,65	11,18	18,13		
5	31,10	21,65	9,45	17,26		
6	31,30	21,20	10,10	17,58		
7	31,44	20,90	10,54	17,81		
8	30,96	21,24	9,72	17,39		
9	32,03	20,83	11,20	18,13		
10	33,16	21,48	11,68	18,36		
11	32,38	21,93	10,45	17,77		
12	31,58	21,65	9,93	17,50		
13	31,51	21,65	9,86	17,47		
14	31,30	21,55	9,75	17,41		
15	31,03	21,45	9,58	17,33		
16	30,80	21,93	9,87	17,47		
17	31,44	20,83	10,61	17,84		
18	31,86	20,76	11,10	18,08		
19	31,76	21,03	10,73	17,90		
20	31,17	21,31	9,86	17,46	Mean of 23	
21	30,76	21,45	9,31	17,19	+ 17",75	
22	30,59	23,10	7,49	16,28		
23	30,08	23,69	6,39	15,73		
24	30,49	23,59	6,90	15,99		
25	30,59	23 38	7,21	16,14		
26	30,46	22,41	8,05	16,56		
27	30,62	23,72	6,90	15,99		
28	30,62	22,41	8,21	16,64		
29	30,25	22,55	7,70	16,33		
30	30,93	23,03	7,90	16,48		
31	29,99	22,20	7,79	16,43	Mean of 11	
April	30,42	22,34	8,08	16,58	+ 16",29	
2	28,43	24,43	4,00	14,53		
3	27,02	26,56	0,46	12,77		
4	26,77	27,05	0,28	12,39	Re-examined and found correct.
5	28,01	25,16	+ 2,85	13,96		
6	29,82	23,66	6,16	15,61		
7	29,82	23,93	5,89	15,48		
8	30,16	23,38	6,78	15,92		
9	30,49	23,03	7,46	16,26		
10	30,66	24,64	6,02	15,54		
11	30,28	24,13	6,15	15,61		
12	29,58	23,96	5,62	15,34		
13	29,20	24,71	4,49	14,78		
14	29,34	24,09	5,25	15,16		
15	29,68	23,69	5,99	15,53		
16	29,68	24,09	5,59	15,33		
18	30,49	24,85	5,64	15,35		
19	30,53	25,06	5,47	15,27		
20	30,07	25,12	4,95	15,01		
21	30,53	24,64	5,89	15,48		
22	30,16	24,92	5,24	15,15		
23	30,23	24,82	5,41	15,24		

OBSERVATIONS FOR LEVEL.

1832				1832					
	D. H.	Illuminating Pivot.	Error from Level.	REMARKS.		D. H.	Illuminating Pivot.	Error from Level.	REMARKS.
Jan.	2 1	East	5,24 E.		April	3 1	East	4,47 E.	
	4 1	5,26 ..			5 1	4,45 ..	
	6 1	5,50 ..			7 1	4,86 ..	
	8 1	5,18 ..			9 1	4,60 ..	
	10 1	4,86 ..			11 1	4,50 ..	
	12 1	5,05 ..			13 1	5,04 ..	
	14 1	5,14 ..			15 1	5,12 ..	
	16 1	5,43 ..			17 1	4,46 ..	
	18 1	5,43 ..			19 1	4,55 ..	
	20 1	5,50 ..			23 1	4,14 ..	
	24 1	5,71 ..			25 1	3,59 ..	
	26 1	5,97 ..			27 1	4,33 ..	
	28 1	6,17 ..			30 1	4,11 ..	
	30 1	6,31 ..		May	2 1	5,26 ..	
Feb.	1 1	5,52 ..			4 1	4,80 ..	
	3 1	6,06 ..			7 1	5,25 ..	
	5 1	5,58 ..			9 1	4,58 ..	
	7 1	6,20 ..			11 1	3,85 ..	
	9 1	5,61 ..			13 1	4,80 ..	
	11 1	5,63 ..			15 1	4,43 ..	
	13 1	5,67 ..			17 1	3,92 ..	
	16 1	4,89 ..			19 1	4,27 ..	
	18 1	4,47 ..			21 1	4,62 ..	
	20 1	4,81 ..			23 1	3,36 ..	
	22 1	4,78 ..			25 1	4,94 ..	
	24 1	4,48 ..			27 1	4,73 ..	
	27 1	5,10 ..			29 1	4,20 ..	
	29 1	5,06 ..			31 1	4,21 ..	
March	2 1	5,69 ..		June	2 1	3,42 ..	
	4 1	5,02 ..			4 1	2,92 ..	
	6 1	5,24 ..			6 1	2,28 ..	
	8 1	3,97 ..			8 1	2,24 ..	
	10 1	3,41 ..			10 1	2,40 ..	
	12 1	4,56 ..			12 1	1,92 ..	
	14 1	4,35 ..			14 1	2,11 ..	
	16 1	5,25 ..			16 1	3,35 ..	
	18 1	4,69 ..			18 1	2,58 ..	After this obser-
	20 1	4,88 ..			22 1	3,10 ..	vation I ad-
	22 1	4,92 ..			24 1	3,13 ..	justed the level.
	24 1	5,56 ..			28 1	2,57 ..	
	26 1	6,78 ..			30 1	2,75 ..	
	28 1	4,07 ..		July	2 1	1,82 ..	
	30 1	4,23 ..			5 1	2,36 ..	Continued cloudy
April	1 1	4,60 ..						weather.

N. B.—The Pages 9, 10, 11, and 12 should have followed after line 20 of Page 53.

OBSERVATIONS FOR LEVEL.

1832				1833				
	1832	Illuminating Pivot.	Error from Level.	REMARKS.	1832	Illuminating Pivot.	Error from Level.	REMARKS.
	D. H.		s.		D. H.		s.	
July	8 1	East	5,37 E.	I adjusted the level before the above reading.	Nov.	27 1	East	1,36 E.
	16 1	7,65 ..		Dec.	1 1	1,19 ..
	20 1	7,67 ..			3 1	1,05 ..
	24 1	6,51 ..			6 1	1,68 ..
	26 1	5,26 ..			8 1	0,50 ..
	28 1	4,30 ..			10 1	0,72 ..
	30 1	4,76 ..			13 1	0,34 ..
Aug.	1 1	5,49 ..			16 1	0,89 ..
	3 1	5,71 ..			18 1	0,71 ..
	5 1	5,77 ..			21 1	1,23 ..
	11 1	6,07 ..		24 1	0,97 ..	
	13 1	5,81 ..		27 1	1,31 ..	
	17 1	6,53 ..		30 1	0,72 ..	
	19 1	6,60 ..					
	21 1	6,21 ..	I re-examined this at 2 P. M. and found it correct.	1833			
	23 1	6,47 ..		Jan.	2 1	East	0,35 E.
	25 1	10,61 ..			4 1	0,30 ..
	27 1	11,75 ..			7 1	1,09 ..
	29 1	11,72 ..			9 1	1,06 ..
	31 1	10,93 ..			11 1	0,66 ..
Sept.	2 1	11,02 ..			14 1	1,45 ..
	4 1	10,43 ..			17 1	3,48 W.
	6 1	11,95 ..			19 1	3,38 ..
	8 1	11,08 ..			21 1	3,23 ..
	10 1	11,55 ..		23 1	3,87 ..	
	12 1	11,24 ..		25 1	4,12 ..	
	15 1	11,50 ..		27 1	4,14 ..	
	20 1	11,48 ..		29 1	4,61 ..	
	22 1	11,97 ..		31 1	5,27 ..	
	24 1	11,43 ..	Feb.	2 1	4,68 ..	
	26 1	11,55 ..		4 1	4,99 ..	
	28 1	11,85 ..		6 1	4,97 ..	
	30 1	12,06 ..		8 1	5,60 ..	
Oct.	2 1	12,32 ..		10 1	5,58 ..	
	4 1	12,14 ..		13 1	5,70 ..	
	6 1	11,88 ..		15 1	5,59 ..	
	9 1	12,05 ..		18 1	5,70 ..	
	12 1	12,46 ..		20 1	5,38 ..	
	14 1	12,18 ..		25 1	5,32 ..	
	19 1	17,48 ..	After the above was registered I lowered the East and 1/2 s.	27 1	5,18 ..	
	19 2	1,12 ..		March	1 1	6,50 ..
	23 1	1,63 ..			4 1	6,17 ..
	26 1	1,54 ..			6 1	6,28 ..
	28 1	1,50 ..			8 1	5,81 ..
	30 1	1,57 ..			10 1	5,12 ..
Nov.	1 1	0,60 ..			12 1	4,74 ..
	3 1	1,18 ..			15 1	5,29 ..
	5 1	0,86 ..			18 1	5,63 ..
	8 1	0,41 ..			20 1	5,10 ..
	10 1	0,32 ..		23 1	6,48 ..	
	12 1	0,81 ..		25 1	6,55 ..	
	14 1	1,02 ..		27 1	7,16 ..	
	16 1	1,06 ..		29 1	6,84 ..	
	21 1	1,77 ..	April	1 1	7,50 ..	
	23 1	1,35 ..		3 1	8,37 ..	

OBSERVATIONS FOR LEVEL.

11

1833	Illuminating Pivot.	Error from Level.	REMARKS.	1833	Illuminating Pivot.	Error from Level.	REMARKS.
	D. H.	s.			D. H.	s.	
April	5 1 East	8,19 W.		Sept.	3 1 East	3,97 E.	
	7 1	7,67 ..			6 1	4,36 ..	
	9 1	8,93 ..			9 1	4,56 ..	
	15 1	8,35 ..			11 1	5,81 ..	
	17 1	8,68 ..			14 1	7,30 ..	
	19 1	8,33 ..			16 1	7,84 ..	
	21 1	8,19 ..			18 1	7,18 ..	
	23 1	8,17 ..			21 1	8,31 ..	
	26 1	8,33 ..			25 1	8,64 ..	
	29 1	8,02 ..			27 1	8,95 ..	I raised the West end 10s.
May	2 1	7,45 ..			28 1	2,96 W.	
	4 1	7,66 ..			30 1	4,06 ..	
	6 1	8,46 ..		Oct.	3 1	4,29 ..	
	8 1	7,22 ..			5 1	3,50 ..	
	11 1	7,57 ..			7 1	4,63 ..	
	13 1	6,38 ..			9 1	4,25 ..	
	18 1	6,57 ..			11 1	3,50 ..	
	23 1	7,29 ..			14 1	3,65 ..	
	25 1	7,59 ..			16 1	3,45 ..	
June	4 1	7,95 ..			19 1	4,07 ..	
	8 1	6,96 ..			21 1	3,91 ..	
	10 1	6,76 ..			23 1	3,54 ..	Rainy weather
	12 1	6,10 ..			31 1	2,71 ..	no observations.
	14 1	5,74 ..		Nov.	7 1	0,20 E.	
	20 1	4,61 ..			9 1	0,09 ..	
	22 1	6,39 ..			13 7	Inverted the axis 6 times.
	24 1	4,65 ..			14 1	Do. 4 times.
	28 1	4,05 ..			14 2	0,25 E.	
	30 1	4,55 ..			18 1	0,85 W.	
July	4 1	3,37 ..			20 1	0,70 ..	
	8 1	3,01 ..			22 1	0,28 ..	
	14 1	2,79 ..			24 1	1,15 ..	
	17 1	1,96 ..			29 1	1,71 ..	
	20 1	1,10 ..		Dec.	2 1	1,89 ..	
	24 1	1,11 ..			4 1	1,98 ..	
	26 1	1,76 ..			7 1	1,84 ..	
	28 1	1,60 ..			9 1	0,67 ..	
Aug.	1 1	1,23 ..			11 1	2,22 ..	
	3 1	1,31 ..			13 1	1,70 ..	
	5 1	1,06 ..			15 1	1,96 ..	
	7 1	0,12 ..			19 1	1,78 ..	
	10 1	0,32 E.			20 1	Inverted the axis 20 times.
	13 1	0,94 ..			21 1	1,08 ..	
	15 1	0,88 ..			23 1	1,34 ..	
	19 1	1,78 ..			25 0	Inverted the axis several times.
	21 1	2,97 ..	Heavy rain with thunder and lightning.				
	24 1	3,86 ..					
	30 1	4,55 ..					

Since the above was put to Press it has occurred to me that some notion may be formed of the figure of the Pivots by noting the indications of the spirit level when applied to the axis under the various directions which the Telescope is capable of assuming when supporting the spirit level; from the

mean of four very careful readings agreeing very well *inter se*, the following were obtained.

Position of the Telescope. °	Error of Level. "		Position of the Telescope. °	Error of Level. "
290 N.P.D.	1,95 W.	110 N.P.D.	2,70 W.
300 —	2,00 —	120 —	2,75 —
310 —	1,80 —	130 —	2,20 —
320 —	1,45 —	140 —	1,75 —
330 —	1,00 —	150 —	1,50 —
340 —	1,05 —	160 —	1,80 —
350 —	2,55 —	170 —	2,05 —
360 —	2,60 —	180 —	2,45 —
10 —	2,50 —	190 —	2,70 —
20 —	2,50 —	200 —	2,35 —
30 —	2,35 —	210 —	2,20 —
40 —	2,10			

It will readily be understood that for the degrees of North Polar Distance intermediate between 40° and 110°, and between 210° and 290°, the spirit level cannot be applied; the results we have obtained, on the whole, are as accordant as might be expected; for assuming the mean of the above (2",10) as the true Error of Level, the greatest error amounts to 1',1.



ERROR OF COLLIMATION.

1832	Azimuth of		N. + S.	N + S + 25",07 2	Mean.	REMARKS.
	North mark.	South mark.				
	"	"	"	"		
April	24 +	30,20	- 24,78	+ 5,42	+ 15,25	
	25	30,25	24,71	5,54	15,30	
	26	30,49	24,61	5,88	15,47	
	27	30,35	24,99	5,36	15,21	
	28	30,32	24,99	5,33	15,20	
	29	30,53	25,12	5,41	15,24	
May	30	30,16	25,09	5,07	15,07	
	1	30,35	25,19	5,16	15,11	
	2	30,28	25,41	4,87	14,97	
	3	30,52	24,92	5,60	15,33	
	4	30,32	25,12	5,20	15,13	
	5	30,42	25,26	5,16	15,11	
	6	30,45	25,34	5,11	15,09	
	7	29,98	25,26	4,72	14,89	
	8	30,15	25,37	4,78	14,92	
	9	29,65	25,44	4,21	14,64	
	10	30,13	25,44	4,69	14,88	Mean of 39
11	30,22	25,51	4,71	14,89	+ 15",06	
12	31,10	23,65	7,45	16,26		
13	32,10	21,82	10,28	17,67		
14	32,17	21,89	10,28	17,67		
15	31,86	22,40	9,42	17,25		
16	31,79	22,09	9,70	17,38		
17	32,21	22,01	10,20	17,63		
18	33,48	22,41	11,07	18,07	
19	33,13	22,02	11,11	18,09	The Micrometer was taken out	
20	33,16	21,96	11,20	18,13	and cleaned, but no derange-	
21	32,82	22,34	10,48	17,78	ment appears to have thence	
22	32,34	22,48	9,86	17,47	resulted.	
23	31,62	23,03	8,59	16,83		
24	31,69	22,68	9,01	17,04		
25	31,44	22,85	8,59	16,83		
26	31,76	22,71	9,05	17,06		
27	31,69	22,51	9,18	17,12		
28	31,93	22,68	9,25	17,16		
29	31,90	22,55	9,35	17,21		
30	31,90	22,48	9,42	17,24		
31	32,13	22,02	10,11	17,59	Mean of 21	
June	1	32,52	21,72	10,80	17,93	+ 17",40
2	32,52	21,59	10,93	18,00		
3	32,62	21,45	11,17	18,12		
4	32,65	21,38	11,27	18,17		
5	32,72	21,45	11,27	18,17		
6	32,86	21,72	11,14	18,10		
7	33,03	21,13	11,90	18,48		
8	33,20	21,13	12,07	18,57		
9	33,02	21,52	11,50	18,28		
10	33,06	21,45	11,61	18,34		
11	33,06	21,20	11,86	18,47		
12	32,75	20,96	11,79	18,43		
13	32,41	21,03	11,38	18,22		
14	32,72	21,20	11,52	18,29		

ERROR OF COLLIMATION.

1832	Azimuth of		N. + S.	N + S +	Mean.	REMARKS.
	North mark.	South mark.		25",07		
	"	"	"	"		
June	16 + 33,16	- 21,13	+ 12,03	+ 18,55		
	17 33,06	21,20	11,86	18,47		
	18 32,52	20,96	11,56	18,31		
	19 32,79	21,41	11,38	18,22		
	20 32,65	20,45	12,20	18,63		
	21 32,82	21,13	11,69	18,38		
	22 32,86	21,31	11,55	18,31		
	24 32,72	20,99	11,73	18,40		
	25 32,68	20,56	12,12	18,59		
	27 32,72	20,96	11,76	18,41		
	28 32,62	20,69	11,93	18,50		
	29 32,62	20,69	11,93	18,50	Mean of 26	
	30 32,62	20,59	12,03	18,55	+ 18",37	
July	1 32,79	20,65	12,14	18,60		
	2 33,55	19,76	13,79	19,43		
	3 33,23	19,83	13,40	19,23		
	4 32,99	20,56	12,43	18,75		
	5 33,34	20,28	13,06	19,07		
	6 33,37	20,00	13,37	19,22		
	7 32,79	19,76	13,03	19,05		
	8 33,23	19,96	13,27	19,17	Mean of 8	
	14 33,06	+ 19",07	
	15	I adjusted the Instrument more nearly to the Meridian.
	16 67,59	32,67	34,92	30,00		
	17 67,04	33,36	33,68	29,37		
	24 67,56	32,70	34,86	29,97		
	25 67,56	33,01	34,55	29,81		
	27 67,21	33,36	33,85	29,46		
	29 67,28	32,88	34,40	29,73		
	30 67,90	32,67	35,23	30,15		
	31 67,56	33,84	33,72	29,39		
August	1 67,38	33,01	34,37	29,72	Mean of 10	
	7 66,59	32,39	34,20	29,63	+ 29",72	
	9	Found the first wire bent I put in a new set.
	13 36,09	- 67,73	- 31,64	- 3,28		
	14 36,26	66,00	29,74	2,33		
	15 36,61	66,35	29,74	2,33		
	17 36,44	65,72	29,28	2,11		
	18 36,47	66,07	29,60	2,26		
	19 35,54	66,17	30,63	2,78		
	20 35,91	66,35	30,44	2,68		
	21 35,77	66,03	30,26	2,59		
	22 35,96	65,72	29,76	2,35		
	23 35,91	65,33	29,42	2,17	Mean of 10	
	24 36,16	64,86	28,70	1,82	- 2",34	
	25 39,49	68,40	28,91	1,92		
	26 39,76	69,05	29,29	2,11		
	27 39,73	69,64	29,91	2,42		
	28 39,89	69,11	29,22	2,08		

ERROR OF COLLIMATION.

1832	Azimuth of		N. + S.	N + S + 25'' 07 2	Mean.	REMARKS.
	North mark.	South mark.				
	"	"	"	"		
August	29 + 39,69	- 68,91	- 29,22	- 2,08		
	30 39,96	68,77	28,81	1,87		
	31 40,21	67,49	27,28	1,11		
Sept.	1 40,21	67,50	27,29	1,11	Mean of 11	
	2 40,07	67,15	27,08	1,01	- 1'',93	
	3 40,00	70,43	30,43	2,08		
	4 39,59	70,29	30,70	2,81		
	5 38,70	70,46	31,76	3,35		
	6 38,52	70,94	32,42	3,67		
	7 38,70	70,87	32,17	3,55		
	8 38,81	71,11	32,30	3,61	Mean of 5	
	9 38,87	71,18	32,31	3,62	- 3'',56	
	13 39,42	70,46	31,04	2,98		
	14 40,38	Some trees obscured the South Mark.
	15 39,15					
	16 39,69					
	18 39,32					
	19 39,28					
	20 39,42	70,77	31,35	3,14		
	21 39,76	70,16	30,70	2,81		
	22 39,83	70,67	30,84	2,88		
	23 39,49	70,46	30,97	2,95		
	24 39,15	70,29	31,14	3,03		
	25 39,08	70,36	31,28	3,11		
	26 39,35	70,53	31,18	3,05		
	27 39,52	70,77	31,25	3,09		
	28 39,87	70,63	30,76	2,85		
	29 39,84	70,60	30,76	2,85		
	30 39,35	70,19	30,84	2,89		
October	1 39,76	70,46	30,70	2,81		
	3 39,89	70,87	30,98	2,95		
	4 40,03					
	5 39,76					
	6 39,55	70,15	30,60	2,76		
	7 39,98	70,49	30,56	2,75		
	8 39,79	70,46	30,67	2,80		
	9 39,76	70,63	30,87	2,90		
	10 40,18	70,12	29,94	2,43		
	11 40,18	69,66	29,48	2,21		
	12 39,93	70,15	30,22	2,57		
	13 39,59	70,35	30,76	2,85		
	14 39,69	70,19	30,50	2,71	Mean of 24	
	17 39,69	70,19	30,50	2,71	12'',79	
	19 39,15	72,80	33,65	4,29	Inverted the Axis several times.
	20 39,21	72,94	33,73	4,33		
	21 39,23	72,87	33,64	4,28		
	22 39,45	72,90	33,45	4,19		
	23 39,18	72,53	33,35	4,14		
	24 38,39	72,08	33,69	4,31		
	25 38,56	72,01	33,45	4,19		
	26 38,52	71,86	33,34	4,13		

ERROR OF COLLIMATION.

1832	Azimuth of		N. + S.	N + S +	Mean.	REMARKS.
	North mark.	South mark.		25",07		
	"	"	"	"		
October	27 + 38,15	- 71,97	- 33,82	- 4,38		
	28 37,98	71,83	33,85	4,39		
	29 38,22	71,48	33,26	4,10		
	30 38,66	72,00	33,34	4,13		
	31 38,49	72,00	33,51	4,22		
Nov.	1 38,39	71,86	33,47	4,20		
	2 38,25	71,14	32,89	3,91		
	3 38,56	71,48	32,92	3,92		
	4 38,12	71,65	33,53	4,23		
	5 37,97	71,38	33,41	4,17		
	6 38,32	71,48	33,16	4,04		
	7 38,12	71,14	33,02	3,98	Mean of 21	
	8 38,32	71,48	33,16	4,04	- 4",13	
	9 41,08	67,48	26,40	0,66	Found the moveable wire loose, which appeared to arise from the gum securing it, having swelled so much from the moist state of the air as to bring it in contact with the plate securing the fixed wires; to remedy this, I removed all the wires, and filed grooves at either end to contain the varnish necessary to secure their ends.
	10 41,21	67,32	26,11	0,52		
	11 41,58	67,66	26,08	0,51		
	12 41,04	67,56	26,52	0,72		
	13 40,77	67,18	26,41	0,67		
	14 41,01	67,25	26,24	0,58		
	15 40,60	67,18	26,58	0,75	Mean of 8	
	16 40,52	67,04	26,52	0,72	- 0",64	
	17 39,69	64,89	25,20	0,06		
	18 39,52	64,78	25,26	0,10		
	19 39,80	65,65	25,85	0,39		
	20 40,04	65,82	25,78	0,36		
	21 40,42	66,04	25,62	0,28		
	22 40,04	64,96	24,92	+ 0,07		
	23 40,21	64,86	24,65	+ 0,21		
	24 40,04	64,78	24,74	+ 0,16		
	25 39,69	64,96	25,27	- 0,10		
	26 39,97	65,33	25,36	0,14		
	27 39,73	65,48	25,75	0,34		
	28 39,86	65,62	25,76	0,35		
	29 39,73	65,47	25,74	0,34		
	30 40,00	64,89	24,89	+ 0,09		
Dec.	1 39,86	64,82	24,96	+ 0,05		
	2 40,53	64,75	24,22	+ 0,43		
	3 40,84	65,30	24,46	+ 0,31		
	4 40,49	64,96	24,47	0,30		
	5 40,21	64,61	24,40	0,33		
	6 40,04	64,55	24,51	0,28		
	7 40,17	64,48	24,31	0,38		
	8 39,73	64,04	24,31	0,38		
	9 39,70	64,21	24,51	0,28		
	10 39,76	64,41	24,65	0,21		
	11 40,04	64,59	24,55	0,26		
	12 40,31	64,61	24,30	0,38		
	13 40,21	64,71	24,50	+ 0,28		
	14 39,86	64,96	25,10	- 0,01		
	15 38,40	63,66	25,26	- 0,09		
	16 38,06	63,38	25,32	- 0,12		
	17 39,49	63,87	24,38	+ 0,35		

ERROR OF COLLIMATION.

1832	Azimuth of		N. + S.	N + S +	Mean.	REMARKS.
	North mark.	South mark.		25",07		
	"	"	"	2		
Dec.	18	+ 39,36	- 64,07	- 24,71	+ 0,18	
	19	39,18	63,76	24,58	+ 0,25	
	20	39,06	63,93	24,87	+ 0,10	
	21	38,84	64,21	25,37	- 0,15	
	22	38,49	63,83	25,34	0,13	
	23	38,56	63,93	25,37	0,15	
	24	38,25	63,59	25,34	0,13	
	25	38,36	63,88	25,52	0,22	
	26	38,40	63,11	24,71	+ 0,18	
	27	38,56	63,07	24,51	+ 0,28	
	28	38,33	63,93	25,60	- 0,26	
	29	38,15	64,78	26,63	0,78	
30	38,84	64,14	25,30	0,11		
31	39,22	64,34	25,12	0,02		
1833						
January	1	+ 38,84	- 64,61	- 25,77	0,35	
	2	38,73	64,00	25,27	0,10	
	3	38,40	63,90	25,50	0,21	
	4	38,84	63,59	24,75	+ 0,16	
	5	38,88	63,96	25,08	- 0,00 Inverted the Axis several times.
	6	38,70	64,00	25,30	0,11	
	7	38,49	64,24	25,75	0,34	Mean of 53
	8	38,56	64,18	25,62	0,27	- 0",01
	9	39,63	62,93	23,30	+ 0,88	
	10	40,63	62,42	21,79	+ 1,64	
	11	40,46	62,21	21,75	1,66	
	12	40,56	62,55	21,99	1,54	
	13	39,52	61,86	22,34	1,36	
	14	39,52	61,86	22,34	1,36	Mean of 6
	15	39,86	62,38	22,52	1,27	+ 1",47 Inverted the Axis several times.
17	40,97	60,21	19,24	+ 2,91		
18	42,52	61,34	18,82	3,12 Inverted the Axis several times.	
19	42,62	61,01	18,39	3,34		
20	42,45	61,59	19,14	2,96		
21	42,27	61,89	19,62	2,72		
22	42,62	61,17	18,55	3,26		
23	42,85	61,41	18,56	3,25		
24	42,34	60,86	18,52	3,27		
25	42,07	60,62	18,55	3,26		
26	41,58	60,21	18,63	3,22		
27	41,76	60,17	18,41	3,33	Mean of 12	
28	41,65	60,65	19,00	3,03	+ 3",14	
29	42,80	59,10	16,30	+ 4,38		
30	44,24	58,07	13,83	5,62		
31	44,34	58,42	14,08	5,49		
February	1	44,51	58,07	13,56	5,75	
	2	43,82	57,73	13,91	5,58	
	3	44,00	57,94	13,94	5,56	
	4	44,00	58,07	14,07	5,50	

ERROR OF COLLIMATION.

1833	Azimuth of		N. + S.	N + S + 25",07 2	Mean.	REMARKS.
	North mark.	South mark.				
	"	"	"	"		
February	5 + 44 21	- 58,42	- 14,21	+ 5,43		
	6 44,41	58,93	14,52	5,27		
	7 44,03	58,63	14,60	5,23		
	8 44,75	58,76	14,01	5,53		
	9 44,58	58,52	13,94	5,56		
	10 44,62	58,76	14,14	5,46		
	11 44,48	59,45	14 97	5,05		
	12 43,93	58,45	14,52	5,27		
	13 44,34	58,66	14,32	5,37		
	14 44,10	59,00	14,90	5,08		
	15 44,34	59,34	15,00	5,03		
	16 44,28	58,93	14,65	5,21		
	17 44,07	59,31	15,24	4,91		
	18 44,34	59,17	14,83	5,12		
	19 44,28	58,59	14,31	5,38		
	20 43,76	58,93	15,17	4,95		
	21 44,24	59,17	14,93	5,07		
	22 44,51	59,59	15,08	4,99		
	23 44,75	58,90	14,15	5,46		
	24 44,51	59,10	14,59	5,24		
	25 45,19	59,00	13,81	5,63		
	26 44,92	58,83	13,91	5,58		
	27 44,75	58,66	13,91	5,58		
	28 44,92	58,83	13,91	5,58		
March	1 44,51	58,59	14,08	5,49		
	2 44,68	58,73	14,05	5,51		
	3 44,92	58,86	13,94	5,56		
	4 45,19	59,10	13,91	5,58		
	5 44,89	58,59	13,70	5,68		
	6 44,68	58,70	14,02	5,52		
	7 45,05	58,91	13,86	5,60		
	8 45,58	59,17	13,59	5,74		
	9 44,03	58,59	14,56	5,25	Mean of 41	
	10 44,65	58,59	13,94	5,46	+ 5",87	
	11 42,44	61,00	18,56	3,25		
	12 42,30	60,97	18,67	3,20		
	13 42,73	60,60	17,87	3,60		
	14 42,90	60,55	17,65	3,71		
	15 42,35	60,48	18,13	3,47		
	16 43,03	61,04	18,01	3,53		
	17 43,27	61,62	18,35	3,36		
	18 42,51	60,86	18,35	3,36		
	19 42,51	61,00	18,49	3,29		
	20 42,48	60,48	18,00	3,53		
	21 42,96	60,51	17,55	3,76		
	22 43,34	60,48	17,14	3,96		
	23 42,99	60,55	17,56	3,75		
	24 42,44	60,14	17,70	3,68		
	25 42,62	60,17	17,55	3,76		
	26 42,27	60,14	17,87	3,60		
	27 42,17	61,17	19,00	3,03		
	28 42,30	60,83	18,53	3,27		
	29 42,27	60,93	18,66	3,20		

ERROR OF COLLIMATION.

1833	Azimuth of		N. + S.	N + S +	Mean.	REMARKS.
	North mark.	South mark.		25",07		
	"	"	"	2		
March	30	+ 42,30	- 61,04	- 18,74	+ 3,16	
	31	42,62	61,31	18,69	3,19	
April	1	42,48	60,69	18,21	3,43	
	2	42,58	60,83	18,25	3,41	
	3	42,51	60,83	18,32	3,37	
	4	43,06	60,86	17,80	3,63	
	5	42,79	60,83	18,04	3,51	
	6	42,83	60,93	18,10	3,48	
	7	42,83	61,14	18,31	3,38	
	8	42,27	60,90	18,63	3,22	
	9	42,44	61,04	18,60	3,23	
	10	42,37	61,14	18,77	3,15	
	11	42,96	61,38	18,42	3,32	
	12	42,58	60,83	18,25	3,41	
	13	42,30	60,62	18,32	3,37	
	14	42,34	60,83	18,49	3,29	
	15	42,62	60,69	18,07	3,50	
	16	42,37	60,97	18,60	3,23	
	17	42,34	60,90	18,56	3,25	
	18	42,41	60,58	18,17	3,45	
	19	42,41	60,61	18,20	3,43	
20	42,55	60,48	17,93	3,57		
21	42,55	60,31	17,76	3,65		
22	42,72	60,83	18,11	3,48		
23	42,30	60,86	18,56	3,25		
24	42,41	60,86	18,45	3,31		
25	42,58	61,11	18,53	3,27		
26	42,21	60,93	18,72	3,17		
27	42,72	60,90	18,18	3,44		
28	42,55	60,79	18,24	3,41		
29	42,48	60,83	18,35	3,36		
30	42,48	61,14	18,66	3,20		
May	1	41,62	60,48	18,86	3,10	
	2	42,10	60,83	18,73	3,17	
	3	42,10	60,65	18,55	3,26	
	4	42,48	60,55	18,07	3,50	
	5	42,48	61,04	18,56	3,25	
	6	42,10	61,00	18,90	3,08	
	7	42,13	60,62	18,49	3,29	
	8	42,17	61,04	18,87	3,10	
	9	42,44	60,72	18,28	3,39	
	10	42,14	60,69	18,55	3,26	Mean of 62
	11	42,14	60,83	18,69	3,19	+ 3",38
	12	40,56	63,45	22,89	1,09	
13	40,56	63,41	22,85	1,11		
14	39,80	63,90	24,10	0,48		
15	39,66	63,83	24,17	0,45		
16	39,86	64,58	24,72	0,18		
17	39,73	64,37	24,64	0,21	Mean of 7	
18	40,00	64,64	24,64	+ 0,21	+ 0",53	
19	40,49	65,58	25,09	- 0,01		
20	39,69	65,47	25,78	0,35		

The centre wire appears to have shifted its position no doubt from the action of the hot land wind; fearing it might have become loose I applied fresh varnish (tincture of Opium) without disturbing the Plate.

ERROR OF COLLIMATION.

1833	Azimuth of		N. + S.	N + S + 25",07 2	Mean.	REMARKS.
	North mark.	South mark.				
	"	"	"	"		
May	21 + 38,56	- 64,85	- 26,29	- 0,61		
	22 38,49	64,44	25,95	0,44		
	23 38,18	64,75	26,57	0,75		
	24 38,59	64,68	26,09	0,51		
	25 38,49	64,78	26,29	0,61		
	26 38,39	64,58	26,19	0,56		
	27 38,32	64,61	26,29	0,61		
	28 38,25	64,89	26,64	0,78		
	29 38,46	64,99	26,53	0,73		
	30 38,56	64,61	26,05	0,49		
	31 38,15	65,30	27,15	1,04		
June	1 38,84	64,34	25,50	0,21		
	2 38,43	64,61	26,18	0,56		
	3 38,70	64,27	25,57	0,25		
	4 38,67	64,34	25,67	0,30		
	5 38,73	64,34	25,61	0,27		
	6 38,52	64,55	26,03	0,48		
	7 38,18	64,14	25,96	0,44		
	8 38,56	64,61	26,05	0,49		
	9 38,70	63,96	25,26	0,09		
	10 38,81	64,14	25,33	0,13		
	11 38,98	64,00	25,02	+ 0,02		
	12 38,91	64,03	25,12	- 0,02		
	13 38,87	64,03	25,16	0,05		
	14 38,73	64,48	25,75	0,34		
	19 39,89	65,30	25,41	0,17		
	20 39,66	65,37	25,71	0,32		
	21	65,37				
	22 39,49	65,37	25,88	0,40		
	23 39,01	64,83	25,82	0,38		
	24 38,77					
	25 39,08	65,13	26,05	0,49		
	26 38,77	64,99	26,22	0,57		
	27 38,56	64,78	26,22	0,57		
	28 38,43	64,78	26,35	0,64		
	29 38,39					
	30 38,49					
July	1 38,59	64,51	25,92	0,42		
	2 39,09	64,41	25,32	0,12		
	3 39,21	64,10	24,89	+ 0,09		
	4 38,87	64,48	25,61	0,27		
	5 39,09	64,24	25,15	0,04		
	6 38,15	64,14	25,99	0,46		
	7 38,87	64,27	25,40	0,16		
	8 38,73	64,51	25,78	0,36		
	9 38,70	64,45	25,75	0,34		
	10 38,22	64,99	26,77	0,85		
	11 38,56	64,51	25,95	0,44		
	12 38,80	65,03	26,23	0,58		
	13 38,56	64,71	26,15	0,54		
	14 38,52	64,68	26,16	0,54		
	15 38,36	64,68	26,32	0,62		
	16 38,12	63,93	25,81	0,37		
	17 38,29	64,37	26,08	0,50		

ERROR OF COLLIMATION.

1833	Azimuth of		N. + S.	N + S + 25",07 2	Mean.	REMARKS.
	North mark.	South mark.				
	"	"	"	"		
July	18 + 37,91	- 64,21	- 26,30	- 0,62		
	19 38,22	64,00	25,78	0,35		
	20 38,32	64,58	26,26	0,59		
	21 38,22					
	22 38,18	64,75	26,57	0,75		
	23 38,09	64,27	26,18	0,56		
	24 38,25	64,71	26,46	0,69		
	25 38,32	64,45	26,13	0,53	Mean of 60	
	26 38,25	64,58	26,33	0,63	- 0",44	
	27 38,15					
	28 37,98	64,68	26,70	0,81		
	29 37,98					
	30 37,67	64,58	26,91	0,92		
	31 37,40	64,55	27,15	1,04		
August	1 37,02	65,03	28,01	1,47		
	2 37,56	64,21	26,65	0,79		
	3 37,49	63,77	26,28	0,60		
	4 37,60	64,51	26,91	0,92		
	5 37,98	64,68	26,70	0,81		
	6 37,63	64,44	26,81	0,87		
	7 37,84	64,71	26,87	0,90		
	8 37,29	64,27	26,98	0,95		
	9 37,12	64,21	27,09	1,01		
	10 37,49	64,27	26,78	0,85		
	11 37,56	64,52	26,96	0,94		
	12 37,49	64,34	26,85	0,89		
	13 37,81	64,45	26,64	0,78		
	14 37,33	64,21	26,88	0,90		
	15 37,12	64,52	27,40	1,16		
	16 37,06	64,27	27,21	1,07		
	17 37,03	64,11	27,08	1,01		
	18 37,43					
	19 37,49	64,27	26,78	0,85		
	20 37,46	64,27	26,81	0,87		
	21 37,81	64,89	27,08	1,01		
	22 37,37	64,61	27,24	1,08	Mean of 25	
	23 37,12	64,61	27,49	1,21	- 0",95	In consequence of heavy rain last night having leaked through the roof, a few drops had fallen upon the eye piece and bent one of the wires ; I put in a new set.
	24 36,85	65,72	28,87	1,90		
	25 36,78	65,72	28,94	1,93		
	26 37,46	67,04	29,58	2,25		
	27 37,46	66,42	28,96	1,94		
	28 38,32	67,80	29,48	2,20		
	30 38,66	66,79	28,13	1,53		
	31 38,43	66,66	28,23	1,58		
Sept.	1 38,52	66,86	28,34	1,63		
	2 38,73	67,04	28,31	1,62	Mean of 9	
	3 38,73	- 1",84	
	4 38,84					
	8 39,52	65,79	26,27	0,60		
	9 39,69	66,24	26,55	0,74		
	10 39,28	66,24	26,96	0,94		

ERROR OF COLLIMATION.

1833	Azimuth of		N. + S.	N + S + 25",07 2	Mean.	REMARKS.
	North mark.	South mark.				
Sept.	11	+ 39,52	- 66,69	- 27,17	- 1,04	Mean of 5 — 0",82
	12	39,08	65,82	26,74	0,83	
	13	37,02	67,38	30,36	2,64	
	14	36,95	67,87	30,92	2,92	
	15	36,95	67,59	30,64	2,78	
	16	37,03	67,69	30,66	2,79	
	17	37,35	67,59	30,23	2,58	
	18	36,95	67,76	30,81	2,87	
	19	36,72	67,97	31,25	3,09	
	20	37,12	68,14	31,02	2,97	
	21	37,19	67,90	30,71	2,82	
	22	36,44				
	23	36,85	68,07	31,22	3,07	
	24	36,78	68,14	31,36	3,14	
	25	36,72	68,01	31,29	3,11	
	26	36,69	68,01	31,32	3,12	
	27	37,81	68,43	30,62	2,78	
	28	38,49	68,68	30,19	2,56	
	29	38,56	69,02	30,46	2,69	
	30	38,94	67,76	28,82	1,87	
October	1	38,94	68,47	29,53	2,23	Mean of 20 — 2",75
	2	39,42	69,60	30,18	2,56	
	3	38,84	68,95	30,11	2,52	
	4	38,59	68,64	30,05	2,49	
	5	38,56	68,68	30,12	2,52	
	6	38,43	68,74	30,31	2,62	
	7	38,53	68,64	30,11	2,52	
	8	38,49	68,74	30,25	2,59	
	9	39,01	68,24	29,23	2,08	
	10	38,91				
	11	38,87	68,40	29,53	2,23	
	12	38,91				
	13	38,66	68,31	29,65	2,29	
	14	38,05	68,14	30,09	2,51	Mean of 10 — 2",43
	15	38,32	67,97	29,65	2,29	
	16	38,39	67,73	29,34	2,13	
	17	39,08	67,38	28,30	1,61	
	18	38,91	67,18	28,27	1,60	
	19	38,77				
	20	38,87				
	21	38,46	68,31	29,85	1,39	
	22	38,63	68,01	29,38	2,15	
	23	38,91	68,01	29,10	2,01	
	24	38,91	68,07	29,16	2,04	
	25	39,18	67,35	28,17	1,55	
	26	39,18	Trees obscured the South Mark.
	27	39,25	
Nov.	1	37,81	67,38	29,57	2,25	
	2	38,15				
	3	38,77	67,11	28,34	1,63	
	4	38,84				

ERROR OF COLLIMATION.

1833	Azimuth of		N. + S.	N + S + 25",07 2	Mean.	REMARKS.
	North mark.	South mark.				
	"	"	"	"		
Nov.	5 + 38,77	- 67,04	- 28,27	- 1,60		
	6 39,42					
	7 39,39	67,38	27,99	1,46		
	8 38,87					
	9 38,46	67,11	28,65	1,79		
	11 37,81					
	12 36,09					
	13 36,72					
	15 35,57	64,96	29,39	2,16	Inverted the Axis several times.
	16 36,09	64,68	28,59	1,76		
	17 35,84	64,71	28,87	1,90		
	18 35,30	63,96	28,66	1,79		
	19 35,30	63,90	28,60	1,76		
	20 35,20	64,27	29,07	2,00		
	21 34,71	64,00	29,29	2,11		
	22 35,06	64,27	29,21	2,07		
	23 35,27	63,27	28,00	1,47		
	24 35,23	63,59	28,36	1,64		
	25 35,23	63,24	28,01	1,47	Mean of 25	
	26 35,13	63,07	27,94	1,43	- 1",83	
	27 35,09	62,86	27,77	1,35		
	28 34,74	62,21	27,47	1,20		
	29 34,58	62,10	27,52	1,22		
	30 34,74	62,14	27,40	1,16		
Dec.	1 34,99	62,28	27,29	1,11		
	2 34,71	62,00	27,29	1,11		
	3 34,78	62,17	27,39	1,16		
	4 34,58	61,93	27,35	1,14		
	5 34,71	62,03	27,32	1,12		
	6 34,78	62,14	27,36	1,14		
	7 35,13	62,21	27,08	1,01		
	8 35,06	62,28	27,22	1,07		
	9 34,92	62,35	27,43	1,18		
	10 34,37	61,73	27,36	1,14		
	11 34,03	61,86	27,83	1,38		
	12 34,10	61,93	27,83	1,38		
	13 34,69	62,14	27,45	1,19		
	14 34,44	61,89	27,45	1,19		
	15 34,54	62,03	27,49	1,21		
	16 34,68	62,21	27,53	1,23		
	17 35,27	62,58	27,31	1,12		
	18 34,40	61,93	27,53	1,23		
	19 34,58	62,10	27,52	1,22	Mean of 24	
	20 33,90	62,03	28,13	1,53	- 1",21	Inverted the Axis several times.
	23 32,31					
	24 31,86	62,83	30,97	2,95	Inverted the Axis several times.
	25 33,24	61,52	28,28	1,60		
	26 34,03	62,55	28,52	1,72		
	27 33,66	61,55	27,89	1,41		
	28 33,59	61,86	28,27	1,60	Mean of 7	
	29 33,59	61,86	28,27	1,60	- 1",62	
	30 33,16	61,86	28,70	1,81		
	31 33,37	61,53	28,16	1,54		

ERROR OF AZIMUTH.

From the foregoing pages, it appears that the Angular distance between the North and South marks has varied from $26^{\circ},97'$, to $25^{\circ},07'$, in the interval between April 1831, and October 1832, it will consequently be our first step to enquire which of the Marks, or if both of them, have moved? for this purpose we will now consult the observations of the Pole Star: correcting the observed transit for Error of the Clock, Error of Collimation, and the Error for Level as modified by the wear of the pivots (already explained at Pages 7 and 8); we obtain the apparent place *affected by the Azimuthal Error*; applying to this the Equations for aberration, nutation, and precession; we obtain the Mean place at the beginning of the year as *affected by the Azimuthal Error*; selecting now the consecutive observations *above and below* the pole, we can determine the values of a^I a^{II} &c. the errors in Azimuth of the center wire as follows:

1832	Observed Transit.	Error of Clock.	Correction for		Aberration, &c.	Mean Right Ascension January 1, 1832.	Resulting values of a^I , a^{II} , &c.
			Level.	Collimation.			
Dec. 7	<i>h. m. s.</i> 0 59 46,17	<i>m. s.</i> + 0 50,04	<i>s.</i> + 1,20	<i>s.</i> - 0,68	<i>s.</i> - 43,28	<i>h. m. s.</i> 0 59 53,45	$a^I = 4,70$ $a^{II} = 4,55$ $a^{III} = 4,08$ $a^{IV} = 3,97$ $a^V = 3,98$ $a^{VI} = 4,08$ $a^{VII} = 3,47$ $a^{VIII} = 3,92$ $a^{IX} = 3,50$ $a^{X} = 3,77$ $a^{XI} = 3,40$ $a^{XII} = 4,39$ $a^{XIII} = 3,28$ $a^{XIV} = 3,28$
7 S.P.	13 0 7,00	+ 0 52,06	- 1,40	+ 0,68	- 42,96	13 0 15,38	
8 S.P.	12 59 59,67	+ 0 56,56	- 1,50	+ 0,68	- 42,32	13 0 13,09	
9	0 59 34,17	+ 0 58,70	+ 1,74	- 0,68	- 42,00	0 59 51,93	
9 S.P.	12 59 52,00	+ 1 0,79	- 1,49	+ 0,68	- 41,69	13 0 13,29	
10	0 59 31,50	+ 1 2,95	+ 1,91	- 0,68	- 41,37	0 59 54,31	
12	0 59 18,20	+ 1 11,71	+ 1,99	- 0,68	- 40,06	0 59 51,16	
12 S.P.	12 59 36,00	+ 1 14,27	- 1,56	+ 0,68	- 41,44	13 0 9,67	
13	0 59 12,40	+ 1 16,79	+ 2,06	- 0,68	- 39,38	0 59 51,19	
13 S.P.	12 59 30,00	+ 1 19,08	- 1,60	+ 0,68	- 39,02	13 0 9,88	
15	0 59 2,30	+ 1 26,47	+ 1,85	- 0,68	- 37,98	0 59 51,86	
15 S.P.	12 59 20,67	+ 1 28,57	- 1,45	+ 0,68	- 37,63	13 0 10,84	
16	0 59 0,60	+ 1 30,67	+ 1,73	- 0,68	- 37,28	0 59 55,04	
16 S.P.	12 59 16,30	+ 1 32,93	- 1,36	+ 0,68	- 36,93	13 0 11,62	
17	0 58 54,14	+ 1 34,59	+ 1,78	- 0,68	- 36,58	0 59 53,25	
18 S.P.	12 59 5,40	+ 1 42,06	- 1,45	+ 0,68	- 35,50	13 0 11,19	
19	0 58 44,50	+ 1 44,45	+ 1,73	- 0,68	- 35,14	0 59 54,86	
20	0 58 36,00	+ 1 49,12	+ 1,89	- 0,68	- 34,42	0 59 51,91	
20 S.P.	12 58 53,00	+ 1 51,20	- 1,28	+ 0,68	- 34,06	13 0 9,54	
21	0 58 32,60	+ 1 53,98	+ 1,52	- 0,68	- 33,71	0 59 53,71	
23 S.P.	12 58 41,10	+ 2 5,99	- 1,40	+ 0,68	- 31,86	13 0 14,51	
24	0 58 13,70	+ 2 8,65	+ 1,67	- 0,68	- 31,46	0 59 51,88	
24 S.P.	12 58 31,10	+ 2 10,90	- 1,36	+ 0,68	- 31,11	13 0 10,21	
25	0 58 11,10	+ 2 13,55	+ 1,65	- 0,68	- 30,71	0 59 54,91	
26	0 58 7,00	+ 2 18,08	+ 1,59	- 0,68	- 29,19	0 59 56,03	
26 S.P.	12 58 21,30	+ 2 20,19	- 1,30	+ 0,68	- 29,60	13 0 11,27	

1833	Observed Transit.	Error of Clock.	Correction for		Aberration, &c.	Mean Right Ascension January 1, 1832.		Resulting values of $a^1, a^2, \&c.$	
			Level.	Collimation.		h. m. s.	h. m. s.		
Jan.	2 S.P.	12 57 45,00	+ 2 50,39	- 1,66	+ 1,23	- 24,12	13 0 10,84	- 2,351 a^{xv}	} $a^{xv} = 3,84$
	3	0 57 24,00	+ 2 52,56	+ 2,12	- 1,23	- 23,72	0 59 53,73	+ 2,316 a^{xv}	
	3 S.P.	12 57 42,00	+ 2 54,22	- 1,67	+ 1,23	- 23,32	13 0 12,46	- 2,351 a^{xvi}	} $a^{xvi} = 4,36$
	4	0 57 17,80	+ 2 55,92	+ 2,14	- 1,23	- 22,92	0 59 51,71	+ 2,316 a^{xvi}	
	4 S.P.	12 57 37,00	+ 2 57,57	- 1,69	+ 1,23	- 22,52	13 0 11,59	- 2,351 a^{xvii}	} $a^{xvii} = 3,84$
	5	0 57 12,40	+ 2 59,23	+ 1,98	- 1,23	- 22,12	0 59 52,26	+ 2,316 a^{xvii}	
	5 S.P.	12 57 11,00	+ 3 0,76	- 1,55	+ 1,23	- 21,73	13 0 8,71	- 2,351 a^{xviii}	} $a^{xviii} = 3,96$
	6	0 57 30,30	+ 3 2,28	+ 1,83	- 1,23	- 21,33	0 59 52,85	+ 2,316 a^{xviii}	
	6 S.P.	12 57 31,00	+ 3 4,02	- 1,37	+ 1,23	- 20,93	13 0 13,95	- 2,351 a^{xix}	} $a^{xix} = 4,56$
	7	0 57 7,00	+ 3 5,76	+ 1,66	- 1,23	- 20,53	0 59 52,66	+ 2,316 a^{xix}	
	8	0 57 4,30	+ 3 9,37	+ 1,67	- 1,23	- 19,74	0 59 54,37	+ 2,316 a^{xx}	} $a^{xx} = 3,51$
	8 S.P.	12 57 19,10	+ 3 11,11	- 1,31	+ 1,23	- 19,35	13 0 10,78	- 2,351 a^{xx}	
	10	0 56 46,20	+ 3 17,11	+ 1,92	+ 2,88	- 18,16	0 59 49,95	+ 2,316 a^{xxi}	} $a^{xxi} = 5,02$
	10 S.P.	12 57 16,60	+ 3 18,87	- 1,45	- 2,88	- 17,77	13 0 13,37	- 2,351 a^{xxi}	
	11	0 56 43,70	+ 3 20,64	+ 2,04	+ 2,88	- 17,37	0 59 51,89	+ 2,316 a^{xxii}	} $a^{xxii} = 4,61$
	11 S.P.	12 57 12,50	+ 3 22,27	- 1,55	- 2,88	- 16,97	13 0 13,37	- 2,351 a^{xxii}	
	14 S.P.	12 56 56,40	+ 3 30,85	- 1,17	- 2,88	- 14,58	13 0 8,62	- 2,351 a^{xxiii}	} $a^{xxiii} = 3,44$
15	1 0 32,60	- 0 28,88	+ 2,44	+ 2,88	- 14,18	0 59 54,86	+ 2,316 a^{xxiii}		
15 S.P.	13 1 4,90	- 0 28,98	- 1,96	- 6,90	- 13,79	13 0 13,27	- 2,351 a^{xxiv}	} $a^{xxiv} = 2,11$	
16	1 0 33,00	- 0 29,35	+ 3,44	+ 6,90	- 13,40	1 0 0,59	+ 2,316 a^{xxiv}		
16 S.P.	13 0 59,50	- 0 29,24	- 2,75	- 6,90	- 13,00	13 0 7,61	- 2,351 a^{xxv}	} $a^{xxv} = 3,40$	
17	1 0 24,60	- 0 30,01	+ 4,43	+ 6,90	- 12,61	0 59 53,30	+ 2,316 a^{xxv}		
17 S.P.	13 1 3,80	- 0 30,38	- 3,54	- 6,90	- 12,21	13 0 10,77	- 2,351 a^{xxv}		

If we now correct the *observed* Azimuth of the North and South Marks for the Error of Collimation, with the assistance of the above values of a we can determine their *true* Azimuth from the meridian; thus:

1832	Azimuth of the centre wire from Meridian.	Azimuth from centre wire of		Azimuth from the Meridian of		REMARKS.	
		North mark.	South mark.	North mark.	South mark.		
December	7 a^1	= N. 4,70 E.	N. 39,79 W.	S. 64,86 E.	N. 35,09 E.	S. 60,16 E.	
	9 a^{ii}	= - 4,55	.. 39,42	.. 64,49	.. 35,37	.. 60,44	..
	10 a^{iii}	= - 4,08	.. 39,55	.. 64,62	.. 35,47	.. 60,54	..
	12 a^{iv}	= - 3,97	.. 39,93	.. 65,00	.. 35,96	.. 61,03	..
	13 a^v	= - 3,98	.. 39,93	.. 65,00	.. 35,95	.. 61,02	..
	15 a^{vi}	= - 4,08	.. 38,50	.. 63,57	.. 34,42	.. 59,49	..
	16 a^{vii}	= - 3,47	.. 38,18	.. 63,25	.. 34,71	.. 59,78	..
	17 a^{viii}	= - 3,92	.. 39,15	.. 64,22	.. 35,23	.. 60,30	..
	19 a^x	= - 3,50	.. 38,94	.. 64,01	.. 35,44	.. 60,51	..
	20 a^c	= - 3,77	.. 38,96	.. 64,03	.. 35,09	.. 60,16	..
	21 a^{xi}	= - 3,40	.. 38,99	.. 64,06	.. 35,59	.. 60,66	..
	24 a^{xii}	= - 4,39	.. 38,38	.. 63,45	.. 33,99	.. 59,06	..
	25 a^{xiii}	= - 3,28	.. 38,58	.. 63,65	.. 35,30	.. 60,37	..
	26 a^{xiv}	= - 3,28	.. 38,22	.. 63,29	.. 34,94	.. 60,01	..
1833 January	3 a^{xv}	= - 3,84	.. 38,61	.. 63,68	.. 34,77	.. 59,84	..
	4 a^{xvi}	= - 4,36	.. 38,68	.. 63,75	.. 34,32	.. 59,39	..

ERROR OF AZIMUTH.

1833	Azimuth of the centre wire from Meridian.	Azimuth from centre wire of		Azimuth from the Meridian of		REMARKS.
		North mark.	South mark.	North mark.	South mark.	
January	5 a^{xvii} = N. 3,84 E.	N. 38,86 W.	S. 63,96 E.	N. 35,04 E.	S. 60,12 E.	
	6 a^{xviii} = — 3,96 ..	38,81 ..	63,88 ..	34,85 ..	59,92 ..	
	7 a^{xix} = — 4,56 ..	38,83 ..	63,90 ..	34,27 ..	59,34 ..	
	8 a^{xx} = — 3,51 ..	38,85 ..	63,92 ..	35,34 ..	60,41 ..	
	10 a^{xxi} = — 5,02 ..	39,00 ..	64,07 ..	33,98 ..	59,05 ..	
	11 a^{xxii} = — 4,61 ..	38,80 ..	63,87 ..	34,19 ..	59,26 ..	
	15 a^{xxiii} = — 3,44 ..	38,59 ..	63,66 ..	35,15 ..	60,22 ..	
	16 a^{xxiv} = — 2,11 ..	38,59 ..	63,66 ..	36,48 ..	41,55 ..	
	17 a^{xxv} = — 3,40 ..	38,06 ..	63,13 ..	34,66 ..	59,73 ..	

Taking the mean of 25 we have 35°,02 and 60°,09 for the true Azimuths. If N and S represent the *observed* Azimuth of the Centre wire from the North and South Marks and, C the Error of Collimation.

$$\begin{aligned} \text{The Azimuth of the Centre wire from the Meridian} &= 35^{\circ},02 \pm C - N. \\ &= 60^{\circ},09 \mp C - S. \\ \text{Taking half the sum} &= 95^{\circ},11 - N - S. \end{aligned}$$

Similarly we have for the Year 1833.

1833	Observed Transit.	Error of Clock.	Correction for		Aberration, &c.	Mean Right Ascension, Jan. 1, 1832.	Resulting values of $a^1, a^2, \&c.$			
			Level.	Collimation.						
Nov.	19	1 1 27,75	— 0 29,87	+ 3,89	— 5,13	— 46,03	h. m. s. 0 59 54,83 + a^1	} $a^1 = 2,63$		
	19 S.P.	13 1 35,75	— 0 28,76	— 3,05	+ 5,13	— 45,79	13 0 7,50 — a^2			
	20	1 1 25,70	— 0 27,78	+ 3,85	— 5,13	— 45,55	0 59 55,31 + a^1			
	20 S.P.	13 1 36,43	— 0 27,06	— 3,01	+ 5,13	— 45,31	13 0 10,40 — a^2			
	23	1 1 17,25	— 0 23,03	+ 3,85	— 5,13	— 44,11	0 59 53,05 + a^3			
	23 S.P.	13 1 30,14	— 0 22,32	— 3,02	+ 5,13	— 43,86	13 0 10,29 — a^3			
	Dec.	5	1 0 56,80	— 0 7,70	+ 4,63	— 3,63	— 37,47		0 59 56,85 + a^4	} $a^4 = 2,63$
		5 S.P.	13 1 10,90	— 0 7,64	— 3,70	+ 3,63	— 37,17		13 0 10,24 — a^4	
		6	1 0 57,25	— 0 6,88	+ 4,59	— 3,63	— 36,85		0 59 58,70 + a^4	
		6 S.P.	13 1 5,27	— 0 6,74	— 3,62	+ 3,63	— 36,54		13 0 6,22 — a^5	
7		1 0 54,00	— 0 6,06	+ 4,58	— 3,63	— 36,21	0 59 56,90 + a^5			
8		1 0 54,25	— 0 5,76	+ 4,18	— 3,63	— 35,58	0 59 57,68 + a^6			
8 S.P.		13 1 4,50	— 0 4,94	— 3,30	+ 3,63	— 35,27	13 0 8,84 — a^6			
9		1 0 54,50	— 0 4,68	+ 3,86	— 3,63	— 34,95	0 59 59,32 + a^6			
9 S.P.		13 1 2,64	— 0 4,23	— 3,05	+ 3,63	— 34,64	13 0 8,57 — a^7			
10		1 0 51,75	— 0 3,60	+ 4,34	— 3,63	— 34,31	0 59 58,77 + a^7			
10 S.P.	13 0 59,17	— 0 3,09	— 3,45	+ 3,63	— 33,97	13 0 6,51 — a^8	} $a^8 = 1,76$			
11	1 0 50,25	— 0 2,48	+ 4,87	— 3,63	— 33,64	0 59 59,59 + a^8				
11 S.P.	13 0 59,57	— 0 2,16	— 3,82	+ 3,63	— 33,30	13 0 8,14 — a^9	} $a^9 = 1,78$			
12	1 0 51,00	— 0 1,60	+ 4,71	— 3,63	— 32,97	1 0 1,73 + a^9				
12 S.P.	13 0 56,50	— 0 1,03	— 3,69	+ 3,63	— 32,63	13 0 7,00 — a^{10}	} $a^{10} = 1,11$			
18	1 0 29,50	+ 0 10,04	+ 4,63	— 3,63	— 28,76	0 59 56,00 + a^{11}				
18 S.P.	13 0 32,00	+ 0 11,06	— 3,63	+ 3,63	— 28,43	13 0 4,85 — a^{11}	} $a^{11} = 1,50$			
19	1 0 29,62	+ 0 12,70	+ 4,60	— 3,63	— 28,04	0 59 59,47 + a^{11}				

1833	Observed Transit.	Error of Clock.	Correction for		Aberration, &c.	Mean Right Ascension, Jan. 1, 1832.			Resulting values of $a^1, a^2, \&c.$
			Level.	Collimation.		<i>h.</i>	<i>m.</i>	<i>s.</i>	
Dec. 24	S.P. 13 0 18,00	+ 0 26,41	- 4,35	+ 7,43	- 24,00	13 0 7,71	-	a^{13}	} $a^{12} = 1,32$
25	1 0 11,00	+ 0 27,77	+ 6,70	- 4,62	- 23,62	1 0 1,45	+ a^{12}		
25	S.P. 13 0 17,80	+ 0 28,77	- 5,27	+ 4,62	- 23,24	13 0 6,90	-	a^{13}	} $a^{13} = 1,55$
26	1 0 3,25	+ 0 30,28	+ 6,41	- 4,62	- 22,85	0 59 57,69	+ a^{13}		
26	S.P. 13 0 16,00	+ 0 31,25	- 5,08	+ 4,62	- 22,49	13 0 8,52	-	a^{14}	} $a^{14} = 2,07$
27	1 0 3,89	+ 0 32,24	+ 6,11	- 4,62	- 22,09	0 59 59,75	+ a^{14}		
27	S.P. 13 0 12,00	+ 0 32,91	- 4,84	+ 4,62	- 21,70	13 0 7,21	-	a^{15}	} $a^{15} = 1,58$
28	S.P. 13 0 7,50	+ 0 35,15	- 4,74	+ 4,62	- 20,92	13 0 5,83	-	a^{16}	
29	0 59 56,37	+ 0 36,66	+ 5,81	- 4,62	- 20,52	0 59 57,92	+ a^{16}		} $a^{16} = 1,68$
30	0 59 57,12	+ 0 38,30	+ 5,67	- 4,62	- 19,74	1 0 0,95	+ a^{17}		
30	S.P. 12 59 59,93	+ 0 39,65	- 4,49	+ 4,62	- 19,35	13 0 4,58	-	a^{17}	} $a^{17} = 0,82$
31	0 59 53,00	+ 0 41,22	+ 5,52	- 4,62	- 18,95	1 0 0,39	+ a^{17}		

Comparing these values of $a, a^1, \&c.$ as before, with the Azimuthal readings corrected for Error of Collimation, we obtain as follows :

1833	Azimuth of the centre wire from Meridian.	Azimuth from centre wire of		Azimuth from the Meridian of		REMARKS.
		North mark.	South mark.	North mark.	South mark.	
November 19	$a^1 = N. 2,63 E.$	<i>s.</i> N. 37,06	<i>s.</i> W. S. 62,13	<i>s.</i> N. 34,43	<i>s.</i> W. S. 59,50	E.
20	$a^2 = - 3,19$.. 37,20	.. 62,07	.. 34,01	.. 59,08	..
23	$a^3 = - 3,64$.. 36,73	.. 61,80	.. 33,09	.. 58,16	..
December 5	$a^4 = - 2,63$.. 35,82	.. 60,89	.. 33,19	.. 58,26	..
6	$a^5 = - 1,77$.. 35,92	.. 60,99	.. 34,15	.. 59,22	..
8	$a^6 = - 2,19$.. 36,13	.. 61,20	.. 33,94	.. 59,01	..
9	$a^7 = - 2,02$.. 36,10	.. 61,17	.. 34,08	.. 59,15	..
10	$a^8 = - 1,76$.. 35,52	.. 60,59	.. 33,76	.. 58,83	..
11	$a^9 = - 1,78$.. 35,41	.. 60,48	.. 33,63	.. 58,70	..
12	$a^{10} = - 1,11$.. 35,41	.. 60,48	.. 34,30	.. 59,37	..
18	$a^{11} = - 1,50$.. 35,63	.. 60,70	.. 34,13	.. 59,20	..
24	$a^{12} = - 1,32$.. 34,81	.. 59,88	.. 33,49	.. 58,56	..
25	$a^{13} = - 1,55$.. 34,84	.. 59,91	.. 33,29	.. 58,36	..
26	$a^{14} = - 2,07$.. 35,75	.. 60,82	.. 33,68	.. 58,75	..
27	$a^{15} = - 1,58$.. 35,07	.. 60,14	.. 33,49	.. 58,56	..
28	$a^{16} = - 1,68$.. 35,19	.. 60,26	.. 33,51	.. 58,58	..
30	$a^{17} = - 0,82$.. 35,19	.. 60,26	.. 34,37	.. 59,44	..
Mean of 17 = N. 33''79 W. S. 58''86 E.						

Whence it appears that the North and South Marks have each deviated 1^m23 , following the same direction in which they had first moved ; for the observations of 1832, we have already found (Page 30.)

The Error in Azimuth..... = $95^m,11 - N - S.$

Whereas for the observations towards the end of 1833, we now find it..... = $92^m,65 - N - S.$

As no circumstances offer to shew, if the present alteration took place gradually, or on a sudden; we will for the present suppose the latter number to take effect from the first of July 1833; and for the previous 6 months, employ the mean of the determinations for 1832 and 1833 or $93^{\circ},87' - N - S$

2

with these we will now compute the observations of the Pole Star, δ Ursæ Minoris, and 76 Draconis, which have been observed both above and below the Pole in 1832 and 1833.

POLARIS AT SUPERIOR CULMINATION.								
1832	Observed A.R. correct- ed for Error of Clock.		Correction for			Aberration, &c.	Mean A.R. 1832.	
			Level.	Azimuth.	Collimation.			
	1h.						1h.	
	m.	s.	s.	s.	s.	s.	m.	s.
October	19	0 53,17	+ 1,38	+ 18,91	- 10,76	- 61,71	0	0,99
	21	0 54,20	+ 1,22	- 61,48	0	2,09
	22	0 55,00	+ 1,16	- 61,36	0	2,95
	23	0 53,56	+ 1,06	- 61,24	0	1,53
	24	0 53,74	+ 1,08	- 61,08	0	1,89
	25	0 54,49	+ 1,10	- 60,92	0	2,82
	26	0 55,57	+ 1,12	- 60,76	0	4,08
	27	0 55,17	+ 1,15	- 60,61	0	3,86
	28	0 55,81	+ 1,16	+ 17,22	- 60,40	0	3,03
	29	0 54,76	+ 0,61	- 60,19	0	1,64
	30	0 56,73	+ 0,00	- 59,98	0	4,21
November	31	0 55,31	+ 1,46	- 59,77	0	3,46
	1	0 55,15	+ 1,75	- 59,51	0	3,85
	2	0 53,47	+ 1,57	- 59,24	0	2,26
	3	0 53,43	+ 1,41	- 58,98	0	3,32
	4	0 51,32	+ 1,49	- 58,71	0	1,56
	5	0 53,04	+ 1,59	- 58,34	0	2,75
	12	0 45,42	+ 1,62	+ 15,34	- 2,21	- 56,06	0	4,20
	13	0 44,80	+ 1,56	- 55,66	0	3,83
	15	0 45,71	+ 1,48	- 54,86	0	5,46
	16	0 44,76	+ 1,46	- 54,46	0	4,89
	17	0 42,52	+ 1,43	+ 11,48	- 0,68	- 54,01	0	0,74
	18	0 42,99	+ 1,34	- 53,56	0	1,57
	19	0 41,42	+ 1,26	- 53,11	0	0,37
21	0 42,96	+ 1,08	- 52,16	0	2,68	
22	0 41,97	+ 1,21	- 51,69	0	2,29	
23	0 42,49	+ 1,41	- 51,19	0	3,51	
25	0 41,51	+ 1,41	- 50,14	0	3,58	
29	0 38,84	+ 1,43	- 48,01	0	3,06	
30	0 38,50	+ 1,46	- 47,44	0	3,32	
December	3	0 36,88	+ 1,57	- 45,64	0	3,61
	4	0 35,79	+ 1,45	- 45,07	0	2,97
	6	0 35,91	+ 1,33	- 43,92	0	4,12
	7	0 36,21	+ 1,20	- 43,28	0	4,93
	9	0 32,87	+ 1,74	- 42,00	0	3,41
	10	0 34,45	+ 1,91	- 41,37	0	5,79
	11	0 32,15	+ 1,84	- 40,72	0	4,07
	12	0 29,91	+ 1,99	- 40,06	0	2,64
13	0 29,19	+ 2,06	- 39,38	0	2,67	
15	0 28,77	+ 1,85	+ 8,67	- 37,98	0	0,63	
16	0 31,27	+ 1,73	- 37,28	0	3,71	

POLARIS AT SUPERIOR CULMINATION.							
1832	Observed A.R. correct- ed for Error of Clock.	Correction for			Aberration, &c.	Mean A.R. 1832.	
		Level.	Azimuth.	Collimation.			
	1h. m. s.	s.	s.	s.	s.	1h. m. s.	
December	17	0 28,73	+ 1,78	+ 8,67	- 0,68	- 36,59	0 1,92
	19	0 28,95	+ 1,73	- 35,14	0 3,53
	20	0 25,12	+ 1,89	- 34,42	0 0,58
	21	0 26,58	+ 1,52	- 33,71	0 2,38
	22	0 24,66	+ 1,58	- 32,96	0 1,27
	24	0 22,35	+ 1,67	- 31,46	0 0,55
	25	0 24,45	+ 1,65	- 30,71	0 3,38
	26	0 25,08	+ 1,59	- 29,96	0 4,70
	27	0 24,97	+ 1,52	- 29,19	0 5,29
1833							
January	2	0 17,00	+ 2,11	+ 10,28	- 1,23	- 8,74	0 19,42
	3	0 16,56	+ 2,12	- 7,94	0 19,79
	4	0 13,72	+ 2,14	- 7,14	0 17,67
	5	0 11,63	+ 1,98	- 6,34	0 16,32
	6	0 13,58	+ 1,83	- 5,55	0 18,91
	7	0 12,76	+ 1,66	- 4,75	0 18,72
	8	0 13,67	+ 1,67	- 3,96	0 20,43
	10	0 3,31	+ 1,92	+ 2,88	- 2,38	0 16,01
	11	0 4,34	+ 2,04	- 1,59	0 17,95
	15	0 3,72	+ 2,44	+ 1,60	0 20,92
16	0 3,65	+ 3,44	+ 2,38	0 22,63	
17	59 54,59	+ 4,42	+ 6,90	+ 3,17	0 19,36	
June	27	0 10,11	+ 5,55	+ 11,19	- 1,76	- 1,20	0 23,89
	28	0 8,16	+ 5,48	- 1,97	0 21,10
July	7	0 19,17	+ 4,86	+ 12,07	- 8,90	0 25,44
	12	0 17,92	+ 4,77	- 12,73	0 20,27
	14	0 19,37	+ 4,73	- 14,23	0 20,18
November	19	0 57,88	+ 3,89	+ 7,32	- 5,13	- 46,03	0 17,93
	20	0 57,92	+ 3,85	- 45,55	0 18,41
	23	0 54,22	+ 3,85	- 44,01	0 16,25
	27	0 54,33	+ 4,34	+ 4,86	- 3,63	- 42,35	0 17,55
December	2	0 50,50	+ 4,64	- 39,28	0 17,09
	3	0 50,84	+ 4,63	- 38,68	0 18,02
	4	0 51,06	+ 4,66	- 38,08	0 18,87
	5	0 49,10	+ 4,63	- 37,47	0 17,49
	6	0 50,37	+ 4,59	- 36,85	0 19,34
	7	0 47,94	+ 4,58	- 36,21	0 17,54
	8	0 48,49	+ 4,18	- 35,58	0 18,32
	9	0 49,82	+ 3,86	- 34,95	0 19,96
	10	0 48,15	+ 4,34	- 34,31	0 19,41
	11	0 47,77	+ 4,87	- 33,64	0 20,23
	12	0 49,40	+ 4,71	- 32,97	0 22,37
	14	0 44,95	+ 4,79	- 31,63	0 19,35
	18	0 39,54	+ 4,63	- 28,76	0 16,64
	19	0 42,32	+ 4,60	- 28,04	0 20,11
	22	0 37,87	+ 4,25	- 25,86	0 17,44
23	0 38,94	+ 4,34	- 25,13	0 19,34	
24	0 47,22	+ 5,52	+ 2,38	- 7,99	- 24,38	0 22,75	
25	0 38,77	+ 6,70	+ 3,32	- 4,62	- 23,62	0 20,55	
26	0 34,53	+ 6,41	- 22,85	0 16,79	
27	0 36,13	+ 6,11	- 22,19	0 18,75	
29	0 33,03	+ 5,81	- 20,52	0 17,02	
30	0 35,42	+ 5,67	- 19,74	0 20,05	
31	0 34,22	+ 5,52	- 18,95	0 19,49	

ERROR OF AZIMUTH.

POLARIS AT INFERIOR CULMINATION.									
1832	Observed A.R. correct- ed for Error of Clock.	Correction for			Aberration, &c.	Mean A.R. 1832.			
		Level.	Azimuth.	Collimation.					
	12h.					12h.			
	m. s.	s.	s.	s.	s.	m. s.			
February	13	59 31 02	+ 1,75	+ 51,62	- 41,30	+ 15,74	59 58,83		
	22	59 31,04	+ 1,33	+ 21,20	0 3,89		
March	11	59 21,78	+ 0,86	+ 48,95	+ 29,30	59 59,09		
	14	59 20,84	+ 1,04	+ 30,24	59 59,77		
	20	59 21,17	+ 1 30	+ 31,74	0 1,86		
	28	59 18,08	+ 0,86	- 38,50	+ 32,90	0 2 29		
	29	59 17,85	+ 0,90	+ 32,96	0 2,16		
	30	59 17 36	+ 0,94	+ 33,03	0 1,78		
	31	59 16 49	+ 1,03	+ 33,09	0 1,06		
April	1	59 11,55	+ 1,12	- 34,30	+ 33,14	0 0,46		
	2	59 6,99	+ 1,08	- 31,06	+ 33,13	59 59,09		
	4	59 5,76	+ 1,05	- 30,33	+ 33,12	59 58,55		
	5	59 7,39	+ 1,05	- 32 90	+ 33,11	59 57,60		
	6	59 9,38	+ 1,15	- 35,51	+ 33,08	59 57,05		
	7	59 10 72	+ 1,25	+ 33,04	59 58,45		
	10	59 18,07	+ 0,99	- 36,72	+ 32,89	0 4,18		
	11	59 18,05	+ 1,03	- 36 87	+ 32,76	0 3,92		
	12	59 17,22	+ 1,16	- 36,21	+ 32,62	0 3 74		
	13	59 18,96	+ 1,29	- 35,51	+ 32,46	0 6,15		
	14	59 15,01	+ 1,31	+ 32,27	0 2,03		
	15	59 15,86	+ 1,33	+ 32 07	0 2,70		
	17	59 17,84	+ 1,02	+ 46,73	+ 31,69	0 1,77		
	18	59 17 70	+ 1,03	+ 31,49	0 1,44		
	19	59 17,14	+ 1,05	+ 31,27	0 0,68		
	21	59 16,50	+ 0,91	+ 30,80	59 59,43		
	26	59 21,02	+ 0,73	+ 29,35	0 2,32		
	27	59 21,51	+ 0,90	+ 29,00	0 2,63		
	30	59 19 99	+ 0,80	+ 27,93	59 59,92		
May	1	59 20,99	+ 1 08	+ 27,58	0 0,87		
	3	59 22 75	+ 1,25	+ 26 68	0 1,90		
	11	59 25,42	+ 0,68	+ 23,06	0 0,40		
	12	59 26,02	+ 0,90	+ 22,57	0 0,71		
	14	59 35,08	+ 1,04	+ 47,82	- 41,16	+ 22,06	0 4 84		
	15	59 33 48	+ 0,90	+ 21 03	0 2,07		
	16	59 36,85	+ 0,79	+ 20,48	0 4,78		
	17	59 34 60	+ 0,68	+ 19,92	0 1,86		
	18	59 33,13	+ 0 76	+ 19 35	59 59 90		
	19	59 38 13	+ 0,84	- 42,76	+ 18,78	0 2,81		
	20	59 38 66	+ 0,92	- 42 85	+ 18,19	0 2,74		
	22	59 33,58	+ 0,71	- 41,16	+ 16,68	59 57,93		
	24	59 34,28	+ 0,78	+ 15,76	59 57,48		
	26	59 33,35	+ 1,12	- 39,75	+ 14,49	59 57,03		
	27	59 37,69	+ 1,02	- 41,16	+ 13 68	59 59,22		
	28	59 36,21	+ 0,88	+ 13,21	59 56,96		
	29	59 37,45	+ 0,76	+ 12,87	59 57,44		
	30	59 36,50	+ 0,76	+ 11,92	59 55,84		
	31	59 37,60	+ 0,76	+ 11,27	59 56,29		
June	5	59 48,05	- 0,02	+ 7,68	0 0,19		
	9	59 49,27	- 0,14	+ 4,80	59 58,33		
	10	59 52,53	- 0,14	- 43,41	+ 4,07	0 0,87		
	11	59 57,32	- 0,26	+ 3,34	0 4,81		
	15	59 59,68	- 0,01	+ 48,85	+ 0,40	0 5,51		
December	7	0 59 06	- 1,40	- 11,46	+ 0,68	- 42,96	0 4,92		
	8	0 56,23	- 1,50	- 42,32	0 1,63		
	9	0 52,79	- 1,49	- 41,69	59 59,93		

POLARIS AT INFERIOR CULMINATION.								
1832	Observed A.R. correct- ed for Error of Clock.	Correction for			Aberration, &c.	Mean A.R. 1832.		
		Level.	Azimuth.	Collimation.				
		13h.				12h.		
		m.	s.	s.	s.	m.	s.	
December	12	0	50,27	- 1,56	- 11,46	+ 0,68	- 39,72	59 58 21
	13	0	49,08	- 1,60	- 39,02	59 57,68
	14	0	51,80	- 1,53	- 38,33	0 1,46
	15	0	49,24	- 1,45	- 8,64	- 37,63	0 2,20
	16	0	49,23	- 1,36	- 36,93	0 2,92
	18	0	47,46	- 1,45	- 35,50	0 2 55
	20	0	44,20	- 1,28	- 34,06	0 0,90
	23	0	47,09	- 1,40	- 31,86	0 5,87
	24	0	42,00	- 1,36	- 31,11	0 1,57
	26	0	41,49	- 1,30	- 29,60	0 2,63
1833								
January	2	0	35,39	- 1,66	- 10,22	+ 1,23	- 8,34	0 12,94
	3	0	36,22	- 1,67	- 7,54	0 14,55
	4	0	34,57	- 1,69	- 6,74	0 17,15
	5	0	30,76	- 1,55	- 5,95	0 14,26
	6	0	35,02	- 1,37	- 5,15	0 19,51
	8	0	30,21	- 1,31	- 3,57	0 16,34
	10	0	35,47	- 1,45	- 2,88	- 1,99	0 18,93
	11	0	34,77	- 1,55	- 1,19	0 18,93
	13	0	26,66	- 1,35	+ 0,40	0 12,61
	14	0	27,25	- 1,17	+ 1,20	0 14,18
	15	0	35,22	- 1,96	- 6,90	+ 1,99	0 18,83
	16	0	30,26	- 2,75	+ 2,78	0 13,17
	17	0	33,42	- 3,54	+ 3,57	0 16,33
November	19	1	6,99	- 3,05	- 7,24	+ 5,13	- 45,79	0 16,04
	20	1	9,37	- 3,01	- 45,31	0 18,94
	21	1	7,54	- 2,92	- 44,83	0 17,68
	22	1	6,66	- 2,81	- 44,35	0 17,39
	23	1	7,81	- 3,02	- 43,86	0 18,82
	28	1	7,25	- 3,49	- 4,81	+ 3,63	- 41,78	0 20,80
December	1	1	3,92	- 3,60	- 39,58	0 19,56
	5	1	3,26	- 3,70	- 37,17	0 21,21
	6	0	58,53	- 3,02	- 36,54	0 17,19
	8	0	59,56	- 3,30	- 35,27	0 19,81
	9	0	58,41	- 3,05	- 34,64	0 19,44
	10	0	56,08	- 3,45	- 33,79	0 17,48
	11	0	57,41	- 3,82	- 33,30	0 19,11
	12	0	55,47	- 3,69	- 32,63	0 17,97
	13	0	53,10	- 3,57	- 31,96	0 16,39
	18	0	49,06	- 3,63	- 28,43	0 15,82
	24	0	44,01	- 4,35	- 2,60	+ 7,43	- 24,00	0 20,89
	25	0	46,57	- 5,27	- 3,29	+ 4,62	- 23,24	0 19,39
	26	0	47,25	- 5,08	- 22,49	0 21,01
	27	0	44,91	- 4,84	- 21,70	0 19,70
	28	0	42,65	- 4,74	- 20,92	0 18,32
	30	0	39,58	- 4,49	- 19,35	0 17,07

ERROR OF AZIMUTH.

δ URSAE MINORIS AT SUPERIOR CULMINATION.									
1832		Observed A.R. correct- ed for Error of Clock.		Correction for			Aberration, &c.	Mean A.R. 1832.	
				Level.	Azimuth.	Collimation.			
		18h.						18h.	
		m.	s.	s.	s.	s.	s.	m.	s.
March	29	26	37,13	— 0,59	— 22,42	+ 17,88	— 2,50	26	29,50
	31	26	40,17	— 0,68	— 3,21	26	31,74
April	9	26	42,11	— 0,72	— 6,37	26	30,48
August	27	26	34,63	— 2,73	+ 7,09	— 2,49	— 5,37	26	31,13
	31	26	32,15	— 2,47	— 3,89	26	30,39
September	4	26	29,16	— 2,30	+ 7,98	— 2,31	26	30,04
	5	26	30,16	— 2,56	— 1,91	26	31,18
	9	26	29,77	— 2,60	— 4,38	— 0,33	26	30,44
	10	26	28,75	— 2,64	— 3,52	+ 0,07	26	30,64
	11	26	28,73	— 2,63	+ 0,46	26	31,03
	22	26	25,27	— 2,76	+ 4,98	26	31,95
	23	26	24,35	— 2,68	+ 5,40	26	31,53
	26	26	21,89	— 2,64	+ 6,66	26	30,36
	27	26	21,32	— 2,68	+ 7,08	26	30,18

δ URSAE MINORIS AT INFERIOR CULMINATION.

δ URSAE MINORIS AT INFERIOR CULMINATION.									
1832		Observed A.R. correct- ed for Error of Clock.		Correction for			Aberration, &c.	Mean A.R. 1832.	
				Level.	Azimuth.	Collimation.			
		6h.						6h.	
		m.	s.	s.	s.	s.	s.	m.	s.
January	4	25	55,71	+ 0,65	+ 23,86	— 6,99	+ 17,23	26	30,46
	12	25	55,52	+ 0,60	— 8,90	+ 17,19	26	28,27
	14	25	54,80	+ 0,62	+ 17,03	26	27,41
	15	25	55,16	+ 0,63	+ 16,94	26	27,69
	24	26	7,00	+ 0,70	+ 24,38	— 19,10	+ 16,01	26	28,99
	29	26	7,70	+ 0,80	+ 15,29	26	29,07
February	11	26	9,69	+ 0,67	+ 12,65	26	29,31
	13	26	10,60	+ 0,67	+ 12,15	26	28,70
	14	26	7,88	+ 0,69	+ 11,90	26	25,75
	15	26	10,61	+ 0,70	+ 11,64	26	28,23
	18	26	14,48	+ 0,46	+ 10,83	26	30,59
	19	26	11,37	+ 0,49	+ 10,55	26	27,69
	20	26	10,24	+ 0,52	+ 10,27	26	26,31
	22	26	12,53	+ 0,51	+ 9,71	26	28,03
	24	26	10,54	+ 0,45	+ 9,13	26	26,40
	25	26	11,84	+ 0,48	+ 8,82	26	26,42
	26	26	14,11	+ 0,53	+ 8,51	26	28,97
	28	26	17,79	+ 0,56	+ 23,10	— 19,88	+ 7,88	26	29,45
March	1	26	17,54	+ 0,61	+ 7,24	26	28,61
	2	26	17,20	+ 0,65	+ 6,91	26	27,98
	3	26	19,84	+ 0,59	+ 6,58	26	30,23
	4	26	18,03	+ 0,54	+ 6,24	26	28,03
	5	26	18,11	+ 0,56	+ 5,90	26	27,78
	13	26	21,96	+ 0,51	+ 3,19	26	28,88
	14	26	22,45	+ 0,45	+ 2,87	26	28,99
	15	26	22,26	+ 0,49	+ 2,55	26	28,52
	17	26	22,38	+ 0,51	+ 1,89	26	28,00
	19	26	23,94	+ 0,48	+ 1,11	26	28,75
	20	26	22,93	+ 0,51	+ 0,73	26	27,39
	21	26	24,47	+ 0,51	+ 0,34	26	28,54

δ URSE MINORIS AT INFERIOR CULMINATION.								
1833	Observed A.R. correct- ed for Error of Clock.	Correction for			Aberration, &c.	Mean A.R. 1833.		
		Level.	Azimuth.	Collimation.				
		6h.				6h.		
		m.	s.	s.	s.	m.	s.	
January	12	26	1,87	— 0,55	— 4,84	— 1,33	+ 16,17	26 11,32
	15	26	0,76	— 0,76	— 1,33	+ 15,97	26 9,80
	16	26	1,42	— 1,06	— 3,18	+ 15,90	26 8,24
	29	26	9,32	— 1,60	— 6,68	+ 14,30	26 10,75
	31	26	8,32	— 1,73	+ 13,94	26 9,01
February	1	26	9,68	— 1,67	+ 13,76	26 10,25
	3	26	8,26	— 1,64	+ 13,39	26 8,49
	9	26	9,24	— 1,78	+ 12,11	26 8,05
	11	26	9,86	— 1,80	+ 11,83	26 8,17
	13	26	12,27	— 1,82	— 5,22	+ 11,14	26 9,69
	28	26	16,47	— 1,55	+ 6,80	26 9,52
March	3	26	16,16	— 1,94	+ 5,81	26 8,13
	6	26	20,43	— 1,94	+ 4,81	26 11,40
	7	26	18,01	— 2,30	+ 4,48	26 8,39
	8	26	16,66	— 1,67	+ 4,15	26 8,24
	9	26	16,25	— 1,79	+ 3,82	26 6,37
	10	26	18,45	— 1,75	+ 3,48	26 8,28
	11	26	19,44	— 1,71	— 4,72	+ 3,14	26 10,93
	13	26	17,52	— 1,70	+ 2,45	26 8,33
	14	26	17,54	— 1,75	+ 2,11	26 7,96
	15	26	17,64	— 1,78	+ 1,76	26 7,58
	16	26	18,89	— 1,80	+ 1,41	26 8,56
	17	26	18,00	— 1,82	+ 1,06	26 7,30
	18	26	18,97	— 1,84	+ 0,71	26 7,90
	19	26	18,87	— 1,80	+ 0,36	26 7,49
	21	26	18,64	— 1,83	— 0,35	26 6,52
	22	26	20,81	— 1,90	— 0,71	26 8,26
	23	26	20,24	— 2,00	— 1,06	26 7,14
	25	26	22,17	— 2,02	— 1,78	26 8,43

76 DRACONIS AT SUPERIOR CULMINATION.

1832	Observed A.R. correct- ed for Error of Clock.	Correction for			Aberration, &c.	Mean A.R. 1832.		
		Level.	Azimuth.	Collimation.				
		20h.				20h.		
		m.	s.	s.	s.	m.	s.	
September	19	54	22,03	— 1,43	+ 3,28	— 1,30	— 6,20	54 16,38
	20	54	21,21	— 1,43	— 1,23	— 6,08	54 15,75
	22	54	20,98	— 1,49	— 5,82	54 15,72
	24	54	20,32	— 1,40	— 5,56	54 15,41
	25	54	19,95	— 1,43	— 5,44	54 15,13
								1833
October	21	54	9,80	+ 1,27	+ 2,99	— 0,86	— 0,76	54 12,44
	22	54	9,99	+ 1,28	— 0,60	54 12,80
	23	54	9,60	+ 1,19	— 0,44	54 12,48
	25	54	8,92	+ 1,18	— 0,11	54 12,12
	30	54	9,47	+ 1,09	+ 0,72	54 13,21
November	6	54	6,87	+ 0,57	+ 1,00	54 11,47

76 DRAGONIS AT INFERIOR CULMINATION.								
1833	Observed A.R. correct- ed for Error of Clock.	Correction for			Aberration, &c.	Mean A.R. 1833.		
		Level.	Azimuth.	Collimation.				
		8h.				6h.		
		m.	s.	s.	s.	m.	s.	
February	23	54	8,60	— 0,31	— 2,22	— 0,51	+ 6,88	54 12,44
March	11	54	10,73	— 0,31	— 1,45	+ 5,72	54 12,47
	12	54	8,49	— 0,28	+ 0,27	+ 5,63	54 11,89
	17	54	8,29	— 0,31	+ 5,12	54 11,15
	18	54	8,76	— 0,32	+ 5,00	54 11,49
	19	54	9,27	— 0,31	+ 4,89	54 12,90
April	25	54	11,53	— 0,36	— 1,45	+ 4,22	54 11,72
	21	54	15,55	— 0,43	+ 0,48	54 11,93

Taking the means, and applying to the observations of 1833, the Annual variations to reduce them to the beginning of 1832, we have:

		Mean A.R. Jan. 1, 1832.		
		h.	m.	s.
I	POLARIS.....S.P.	52 Observations in the first six months of 1832...	13	0 0,86
II	Ann. Var. 17",86	{ 26 Observations towards the end of 1832, and in January 1833.	13	0 0,87
III	Ann. Var. 17",86	{ 22 Observations in the months of November and December 1833.	13	0 2,85
IV	POLARIS.....	{ 62 Observations in November and December 1832, and in January 1833.	1	0 2,93
V	Ann. Var. 3",72	5 Observations in June and July 1833.....	1	0 6,39
VI	Ann. Var. 3",72	{ 27 Observations in the months of November and December 1833.	1	0 3,00
VII	URSAE MIN. S.P.	30 Observations in the three first months of 1832...	6 26	28,35
VIII	Ann. Var. 19",19	{ 29 Observations in the three first months of 1833.	6 26	27,83
IX	URSAE MIN....	14 Observations towards the middle of 1832.....	18 26	30,75
X	76 DRAGONIS S.P.	{ 8 Observations in March 1833.....	8 54	15,72
XI	76 DRAGONIS....	5 Observations in September 1832.....	20 54	15,68
XII	Ann. Var. 3",72	6 Observations in October and November 1833...	20 54	16,17

Examining these results attentively; we notice, from the near agreement of No. I with No. II and of No. VII with No. VIII, that any error of Azimuth affecting the observations at the beginning of 1832, affect equally those towards the end of that year and for the three first months of 1833: let this error be represented by a . No. V shews us that some larger error which we will call a' exists in the months of June and July 1833. To No. XII, or the observations for October and the early part of November 1833, we will assign an error a'' . Finally, comparing No. III with No. VI we find that an

error of Azimuth of no consequence is attached to the observations between the 19th November and the end of the year 1833.

Taking the mean of I and II and of VII and VIII we obtain the following Equations.

$$\begin{aligned}
 s. \quad 0,86 + 2,35 a \pm \frac{e}{\sqrt{n}} &= s. \quad 2,93 - 2,32 a \pm \frac{e}{\sqrt{n^1}} \\
 28,09 + 1,10 a \pm \frac{e^1}{\sqrt{n^{11}}} &= 30,75 - 1,07 a \pm \frac{e^1}{\sqrt{n^{11}}} \\
 15,72 + 0,47 a \pm \frac{e^{11}}{\sqrt{n^{11}}} &= 15,68 - 0,44 a \pm \frac{e^{11}}{\sqrt{n^{11}}}
 \end{aligned}$$

where $e, e^1, \&c.$ represent the probable errors of a single observation and $n, n^1, \&c.$ the number of observation constituting each result. Considering the low altitude at which Stars below the pole are seen in this latitude, it must be expected that the unsteadiness consequent thereto will give rise to large errors of observation; in the case of the Pole Star, I propose to assume the mean error of a single observation to be two seconds; for δ Ursæ Minoris, one second; and for 76 Draconis seven tenth of a second—substituting these values, we determine:

$$\begin{aligned}
 4,67 a = 2,07 \pm ,32 \text{ or } a = 0,44 \pm ,07 \\
 2,17 a = 2,66 \pm ,28 - a = 1,22 \pm ,13 \\
 0,91 a = 0,04 \pm ,40 - a = -0,05 \pm ,44
 \end{aligned}$$

giving to each of these results a weight in the inverse ratio of the probable error, we find $a = 0',63$; hence the Azimuth of the North and South Marks for the year 1832, and for the first 3 months of 1833; (instead of the results found at Page 30) will be N. $35',51$ W. and S. $60',69$ E. Computing now the observations of the Pole Star with these newly found errors of Azimuth and taking the mean we find:

		Mean Right Ascension reduced to January 1, 1832.					
		h. m. s.			h. m. s.		
78	Observations <i>below</i> the Pole in 1832 and in January 1833.....	13	0	2,34	13	0	2,45
22	Observations <i>below</i> the Pole towards the end of 1833.....			2,85			
62	Observations <i>above</i> the Pole in 1832 and in January 1833.....	1	0	1,47	1	0	1,93
27	Observations <i>above</i> the Pole towards the end of 1833.....			3,00			

With the mean of these *1h. Om. 2,19s.* We will now proceed to compute a^1 or, since there are only five observations, it will perhaps be better to compute from these the Azimuth of the North and South Marks as follows:

ERROR OF AZINUTH:

1833	Observed A.R. correct- ed for Clock Error.	Correction for			Mean A.R. January 1, 1832.
		Level.	Collimation.	Aberration, &c.	
	<i>h. m. s.</i>	<i>s.</i>	<i>s.</i>	<i>s.</i>	<i>h. m. s.</i>
June 27	1 0 10,11	+ 5,55	- 1,76	- 1,20	0 59 56,92 + 2,32 a^I
28	1 0 8,16	+ 5,48	- 1,97	0 59 54,13 + 2,32 a^{II}
July 7	1 0 19,17	+ 4,86	- 8,90	0 59 57,79 + 2,32 a^{III}
12	1 0 17,92	+ 4,77	- 12,73	0 59 52,42 + 2,32 a^{IV}
14	1 0 19,37	+ 4,73	- 14,23	0 59 52,33 + 2,32 a^V

	<i>s.</i>			<i>s.</i>
Hence	5,27	=	2,32 a^I	or a^I = 2,27
—	8,06	=	2,32 a^{II}	— a^{II} = 3,47
—	4,40	=	2,32 a^{III}	— a^{III} = 1,90
—	9,77	=	2,32 a^{IV}	— a^{IV} = 4,21
—	9,86	=	2,32 a^V	— a^V = 4,29

employing these in conjunction with the registered variation of the centre wire from the North and South Marks at Page 24, when corrected for Collimation, we obtain the

AZIMUTH OF THE					
North Mark.			South Mark.		
	<i>s.</i>			<i>s.</i>	
N.	36,86	W.	S.	61,93 E.
—	35,60	—	—	60,67 —
—	37,13	—	—	62,29 —
—	35,17	—	—	60,24 —
—	34,77	—	—	59,84 —
Mean N.			35,91 W.	S. 60,98 E.

To compute a^{II} we must now with the value found above for a correct the place of 76 Draconis given at Pages 37 and 38, we have from

	Mean A.R. 1832.		
	<i>h.</i>	<i>m.</i>	<i>s.</i>
5 Observations in September, 1832 above the Pole.....	20	54	15,40
8 Observations in March 1833 below the Pole.....	8	54	16,02

Taking the mean of these and putting it equal to No. XII, we have

$$12 \text{ } \begin{matrix} \text{h.} & \text{m.} & \text{s.} \\ \text{+} & 8 & 54 & 15,71 \end{matrix} = \begin{matrix} \text{h.} & \text{m.} & \text{s.} \\ 20 & 54 & 16,17 \end{matrix} - 0,44 \text{ } a^{II}$$

$$\text{or } a^{II} = 0'',98$$

applying this to the numbers found at Page 32, which it will be recollected

have been employed from the 1st of July 1833; we obtain the Azimuths of the North and South Marks for the month of October, and for the early part of November 1833; and recapitulating, we have found altogether as follows.

	AZIMUTH OF THE MARK			
	to the North.		to the South.	
	s.		s.	
Observations of 1831.....	N. 35,34	W.	S. 62,31	E.
1832, and January, February, and March of 1833....	— 35,51	—	— 60,69	—
June and July 1833.....	— 35,91	—	— 60,98	—
October and the early part of November 1833.....	— 34,77	—	— 59,84	—
November and December 1833.....	— 33,79	—	— 58,86	—

The results of 1832 from the number of observations employed are probably very near the truth; comparing them with those of 1831 it appears probable that the North Mark has remained firm, and that an alteration has taken place in the position of the South Mark to the amount of 1",6. The results for the months of June and July 1833, being determined from five observations of the Pole Star only, cannot be supposed very accurate; their tendency is however to support the values found for 1832. The results of October and the early part of November 1833, determined from five observations of 76 Draconis, possess still less claims to accuracy, their tendency is however rather to support the results of 1832 than those of November and December 1833, these last from the number of observations employed we may presume to be a good determination; from these considerations we are led to conclude, that the position of the North Mark has remained unchanged from the time of the erection of the Transit Instrument (January 1831) up to the early part of November 1833 when it moved 1",4 towards the East = 0',64 Inches; that the South Mark shifted to the Westward towards the end of 1831 or the beginning of 1832 to the amount 1",61, and that a further alteration in the same direction took place to the amount 2",42 towards the beginning of November 1833: to determine the precise date of the former alteration we unfortunately have not a sufficient number of observations, but the uncertainty thereby introduced into one or two months observations (from an error of eight tenths of a second of space) will on inspecting the table shewing the correction for 1" error of Azimuth be found not to exceed 0",14 of time. For the more accurate determination of the date of the latter alteration, we will now examine the daily observations of the Azimuth of the centre wire from the North and South Marks for 1833: to get rid of the error of Collimation with which each of these is affected, we will examine the sum, or N + S; here we find the utmost regularity to exist up to the 26th of September when

an alteration to the amount of two seconds occurs, but in a direction contrary to the alteration for which we desire to account; we must consequently attribute it to some alteration of the Instrument itself: from this time up to the end of the year, one alteration only occurs deserving of notice; it is on the 11th of November, being in amount such as very nearly agrees with the alteration which we found above to have taken place towards the early part of November; I should hesitate to assume this day as the date of the alteration on these grounds were it not that another circumstance (the fall of four or five Inches of rain) which took place at this time, seems to render it probable that the foundation of the buildings forming the North and South Marks may have given way; to satisfy myself on this head, on a late occasion I paid a visit to the South Mark, which is the end of a very substantially built brick dwelling House, situated at about one and a half miles distant from the Observatory; here I found a crack in the wall about 2 Inches wide, which is just sufficient to account for the alterations above found; the North Mark is a square, (brick and chunam) pyramid, of weather beaten and rough exterior, thereby offering no means of detecting a small alteration of the foundation, but from the nature of the soil (a bed of loose sand) it is easy to believe that a considerable fall of rain might have produced the alteration in question; consequently, in computing the corrections for Azimuth for 1832 and for 1833 up to the 12th of November, we must employ the number

$$\frac{96'',20 - N - S}{2}$$

and from the 12th November 1833, to the end of the year

$$\frac{92'',65 - N - S}{2}$$

employing these with the observed Azimuth of the North and South Marks from the centre wire we obtain as follows:

1832	N. — S.	$\frac{96'',20 - N - S}{2}$	REMARKS.	1832	N. — S.	$\frac{96'',20 - N - S}{2}$	REMARKS.
Jan.				Jan.			
1	+ 48,14	+ 24,03		7	+ 54,80	+ 20,70	
2	51,69	22,26		8	51,35	22,42	
3	51,97	22,11		9	51,94	22,13	
5	52,41	21,89		10	54,40	20,90	
6	51,83	22,18		11	52,48	21,86	

ERROR OF AZIMUTH.

1832	N. — S.	96'' 20 — N — S.		REMARKS.	1832	N. — S.	96'' 20 — N — S.		REMARKS.
		s.	s.				s.	s.	
Jan. 12	+ 52,55	+ 21,82			March 15	+ 52,48	+ 21,86		
13	52,48	21,86			16	51,72	22,23		
14	51,12	22,54			17	52,27	21,96		
15	50,84	22,68			18	52,62	21,79		
16	51,70	22,25			19	52,72	21,70		
17	50,89	22,65			20	52,48	21,86		
18	52,31	21,94		Mean of 18 = + 22'',16.	21	52,21	21,99		
19	50,86	22,67		New wires.	22	53,69	21,26		
Feb. 3	53,20	21,50		New wires.	23	53,77	21,21		
4	48,82	23,69			24	54,08	21,06		
5	50,86	22,67			25	53,97	21,11		
6	50,47	22,86			26	52,87	21,67		
7	48,57	23,81			27	54,34	20,93		
8	49,95	23,12			28	53,08	21,58		
9	49,91	23,14			29	52,80	21,70		
10	48,86	23,67			30	53,96	21,12		
11	49,28	23,46			31	52,10	22,00		
12	51,30	22,45			April 1	52,78	21,72		
13	50,88	22,66			2	52,86	21,67		
14	50,82	22,69			3	53,58	21,31		
15	51,37	22,41			4	53,82	21,19		
16	51,26	22,47			5	53,17	21,51		
17	51,33	22,43			6	53,48	21,36		
18	52,29	21,95			7	53,78	21,22		
19	50,39	22,90			8	53,54	21,33		
20	51,13	22,53			9	53,52	21,34		
21	51,65	22,27			10	53,30	20,45		
22	51,41	22,39			11	54,41	20,89		
23	51,57	22,31			12	53,54	21,33		
24	51,57	22,31			13	53,94	21,14		
25	51,69	22,25			14	53,43	21,38		
26	52,17	22,01		Mean of 24 = + 22'',69	15	53,37	21,41		Mean of 49 = + 21'',46
27	52,14	22,03			16	53,77	21,21		
28	53,68	21,26			18	55,34	20,43		
29	53,44	21,38			19	55,59	20,30		
March 1	53,23	21,48			20	55,19	20,50		
2	52,90	21,62			21	55,17	20,51		
3	53,20	21,50			22	55,08	20,56		
4	52,48	21,86			23	55,05	20,57		
5	52,75	21,72			24	54,98	20,61		
6	52,50	21,85			25	54,96	20,62		
7	52,34	21,93			26	55,10	20,55		
8	52,20	22,00			27	55,34	20,43		
9	52,86	21,67			28	55,31	20,44		
10	54,64	20,78			29	55,65	20,27		
11	54,31	20,94			30	55,25	20,47		
12	53,23	21,48			May 1	55,54	20,33		
13	53,16	21,52			2	55,69	20,26		
14	52,85	21,67			3	55,44	20,38		
					4	55,44	20,38		
					5	55,62	20,26		

ERROR OF AZIMUTH.

1832	N. — S.	96'' 20 — N — S.	REMARKS.	1832	N. — S.	96'' 20 — N — S.	REMARKS.		
May	6	+ 55,79	+ 20,20	<p>Mean of 40 = + 20'',56</p> <p>Mean of 22 = + 21'',04</p>	June	28	+ 53,31	+ 21,44	
	7	55,24	20,48			29	53,31	21,44	
	8	55,52	20,34			30	53,21	21,49	
	9	55,09	20,56		July	1	53,44	21,38	
	10	55,57	20,31			2	53,31	21,44	
	11	55,73	20,23			3	53,06	21,57	
	12	54,75	20,72			4	53,55	21,32	
	13	53,92	21,14			5	53,62	21,29	
	14	54,06	21,07			6	53,37	21,41	
	15	54,30	20,95			7	52,55	21,82	
	16	53,88	21,16			8	53,19	21,50	
	17	54,22	20,99		<p>Mean of 17 = + 21'',40</p> <p>Adjusted the Instrument.</p>	16	100,26	2,03	
	18	55,89	20,16			17	100,40	2,10	
	19	55,15	20,52			24	100,26	2,03	
	20	55,12	20,54			25	100,57	2,18	
	21	55,16	20,52			27	100,57	2,18	
	22	54,82	20,69			29	100,16	1,98	
	23	54,65	20,77			30	100,57	2,18	
	24	54,37	20,91			31	101,40	2,60	
	25	54,29	20,95			Aug.	1	100,39	2,09
	26	54,47	20,86				7	98,98	1,39
	27	54,20	21,00		<p>Mean of 10 = — 2'',08</p> <p>A new set of lines.</p>	13	103,82	3,81	
	28	54,61	20,79			14	102,26	3,03	
	29	54,45	20,87			15	102,96	3,38	
	30	54,38	20,91			17	102,16	2,98	
	31	54,15	21,02			18	102,54	3,17	
	June	1	54,24			20,98	19	101,71	2,75
2		54,11	21,04			20	102,26	3,03	
3		54,07	21,06			21	101,80	2,80	
4		54,03	21,08			22	101,68	2,74	
5		54,17	21,01			23	101,24	2,52	
6		54,58	20,81			24	101,02	2,41	
7		54,16	21,02		<p>Mean of 11 = — 2'',97</p>	25	107,89	5,84	
8		54,33	20,93			26	108,81	6,30	
9		54,54	20,83			27	109,37	6,58	
10		54,51	20,84			28	109,00	6,40	
11		54,26	20,97			29	108,60	6,20	
12		53,71	21,24			30	108,73	6,26	
13		53,44	21,38			31	107,79	5,79	
14		53,92	21,14			Sept.	1	107,71	5,75
15		53,92	21,14	2			107,22	5,51	
16		54,29	20,95	3			110,43	7,11	
17		54,26	20,97	4	109,88		6,84		
18		53,48	21,36	5	109,16		6,48		
19		54,20	21,00	6	109,46		6,63		
20		53,10	21,55	7	109,57		6,68		
21		53,95	21,12	8	109,92		6,86		
22		54,17	21,01	9	110,05		6,92		
23	53,71	21,24							
24	53,24	21,48							
25	53,24	21,48							
26	53,68	21,26							
27									

1882		N. — S.	96',30 — N — S. s.	REMARKS.	1882		N. — S.	96',30 — N — S. s.	REMARKS.
Sept.	13	+109,68	6,74		Nov.	13	+107,95	5,87	
	20	110,19	6,99			14	108,26	6,03	
	21	110,22	7,01			15	107,78	5,79	Mean of 8 = — 5",99
	22	110,50	7,15			16	107,56	5,68	
	23	109,95	6,87			17	104,58	4,19	
	24	109,44	6,62			18	104,30	4,05	
	25	109,44	6,62			19	105,45	4,62	
	26	109,88	6,84			20	105,86	4,83	
	27	110,29	7,04			21	106,46	5,13	
	28	110,50	7,15			22	105,00	4,40	
29	110,44	7,12		23	105,07	4,43			
Oct.	1	110,22	7,01		24	104,82	4,31		
	3	110,76	7,28		25	104,65	4,22		
	6	109,70	6,75		26	106,30	4,55		
	7	110,42	7,11		27	105,21	4,50		
	8	110,25	7,02		28	105,48	4,64		
	9	110,39	7,09		29	105,20	4,50		
	10	110,30	7,05		30	104,82	4,24		
	11	109,83	6,81		Dec.	1	104,82	4,24	
	12	110,08	6,94			2	105,28	4,54	
	13	109,94	6,87			3	106,14	4,97	
14	110,04	6,92	Mean of 31 = — 6",90	4		105,45	4,62		
17	109,88	6,84		5		104,82	4,31		
				6		104,59	4,19		
19	111,95	7,87	Inverted the axis several times.	7		104,65	4,22		
20	112,15	7,97		8		103,77	3,78		
21	112,10	7,95		9		103,91	3,85		
22	112,35	8,07	Mean of 5 = — 7",92	10		104,17	3,98		
23	111,71	7,75		11	104,63	4,21			
				12	104,92	4,36			
24	110,47	7,13		13	104,92	4,36	Mean of 28: = — 4",38		
25	110,57	7,18		14	104,82	4,31			
26	110,38	7,09							
27	110,12	6,96		15	102,06	2,93			
28	109,81	6,80		16	101,44	2,62			
29	109,70	6,75		17	103,36	3,58			
30	110,66	7,23		18	103,43	3,61			
31	110,49	7,14		19	102,94	3,37			
Nov.	1	110,25	7,02		20	102,99	3,39		
	2	109,39	6,59		21	103,05	3,42		
	3	110,04	6,92		22	102,32	3,06		
	4	109,77	6,78		23	102,40	3,14		
	5	109,35	6,58		24	101,84	2,82		
	6	109,80	6,80		25	102,24	3,02		
	7	109,26	6,53	Mean of 16 = — 6",89	26	101,51	2,66		
	8	109,80	6,80		27	101,63	2,71		
					28	102,26	3,03		
	9	108,56	6,18	A new set of lines put in.	29	102,93	3,36		
10	108,53	6,16		30	102,98	3,39	Mean of 17 = — 3",16		
11	108,24	6,02		31	103,56	3,68			
12	108,60	6,20							

ERROR OF AZIMUTH.

1833	N. — S.	96° 20' — N — S.		REMARKS.	1833	N. — S.	96° 20' — N — S.		REMARKS.	
		s.	s.				s.	s.		
Jan.	1	+103,45	3,62	Inverted the axis.	Feb.	19	+102,87	3,54		
	2	102,73	3,26			20	102,69	3,24		
	3	102,30	3,05			21	103,41	3,60		
	4	102,43	3,11			22	104,10	3,95		
	5	102,84	3,32			23	103,65	3,73		
	6	102,70	3,25			24	103,61	3,70		
	7	102,73	3,26			25	104,19	3,99		
	8	102,74	3,27			26	103,75	3,77		
	9	102,56	3,18			27	103,41	3,60		
	10	103,05	3,42			28	103,75	3,77		
	11	102,67	3,23			March	1	103,10		3,45
	12	103,11	3,45				2	103,41		3,60
	13	101,38	2,59				3	103,78		3,79
	14	101,28	2,54				4	104,29		4,04
15	102,24	3,02	5	103,48	3,64					
17	101,18	2,49	6	103,38	3,59					
18	103,86	3,83	7	103,96	3,88					
19	103,63	3,71	8	104,75	4,28					
20	104,04	3,92	9	102,62	3,21					
21	104,16	3,98	10	103,24	3,52					
22	103,79	3,79	11	103,44	3,62					
23	104,26	4,03	12	103,27	3,53					
24	103,20	3,50	13	103,33	3,56					
			14	103,45	3,62					
			15	102,83	3,31					
			16	104,07	3,93					
			17	104,89	4,34					
			18	103,37	3,58					
			19	103,51	3,65					
			20	102,96	3,38					
			21	103,47	3,63					
			22	103,82	3,81					
			23	103,54	3,67					
			24	102,58	3,19					
			25	102,79	3,29					
			26	102,41	3,10					
			27	103,34	3,57					
			28	103,13	3,46					
			29	103,20	3,50					
			30	103,34	3,57					
			31	103,93	3,86					
			April	1	103,17	3,48				
				2	103,41	3,60				
				3	103,34	3,57				
				4	103,92	3,86				
				5	103,62	3,71				
				6	103,76	3,78				
				7	103,97	3,88				
				8	103,17	3,49				
				9	103,48	3,64				
				10	103,51	3,63				
				11	104,34	4,07				

Inverted the axis.

Mean of 14
= — 3",18

Inverted the axis.

Inverted the axis.

Mean of 7
= — 3",82

Mean of 12
= — 3",00

ERROR OF AZIMUTH.

1833			REMARKS	1833			REMARKS
N. — S.	96'' 20 — N — S.	2		N. — S.	96'' 20 — N — S.	2	
April 12	+103,41	3,60		June 1	+103,18	3,49	
13	102,92	3,36		2	103,04	3,42	
14	103,17	3,48		3	102,97	3,38	
15	103,31	3,55		4	103,01	3,40	
16	103,34	3,57		5	103,07	3,43	
17	103,24	3,52		6	103,07	3,43	
18	102,99	3,39		7	102,32	3,06	
19	103,02	3,41		8	103,17	3,48	
20	103,03	3,41		9	102,66	3,23	
21	102,86	3,33		10	102,95	3,37	
22	103,55	3,67		11	102,98	3,39	
23	103,16	3,48		12	102,94	3,37	
24	103,27	3,53		13	102,90	3,35	Mean of 25
25	103,69	3,74		14	103,21	3,50	= — 3'',42
26	103,14	3,47					
27	103,62	3,71		19	105,19	4,49	
28	103,34	3,57		20	105,03	4,41	
29	103,31	3,55		22	104,86	4,33	
30	103,62	3,71		23	104,86	4,33	Mean of 5
May 1	102,10	2,95		25	104,21	4,00	= — 4'',31
2	102,93	3,36					
3	102,75	3,28		26	103,76	3,78	
4	103,03	3,41		27	103,34	3,57	
5	103,52	3,66		28	103,21	3,50	
6	103,10	3,45		July 1	103,10	3,45	
7	102,75	3,27		2	103,50	3,65	
8	103,21	3,50	Mean of 94	3	103,31	3,55	
9	103,16	3,48	= — 3'',56	4	103,35	3,57	
10	102,83	3,31	I applied fresh varnish to the wires.	5	103,33	3,56	
				6	103,29	3,54	
11	103,55	3,67		7	103,14	3,47	
12	104,01	3,90		8	103,24	3,52	
13	103,97	3,88		9	103,15	3,47	
14	103,70	3,75		10	103,21	3,50	
15	103,49	3,64		11	103,07	3,43	
16	104,44	4,12		12	103,83	3,81	
17	104,10	3,95		13	103,27	3,53	
18	104,64	4,22		14	103,20	3,50	
19	106,07	4,93	Mean of 10	15	103,04	3,42	
20	105,16	4,48	= — 4'',05	16	102,05	2,92	
				17	102,66	3,23	
21	103,41	3,60		18	102,12	2,96	
22	102,93	3,36		19	102,22	3,01	
23	102,93	3,36		20	102,90	3,35	
24	103,27	3,53		22	102,93	3,37	
25	103,27	3,53		23	103,36	3,58	
26	102,97	3,38		24	102,96	3,38	
27	102,93	3,36		25	102,77	3,29	
28	103,14	3,47		26	102,83	3,31	
29	103,45	3,62		28	102,66	3,23	
30	103,17	3,48		30	102,25	3,02	
31	103,45	3,62		31	101,95	2,87	

ERROR OF AZIMUTH.

1833	N. — S.	96',20 — N — S. 2	REMARKS.	1833	N. — S.	96',20 — N — S. 2	REMARKS.
Aug.	1 + 102,05	2,92		Sept.	28 + 107,17	5,48	
	2 101,77	2,78			29 107,58	5,69	
	3 101,26	2,53			30 106,70	5,25	
	4 102,11	2,95		Oct.	1 107,41	5,60	
	5 102,66	3,23			3 107,79	5,80	
	6 102,07	2,93			4 107,23	5 51	
	7 102,55	3,17			5 107,24	5,52	
	8 101,56	2,68			6 107,17	5,49	
	9 101,33	2,57			7 107,17	5,49	
	10 101,76	2,78			8 107,23	5,51	
	11 102,08	2,94			9 107,25	5,52	
	12 101,83	2,81			11 107,27	5,53	
	13 102,26	3,03			13 106,97	5,38	
	14 101,54	2,67			14 106,19	5,00	
	15 101,64	2,72			15 106,29	5,04	
	16 101,33	2,57			16 106,12	4,96	
	17 101,14	2,47			17 106,46	5,13	
	19 101,76	2,78			18 106,09	4,94	
	20 101,73	2,77			21 106,77	5,28	
	21 102,70	3,25			22 106,64	5,22	
	22 101,98	2,89			23 106,92	5,36	
	23 101,73	2,77			24 106,98	5,39	
	24 102,57	3,18	Mean of 55 = — 3",18		25 105,53	4,67	
	25 102,50	3,15		Nov.	1 105,19	4,50	
	26 104,50	4,15			3 105,88	4,84	
	27 103,88	3,84			5 105,81	4,81	
	28 106,12	4,96			7 106,77	5,28	Mean of 29 = — 5",24
	30 105,45	4,62			9 105,57	4,68	
	31 105,09	4,44			15 100 53	3,94	
Sept.	1 105,38	4,59			16 100,77	4 06	
	2 105,77	4 78			17 100,55	3,95	
	8 105,31	4,55			18 99,26	3,30	
	9 105 93	4,86			19 99,20	3,27	
	10 105,52	4,66			20 99,47	3,41	
	11 106,21	5,00			21 98,71	3,03	
	12 104,90	4 35			22 99,33	3,34	
	13 104,40	4,10			23 98,54	2,94	
	14 104,83	4 31			24 98,82	3,08	
	15 104,54	4,17			25 98,47	2,91	
	16 104,72	4,26			26 98,20	2,77	Mean of 13 = — 3",28
	17 104,95	4,37			27 97,95	2,65	
	18 104,71	4,25			28 96,95	2,15	
	19 104,69	4,24			29 96,68	2,01	
	20 105,26	4,53			30 96,88	2,12	
	21 105,09	4,44		Dec.	1 97,27	2,31	
	23 105,92	4,86			2 96,71	2,03	
	24 104,92	4,36			3 96,95	2,15	
	25 104,73	4,26	Mean of 25 = — 4",45		4 96,51	1,93	
	26 104,70	4,25			5 96,74	2,04	
	27 106,24	5,02			6 96,92	2,13	

1833	N. — S.	92'',65 — N — S.	REMARKS.	1833	N. — S.	92'',65 — N — S.	REMARKS.
Dec. 7	+ 97,34	— 2,34		Dec. 19	+ 96,68	— 2,01	Mean of 23
8	97,34	2,34		20	95,93	1,64	= — 2'',43
9	97 27	2,31		24	94,69	1,02	Inverted the axis.
10	96,10	1,72		25	94,76	1,05	
11	95,89	1,62		26	96,58	1,96	Inverted the axis.
12	96,03	1,69		27	95,21	1,28	
13	96 83	2,09		28	95,45	1,40	
14	96,33	1,84		29	95,45	1,40	
15	96,57	1,96		30	95,02	1,18	Mean of 8
16	96,89	2,12		31	94,90	1,12	= — 1'',30
17	97,85	2,60					
18	96,33	1,84					

The following table exhibits the amount of error caused by an uncertainty of one second in the position of the Instrument for *unreduced* observations ; and the amount of error *after reduction*, or the error which may be expected to attach to the places of the Sun, Moon, Planets, and fixed Stars which hereafter follow, in case an error of Azimuth to this amount has been committed.

North Polar Distance of the Object.	Correction for 1s. Error of Azimuth.	Error of the computed Result.
20	+ 0,1634	+ 0,178
25	+ 0,1245	+ 0,140
30	+ 0,0974	+ 0,112
35	+ 0,0776	+ 0,093
40	+ 0,0623	+ 0,077
45	+ 0,0499	+ 0,065
50	+ 0,0394	+ 0,054
55	+ 0,0304	+ 0,045
60	+ 0,0224	+ 0,037
65	+ 0,0152	+ 0,030
70	+ 0,0084	+ 0,023
75	+ 0,0023	+ 0,017
80	— 0,0036	+ 0,011
85	— 0,0095	+ 0,006
90	— 0,0151	0,000
95	— 0,0208	— 0,006
100	— 0,0265	— 0,011
105	— 0,0324	— 0,017
110	— 0,0387	— 0,024

North Polar Distance of the Object.		Correction for 1s. Error of Azimuth,		Error of the computed Result.
		"		"
115	— 0,0455	— 0,030
120	— 0,0526	— 0,037
125	— 0,0609	— 0,046
130	— 0,0696	— 0,055
135	— 0,0801	— 0,065
140	— 0,0922	— 0,077
145	— 0,1077	— 0,093
150	— 0,1276	— 0,112
155	— 0,1544	— 0,139
160	0,1934	— 0,178

REDUCTIONS EMPLOYED.

In the reductions of the Observations for 1832 and 1833, I have continued to employ the numbers *a*, *b*, *c*, *d*, &c. given in the Catalogue of the Royal Astronomical Society, and for the numbers of A, B, C, D, I have availed myself of the values given in the Supplements to the Nautical Almanac, which I have reduced to nine o'clock in the evening for the Meridian of Madras; in the case of the Pole Star, and δ Ursæ Minoris the computations have been made for the moment of Transit.

ON THE CLOCK ERRORS AND CLOCK RATES.

In the result of Observations for 1831 Vol. I. I have explained at some length the method employed for the determination of the error and rate of the clock, and have exhibited the degree of accuracy to which the observa-

tions lay claims; on the present occasion I have therefore thought it sufficient to refer to these, and to state, that the reduction of the Observations for 1832 and 1833, have been effected agreeably to the plan there laid down with but one slight exception, namely; in the reductions for 1831, I had employed the Greenwich Catalogue of 720 Stars, whereas in the reduction for 1832 and 1833; those Stars only of this Catalogue have been employed, which are situated between the limits of 65° and 115° of North Polar Distance; by this arrangement we are enabled to correct the Right Ascension of a Star for any small error of Level, Azimuth, or Collimation which may have been committed in the reduction, from inequality of the Pivots, or from a wrong assumption of the position of either of the Marks with regard to the meridian: independent of this consideration, the more rapid motion of Equatorial Stars through the field of view recommends them to preference where *general* accuracy only is our aim.

With regard to the accuracy of the determination of the Clock Errors, I may very safely claim for them an increased degree above that of 1831, and considering that with one exception only the same observers have been employed, this of course could only be expected; the exception I allude to is the exclusion of the Assistant S from making further observations; it will be recollected that towards the beginning of 1831 the observers S. M. A. R. and T. or my four Assistants and self agreed to two or three tenths of a second of time in estimating the time of Transit of a Star, whereas towards the end of that year, the Assistant S had acquired a habit of observing which gave rise to a difference of two seconds of time from the other three Assistants; in consequence of my absence from Madras at this time, (being otherwise employed in Calcutta) the evil was allowed to exist up to the middle of the year 1832, since which time I have not allowed the Assistant S to make any observations, and (agreeable to the plan followed in 1831) have employed only those observations of his before this time, which are situated in the vicinity of *known* Stars, and have rejected the rest; with regard to the Assistants M. R. and A. they continue up to the present time steadily to observe within two tenths of a second of myself and with about the same degree of accuracy. With a view to discover the cause of the difference above found, I lately tried the effect of pressure upon the Telescope whilst observing, this being the only means by which so large a discordance as two seconds might be accounted for, the result was, that a pressure of 5 pounds upon the end of the Telescope did not produce a deviation to the amount of 10 seconds of space; a fact, which (although it leaves us unsatisfied as to the present enquiry) speaks very satisfactorily with regard to the stability of the Telescope. With regard to

the going of the Clock ; it will be remarked that its irregularities are both large and frequent ; this is partly due to an ill constructed click, which I have not been able to get remedied at Madras, whereby the Clock has stopt, or tript, on the days of winding, (on the 1st and 15th of each month) ; and partly from the decayed state of the Clock case, which has allowed spiders to creep into the works ; the latter cause has I hope now been removed by a new plank which I lately caused to be screwed to the back of the case, and the former it must be recollected does not affect the reduced places of the Sun or Stars ; for the irregularities with which we *have* to contend, I may remark, that the method of reduction (the employment of the places of several known Stars, and separating the results into sets occupying two or three hours of A. R. only) keeps so severe a check upon the error of the Clock ; that an error of one tenth of a second of time from this cause is of unfrequent occurrence ; in a few cases however where uncertainty to the amount of 3 or 4 tenths has occurred, I have rejected the observations altogether.

1832	Clock Rate by		Difference.	REMARKS.	1832	Clock Rate by		Difference.	REMARKS.
	Sun.	Stars.				Sun.	Stars.		
Jan.	<i>s.</i> 3 + 2,43	<i>s.</i> + 2,21	<i>s.</i> 0,22		Feb. 19	<i>s.</i>	<i>s.</i> + 2,98	<i>s.</i>	
	5.....	+ 2,51			20	+ 3,09	+ 3,15	0,06	
	6.....	+ 2,03			21	+ 2,98	+ 2,87	0,11	
	10 + 2,07	+ 1,92	0,15		22	+ 2,74	+ 2,75	0,01	
	12 + 1,72	+ 1,79	0,07		23	+ 3,03	+ 2,72	0,31	
	13.....	+ 2,84			24	+ 2,50	+ 2,72	0,22	
	14 + 2,27	+ 2,00	0,27		25	+ 3,28	+ 2,70	0,58	
	16.....	+ 2,66			26	+ 2,94		
	26 + 2,50	+ 2,77	0,27		27	+ 2,90		
	27 + 2,02	+ 2,27	0,25		28	+ 2,70		
	29 + 2,07	+ 2,20	0,13		29	+ 3,13	Stopt(I presume)
	30.....	+ 2,31			March 1	+ 0,36		two seconds in
	31 + 3,10	+ 3,02	0,08		2	+ 2,30		winding.
Feb.	1.....	The Clock stopt	3	+ 2,48	+ 2,40	0,08	
	2.....	+ 2,57		in winding.	5	+ 2,64	+ 2,55	0,09	
	3.....	Stopt the Clock	6	+ 2,45	+ 2,26	0,19	
	4 + 2,39	+ 2,37	0,02	four minutes.	7	+ 1,99	+ 2,00	0,01	
	5 + 2,39	+ 2,24	0,15		8	+ 2,14	+ 2,44	0,30	
	6 + 2,62	+ 2,63	0,01		9	+ 2,12	+ 2,08	0,04	
	7 + 2,18	+ 2,04	0,14		10	+ 2,31	+ 2,50	0,19	
	8 + 2,19	+ 2,13	0,06		11	+ 2,50		
	9 + 2,22	+ 2,10	0,12		12	+ 2,46		
	10 + 1,82	+ 1,88	0,06		13	+ 2,41	+ 2,22	0,19	
	11 + 2,15	+ 1,95	0,20		14	+ 2,52	+ 2,52	0,00	
	12.....	+ 2,08			15	+ 1,87	+ 2,23	0,36	
	13.....	+ 2,01			16	+ 2,29	+ 2,25	0,04	
	14.....	+ 1,94			17	+ 2,27	+ 2,60	0,33	
	15 + 1,89	+ 2,38	0,49		18	+ 2,13	+ 2,39	0,26	
	18.....	+ 2,99			19	+ 2,07	+ 2,13	0,06	

1832	Clock Rate by		Difference.	REMARKS.	1832	Clock Rate by		Difference.	REMARKS.
	Sun.	Stars.				Sun.	Stars.		
	<i>s.</i>	<i>s.</i>	<i>s.</i>			<i>s.</i>	<i>s.</i>	<i>s.</i>	
March 20	+ 2,26	+ 2,10	0,16		May 14	+ 0,92	
21	+ 1,81	+ 2,05	0,24		15	+ 1,14	+ 4,68	The Clock tript
22	+ 2,04	+ 1,92	0,12		16	+ 3,29	in winding.
23	+ 2,23			17	+ 9,62	I removed the
24	+ 2,24	+ 2,14	0,10		18	- 0,20		head of the Clock
25	+ 2,02	+ 2,12	0,10		19	- 1,57		and cleaned the
26	+ 2,13	+ 2,28	0,15		20	- 1,64		scapement, &c.
27	+ 2,27	+ 2,40	0,13		21	- 1,67		and applied fresh
28	+ 2,72	+ 3 05	0,33		22	- 2,00		oil.
29	+ 3,05			23	- 1,66		
30	+ 3,43			25	- 1,22			
31	+ 3,65	+ 3,83	0,18		26	- 0,64	- 1 05	0,41	
April 1	+ 3 21			27	- 1,08		
2	+ 3,68	+ 3,73	0,05		28	- 1,02		
3	+ 3,53	+ 3,33	0,20		29	- 0,72		
4	+ 2,99	+ 3,05	0,06		30	- 0,81		
5	+ 3,02	+ 2,97	0,05		31	- 0 87		
6	+ 3,00	+ 2,90	0,10		June 1	- 0,80	- 1,05	0,25	
7	+ 2,83	+ 2,83	0,00		2	- 1,83			
9	+ 2,71	+ 3,15	0,44		4	- 1,89	- 1,65	0,24	
10	+ 3,29	+ 3,13	0,16		5	- 2,37	- 2,44	0,07	
11	+ 3,09	Stopt the Clock	7	- 2,22			
12	+ 3,40	four minutes and	8	- 2,61			
13	+ 2,57	lengthened the	9	- 2,27	- 2,30	0,03	
14	+ 2,80	pendulum.	10	- 2,54		
15	+ 0,21	The Click re-	11	- 2,50	- 2,54	0,04	
16	0,00	fused to do its duty	12	- 2,72	- 2,36	0,36	
17	- 0,24	in consequence	13	- 2,48	- 2,82	0,34	
18	- 0,43	- 0,36	0,07	the Clock stopt for	14	- 3,27	- 3,18	0,09	
19	- 0,46	- 0,43	0,03	several seconds in	15	- 3,00	- 2,86	0,14	
20	- 0,23			winding.	16	- 2,78	- 2,90	0,12	
21	- 0,65	- 0,98	0,33		18	- 2,71			
22	- 0,10	- 0,45	0,35		23	- 2,88			
23	- 0,36	- 0,54	0,18		27	+ 3,48			
24	- 0,24				28	+ 4,99			
25	- 0 85				July 2	+ 2,31			
26	- 0,03	- 0,38	0,35		3	+ 0,65			
27	+ 0,20	0,00	0,20		5	+ 6,34			
28	- 0,56	- 0,46	0,10		7		The Clock was
29	- 0,14	- 0,29	0,15		16	- 5,00		cleaned by
30	+ 0,02	+ 0,05	0,03		20	- 4,61			Mr. Law.
May 1	- 0,06	0,00	0,06		25	- 5,27	- 5,05	0,22	
2	+ 0,14	- 0,37	0,51		26	- 5,90			
3	- 0,44	+ 0,11	0,55		28	- 5,54			
4	- 0,03	- 0,27	0,24		29	- 5,43		
5	- 0,46	- 0,27	0,19		30	- 5,48			
6	- 0,55	- 0,19	0,36		Aug. 1	+ 7,54			
7	- 0,12	- 0,41	0,29		2	+ 6,88			
8	- 0,09	- 0,16	0,07		3	+ 7,46			
9	+ 0,21	- 0,09	0,30		5	+ 8,49			
11	+ 4 +	The Clock pro-	7		The Clock was
12	- 0,35	0,00	0,35	bably tript 4					inspected by
13	+ 0,65		seconds.					Mr. Law.

ON THE CLOCK ERRORS AND CLOCK RATES.

1832	Clock Rate by		Difference.	REMARKS.	1832	Clock Rate by		Difference.	REMARKS.		
	Sun.	Stars.				Sun.	Stars.				
Aug.	8	+15,81	Regulated.	Oct.	23	+ 1,07	+ 1,01	0,06	
	14	- 4,51	- 5,00	0,49			24	+ 0,91	+ 0,87	0,04	
	17	- 5,14	- 5,00	0,14			25	+ 0,14	Stopt the Clock
	18	- 5,14	- 5,28	0,14			26	+ 0,55	+ 0,29	0,26	two minutes.
	19	- 5,38					27	+ 0,32	+ 0,37	0,05	
	20	- 5,13	- 5,19	0,06			28	+ 0,41		
	21	- 5,46					29	+ 1,31		
	22	- 5,09					30	+ 1,46	+ 1,27	0,19	
	23	- 5,49	- 5,27	0,22			31	+ 0,77	+ 0,42	0,35	
	24	- 5,39			Nov.	1	+ 0,40	+ 0,20	0,20	
	25	- 5,34					2	+ 0,54		
	27	- 5,46				3	+ 0,42	+ 1,07	0,65	
	28	- 4,85	- 5,08	0,23			4	+ 5,26	+ 8,91		
	29	- 5,33					5	+ 8,42		
	30	- 5,36	- 5,27	0,09			7	cleaned the
	31	- 5,05	- 3,82				9	- 4,39		Clock.
Sept.	1	+ 0,23				10	- 4,55	- 4,49	0,06	
	4	+ 1,70	+ 1,13	0,57			12	- 4,57		
	6	+ 1,17				13	- 4,47	- 4,47	0,00	
	7	+ 1,20	+ 1,45	0,25			15	- 4,61	- 4,52	0,09	
	9	+ 1,88	+ 2,25	0,37	Advanced the		16	- 4,51	- 4,33	0,18	
	11	+ 0,33		Clock 3 mi-		17	- 4,06	- 4,20	0,14	
	12	+ 0,64		minutes.		18	- 4,25	- 4,06	0,19	
	14	+ 0,58				19	- 4,38	- 4,22	0,16	
	15	+ 0,56	+ 1,01	0,45			21	- 3,79			
	19	+ 1,76				22	- 4,53	- 3,90	0,63	
	21	- 5,07					23	- 4,11	- 4,23	0,12	
	22	- 5,11	- 5,06	0,05			24	- 4,26	- 4,12	0,14	
	23	- 5,05	- 5,11	0,06			25	- 4,49	- 4,56	0,07	
	24	- 5,55	- 5,09	0,46			26	- 4,10			
	25	- 4,97					27	- 4,44			
	26	- 4,37	- 4,43	0,06			29	- 4,56		
	27	- 4,24	- 4,39	0,15			30	- 4,58	- 4,53	0,05	
	28	- 4,25				Dec.	1	- 4,87			
	30	- 1,39				4	- 4,65		
Oct.	1	+ 10,23	Wound up the		5	- 4,42	- 4,40	0,02	
	2	+ 12,59	+ 14,53		Clock.		6	- 4,47		
	3	+ 14,04				7	- 4,41	- 4,47	0,06	
	4	+ 12,62				8	- 3,93	- 4,20	0,27	
	5	+ 8,97				9	- 4,56	- 4,15	0,41	
	6	+ 7,94	+ 7,84	0,10			10	- 4,40	- 4,24	0,16	
	7	+ 10,07					11	- 4,40		
	8	+ 6,85	+ 5,56				12	- 4 81		
	9	+ 4,36	+ 4,42	0,06			14	- 4,61	- 4,67	0,06	
	11	+ 4,17	+ 4,19	0,02			15	- 4,82	- 4,84	0,02	
	12	+ 3,70	+ 4,76				16	- 4,24		
	13	+ 4,43	+ 4,22	0,21			17	- 4 12		
	14	+ 5,11				18	- 4,79		
	15	+ 4,77	Wound up the		19	- 4,02	- 4,43	0,41	
	19	- 0,10		Clock.		20	- 4,76	- 4,46	0,30	
	20	- 1,18					21	- 4,61	- 4,89	0,28	
	21	+ 0,32	+ 1,00				22	- 5,17			
	22	+ 0,95	+ 1,26	0,31			23	- 5,14			

1832				REMARKS.	1833						
Clock Rate by			Difference.		Clock Rate by			Difference.	REMARKS.		
Sun.	Stars.				Sun.	Stars.					
Dec.	24	- 4,76	- 4,83	0,07	Feb.	17	+ 0,89	+ 0,73	0,16		
	25	- 4,51	- 4,70	0,19		18	+ 0,67	+ 0,64	0,03		
	26	- 4,91	- 4,53	0,38		19	+ 0,81	+ 0,74	0,07		
	27	- 4,51	- 4,93	0,42		20	+ 0,58	+ 0,27	0,31		
	28	- 4 63				21	+ 0,62				
	29	- 4,62				22	+ 1,69				
						23	The Clock tript	
						24				9 seconds; I ap-	
						25				plied fresh oil to	
1833						26	+ 0,27	+ 0,46	0,19	the escapement.	
Jan.	3	- 3,74	- 3,56	0,18		27	+ 0,79	+ 0,73	0,06		
	4	- 3,41	- 3,37	0,04		28	+ 2,08				
	5	- 3,52	- 3,17	0,35		March	1	+ 1,70	+ 1,18	0,52	
	6	- 3,29	- 3,28	0,01			2	+ 0,96	+ 1,05	0,09	
	7	- 3,40					3	+ 0,80	+ 0,52	0,28	
	8	- 3,74	- 3,63	0,11			4	+ 0,32	+ 0,63	0,31	
	9	- 3,34	- 3,50	0,16			5	+ 0,67	+ 0,88	0,21	
	10	- 3,73	- 3,99	0,26			6	+ 0,98	+ 0,65	0,33	
	11	- 3,74	- 3,56	0,18			7	+ 0,38	+ 0,52	0,14	
	12	- 3,44	- 2,98	0,46			8	+ 0,46	+ 0,74	0,28	
	14	- 2,80	- 2,93	0,13			9	+ 1,23	+ 0,90	0,33	
	15	+ 0,25	Regulated the	10	+ 0,79			
	16	+ 0,20	+ 0,25	0,05	Clock.	11	+ 0,60	+ 1,00	0,40		
	17	+ 0,74	+ 0,64	0,10		12	+ 0,67	+ 0,46	0,21		
	18	+ 0,86	+ 0,62	0,24		13	+ 0,40	+ 0,46	0,06		
	19	+ 0,44	+ 0,50	0,06		14	+ 0,41	+ 0,46	0,05		
	20	+ 0,68	+ 0,20	0,48		15	+ 1,32	+ 1,86	0,54		
	21	- 0,04	+ 0,06	0,10		16	+ 3,86			
	22	+ 0,14	+ 0,18	0,04		17	+ 3,97			
	23	- 0,05			18	+ 1,80			
	24	- 0,23			19	+ 1,17	+ 1,61	0,44		
	25	- 0,14			20	+ 2,00	+ 2,00	0,00		
	26	- 0,20	0,00	0,20		21	+ 1,42	+ 0,80	0,62		
	27	- 0,42			22	+ 0,58	+ 0,84	0,26		
	28	- 0,12	- 0,48	0,36		23	+ 0,65	+ 0,65	0,00		
	29	- 0,21	- 0,46	0,25		24	+ 1,93			
	30	- 0,13	0,00	0,13		25	+ 1,27	+ 0,63			
	31	- 0,01	+ 0,04	0,05		26	+ 0,69	+ 0,95	0,26		
Feb.	1	+ 0,26	+ 0,16	0,10		27	+ 0,80	+ 0,88	0,08		
	2	+ 0,31	+ 0,12	0,19		28	+ 0,83	+ 0,65	0,18		
	3	+ 0,17	+ 0,15	0,02		29	+ 0,70	+ 0,62	0,08		
	4	+ 0,11	- 0,11	0,22		30	+ 0,47	+ 0,43	0,04		
	5	- 0,32	- 0,12	0,20		31	+ 0,50	+ 0,54	0,04		
	6	- 0,10	0,00	0,10		April	1	+ 0,33	+ 0,60	0,27	
	7	+ 0,02				2	+ 0,83	+ 0,60	0,23	
	8	+ 0,20	+ 0,05	0,15			3	+ 0,39	+ 0,35	0,04	
	9	+ 0,56	+ 0,74	0,18			4	+ 0,60	+ 1,46		
	10	+ 0,64				5	+ 1,87		
	11	+ 0,63	+ 0,39	0,24			6	+ 0,74		
	12	+ 0,74	+ 0,78	0,04			7	+ 1,61		
	13	+ 0,81	+ 0,89	0,08			8	+ 2,51		
	14	+ 0,50	+ 0,85	0,35			9	+ 2,72	+ 2,88	0,16	
	15	+ 0,59	+ 0,58	0,01			10	+ 3,23			
	16	+ 0,59	+ 0,65	0,06							

1833	Clock Rate by		Difference.	REMARKS.	1833	Clock Rate by		Difference.	REMARKS.
	Sun.	Stars.				Sun.	Stars.		
	<i>s.</i>	<i>s.</i>	<i>s.</i>			<i>s.</i>	<i>s.</i>	<i>s.</i>	
April	11	+ 1,89			June	11 - 4 51		
	13 + 2,80				12	- 4,66 - 4,62	0,04	
	14 + 3,23				13	- 4,63 - 4,81	0,18	
	15	+ 6,53 + 8,14				14	- 4,70 - 4,50	0,20	
	16	+ 7,57 + 5,83	I found a fine		19	Regulated the
	17	+ 2,68 + 2,25	0,43	cobweb attached		21 - 0,02		Clock.
	18	+ 2,11 + 2,37	0,26	to the escapement		22		
	19	+ 2,69 + 3,00	0,31	which I removed.		23 - 0,20		
	20			26	- 1,52		
	21 - 2,01		Regulated the		27	- 2,14		
	22	- 1,84 - 2,05	0,21	Clock.		28	- 2,44		
	23	- 2,12 - 2,40	0,28			29	- 3,01 - 3,31	0,30	
	24	- 2,48 - 2,37	0,11			30 - 3,78		
	25	- 2,16 - 2,35	0,19		July	1	- 3,91 - 4,52		
	26 - 2,12				2	- 4,21 - 4,20	0,01	
	27	- 2,27 - 2,24	0,03			3	- 4,52		
	28	- 1,56 - 1,28	0,28			5	- 4,45 - 4,46	0,01	
	29	- 2,32 - 2,91	0,59			7 - 4,61		
	30	- 3,10 - 3,25	0,15			8 - 5,48		
May	1	- 3,44 - 3,43	0,01			9	- 5,59		
	2	- 3,44 - 3,28	0,06			12	Regulated the
	3	- 3,42 - 3,19	0,23			13	+ 0,60 + 0,42	0,18	Clock.
	4	- 3,00 - 2,76	0,24			15	+ 0,44 + 0,02	0,42	
	5 - 2,70				16		
	6	- 2,72 - 2,70	0,02			17	+ 0,10		
	7	- 2,83				18	+ 0,09		
	8	- 3,32 - 3,38	0,06			19	- 0,10		
	9	- 3,76 - 3,63	0,13			20	- 0,26 - 0,24	0,02	
	10	- 3,41 - 3,51	0,10			21	- 0,80		
	11	- 3,36 - 3,50	0,14			23 - 0,94		
	12	- 3,55 - 3,35	0,20			25	- 0,66 - 0,66	0,00	
	13	- 3,65 - 4,07				26 - 0,67		
	14	- 3,77				27	- 0,80 - 0,85	0,05	
	19	- 3,08				28	- 1,06 - 0,76	0,30	
	20	- 3,62				29	- 1,10		
	22	- 3,52				30	- 1,00		
	23	- 5,02 - 4,53	0,49		Aug.	1	- 0,66 - 0,63	0,03	
	24	- 4,46 - 4,67	0,21			2	- 0,26 - 0,85	0,59	
	25	- 4,95				3	- 0,71		
	26	- 5,31				4	- 0,53 - 0,69	0,16	
	28	- 4,81				5	- 0,29 - 0,48	0,19	
	29	+ 4,07				6	- 1,33 - 1,20	0,13	
	30	+ 4,97 + 5,54				7	- 0,89 - 1,14	0,25	
	31	+ 5,90 + 6,27	0,37			8	- 0,54 - 0,94	0,40	The minute hand
June	1	+ 6,45	Wound up the		10 - 2,51		of the Clock stopt,
	2	- 3,55 - 4,75		Clock.		11	- 2,68		in consequence of
	3	- 4,41 - 3,40				12	- 2,57 - 2,75	0,18	having become
	4	- 3,39				13	- 1,84		loose.
	5	- 4,03				14 - 1,68		
	6	- 4,16				15	- 2,09 - 1,77	0,32	
	7	- 4,50				16	- 1,94		
	8	- 4,38 - 4,54	0,16			17	- 0,91 - 0,74	0,17	
	9 - 4,66				18 - 1,41		

1833	Clock Rate by		Difference.	REMARKS.	1833	Clock Rate by		Difference.	REMARKS.
	Sun.	Stars.				Sun.	Stars.		
	<i>s.</i>	<i>s.</i>	<i>s.</i>			<i>s.</i>	<i>s.</i>	<i>s.</i>	
Aug. 23	- 1,59	- 1,63	0,04		Oct. 20 + 1,92			
24	- 2,01				21 + 0,80			
28	- 1,22				22	+ 1,62 + 1,41	0,21		
29 - 1,44				23	+ 1,27 + 1,21	0,06		
30	- 1,19	- 1,05	0,14		24 + 1,14			
31	- 1,56				31 + 0,85			
Sept. 2	- 1,41				Nov. 1	+ 0,77			
3	- 0,58				5 + 2,10			
5	An alteration of	6 + 0,83			
6	- 0,38			about 30 seconds	7 + 0,41	Forwarded the	
7	- 1,17			in the error of the	16 + 1,06		Clock two mi-	
8	- 0,84	- 1,17	0,33	clock took place	17 + 1,55		nutes.	
9	- 1,37	- 1,00	0,37	between the 3d	18 + 0,96			
10	- 1,36	- 1,65	0,29	and 5th.	19	- 1,37 - 2,05			
11	- 1,52	- 1,08	0,44		20 - 1,99			
12	- 1,45	- 1,50	0,05		21 - 1,88			
13 - 1,49				22	- 1,46			
14 - 1,27				23 - 1,67			
15	- 1,16	- 1,39	0,23		29	- 1,73			
16	- 1,25				Dec. 1	- 1,27			
17 - 0,75				2	- 1,06 - 1,49	0,43		
18	- 0,96	- 1,46	0,50		3	- 1,08 - 0,83	0,25		
19	- 1,36				4 - 0,34			
21 - 1,64				5	- 0,50 - 0,51	0,01		
26	- 1,53	- 1,26	0,27		6 - 0,88			
27	- 0,36				7	- 0,92 - 0,86	0,06		
28	- 1,78				8	- 0,75			
29	- 1,90				9	- 0,85 - 0,89	0,04		
30 - 1,46				10	- 0,74			
Oct. 2 - 1,62				11	- 1,22 - 1,12	0,10		
4	- 1,47	- 1,46	0,01		12	- 0,93 - 1,10	0,17		
5 - 1,20				13 - 1,16			
6	- 1,38	- 1,50	0,12		14	- 1,19			
7 - 1,68				17 - 2,40			
8 - 1,61				18	- 2,15 - 2,28	0,13		
9 - 0,77				19	- 2,61 - 2,72	0,11		
10 - 0,93				20	- 2,21 - 1,93	0,28		
11 - 1,18				22	- 2,62			
12 - 1,24				23	- 3,02 - 2,77			
13 - 0,94				24	- 2,35 - 2,49	0,14		
14 + 0,31				25 - 2,13			
15	+ 0,06				26	- 2,42 - 2,46	0,04		
16	+ 0,85 + 0,78		0,07		27	- 2,33			
17 + 2,12				29 - 1,64			
18	+ 2,30				30	- 1,99 - 2,13	0,14		
19	+ 2,56 + 2,65		0,09		31	- 2,96			

OF THE MURAL CIRCLE.

This Instrument having been already sufficiently described in Vol. I, it is only necessary for me here to remark that I have continued to employ the full aperture ($3\frac{3}{4}$ Inches) and the same power (about 140) as heretofore. Towards the end of the year 1832, being desirous of ascertaining the amount of error of some of the divisions, I availed myself of the cloudy evenings which then occurred, to measure the angular distance between two *collimators*, which I had previously adjusted to subtend an angle of 90° ; by this means I was put in possession of the error of the points 90° 180° 270° .

I now placed two collimators so as to subtend an angle of 30° and thus obtained the error of the points 30° , 60° , 120° , &c. and subdividing these, eventually arrived at the errors of every fifth degree; the particulars of these measurements having been transmitted to England for publication, it is only necessary for me to remark, that the largest error which would be committed by the employment of any division together with that situated at 180° distance, did not exceed $2''.5$; and this is probably too large, being subject to the errors of observation. On the occasion of making these observations it was necessary to unclamp the Telescope from the circle, and on again clamping it to readjust the Telescope for Level, &c.; but at no other time during the years 1832 and 1833, has any adjustment been found necessary. On inspecting the rough observations a consistency is found to exist among the microscope readings which speaks in a manner highly creditable for the stability of the axis. With regard to the state of the Instrument I may safely assert, that now, after three and a half years of active employ, it is in no respect injured by wear, and but little deteriorated in appearance.

METEOROLOGICAL INSTRUMENTS EMPLOYED.

The Barometer employed at the beginning of 1832 was made by Cary; this

on being compared (see page 59, Vol. I.) with Standard Barometer No 3, by Gilbert, shewed that the indications of the former were in defect 0,152 Inches, hence it is necessary to increase the Barometrical indications set down in the Mural Circle Book by this amount from the 1st January up to the 20th February 1832; for the observations after this date the Standard Barometer No. 3, by Gilbert was employed, which consequently only requires the correction for capillary action + ,027. With a view to discover if the Barometer in question has remained undisturbed, I have occasionally compared it with another Standard No. 6, by Gilbert; the result of these comparisons shew that the same difference (.018) exists between them as found in Calcutta, when they were compared with the other Standards in the Surveyor General's Office. The Thermometers employed at the commencement of 1832 were A and B by Jones, which from comparisons made with the Standard A belonging to the Surveyor General's Office at Calcutta, appear to be 0°,54 and 0°,47 respectively *too low*; hence, (Thermometer A having been employed "*in doors*" for 1832 and 1833) it becomes necessary to add 0°,54 to the in door Thermometer as set down in the Circle Book; for the Thermometer "*without*", the Thermometer B was employed up to 1st March 1832; for which period the indications must consequently be increased 0°,47: after this time and up to the end of 1833, a Standard Thermometer by Troughton was employed, which I selected in Calcutta as agreeing with the Standard A in the Surveyor General's Office; consequently from the 1st March 1832 up to the end of 1833, the *out door* Thermometer as set down in the Mural Circle Book does not require correction.

OBSERVATIONS MADE WITH THE MURAL CIRCLE.

Having found it inconvenient to observe the reflected image of Stars from a basin of quicksilver by reason of the disturbance necessarily produced by the observer at the Transit Instrument, I have during the years 1832 and 1833, given up observing by reflection. In the determination of the Index Error I have continued to employ those Stars of the Greenwich Catalogue which are situated between 25° and 90° of N. P. D. these being the limits between which the uncertainty of refraction is but small.

In the reduction of the Greenwich Catalogue Bradley's table of refraction was employed, whereas in the reduction of the Madras Results I have for reasons explained at Page 61, Vol. I. employed Atkinson's table; I have consequently reduced the Greenwich Catalogue to the tenor of Atkinson's table of refraction before using it in computing the Index Error (see Page 62, Vol. I.)

The table of Index Errors which now follows has been employed in computing the places of the fixed Stars, and the Planets when the centre of the body has been observed; but in the case of the Sun and Moon, and of Planets where the *limb* has been observed, an allowance has been made of 1",2 for the semi-diameter of the wire.

Index Error of the Madras Mural Circle for the years 1832 and 1833.

Date.	No. of Observations.	Index Error.	Mean.	REMARKS.	Date.	No. of Observations.	Index Error.	Mean.	REMARKS.
1832					1832				
Jan.	1	14	2 58,41		Feb.	19	8	3 3,44	
	3	12	2 59,11			21	11	3 4,04	
	5	21	2 59,08			23	10	3 4,62	
	6	9	3 0,19			24	11	3 2,03	
	10	8	2 59,06			26	21	3 18,11	I adjusted the microscopic readings.
	12	4	2 59,40			27	7	3 16,09	
	13	9	3 1,04			29	15	3 16,47	
	14	12	3 4,37		March	1	11	3 16,63	
	18	9	3 1,20			2	10	3 16,37	
	22	17	3 1,40			4	15	3 16,28	} — 3 16,37
	24	9	3 2,00			5	6	3 15,61	
	26	20	3 1,66			7	8	3 15,25	} — 3 15,47
	27	18	3 2,33			10	13	3 15,55	
	28	13	3 2,52			11	8	3 27,81	I took the In- strument down, and applied fresh oil to the axis.
	29	12	3 1,95			12	6	3 28,16	
	30	14	3 2,45			13	8	3 28,48	
	31	9	3 3,93			15	10	3 28,88	
Feb.	1	13	3 3,00			18	12	3 28,80	
	2	12	3 2,79			20	13	3 26,85	
	3	12	3 3,73			23	9	3 27,03	
	4	10	3 3,74			24	10	3 28,05	
	5	10	3 3,51			25	8	3 27,30	
	6	11	3 3,29			26	8	3 27,41	
	7	4	3 3,36			28	9	3 27,59	
	8	9	3 2,30			29	9	3 28,55	
	10	11	3 2,50			30	6	3 28,36	
	11	7	3 2,19			31	10	3 28,57	
	13	9	3 2,70		April	1	12	3 28,13	
	15	15	3 2,14			3	20	3 27,32	— 3 27,61

INDEX ERROR OF THE MURAL CIRCLE FOR 1832 AND 1833. 61

Date.	No. of Observations.	Index Error.	Mean.	REMARKS.	Date.	No. of Observations.	Index Error.	Mean.	REMARKS.
1832					1832				
April	4	10	3 27,21	} — 3 27,61	Sept.	27	11	3 25,11	
	6	14	3 28,04		30	7	3 25,18		
	10	10	3 27,49		Oct.	1	13	3 25,09	
	12	9	3 27,39			2	10	3 24,65	
	13	7	3 27,53			3	15	3 25,88	
	14	6	3 28,42			7	11	3 25,28	
	16	8	3 27,43			9	10	3 24,22	
	20	7	3 26,83			11	9	3 25,12	
	22	10	3 25,17			21	15	3 25,15	
	24	6	3 25,32			24	10	3 24,00	
	26	11	3 24,99	} — 3 24,89	26	10	3 25,00		
	28	14	3 24,51		27	7	3 24,27		
	30	12	3 24,24		29	7	3 26,39		
May	1	4	3 24,64		30	11	3 26,03		
	2	6	3 25,37		31	8	3 25,27		
	4	10	3 26,42	Nov.	2	9	3 25,65		
	9	11	3 27,75		3	9	3 22,42		
	11	7	3 26,60		4	9	3 23,64		
	12	10	3 26,72	} — 3 26,77		8	7	3 24,71	
	16	14	3 26,39			10	7	3 23,06	
	17	5	3 25,46			12	7	3 24,59	
	18	8	3 25,25			16	9	1 19,14	
	20	10	3 25,51			17	9	1 19,74	
	22	10	3 24,86	} — 3 25,23		19	11	1 20,16	
	26	9	3 25,46			21	6	1 20,10	
	28	7	3 24,88			22	8	1 19,84	
	31	7	3 25,00			25	12	1 20,17	
June	5	9	3 25,44			29	11	1 18,84	
	10	7	3 24,04			30	6	1 17,60	
	12	11	3 23,59	} — 3 25,23	Dec.	4	8	4 46,81	
	17	13	3 25,33			6	8	4 47,47	
	22	3	3 25,17			8	13	2 50,41	
July	29	11	3 23,25			12	7	3 2,73	
Aug.	8	10	3 23,22			13	6	3 0,11	
	11	12	3 22,28			16	12	4 24,04	
	15	7	3 21,74			17	8	4 23,57	
	18	12	3 22,85			18	7	4 25,01	
	23	13	3 23,88			19	5	4 24,67	
	24	11	3 23,70			21	11	7 17,71	
	26	11	3 25,02	} — 3 25,23		22	9	7 16,53	
	27	8	3 24,25			24	10	7 17,48	
	28	13	3 24,36			25	9	7 18,36	
	31	10	3 25,39			27	5	3 3,13	
Sept.	7	10	3 24,97						
	9	10	3 24,05						
	11	10	3 23,95						
	19	14	3 23,72						
	22	11	3 24,67						
	23	9	3 25,63						
	24	13	3 24,39						
	25	15	3 24,78						
	26	12	3 24,44						
					1833				
					Jan.	2	11	3 4,30	
						3	17	3 5,36	
						4	17	3 5,85	
						5	9	3 6,05	

Unclamped the Telescope.

Unclamped the Telescope.

Do. do.

Do. do.

Do. do.

Do. do.


I took down the Circle and cleaned the axis.

62 INDEX ERROR OF THE MURAL CIRCLE FOR 1832 AND 1833.

1833				1833					
Date.	No. of Observations.	Index Error.	Mean.	REMARKS.	Date.	No. of Observations.	Index Error.	Mean.	REMARKS.
Jan. 6	11	- 3 4,66	} - 3 4,56		March 10	16	- 3 6,27	}	Unclamped the Telescope.
8	11	3 4,74							
10	9	3 4,08	} - 3 5,12		12	17	1 34,75	} - 1 36,58	
11	8	3 4,74							
14	12	3 5,12	} - 3 5,83		13	19	1 34,35	} - 1 35,36	
15	11	3 5,12							
16	11	3 5,51	} - 3 5,83		14	20	1 35,30	} - 1 36,41	
17	9	3 6,38							
18	8	3 5,56	} - 3 5,57		15	19	1 35,11	} - 1 33,24	
19	10	3 6,08							
20	7	3 5,89	} - 3 6,29		16	17	1 36,48	} - 1 33,02	
21	9	3 5,53							
22	8	3 5,63	} - 3 5,65		17	18	1 36,45	} - 1 31,52	
23	11	3 6,45							
25	17	3 5,90	} - 3 5,65		18	19	1 36,26	} - 1 31,52	
26	14	3 5,76							
27	14	3 5,73	} - 3 5,65		19	16	1 37,09	} - 1 31,52	
28	16	3 5,84							
29	12	3 5,64	} - 3 5,65		20	16	1 36,75	} - 1 31,52	
31	9	3 6,55							
Feb. 2	12	3 6,33	} - 3 5,57		21	15	1 36,53	} - 1 31,52	
4	11	3 5,89							
5	8	3 6,27	} - 3 5,57		22	19	1 36,50	} - 1 31,52	
6	12	3 4,35							
8	13	3 4,36	} - 3 5,57		23	15	1 35,30	} - 1 31,52	
9	16	3 5,06							
10	13	3 5,71	} - 3 5,57		24	15	1 35,49	} - 1 31,52	
11	14	3 5,35							
12	14	3 5,58	} - 3 5,57		25	15	1 34,91	} - 1 31,52	
13	13	3 5,67							
14	13	3 5,53	} - 3 5,57		26	10	1 35,23	} - 1 31,52	
15	14	3 6,25							
16	11	3 6,33	} - 3 5,57		27	11	1 35,72	} - 1 31,52	
17	7	3 5,30							
18	12	3 5,69	} - 3 5,57		28	12	1 35,47	} - 1 31,52	
19	12	3 6,32							
20	11	3 5,21	} - 3 5,57		29	14	1 36,22	} - 1 31,52	
21	10	3 5,58							
22	14	3 5,38	} - 3 5,57		30	7	1 36,39	} - 1 31,52	
23	12	3 5,76							
24	10	3 5,57	} - 3 5,57		31	13	1 35,75	} - 1 31,52	
26	16	3 5,81							
27	12	3 5,98	} - 3 5,57		April 1	10	1 36,60	} - 1 31,52	
28	12	3 5,77							
March 1	13	3 5,52	} - 3 5,57		2	13	1 35,93	} - 1 31,52	
2	11	3 5,04							
4	21	3 5,18	} - 3 5,57		3	17	1 35,40	} - 1 31,52	
5	13	3 5,07							
6	16	3 4,56	} - 3 5,57		4	14	1 35,52	} - 1 31,52	
7	14	3 4,49							
8	14	3 3,94	} - 3 5,57		5	7	1 35,01	} - 1 31,52	
9	15	3 3,46							
					6	11	1 35,28	} - 1 31,52	
					7	9	1 35,19		
					8	9	1 34,36	} - 1 31,52	
					9	6	1 33,45		
					13	9	1 34,31	} - 1 31,52	
					15	8	1 33,28		
					17	9	1 33,26	} - 1 31,52	
					19	9	1 33,19		
					20	6	1 33,90	} - 1 31,52	
					22	9	1 33,94		
					24	12	1 35,44	} - 1 31,52	
					27	12	1 34,07		
					28	8	1 32,80	} - 1 31,52	
					30	8	1 33,80		
					May 3	10	1 32,26	} - 1 31,52	
					4	9	8 1 31,57		
					23	10	1 31,47	} - 1 31,52	
					24	30	8 1 32,06		
					June 2	10	1 30,85	} - 1 31,52	
					7	7	1 30,04		
					8	8	1 28,56	} - 1 31,52	
					9	10	1 27,78		
					11	8	1 28,69	} - 1 31,52	
					12	8	1 28,39		
					20	10	1 26,82	} - 1 31,52	
					28	10	1 26,77		

INDEX ERROR OF THE MURAL CIRCLE FOR 1832 AND 1833. 63

1832				1833					
Date.	No. of Observations.	Index Error.	Mean.	REMARKS.	Date.	No. of Observations.	Index Error.	Mean.	REMARKS.
June 30	9	- 1 27,11	" "		Oct. 13	8	- 1 28,15	" "	
July 2	12	1 25,95			15	11	1 28,29		
5	11	1 27,72			16	7	1 28,99		
8	11	1 26,00			20	13	1 28,73	- 1 28,38	
12, 16	11	1 27,53			23	13	1 27,95		
23	10	1 27,18	- 1 27,37		Nov. 1	15	1 28,70		
26	10	1 27,39			9	7	1 25,22	- 1 25,18	Much rain
Aug. 1	12	1 26,83			17	8	1 25,14		which fell between the 1st and 9th, may have occasioned this change.
3	14	1 25,78			19	14	1 24,44		
5	15	1 26,99			21	14	1 23,94		
7	11	1 25,60	- 1 26,38		22	10	1 24,95	- 1 24,30	
10	13	1 27,06			23	11	1 24,56		
14	14	1 26,23			Dec. 2	15	1 24,33		
16	14	1 26,16			3	13	1 24,99		
29	14	1 27,84			4	13	1 24,59	- 1 24,85	
Sept. 5	14	1 26,66			5	12	1 24,96		
9	12	1 27,66			6	10	1 25,56	- 1 25,52	
10	8	1 26,87			7	8	1 25,48		
11	8	1 28,19			8	10	1 26,57		
13	14	1 28,89			9	8	1 26,90		
15	13	1 28,39	- 1 28,55		10	11	1 27,75		
20	11	1 28,61			14	18	1 28,19		
26	14	1 28,33			18	10	1 25,70		Much rain.
30	7	1 28,55			20	11	1 25,81	- 1 25,86	
Oct. 2	9	1 28,47			23	12	1 25,47		
4	7	1 28,34			25	14	1 28,12	- 1 27,98	
7	11	1 28,71			27	15	1 27,84		
11	15	1 27,33	- 1 28,38		29	11	1 26,89	- 1 27,11	
12	9	1 28,10			31	15	1 27,34		



RESULT OF OBSERVATIONS MADE WITH THE TRANSIT INSTRUMENT AND MURAL CIRCLE, IN THE YEARS 1832 AND 1833.

In the first place we will examine the observations of the Sun ; the observed transit of the first and second limb over the five wires furnishes us with the means of determining the semi-diameter ; for we have \odot mean semi-diameter = $\left(\frac{\odot 2 L - \odot 1 L}{30} \right) \left(1 + \frac{a^1 - a}{48} \right) \sin. \text{N.P.D.} + \text{Log. } \ominus - \odot$. Where a and a^1 represent the Right Ascension of the Sun, at the noon preceding, and at the noon following the day of observation ; in the next place, correcting the observed A.R. for the error of the Clock, and the observed N.P.D. for refraction parallax and Semi-diameter, we obtain results which we will now compare with the places interpolated from the Nautical Almanac, as follows.

Comparison of the observed A.R. and N.P.D. of the Sun, with their places interpolated from the Nautical Almanac, &c.

1832	Observed R.A.			A.R. from Nautical Almanac.		Error of Tables.	Observed N.P.D.	N.P.D. from Nautical Almanac.		Error of Tables.	Mean Semi-diameter.							
	h.	m.	s.	m.	s.			m.	s.		m.	s.						
Jan.	1	18	43	11,27	43	11,40	+	0,13	113	5	10	16	5	16,00	+	5,84	15	58,02
	2				113	0	25	66	0	30,00	+	4,34		
	3	18	52	1,81	52	1,50	-	0,31	112	55	9,19	55	14,00	+	4,81	16	1,44	
	4				112	49	25,77	49	31,00	+	5,23			
	6	19	5	14,50	5	13,90	-	0,60	112	36	43,55	36	44,00	+	0,45	16	1,64	
	7	19	9	38,17	9	37,20	-	0,97	112	29	39,23	29	41,00	+	1,77	15	58,00	
	8	19	14	0,85	14	0,00	-	0,85	112	22	4,23	22	12,00	+	7,77			
	9	19	18	22,72	18	22,30	-	0,42	112	14	11,47	14	17,00	+	5,53	16	0,80	
	10	19	22	44,67	22	44,10	-	0,57	112	5	46,20	5	54,00	+	7,80	16	0,18	
	11	19	27	6,35	27	5,30	-	1,05	111	56	57,08	57	3,00	+	5,92			
	12	19	31	26,78	31	35,80	-	0,98	111	47	47,83	47	46,00	-	1,83	16	2,34	
	13	19	35	46,46	35	45,70	-	0,76	111	38	2,25	38	4,00	+	1,75			
	14	19	40	5,78	40	5,10	-	0,68	111	28	0,62	28	0,00	-	0,62	16	3,33	
	15	19	44	24,63	44	23,50	-	1,13	111	17	30,45	17	32,00	+	1,55	16	3,18	
	18				110	43	35,60	43	38,00	+	2,40			
	19				110	31	28,91	31	32,00	+	3,09	16	3,28	
	21				110	6	10,93	6	10,00	-	0,93			
	22				109	52	58,42	52	55,00	-	3,42			
	24	20	22	39,78	22	38,80	-	0,98	109	25	12,68	25	20,00	+	7,39	16	1,10	
	25	20	26	50,94	26	50,30	-	0,64	109	11	2,58	11	0,00	-	2,58	16	0,67	

RESULT OF OBSERVATIONS IN 1832 AND 1833.

1832	Observed A.R.			A.R. from Nautical Almanac		Error of Tables.	Observed N.P.D.			N.P.D. from Nautical Almanac.		Error of Tables.	Mean Semi-diameter.		
	h.	m.	s.	m.	s.		'	"	'''	'	"		'''	'	"
May	18						70	26	0,85	26	3,00	+	2,15	16 6,48	
	19						70	13	4,39	13	2,00	-	2,39	15 57,80	
	20													16 2,00	
	21	3	51	53,69	51	53,10	- 0,59							16 2,45	
	22	3	55	54,12	55	53,50	- 0,62	69	36	4,10	36	1,00	-	3,10	16 2,33
	23	3	59	55,17	59	54,50	- 0,67	69	21	24,51	24	23,00	-	1,51	16 1,96
	24	4	3	56,77	3	55,90	- 0,87	69	12	58,73	13	5,00	+	6,27	
	25							69	2	7,24	2	8,00	+	0,76	16 1,26
	26							68	51	22,12	51	32,00	+	9,88	16 6,14
	27	4	16	4,58	16	3,70	- 0,88	68	41	19,69	41	19,00	-	0,69	16 0,14
	29	4	24	11,64	24	11,40	- 0,24								
	30	4	28	16,87	28	15,90	- 0,97	68	12	49,40	12	52,00	+	2,60	16 1,72
	June	31	4	32	21,49	32	20,90	- 0,59	68	4	1,99	4	7,00	+	5,01
1		4	36	27,01	36	26,20	- 0,81	67	55	44,58	55	47,00	+	2,42	16 1,42
2		4	40	32,24	40	31,90	- 0,34	67	47	43,95	47	47,00	+	3,05	16 2,96
3		4	44	38,78	44	38,00	- 0,78	67	40	13,26	40	14,00	+	0,74	16 2,52
4		4	48	45,34	48	44,50	- 0,84	67	33	0,66	33	5,00	+	4,34	15 59,30
5		4	52	51,52	42	51,40	- 0,12	67	26	14,34	26	16,00	+	1,66	15 59,04
6		4	56	58,89	56	58,40	- 0,49	67	19	51,10	19	51,00	-	0,10	16 3,98
7		5	1	6,37	1	5,80	- 0,57	67	13	48,35	13	51,00	+	2,65	
8		5	5	13,86	5	13,40	- 0,46	67	8	8,77	8	15,00	+	6,23	
9		5	9	21,89	9	21,40	- 0,49	67	3	4,85	3	3,00	-	1,85	16 2,30
10		5	13	30,16	13	29,60	- 0,56	66	58	8,42	58	15,00	+	6,58	16 1,33
11		5	17	38,91	17	38,00	- 0,91	66	53	50,47	53	51,00	+	0,53	16 2,05
12		5	21	47,19	21	46,60	- 0,59	66	49	48,77	49	52,00	+	3,23	16 3,05
13		5	25	56,03	25	55,20	- 0,83	66	46	15,80	46	18,00	+	2,20	16 2,30
14		5	30	4,78	30	4,20	- 0,58	66	43	5,01	43	7,00	+	1,99	15 59,60
15		5	34	14,08	34	13,40	- 0,68	66	40	19,76	40	20,00	+	0,24	16 0,46
16		5	38	23,34	38	22,50	- 0,84	66	37	56,16	38	1,00	+	4,84	16 1,72
17								66	36	1,30	36	4,00	+	2,70	15 59,93
18	5	46	41,46	46	41,20	- 0,26	66	34	27,43	34	33,00	+	5,57	16 1,61	
22							66	32	29,86	32	32,00	+	2,14	16 2,00	
23							66	33	2,38	33	4,00	+	1,62		
24							66	34	6,15	34	2,00	-	4,15	16 1,72	
26							66	37	12,21	37	10,00	-	2,21	15 58,30	
27							66	39	18,10	39	23,00	+	4,90	16 1,24	
28							66	41	57,75	41	57,00	-	0,75		
July	30						66	48	19,61	48	23,00	+	3,39	16 2,10	
	2						66	56	27,42	56	25,00	-	2,42	16 1,68	
	3						67	1	2,13	1	3,00	+	0,87	16 0,10	
	4						67	6	7,63	6	5,00	-	2,63	16 3,56	
	5						67	11	27,68	11	30,00	+	2,32	16 2,11	
	6						67	17	15,99	17	20,00	+	4,01	15 57,86	
	7						67	23	27,86	23	32,00	+	4,14	15 54,90	
	16						67	36	44,73	38	43,00	-	1,73	15 58,07	
	20						69	18	48,33	18	53,00	+	4,67	15 58,04	
	24	8	14	5,86	14	5,10	- 0,76	69	6	39,55	6	35,00	-	4,55	
	25	8	18	3,34	18	2,80	- 0,54	70	19	27,21	19	21,00	-	6,21	16 0,60
26	8	22	0,19	21	59,80	- 0,39	70	32	35,49	32	28,00	-	7,49	16 1,02	
28	8	29	52,89	29	52,50	- 0,39	70	59	44,48	59	37,00	-	7,48	16 1,86	
30							71	28	3,32	28	3,00	-	0,32	16 3,48	
August	31						71	42	44,17	42	42,00	-	2,17	16 0,58	
	1						71	57	43,37	57	42,00	-	1,37	16 2,13	

RESULT OF OBSERVATIONS IN 1832 AND 1833.

1832	Observed R.A.			A.R. from Nautical Almanac.		Error of Tables.	Observed N.P.D.			N.P.D. from Nautical Almanac.		Error of Tables.	Mean Semi-diameter.		
	h.	m.	s.	m.	s.		°	'	"	°	'		"	'	"
August	2						72	12	59,72	12	57,00	-	2,72		
	3						72	28	35,78	28	29,00	-	6,78	16 2,34	
	5						73	0	30,57	0	28,00	-	2,57	16 0,18	
	6						73	16	49,42	16	51,00	+	1,58	16 3,90	
	7						73	33	29,44	33	29,00	-	0,44	16 2,42	
	8						73	50	25,78	50	24,00	-	1,78	16 2,65	
	11						74	42	48,86	42	42,00	-	6,86	15 58,94	
	12						75	0	47,12	0	38,00	-	9,12	16 0,58	
	13	9	31	23,86	31	23,50	- 0,36	75	18	57,04	18	48,00	-	9,04	16 2,92
	14	9	35	9,95	35	9,10	- 0,85	75	37	16,45	37	13,00	-	3,45	15 58,14
	17	9	46	24,37	46	23,40	- 0,97	76	33	50,58	33	47,00	-	3,58	16 0,48
	18	9	50	7,78	50	7,00	- 0,78	76	53	7,52	53	4,00	-	3,52	16 0,20
	19	9	53	50,86	53	50,20	- 0,66	77	12	34,40	12	34,00	-	0,40	
	20	9	57	33,69	57	32,90	- 0,79	77	32	15,95	32	17,00	+	1,05	16 1,50
	21	10	1	15,59	1	15,30	- 0,29	77	52	9,12	52	11,00	+	1,88	16 2,80
	22	10	4	57,57	4	57,10	- 0,47	78	12	17,81	12	18,00	+	0,19	16 2,01
	23	10	8	38,75	8	38,50	- 0,25	78	32	34,72	32	35,00	+	0,28	15 59,88
	25	10	16	0,87	16	0,20	- 0,67	79	13	50,09	13	43,00	-	7,09	
	27	10	23	20,62	23	20,00	- 0,62	79	55	35,70	55	31,00	-	4,70	16 3,90
	28	10	27	0,20	26	59,30	- 0,90	80	16	40,24	16	40,00	-	0,24	16 0,60
29	10	30	39,24	30	38,40	- 0,84	80	38	4,17	37	59,00	-	5,17		
30	10	34	17,75	34	17,00	- 0,75	80	59	26,61	59	26,00	-	0,61		
31	10	37	56,23	37	55,30	- 0,93	81	21	0,28	21	1,00	+	0,72	16 1,68	
Sept.	2						82	4	34,16	4	38,00	+	3,84	15 59,20	
	4	10	52	25,74	52	25,30	- 0,44	82	48	42,95	48	43,00	+	0,05	16 0,82
	6	10	59	39,23	59	38,80	- 0,43	83	33	16,04	33	17,00	+	0,96	15 59,30
	7	11	3	15,93	3	15,20	- 0,73	83	55	43,26	55	44,00	+	0,74	16 2,54
	9	11	10	27,91	10	27,30	- 0,61	84	41	2,89	40	54,00	-	8,89	16 0,12
	10	11	14	4,16	14	3,10	- 1,06	85	3	47,45	3	38,00	-	9,45	16 3,85
	14							86	35	24,45	35	20,00	-	4,45	
	15	11	32	1,05	32	0,70	- 0,35	86	58	30,78	58	26,00	-	4,78	16 4,16
	16	11	35	36,33	35	36,10	- 0,23	87	21	38,43	21	35,00	-	3,43	16 0,18
	20	11	49	58,33	49	57,90	- 0,43	88	54	42,27	54	42,00	-	0,27	16 0,22
	21	11	53	34,02	53	33,50	- 0,52	89	18	7,92	18	5,00	-	2,92	16 0,95
	22	11	57	12,62	57	9,10									
	23	12	0	45,77	0	45,00	- 0,77	90	5	1,09	4	53,00	-	8,09	15 59,74
	24	12	4	21,46	4	21,00	- 0,46	90	28	22,24	28	19,00	-	3,24	16 1,70
25	12	7	57,55	7	57,20	- 0,35	90	51	48,97	51	45,00	-	3,97	16 2,28	
26	12	11	33,26	11	33,30	+	91	15	13,55	15	11,00	-	2,55	16 3,98	
27	12	15	10,66	15	9,80	- 0,86	91	38	35,55	38	36,00	+	0,45	16 1,61	
28	12	18	46,92	18	46,40	- 0,52	92	2	2,24	2	1,00	-	1,24		
30	12	26	0,94	26	0,20	- 0,74	92	48	47,45	48	48,00	+	0,55	16 2,07	
October	1	12	29	37,70	29	37,80	- 0,10	93	12	2,44	12	8,00	+	5,56	15 59,43
	2	12	33	16,41	33	15,20	- 1,21	93	35	28,25	35	27,00	-	1,25	16 4,17
	4	12	40	32,07	40	31,50	- 0,57	94	21	58,44	21	55,00	-	3,44	
	5							94	45	10,97	45	5,00	-	5,97	16 1,35
	6							95	8	13,54	8	10,00	-	3,54	16 1,72
	7	12	51	28,31	51	28,10	- 0,21	95	31	16,87	31	12,00	-	4,87	16 0,84
	8	12	55	9,28	55	7,90	- 1,38	95	54	12,56	54	10,00	-	2,56	16 1,55
	9	12	58	48,97	58	48,00	- 0,97	96	17	8,14	17	3,00	-	5,14	16 3,85
	11	13	6	10,42	6	9,50	- 0,92	97	2	36,15	2	35,00	-	1,15	16 3,10
	12	13	9	52,25	9	51,30	- 0,95	97	25	17,91	25	12,00	-	5,91	16 4,56
	13	13	13	34,20	13	33,20	- 1,00	97	47	42,99	47	43,00	+	0,01	16 1,51

RESULT OF OBSERVATIONS IN 1832 AND 1833.

1833	Observed A.R.			A.R. from Nautical Almanac.		Error of Tables.	Observed N.P.D.			N.P.D. from Nautical Almanac.		Error of Tables.	Mean Semi-diameter.				
	h.	m.	s.	m.	s.		°	'	"	'	"		"	'	"		
June	8						67	9	35,10	9	33,00	-	2,10	16	1,92		
	9	5	8	19,65	8	19,60	-	0,05	67	4	14,97	4	16,00	+	1,03	16	2,90
	11	5	16	36,61	16	36,30	-	0,31	66	54	51,15	54	52,00	+	0,85		
	12	5	20	45,15	20	44,90	-	0,25	66	50	48,41	50	45,00	-	3,41	16	3,18
	13	5	24	54,22	24	53,80	-	0,42	66	47	6,46	47	6,00	-	0,64	16	1,94
	14	5	29	3,14	29	2,90	-	0,24	66	43	49,60	43	49,00	-	0,60	16	3,16
	19							66	33	37,07	33	35,00	-	2,07	16	0,72	
	20							66	32	45,59	32	47,00	+	1,41	16	3,48	
	21							66	32	23,63	32	24,00	+	0,37	16	2,37	
	22							66	32	28,80	32	26,00	-	2,80	16	1,60	
	23	6	6	29,24	6	28,80	-	0,44	66	32	50,98	32	52,00	+	1,02	15	59,34
	25							66	35	0,36	34	58,00	-	2,36			
	26							66	36	39,12	36	38,00	-	1,12	16	2,25	
	27							66	38	46,12	38	43,00	-	3,12	16	2,16	
	28							66	41	14,50	41	14,00	-	0,50	16	3,58	
29	6	31	24,35	31	23,60	-	0,75	66	44	11,45	44	8,00	-	3,45	16	1,66	
30							66	47	30,50	47	27,00	-	3,50				
July	1						66	51	12,76	51	9,00	-	3,76	16	1,67		
	2	6	43	48,73	43	48,40	-	0,33	66	55	20,33	55	17,00	-	3,33	16	0,94
	3	6	47	56,14	47	56,20	+	0,06	66	59	51,96	59	49,00	-	2,96	16	2,36
	5	6	56	10,97	56	10,80	-	0,17	67	10	2,41	10	4,00	+	1,59	16	1,24
	7							67	22	1,28	21	54,00	-	7,28	16	1,25	
	8	7	8	30,58	8	30,40	-	0,18	67	31	24,40	28	25,00	+	0,60		
	9							67	35	20,93	35	20,00	-	0,93	16	0,26	
	12	7	24	51,43	24	51,40	-	0,03	67	58	15,41	58	21,00	+	5,59		
	13	7	28	55,82	28	55,60	-	0,22	68	6	46,70	6	47,00	+	0,30		
	15	7	37	3,29	37	2,50	-	0,79	68	24	41,77	24	46,00	+	4,23	16	0,02
	16							68	34	21,94	34	18,00	-	3,94	16	3,98	
	17							68	44	12,97	44	14,00	+	1,03	16	1,92	
	19							68	54	32,08	54	31,00	-	1,08	16	3,70	
	19							69	5	7,40	5	8,00	+	0,60	16	4,40	
	20							69	16	9,90	16	8,00	-	1,90	16	1,26	
21							69	27	28,92	27	28,00	-	0,92	15	58,18		
23							69	51	17,19	51	11,00	-	6,19				
25	8	17	6,42	17	5,90	-	0,52										
27	8	24	59,85	24	59,50	-	0,35	70	42	36,84	42	37,00	+	0,16	16	1,94	
28	8	28	55,62	28	55,40	-	0,22	70	56	18,15	56	15,00	-	3,15	16	2,67	
29							71	10	17,36	10	12,00	-	5,36	16	1,94		
30							71	24	27,07	24	29,00	+	1,93	15	59,98		
August	1						71	54	3,50	53	59,00	-	4,50	16	1,58		
	2						72	9	7,58	9	10,00	+	2,42	16	3,65		
	3						72	24	39,17	24	37,00	-	2,17	6	2,23		
	4	8	56	9,79	56	9,70	-	0,09	72	40	20,96	40	24,00	+	3,04	16	3,16
	5	9	0	1,20	0	0,80	-	0,40	72	56	25,53	56	26,00	+	0,47	16	1,40
	6	9	3	51,52	3	51,30	-	0,22	73	12	41,00	12	45,00	+	4,00	15	59,18
	7							73	29	24,87	29	21,00	-	3,87			
	8							73	46	13,25	46	12,00	-	1,25			
	9							74	3	20,28	3	19,00	-	1,28	16	1,68	
	10	9	19	7,60	19	7,40	-	0,20	74	20	46,67	20	42,00	-	4,67	16	3,20
	11	9	22	55,33	22	55,10	-	0,23	74	38	25,82	38	20,00	-	5,82	16	1,24
	12	9	26	42,71	26	42,70	-	0,01	74	56	16,70	56	13,00	-	3,70	15	59,95
	13	9	30	29,74	30	28,90	-	0,84									
	14							74	32	42,15	32	44,00	+	1,85	16	0,00	
	15	9	38	0,76	38	0,30	-	0,46	74	51	21,86	51	19,00	-	2,86	16	2,51

1832	Observed R.A.			A.R. from Nautical Almanac.		Error of Tables.	Observed N.P.D.			N.P.D. from Nautical Almanac.		Error of Tables.	Mean Semi-diameter.				
	h.	m.	s.	m.	s.		s.	°	'	"	'		"	'	"		
Nov.	6	14	44	42,88	44	42,30	-	0,58	105	56	47,82	56	48,00	+	0,18	16	2,43
	14								108	12	10,38	12	15,00	+	4,62	16	3,08
	16								108	42	56,50	43	2,00	+	5,50		
	17	14	29	25,78	29	25,60	-	0,18								16	1,70
	18	14	33	35,32	33	34,70	-	0,62	109	12	23,54	12	29,00	+	5,46	16	0,66
	19	14	37	44,48	37	44,40	-	0,08	109	26	42,72	26	43,00	+	0,28	16	1,15
	20								109	40	35,26	40	35,00	-	0,26		
	22	15	50	19,06	50	18,60	-	0,46	110	7	16,74	7	13,00	-	3,74		
	23	15	54	31,45	54	31,40	-	0,05	110	19	59,62	19	59,00	-	0,62	16	2,53
	24								110	32	27,00	32	23,00	-	4,00		
	25								110	44	21,60	44	23,00	+	1,40		
	28								111	18	3,82	18	4,00	+	0,18	16	0,44
	29								111	28	26,70	28	31,00	+	4,30	16	1,75
Dec.	1								111	48	4,13	48	8,00	+	3,87	16	3,22
	2								111	57	19,23	57	20,00	+	0,77	16	0,77
	3	16	37	21,84	37	21,30	-	0,54	112	6	4,65	6	6,00	+	1,35	16	2,34
	5								112	22	19,90	22	22,00	+	2,10	15	57,60
	6	16	50	26,18	50	25,30	-	0,88	112	29	46,61	29	50,00	+	3,39	15	59,68
	7	16	54	48,07	54	47,70	-	0,37	112	36	50,59	36	50,00	-	0,59	15	57,90
	8	16	59	10,94	59	10,70	-	0,24	112	43	20,30	43	29,00	+	8,70	16	3,36
	9								112	49	31,06	49	37,00	+	5,94	16	2,58
	10								112	55	12,84	55	10,00	-	2,84	16	2,33
	11	17	11	22,42	12	22,20	-	0,22	113	0	29,49	0	34,00	+	4,51	16	3,23
	12	17	16	47,39	16	46,90	-	0,49	113	5	13,94	5	21,00	+	7,06	16	2,60
	14								113	13	25,90	13	33,00	+	7,10		
	15								113	16	51,90	16	57,00	+	5,10	16	0,15
	18								113	24	17,72	24	23,00	+	5,28		
	19	17	47	48,04	47	47,50	-	0,54	113	25	50,71	25	53,00	+	2,29		
	20	17	52	15,12	52	14,00	-	1,12	113	26	53,65	26	56,00	+	2,35	6	0,15
	22								113	27	32,56	27	39,00	+	6,44	16	0,00
	23	18	5	33,90	5	33,60	-	0,30	113	27	13,71	27	19,00	+	5,29	16	2,32
	24								113	26	20,51	26	29,00	+	8,49	16	1,84
	26	18	18	53,37	18	53,20	-	0,17	113	23	18,50	23	24,00	+	5,50	6	1,75
	27	18	23	19,90	23	19,50	-	0,40	113	21	5,75	21	10,00	+	4,25	16	3,56
	29	18	32	12,21	32	11,60	-	0,61	113	15	9,69	15	17,00	+	7,31	16	3,14
	30	18	36	37,59	36	37,30	-	0,29	113	11	32,74	11	39,00	+	6,26	16	1,08
	31	18	41	3,31	41	3,00	-	0,31	113	7	26,81	7	32,00	+	5,19	16	0,46

Taking the mean of the above measures of the Sun's Semi-diameter for 1832 we have from 258 Observations...16' 1",52
 ————— 1833 ————— 257 ————— ...16' 1",30
 differing very little from that found from the Observations of 1831.*

We will now select from the above, those observations which are made near to the Solstices, for the determination of the value of the obliquity of the Ecliptic as follows.

* Vol. I Page 69, ☉s. Mean Semi-diameter for 16' 0",15 read 16' 1",15,

Observations of the Sun made near to the Summer Solstices of 1832 and 1833, applied to the determination of the obliquity of the Eceleptic.

1832	N. P. D.	Reduction.	Sun's Latitude.	Solstitial N.P.D.	Correction for		Mean N.P.D. of the Solstice reduced to January 1.
					D r. Nut.	Gr. Nut + t. 0°, 46 365	
May	30 68 12 49 40	1 40 26,74	—	0,89 66 32 21,77	—	5,45	0,58 66 32 15,74
	31 68 4 1,99	1 31 42,77	—	0,78 66 32 18,44	—	5,44	0,60 66 32 12,40
June	1 67 55 44,58	1 23 22,12	—	0,64 66 32 21,82	—	5,43	0,61 66 32 15,78
	2 67 47 43,95	1 15 24,23	—	0,48 66 32 19,24	—	5,43	0,63 66 32 13,18
	3 67 40 13,26	1 7 49,63	—	0,33 66 32 23,30	—	5,42	0,64 66 32 17,24
	4 67 33 0 66	1 0 38,25	—	0,17 66 32 22,24	—	5,41	0,65 66 32 16,18
	5 67 26 14,34	0 53 50,42	—	0,02 66 32 23,90	—	5,40	0,66 66 32 17,84
	6 67 19 51,10	0 47 26,50	+	0,11 66 32 24,71	—	5,39	0,67 66 32 18 65
	7 67 13 48,35	0 41 26,23	+	0 25 66 32 22,37	—	5,38	0,68 66 32 16,31
	8 67 8 3,77	0 35 50,30	+	0,33 66 32 18,80	—	5,38	0,69 66 32 12,73
	9 67 3 4 85	0 30 38,07	+	0,40 66 32 27,18	—	5,37	0,70 66 32 21,11
	10 66 58 8 42	0 25 50,08	+	0,42 66 32 18 76	—	5,36	0,71 66 32 12,69
	11 66 53 50,47	0 21 26,12	+	0,43 66 32 24,78	—	5,36	0,71 66 32 12,71
	12 66 49 48,77	0 17 27,08	+	0,39 66 32 22,08	—	5,35	0,72 66 32 16,01
	13 66 46 15,80	0 13 52,37	+	0,32 66 32 23,75	—	5,34	0,73 66 32 17,68
	14 66 43 5,01	0 10 41,97	+	0 20 66 32 23,24	—	5,34	0,73 66 32 17 17
	15 66 40 19,76	0 7 56,09	+	0,10 66 32 23,77	—	5,33	0,74 66 32 17,70
	16 66 37 56,16	0 5 35,05	—	0,06 66 32 21,05	—	5,32	0,74 66 32 14 99
	17 66 36 1,30	0 3 38 63	—	0 23 66 32 22,44	—	5,32	0,75 66 32 16,37
	18 66 34 27,43	0 2 6,94	—	0,37 66 32 20,12	—	5,31	0,75 66 32 14,05
	22 66 32 29 86	0 0 7,95	—	0 85 66 32 21,06	—	5,28	0,75 66 32 15,03
	23 66 33 2 38	0 0 40,25	—	0,88 66 32 21,25	—	5,27	0,76 66 32 15 22
	24 66 34 6,15	0 1 37,28	—	0,89 66 32 27 98	—	5,27	0,76 66 32 21,95
	26 66 37 12,21	0 4 45,67	—	0,81 66 32 25 73	—	5,26	0,76 66 32 19,71
27 66 39 18,10	0 6 56,78	—	0,74 66 32 20,38	—	5,25	0,76 66 32 14,37	
28 66 41 57,75	0 9 32,85	—	0,59 66 32 24 31	—	5,24	0,75 66 32 18,32	
30 66 48 19,61	0 15 58,10	—	0,31 66 32 21,20	—	5,23	0,75 66 32 15,22	
July	2 66 56 27,42	0 24 0,90	—	0 00 66 32 26,52	—	5,22	0,74 66 32 20 56
	3 67 1 2,13	0 28 38,57	+	0,15 66 32 23,71	—	5,21	0,74 66 32 17,76
	4 67 6 7,63	0 33 40,30	+	0,26 65 32 27,59	—	5,20	0,73 66 32 21 66
	5 67 11 27,68	0 39 5,92	+	0,36 66 32 22 12	—	5,19	0,72 66 32 16 21
	6 67 17 15 99	0 44 55 45	+	0,42 66 32 20 96	—	5,18	0,71 66 32 15 07
	7 67 23 27,86	0 51 8,68	+	0,46 66 32 19,74	—	5,17	0,70 66 32 13,77
1833							
May	30 68 14 56 42	1 42 38,98	+	0,28 66 32 18,72	—	2,62	0,58 66 32 15 52
	31 68 6 9,50	1 33 49,77	+	0,25 66 32 19,98	—	2,61	0 60 66 32 16,77
June	1 67 57 43 95	1 25 23,25	+	0,16 66 32 20,86	—	2,60	0 61 66 32 17,65
	2 67 49 39 37	1 17 20,23	+	0,06 66 32 19,20	—	2,59	0,63 66 32 15,98
	3 67 41 58 90	1 9 39,97	—	0,05 66 32 18,88	—	2,58	0,64 66 32 15 66
	4 67 34 46,36	1 2 23,22	—	0,12 66 32 23,02	—	2,57	0 65 66 32 19,80
	5 67 27 49,58	0 55 29,80	—	0,30 66 32 19 48	—	2,57	0 66 66 32 16,25
	6 67 21 22,53	0 49 0,02	—	0,44 66 32 22,07	—	2,56	0 67 66 32 18,84
	7 67 15 12,89	0 42 53 90	—	0,55 66 32 18,44	—	2,55	0,68 66 32 15,21
	8 67 9 35,10	0 37 11,80	—	0,66 66 32 22,64	—	2,54	0 69 66 32 19 41
	9 67 4 14,97	0 31 53,58	—	0,77 66 32 20,62	—	2,53	0 70 66 32 17,39
	11 66 54 51,15	0 22 29,62	—	0,85 66 32 20,68	—	2,52	0,71 66 32 17,45
	12 66 50 48,41	0 18 24,09	—	0,86 66 32 23,46	—	2,51	0,72 66 32 20,23
	13 66 47 6,64	0 14 43 08	—	0,85 66 32 22,71	—	2,50	0,73 66 32 19,48
	14 66 43 49,60	0 11 26,56	—	0,78 66 32 22,26	—	2,49	0,73 66 32 19,04

1833	N. P. D.			Reduction.			Sun's Latitude.	Solstitial N.P.D.			Correction for		Mean N.P.D. of the Solstice reduced to January 1.						
											Dr. Nut.	Cr. Nut. t 0",46 + 365							
June	19	66	33	37,07	0	1	13,92	—	0,20	66	32	22,95	—	2,45	—	0,75	66	32	19,75
	20	66	32	45,59	0	0	23,75	—	0,09	66	32	21,75	—	2,44	—	0,75	66	32	18,56
	21	66	32	23,63	0	0	2,44	+	0,06	66	32	21,25	—	2,43	—	0,76	66	32	18,06
	22	66	32	28,80	0	0	3,87	+	0,17	66	32	25,10	—	2,42	—	0,76	66	32	21,92
	23	66	32	50,98	0	0	30,18	+	0,25	66	32	21,05	—	2,41	—	0,76	66	32	17,88
	25	66	35	0,36	0	2	37,15	+	0,31	66	32	23,52	—	2,40	—	0,76	66	32	20,36
	26	66	36	39,12	0	4	17,67	+	0,30	66	32	21,75	—	2,39	—	0,76	66	32	18,60
	27	66	38	46,12	0	6	22,88	+	0,27	66	32	23,51	—	2,38	—	0,76	66	32	20,37
	28	66	41	14,50	0	8	52,72	+	0,20	66	32	21,98	—	2,37	—	0,75	66	32	18,86
	29	66	44	11,45	0	11	47,10	+	0,10	66	32	24,45	—	2,36	—	0,75	66	32	21,34
30	66	47	30,50	0	15	5,81	+	0,04	66	32	24,73	—	2,35	—	0,75	66	32	21,63	
July	1	66	51	12,76	0	18	49,00	—	0,12	66	32	23,64	—	2,34	—	0,74	66	32	20,56
	2	66	55	20,33	0	22	56,44	—	0,25	66	32	23,64	—	2,33	—	0,74	66	32	20,57
	3	66	59	51,96	0	27	27,98	—	0,38	66	32	23,60	—	2,32	—	0,73	66	32	20,55
	5	67	10	2,41	0	37	43,30	—	0,62	66	32	18,49	—	2,30	—	0,72	66	32	15,47
	7	67	22	1,28	0	49	34,80	—	0,78	66	32	25,70	—	2,29	—	0,70	66	32	22,71
	8	67	28	24,40	0	56	4,65	—	0,82	66	32	18,93	—	2,28	—	0,69	66	32	15,96
	9	67	35	20,93	1	2	58,81	—	0,82	66	32	21,30	—	2,27	—	0,68	66	32	18,35

And further we have:

Observations of the Sun made near to the Winter Solstice of 1832 and 1833, applied to the determination of the obliquity of the Ecliptic.

1832	N. P. D.			Reduction.			Sun's Latitude.	Solstitial N.P.D.			Correction for		Mean N.P.D. of the Solstice reduced to January 1.							
											Dr. Nut.	Cr. Nut. t 0",46 + 365								
Jan.	1	112	5	10,16	0	22	17,90	+	0,74	113	27	28,80	+	6,46	+	0,50	113	27	35,76	
	2	113	0	25,66	0	27	5,68	+	0,62	113	27	31,96	+	6,46	+	0,49	113	27	38,91	
	3	112	55	9,19	0	32	20,98	+	0,51	113	27	30,68	+	6,45	+	0,48	113	27	37,61	
	4	112	49	25,77	0	38	3,80	+	0,37	113	27	29,94	+	6,44	+	0,47	113	27	36,83	
	6	112	36	43,55	0	50	50,74	+	0,07	113	27	34,36	+	6,43	+	0,46	113	27	41,25	
	7	112	29	39,23	0	57	54,64	—	0,09	113	27	33,78	+	6,43	+	0,45	113	27	40,66	
	8	112	22	4,23	1	5	25,20	—	0,24	113	27	29,19	+	6,42	+	0,44	113	27	36,05	
	9	112	14	11,47	1	13	22,27	—	0,38	113	27	33,36	+	6,41	+	0,43	113	27	40,20	
	10	112	5	46,20	1	21	45,94	—	0,47	113	27	31,67	+	6,40	+	0,42	113	27	38,49	
	11	111	56	57,08	1	30	34,67	—	0,55	113	27	31,20	+	6,39	+	0,41	113	27	38,00	
	12	111	47	47,83	1	39	49,30	—	0,58	113	27	36,55	+	6,38	+	0,40	113	27	43,33	
	13	111	38	2,25	1	49	29,18	—	0,59	113	27	30,84	+	6,37	+	0,40	113	27	37,61	
	14	111	28	0,62	1	59	34,57	—	0,55	113	27	34,64	+	6,36	+	0,39	113	27	41,39	
	15	111	17	30,45	2	10	3,90	—	0,51	113	27	33,84	+	6,35	+	0,37	113	27	40,56	
	18	110	43	35,60	2	43	58,52	—	0,16	113	27	33,95	+	6,34	+	0,31	113	27	40,61	
	19	110	31	28,91	2	56	4,98	—	0,02	113	27	33,87	+	6,33	+	0,30	113	27	40,50	
	21	110	6	10,93	3	21	25,43	+	0,26	113	27	36,62	+	6,32	+	0,27	113	27	43,21	
	Nov.	21	109	57	12,86	3	30	25,78	+	0,81	113	27	39,45	+	4,14	+	0,66	113	27	44,25
		22	110	10	16,30	3	17	22,52	+	0,75	113	27	39,57	+	4,13	+	0,68	113	27	44,38
		23	110	22	53,92	3	4	41,40	+	0,70	113	27	36,02	+	4,13	+	0,69	113	27	40,84
24		110	35	16,84	2	52	22,67	+	0,58	113	27	40,09	+	4,12	+	0,71	113	27	44,92	
25		110	47	4,37	2	40	26,83	+	0,46	113	27	31,66	+	4,11	+	0,72	113	27	36,49	
26		110	58	39,92	2	28	55,10	+	0,31	113	27	35,33	+	4,11	+	0,74	113	27	40,18	
27		111	9	50,58	2	17	46,50	+	0,17	113	27	37,25	+	4,10	+	0,75	113	27	42,10	

1833	N. P. D.			Reduction.	Sun's Latitude.	Solstitial N.P.D.	Correction for		Mean N.P.D. of the Solstice reduced to January 1.	
							Dr. Nut.	Gr. Nut t. 0",46 + 365		
Dec. 14	113	13	25,90	0 14	7,12	+ 0,17	113 27 33,19	+ 0,97	+ 0,96	113 27 35,12
15	113	16	51,90	0 10	42,95	+ 0,05	113 27 34,90	+ 0,96	+ 0,97	113 27 36,83
18	113	24	17,72	0 3	17,62	- 0,20	113 27 35,14	+ 0,93	+ 0,99	113 27 37,06
19	113	25	50,71	0 1	46,73	- 0,23	113 27 37,21	+ 0,92	+ 0,99	113 27 39,12
20	113	26	53,65	0 0	43,16	- 0,24	113 27 36,57	+ 0,91	+ 1,00	113 27 38,48
22	113	27	32,56	0 0	0,92	- 0,14	113 27 33,34	+ 0,90	+ 1,00	113 27 35,24
23	113	27	13,71	0 0	22,10	- 0,04	113 27 35,77	+ 0,89	+ 1,00	113 27 37,66
24	113	26	20,51	0 1	11,90	+ 0,07	113 27 32,48	+ 0,88	+ 0,99	113 27 34,35
26	113	23	18,50	0 4	16,00	+ 0,29	113 27 34,79	+ 0,86	+ 0,98	113 27 36,63
27	113	21	5,75	0 6	30,25	+ 0,44	113 27 36,44	+ 0,85	+ 0,98	113 27 38,27
29	113	15	9,69	0 12	23,40	+ 0,68	113 27 33,77	+ 0,83	+ 0,97	113 27 35,57

Taking the means we have :

Mean Obliquity January 1, 1832.

From 33 Observations of the Summer Solstice of 1832.....	23 27 43,59
From 33 ————— of 1833 23° 27' 41",29 — 0",46....	23 27 40,83

Obliquity from Summer Solstices of 1832 and 1833 = 23 27 42,21

From 40 Observations at the Winter Solstice of 1832-33.....	23 27 39,20
From 47 ————— of 1833-34 23° 27' 39",07 — 0",46..	23 27 38,52

Obliquity from Winter Solstices of 1832 and 1833 = 23 27 38,83

Finally we have from the means of the whole..... 23 27 40 52

We will now from the Observations of the Sun near to the time of the Equinoxes compare the Right Ascension as determined by the Transit Instrument with that computed from the observed N. P. D.

Observations of the Sun made near to the Vernal Equinox in 1832 and 1833, applied to the determination of the error of the assumed Equinoctial Point.

1832	Reduced N.P.D. of the Sun.			Correc- tion.	N. P. D. reduced on account of Sun's Latitude	Compu'ed A. R.	Observed A.R.	Error of Eq. Point.	REMARKS.
	°	'	"	"	°	'	"		
February 10	104	38	3,00	- 0,69	104 38 2,31	h. m. s.	21 32 2,41	32 1,90	- 0,51
11	104	18	41,46	- 0,64	104 18 40,82	21 35 59,16	35 59,42	+ 0,26	
12	103	58	58,84	- 0,55	103 58 58,29	21 39 56,50	39 56,67	+ 0,17	
14	103	19	0,46	- 0,29	103 19 0,17	21 47 47,29	47 46,51	- 0,78	
15	102	58	40,60	+ 0,18	102 58 40,78	21 51 41,66	51 41,44	- 0,22	

RESULT OF OBSERVATIONS IN 1832 AND 1833.

1832	Reduced N.P.D. of the Sun.			Correc- tion.	N. P. D. reduced on account of Sun's Latitude.	Computed A. R.			Observed A.R.			Error of Eq. Point.	REMARKS.				
	°	'	"	"	°	'	"	<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>m.</i>	<i>s.</i>	<i>s.</i>				
February	17	102	17	24,55	+	0,21	102	17	24,76	21	59	28,26	59	27,62	-	0,64	
	18	101	56	29,40	+	0,23	101	56	29,63	22	3	20,34	3	19,98	-	0,36	
	20	101	14	6,88	+	0,45	101	14	7,33	22	11	2,18	11	2,05	-	0,13	
	21	100	52	35,17	+	0,50	100	52	35,67	22	14	52,60	14	52,05	-	0,55	
	22	100	30	59,96	+	0,53	100	31	0,49	22	18	41,77	18	41,41	-	0,36	
	23	100	9	14,08	+	0,55	100	9	14,63	22	22	30,20	22	30,61	+	0,41	
	24	99	47	16,27	+	0,52	99	47	16,79	22	26	18,47	26	18,67	+	0,20	
	25	99	25	5,41	+	0,46	99	25	5,87	22	30	6,85	30	6,02	-	0,83	
	26	99	2	47,56	+	0,35	99	2	47,91	22	33	54,39	33	53,64	-	0,75	
	March	1	97	32	35,88	-	0,20	97	32	35,68	22	48	56,18	48	55,93	-	0,25
		2	97	9	40,18	-	0,38	97	9	39,80	22	52	41,22	52	40,20	-	1,02
		3	96	46	46,24	-	0,52	96	46	45,72	22	56	24,52	56	24,18	-	0,34
		4	96	23	43,27	-	0,63	96	23	42,64	23	0	7,87	0	7,21	-	0,66
		5	96	0	38,10	-	0,69	96	0	37,41	23	3	50,27	3	50,45	+	0,18
		6	95	37	17,60	-	0,75	95	37	16,85	23	7	33,89	7	33,06	-	0,83
		7	95	14	8,80	-	0,80	95	14	8,00	23	11	14,53	11	15,11	+	0,58
8		94	50	45,89	-	0,81	94	50	45,08	23	14	56,36	14	56,67	+	0,31	
9		94	27	20,70	-	0,76	94	27	19,94	23	18	37,56	18	37,99	+	0,43	
10		94	3	45,61	-	0,69	94	3	44,92	23	22	19,43	22	18,97	-	0,46	
11		93	40	16,45	-	0,59	93	40	15,86	23	25	59,56	26	0,07	+	0,51	
12		93	16	34,74	-	0,46	93	16	34,28	23	29	40,93	29	39,62	-	1,31	
13		92	53	3,09	-	0,30	92	53	2,79	23	33	20,07	33	19,43	-	0,64	
14		92	29	24,00	-	0,17	92	29	23,83	23	36	59,81	36	59,13	-	0,68	
15		92	5	48,99	-	0,03	92	5	48,96	23	40	38,43	40	38,44	+	0,01	
16		91	42	4,68	+	0,11	91	42	4,79	23	44	18,12	44	17,49	-	0,63	
April	17	91	18	26,27	+	0,23	91	18	26,50	23	47	56,55	47	56,47	-	0,08	
	18	90	54	47,50	+	0,35	90	54	47,85	23	51	34,77	51	35,02	+	0,25	
	19	90	31	7,53	+	0,42	90	31	7,95	23	55	13,02	55	13,50	+	0,48	
	20	90	7	25,25	+	0,47	90	7	25,72	23	58	51,53	58	52,11	+	0,58	
	21	89	43	42,84	+	0,47	89	43	43,31	0	2	30,04	2	30,28	+	0,24	
	22	89	20	6,12	+	0,44	89	20	6,56	0	6	7,73	6	8,27	+	0,54	
	23	88	56	24,08	+	0,39	88	56	24,47	0	9	46,38	9	46,31	-	0,07	
	24	88	32	40,98	+	0,29	88	32	41,27	0	13	25,39	13	24,39	-	1,00	
	25	88	9	11,58	+	0,17	88	9	11,75	0	17	2,58	17	2,32	-	0,26	
	26	87	45	38,10	+	0,04	87	45	38,14	0	20	40,75	20	40,24	-	0,51	
	27	87	22	8,80	-	0,11	87	22	8,69	0	24	18,71	24	18,21	-	0,50	
	28	86	58	50,12	-	0,26	86	58	49,86	0	27	55,53	27	56,14	+	0,61	
	30	86	12	1,12	-	0,54	86	12	0,58	0	35	12,77	35	12,60	-	0,17	
	31	85	48	52,19	-	0,67	85	48	51,52	0	38	50,03	38	50,74	+	0,71	
	1	85	25	35,96	-	0,78	85	25	35,18	0	42	29,22	42	29,22	+	0,00	
	2	85	2	39,39	-	0,84	85	2	38,55	0	46	6,21	46	7,69	+	1,48	
3	84	39	36,02	-	0,86	84	39	35,16	0	49	45,22	49	46,12	+	0,90		
4	84	16	35,87	-	0,84	84	16	35,03	0	53	24,77	53	24,51	-	0,26		
5	83	53	45,69	-	0,81	83	53	44,88	0	57	3,84	57	3,39	-	0,45		
6	83	31	1,49	-	0,75	83	31	0,74	1	0	43,15	0	42,48	-	0,67		
15	80	12	31,85	+	0,35	80	12	32,20	1	33	43,40	33	43,40	+	0,00		
16	79	51	10,54	+	0,38	79	51	10,92	1	37	25,33	37	25,75	+	0,42		
1833																	
February	10	104	23	34,23	+	0,56	104	23	34,79	21	34	59,87	34	59,51	-	0,36	
	12	103	44	4,78	+	0,47	103	44	5,25	21	42	53,67	42	53,30	-	0,37	
	13	103	24	1,74	+	0,38	103	24	2,12	21	46	48,98	46	48,61	-	0,37	
	15	102	43	13,18	+	0,18	102	43	13,36	21	54	38,10	54	36,82	-	1,28	
	16	102	22	39,38	+	0,05	102	22	39,43	21	58	29,85	58	30,01	+	0,16	

1833	Reduced N.P.D. of the Sun.			Correc- tion.	N. P. D. Reduced on account of Sun's Latitude			Computed A. R.			Observed A.R.			Error of Eq.Point.	REMARKS.	
	°	'	"		°	'	"	h.	m.	s.	m.	s.	s.			
February	17	102	1	44,63	-	0,09	102	1	44,57	22	2	22,59	2	22,58	-	0,01
	18	101	40	38,74	-	0,22	101	40	38,52	22	6	14,58	6	14,37	-	0,21
	19	101	19	23,09	-	0,34	101	19	22,75	22	10	5,66	10	5,21	-	0,45
	21	100	36	23,80	-	0,51	100	36	23,29	22	17	45,14	17	45,75	+	0,61
	25	99	8	20,59	-	0,53	99	8	20,06	22	32	58,25	32	58,11	-	0,14
	26	98	45	58,69	-	0,48	98	45	58,21	22	36	44,95	36	44,77	-	0,18
	27	98	23	31,78	-	0,38	98	23	31,40	22	40	30,63	40	30,64	+	0,01
March	28	98	0	52,63	-	0,28	98	0	52,35	22	44	16,58	44	16,18	-	0,40
	1	97	38	12,79	-	0,19	97	38	12,60	22	48	0,95	48	1,02	+	0,07
	2	97	15	21,56	-	0,06	97	15	21,50	22	51	45,61	51	45,46	-	0,15
	3	96	52	30,74	+	0,06	96	52	30,80	22	55	28,69	55	29,32	+	0,63
	4	96	29	27,33	+	0,18	96	29	27,51	22	59	12,41	59	12,55	+	0,14
	5	96	6	22,57	+	0,28	96	6	22,85	23	2	55,03	2	55,14	+	0,11
	6	95	43	9,59	+	0,38	95	43	9,97	23	6	37,72	6	37,87	+	0,15
	7	95	19	58,31	+	0,45	95	19	58,76	23	10	19,00	10	19,71	+	0,71
	9	94	33	12,83	+	0,47	94	33	13,30	23	17	42,09	17	42,63	+	0,54
	10	94	9	44,96	+	0,45	94	9	45,41	23	21	23,06	21	23,59	+	0,53
	11	93	46	6,52	+	0,40	93	46	6,92	23	25	4,85	25	4,17	-	0,68
	13	92	58	59,66	+	0,21	92	58	59,87	23	32	24,73	33	24,34	-	0,39
	15	92	11	45,46	-	0,02	92	11	45,44	23	39	43,47	39	43,47	-	0,00
	19	90	36	56,16	-	0,49	90	36	55,67	23	54	19,60	54	19,30	-	0,30
	21	89	49	34,57	-	0,64	89	49	33,93	0	1	36,19	1	36,46	+	0,27
22	89	25	54,83	-	0,65	89	25	54,18	0	5	14,31	5	14,68	+	0,37	
23	89	2	12,24	-	0,65	89	2	11,59	0	8	52,97	8	52,94	-	0,03	
25	88	14	57,74	-	0,68	88	14	57,06	0	16	9,32	16	9,05	-	0,27	
26	87	51	22,43	-	0,50	87	51	21,93	0	19	47,62	19	46,96	-	0,66	
27	87	27	56,06	-	0,40	87	27	55,66	0	23	24,96	23	25,21	+	0,25	
29	86	41	6,18	-	0,17	86	41	6,01	0	30	40,74	30	41,03	+	0,29	
30	86	17	41,50	-	0,05	86	17	41,45	0	34	19,52	34	18,96	-	0,56	
April	1	85	31	19,65	+	0,20	85	31	19,85	0	41	34,97	41	34,99	+	0,02
	2	85	8	19,02	+	0,29	85	8	19,31	0	45	12,33	45	13,33	+	1,00
	3	84	45	17,78	+	0,35	84	45	18,14	0	48	50,76	48	51,63	+	0,87
	4	84	22	15,83	+	0,38	84	22	16,21	0	52	30,29	52	30,08	-	0,21
	5	83	59	24,94	+	0,39	83	59	25,33	0	56	9,18	56	8,63	-	0,55
	6	83	36	42,09	+	0,40	83	36	42,49	0	59	47,97	59	47,96	-	0,01
	7	83	14	5,33	+	0,35	83	14	5,68	1	3	27,03	3	26,81	-	0,22
	8	82	51	36,47	+	0,28	82	51	36,75	1	7	6,16	7	6,06	-	0,10
	9	82	29	14,25	+	0,18	82	29	14,43	1	10	45,61	10	45,34	-	0,27
	14	80	39	28,60	-	0,45	80	39	28,15	1	29	6,19	29	7,01	+	0,82
	17	79	35	18,23	-	0,74	79	35	17,49	1	40	11,64	40	12,38	+	0,74
	19	78	53	18,19	-	0,75	78	53	17,44	1	47	37,84	47	37,39	-	0,45
20	78	32	37,70	-	0,73	78	32	36,97	1	51	20,97	51	20,78	-	0,19	
22	77	51	45,72	-	0,59	77	51	45,13	1	58	49,40	58	48,71	-	0,69	

NOTE.—In the foregoing computations, and in those which follow, the Sun's Latitude has been computed from VINCE'S Tables for the year 1832; and copied from the Nautical Almanac for the year 1833: the values of the obliquity of the Ecliptic employed are those given in the Supplements to the Nautical Almanac.

And further we have:

Observations of the Sun made near to the Autumnal Equinox in 1832 and 1833, applied to the determination of the error of the assumed Equinoctial Point.

1832	Reduced N.P.D. of the Sun.	Correc- tion.	N. P. D. Reduced on account of Sun's Latitude.	Computed A. R.	Observed A.R.	Error of Eq. Point.	REMARKS.
	• / "	"	• / "	h. m. s.	m. s.	s.	
August	19 77 12 34,40	— 0,75	77 12 33,65	9 53 50 10	53 50,86	+ 0,76	
	20 77 32 15,95	— 0,57	77 32 15,38	9 57 32,69	57 33,69	+ 1,00	
	21 77 52 9,12	— 0,50	77 52 8,62	10 1 14,74	1 15,59	+ 0,85	
	22 78 12 17,81	— 0,39	78 12 17,42	10 4 57,07	4 57,57	+ 0,50	
	23 78 32 34,72	— 0,29	78 32 34,43	10 8 38,40	8 38,75	+ 0,35	
	25 79 13 50,09	+ 0,10	79 13 50,19	10 16 1,40	16 0,87	— 0,53	
	27 79 55 35,70	+ 0,37	79 55 36,07	10 22 20,81	23 20,62	— 0,19	
	28 80 16 40,24	+ 0,48	80 16 40,72	10 26 59,46	27 0,20	+ 0,74	
	29 80 38 4,17	+ 0,57	80 38 4,74	10 30 39,46	30 39,24	+ 0,22	
	30 80 59 26,61	+ 0,60	80 59 27,21	10 34 17,32	34 17,75	+ 0,43	
	31 81 21 0,28	+ 0,62	81 21 0 90	10 37 55,15	37 56,23	+ 1,08	
September	4 82 48 42,95	+ 0,31	82 48 43,26	10 52 25,48	52 25,74	+ 0,26	
	6 83 33 16,04	+ 0,06	83 33 16,10	10 59 38,73	59 39,23	+ 0,50	
	7 83 55 43,26	— 0,08	83 55 43,18	11 3 15,17	3 15,93	+ 0,76	
	9 84 41 2,89	— 0,37	84 41 2 52	11 10 28,67	10 27,91	— 0,76	
	10 85 3 47,45	— 0,49	85 3 46,96	11 14 4,63	14 4,16	— 0,47	
	15 86 58 30,78	— 0,63	86 58 30,15	11 32 1,45	32 1,05	— 0,40	
	16 87 21 38,43	— 0,53	87 21 37,90	11 35 36,58	35 36,33	— 0,25	
	20 88 54 42,27	0,00	88 54 42,27	11 49 57,91	49 58,33	+ 0,42	
	21 89 18 7,92	+ 0,16	89 18 8,08	11 53 34,07	53 34,02	— 0,05	
	23 90 5 1,09	+ 0,44	90 5 1,53	12 0 46,25	0 45,77	— 0,48	
	24 90 28 22,24	+ 0,57	90 28 22,81	12 4 21,60	4 21,46	— 0,14	
	25 90 51 48,97	+ 0,65	90 51 49,62	12 7 57,81	7 57,55	— 0,26	
	26 91 15 13,55	+ 0,70	91 15 14,25	12 11 33,85	11 33,26	— 0,59	
	27 91 38 35,55	+ 0,71	91 38 36,26	12 15 9,73	15 10,66	+ 0,93	
	28 92 2 2,24	+ 0,72	92 2 2,96	12 18 46,59	18 46,92	+ 0,33	
	30 92 48 47,45	+ 0,57	92 48 48,02	12 26 0,40	26 0,94	+ 0,54	
October	1 93 12 2,44	+ 0,41	93 12 2,85	12 29 36,85	29 37,70	+ 0,85	
	2 93 35 28,25	+ 0,29	93 35 28,54	12 33 15,61	33 16,41	+ 0,80	
	4 94 21 58,44	0,00	94 21 58,44	12 40 31,94	40 32,07	+ 0,13	
	7 95 31 16,87	— 0,36	95 31 16,51	12 51 28,70	51 28,31	— 0,39	
	8 95 54 12,56	— 0,47	95 54 12,09	12 55 8,03	55 9,28	+ 1,25	
	9 96 17 8,14	— 0,52	96 17 7,62	12 58 48,53	58 48,97	+ 0,44	
	11 97 2 36,15	— 0,54	97 2 35,61	13 6 9,61	6 10,42	+ 0,81	
	12 97 25 17,91	— 0,53	97 25 17,38	13 9 51,89	9 52,25	+ 0,36	
	13 97 47 42,99	— 0,43	97 47 42,56	13 13 32,95	13 34,20	+ 1,25	
	14 98 10 10,93	— 0,37	98 10 10,56	13 17 16,07	17 16,69	+ 0,62	
	15 98 32 28,43	— 0,22	98 32 28,21	13 20 59,15	20 59,66	+ 0,51	
	21 100 43 39,23	+ 0,61	100 43 39,84	13 43 32,00	43 31,90	— 0,10	
	22 101 4 51,71	+ 0,68	101 4 52,39	13 47 18,26	47 19,30	— 1,04	
	23 101 25 59,41	+ 0,76	101 26 0,17	13 51 6,11	51 7,33	+ 1,22	
	24 101 47 2,79	+ 0,80	101 47 3,59	13 54 55,73	54 56,51	+ 0,78	
	25 102 7 52,25	+ 0,80	102 7 53,05	13 58 45,50	58 45,82	+ 0,32	
	26 102 28 29,93	+ 0,75	102 28 30,68	14 2 35,87	2 36,37	+ 0,50	
	27 102 48 55,17	+ 0,67	102 48 55,84	14 6 26,80	6 27,71	+ 0,91	
	28 103 9 6,56	+ 0,55	103 9 7,11	14 10 18,12	10 19,51	+ 1,39	
	29 103 29 9,27	+ 0,42	103 29 9,69	14 14 10,91	14 12,06	+ 1,15	
	30 103 49 0,54	+ 0,24	103 49 0,78	14 18 4,71	18 5,38	+ 0,67	

1833	Reduced N.P.D. of the Sun.	Correc-tion.	N. P. D. Reduced on account of Sun's Latitude.	Computed A. R.	Observed A.R.	Error of Eq.Point.	REMARKS.
August 21	77 47 25,65	+ 0,38	77 47 26,03	h. m. s. 10 0 22,75	m. s. 0 22,75	s. 0,00	
23	78 27 36,67	+ 0,23	78 27 36,90	10 7 44,85	7 46,30	+ 1,45	
September 6	83 27 49,66	- 0,37	83 27 49,29	10 58 46,16	58 46,05	- 0,11	
7	83 50 7,20	- 0,25	83 50 6,95	11 2 21,37	2 22,65	+ 1,28	
8	84 12 40,76	- 0,13	84 12 40,63	11 5 58,00	5 58,18	+ 0,18	
9	84 35 23,52	0,00	84 35 23,52	11 9 34,98	9 35,03	+ 0,05	
10	84 58 10,48	+ 0,12	84 58 10,60	11 13 11,58	13 11,05	- 0,53	
11	85 20 58,43	+ 0,25	85 20 58,68	11 16 47,38	16 46,58	- 0,80	
12	85 43 52,64	+ 0,36	85 43 53,00	11 20 23,30	20 22,64	- 0,66	
13	86 6 42,77	+ 0,45	86 6 43,22	11 23 57,76	23 59,07	+ 1,31	
14	86 29 50,37	+ 0,51	86 29 50,88	11 27 34,22	27 34,41	+ 0,19	
15	86 52 56,52	+ 0,55	86 52 57,07	11 31 9,80	31 10,00	+ 0,20	
16	87 16 6,30	+ 0,54	87 16 6,84	11 34 45,33	34 45,32	- 0,01	
18	88 2 29,35	+ 0,44	88 2 29,79	11 41 55,50	41 56,45	+ 0,95	
26	91 9 35,25	- 0,45	91 9 34,80	12 10 41,60	10 41,70	+ 0,10	
October 2	93 29 47,72	- 0,41	93 29 47,31	12 32 22,37	32 22,92	+ 0,55	
3	93 53 3,65	- 0,31	93 53 3,34	12 36 0,16	36 0,40	+ 0,24	
4	94 16 17,33	- 0,19	94 16 17,14	12 39 38,29	39 38,55	+ 0,26	
6	95 2 38,90	+ 0,08	95 2 38,98	12 46 56,25	46 56,12	- 0,13	
11	96 57 10,39	+ 0,61	96 57 11,00	13 5 16,71	5 17,20	+ 0,49	
13	97 42 21,89	+ 0,65	97 42 22,54	13 12 40,07	12 40,71	+ 0,64	
14	98 4 49,11	+ 0,63	98 4 49,74	13 16 22,69	16 23,46	+ 0,77	
15	98 27 10,50	+ 0,56	98 27 11,06	13 20 5,93	20 6,35	+ 0,42	
16	98 49 23,52	+ 0,48	98 49 24,00	13 23 49,56	23 50,19	+ 0,63	
18	99 33 33,44	+ 0,27	99 33 33,71	13 31 19,73	31 19,47	- 0,26	
19	99 55 20,54	+ 0,16	99 55 20,70	13 35 4,77	35 5,13	+ 0,36	
22	100 59 48,55	- 0,30	100 59 48,25	13 46 23,75	46 25,03	+ 1,28	
23	101 21 4,99	- 0,30	101 21 4,69	13 50 12,53	50 12,57	+ 0,04	
24	101 42 5,42	- 0,37	101 42 5,05	13 54 0,98	54 1,50	+ 0,52	

Taking the means and referring to the Observations of 1831, for the results of that year we have:

ERROR OF THE ASSUMED EQUINOCTIAL POINT.				MEAN.
	s.		s.	s.
1831 From 19 Obs. at Vernal Eq.	+ ,055	from 17 Obs. at Aut. Eq.	+ ,267	+ ,161
1832 — 50 — — — —	,140	48 — — — —	+ ,399	+ ,130.
1833 — 48 — — — —	,046	29 — — — —	+ ,325	+ ,140
General Mean.....	- ,068.		+ ,352	+ ,142

Now the above observed places are derived from the Equinoctial Point assumed by Dr. Maskelyne + 0",20: hence it appears that the place of the true Equinox, is Dr. MASKELYNE + 0",058.

It must here be recollected that the above measures of N. P. D. are derived from a comparison of the observed places of certain fixed Stars, with their places given in the Greenwich Catalogue; the latter depending upon the

assumption that the latitude of the Greenwich Royal Observatory = $51^{\circ} 28' 39''.00$. Now any error in this assumption will necessarily occasion a similar error in the determination of the North Polar Distance of the Sun, Planets, Moon, and fixed Stars; and further, our result of the latitude of the Madras Observatory determined at Page 95, of Vol. I. will be erroneous to the like amount.

If to the above cause we now refer the disagreement between the Solstitial declination of the Sun in Winter and Summer at Page 79, and the disagreement between the Equinoctial point found from the Spring and Autumn Observations as above; we determine as follows.

	Latitude of Greenwich.		
	°	'	"
To reconcile the Summer and Winter Solstices of 1831.....	51	28	38,29
_____ — — — — — 1832.....	51	28	36,81
_____ — — — — — 1833.....	51	28	37,85
_____ — Spring and Autumn Equinoxes — 1831.....	51	28	38,30
_____ — — — — — 1832.....	51	28	37,26
_____ — — — — — 1833.....	51	28	37,80
Giving to each result the same weight and taking the Mean =	51	28	37,72

and the reduced value of the latitude of the Madras Observatory $13^{\circ} 4' 7''.93$: for the present I propose to consider these determinations too small by half a second at least; an opinion which rests on the improbability that the numerous and carefully made Observations at Greenwich can err to this amount on the one hand, and on the other from the general irregularity of the Solar Observations at Madras, the above result cannot be allowed to determine a point of so much importance and to this degree of accuracy.

With regard to the irregularity just noticed I have to remark, that in this climate the edge of the Sun is frequently ill defined and tremulous, which will account for some of the discordances which are found; whether the fierce rays of a vertical Sun which on one occasion may unavoidably remain longer on the Telescope than at another will account for the rest, is a subject to which I propose immediately to turn my attention.

In the next place we come to the Observations of the Planets; these have been reduced to the *apparent* place, as would be viewed by an observer situated at the centre of the Earth; for this purpose the parallaxes employed have been computed from the Horizontal Parallaxes given in the Supplement to the Nautical Almanac.

Apparent Right Ascension and North Polar Distance of MERCURY.

1832	Madras Mean Time of Observations.			Point Observed.	A. R.			Point Observed.	N. P. D.			REMARKS.	
	h.	m.	s.		h.	m.	s.		°	'	"		
February	18	22	52	27,8	Centre.	20	44	22,92	Centre.	109	33	37,96	
March	12	23	49	32,7	—	23	13	5,84	—	97	9	23,59	
April	2	0	52	55,8	—	1	35	30,68	—	
	3	0	55	45,8	—	1	42	17,52	—	78	12	35,78	
	4	0	58	28,0	—	1	48	56,37	—	77	23	9,48	
	5	1	0	58,6	—	1	55	24,09	—	76	35	39,44	
	7	1	5	27,5	—	2	7	47,14	—	75	7	23,45	
	9	1	8	58,3	—	2	19	12,22	—	73	48	43,07	
	10	1	10	24,5	—	2	24	35,39	—	73	13	23,41	
	October	6	23	7	26,8	—	12	10	54,23	—	89	6	16,89
November	5	0	13	34,3	—	15	11	35,35	—	108	41	17,75	
	10	0	25	11,1	—	15	42	56,69	—	111	5	10,69	
	12	0	29	56,0	—	15	55	36,31	—	111	55	33,71	
	15	0	37	12,4	—	16	14	42,77	—	113	2	59,82	
	18	0	44	33,3	—	16	33	55,71	—	112	59	58,15	
	19	0	47	2,4	—	16	40	21,68	—	114	16	34,42	
	23	0	56	51,7	—	17	5	58,04	—	115	10	9,17	
	December	5	1	20	45,8	—	18	17	14,48	—	115	36	52,37
8		1	22	50,6	—	18	31	9,31	—	115	11	51,63	
1833													
March	18	0	59	33,0	—	0	42	2,21	—	84	51	7,51	
	23	1	9	0,8	—	1	11	15,29	—	80	42	17,95	
	25	1	10	58,6	—	1	21	6,36	—	79	17	51,48	
	26	1	11	28,3	—	1	25	32,76	—	78	39	40,16	
	27	1	11	37,4	—	1	29	38,62	—	78	4	34,61	
	28	1	11	24,2	—	1	33	21,79	—	77	32	35,76	
	29	1	10	48,0	—	1	36	42,12	—	77	3	50,33	
	April	1	1	6	36,7	—	1	44	19,53	—	75	58	5,46
May	28	22	36	—	—	—	75	8	57,74	
	31	22	50	36,5	—	3	21	57,46	—	73	21	8,20	
July	17	1	50	—	—	—	74	45	54,00	
October	19	0	21	13,8	—	14	11	17,32	—	103	39	32,36	
	21	0	25	15,4	—	14	23	12,66	—	104	54	19,58	
December	23	22	27	30,1	—	16	37	26,89	—	109	32	19,03	
	25	22	24	44,8	—	16	43	33,15	—	109	58	33,07	

Apparent Right Ascension and North Polar Distance of VENUS.

1832	Madras Mean Time of Observations.			Point Observed.	A. R.			Point Observed.	N. P. D.			REMARKS.	
	h.	m.	s.		h.	m.	s.		°	'	"		
January	24	21	5	59,0	2 L.	17	19	50,61	Centre.	110	47	53,14	
	26	21	8	1,9	—	17	29	47,29	—	111	2	15,27	
	29	21	11	14,3	—	17	44	48,13	—	111	19	47,23	
	30	21	12	19,3	—	17	49	50,22	—	
	31	21	13	23,6	—	17	54	52,60	—	111	28	33,17	
February	1	21	14	30,5	—	17	59	56,21	—	111	32	7,02	
	3	21	16	45,8	—	18	10	4,92	—	111	37	38,93	

Apparent Right Ascension and North Polar Distance of VENUS, continued.

1832	Madras Mean Time of Observations.			Point Observed.	A. R.			Point Observed.	N. P. D.			REMARKS.		
	h.	m.	s.		h.	m.	s.		"	"	"			
February	4	21	17	54,1	2 L.	18	15	9,91	Centre.	111	39	27,33		
	5	21	19	0,4	---	18	20	15,42	---	111	40	42,87		
	7	21	21	21,8	---	18	30	27,30	---	111	41	29,47		
	8	21	22	31,8	---	18	35	34,28	---	111	40	45,50		
	11	21	26	0,9	---	18	50	55,45	---	111	35	24,42		
	12	21	27	12,1	---	18	56	2,61	---	111	32	22,38		
	22	21	38	53,4	---	19	47	10,59	---	110	28	46,09		
	23	21	40		---	---	110	19	7,20		
	24	21	41	10,0	---	19	57	20,64	---	110	8	50,92		
	25	21	42	15,2	---	20	2	24,93	---	109	58	1,77		
	27	21	44	30,6	---	20	12	31,55	---	109	34	36,02		
	29	21	46	41,0	---	20	22	35,13	---	109	8	56,41		
	March	1	21	47	45,5	---	20	27	36,54	---	108	55	11,00	
		2	21	48	48,6	---	20	32	36,44	---	108	40	58,79	
3		21	49	51,4	---	20	37	36,03	---	108	26	14,11		
4		21	50	54,0	---	20	42	34,88	---	108	10	58,20		
5		21	51	55,7	---	20	47	32,66	---	107	55	9,01		
7		21	53	54,6	---	20	57	26,11	---	107	22	4,86		
11		21	57	43,5	---	21	17	1,75	---	106	9	59,08		
12		21	58	39,2	---	21	21	54,02	---	105	50	49,63		
13		21	59	33,5	---	21	26	45,02	---	105	31	12,87		
15		22	1	18,8	---	21	36	23,79	---	104	50	38,67		
17		22	3	0,5	---	21	45	59,34	---	104	8	21,96		
19		22	4	39,9	---	21	55	31,65	---	103	24	30,87		
26		22	9	59,6	---	22	28	27,31	---	100	40	0,73		
August		13	0	24	39,7	---	9	51	31,40	---	75	35	46,22	
	17	0	28	2,5	---	10	10	41,59	---	77	13	34,46		
	20	0	30	24,3	---	10	24	53,40	---	78	32	45,56		
	21	0	31		---	---	78	59	47,08		
September	11	0	52	53,4	---	12	5	49,74	---	89	18	0,54		
	24	0	52	10,2	---	13	4	41,95	---	95	56	45,22		
	26	0	54	14,1	---	13	13	52,42	---	96	59	59,85		
	27	0	54	47,7	---	13	18	25,41	---	97	26	55,75		
October	2	0	57	24,7	---	13	41	30,20	---	99	53	59,97		
	8	1	1	51,2	---	14	9	37,16	---	102	42	49,05		
	12	1	5	8,4	---	14	28	40,84	---	104	29	40,59		
	13	1	6	0,5	---	14	33	29,53	---	104	55	35,06		
	24	1	16	50,4	---	15	27	43,17	---	109	12	55,96		
	25	1	17	56,9	---	15	32	46,23	---	109	33	35,33		
	26	1	19	4,3	---	15	37	50,36	---	109	3	41,23		
	27	1	20	13,3	---	15	42	56,33	---	110	13	15,54		
	28	1	21	23,3	---	15	48	3,08	---	110	32	10,97		
	29	1	22	34,0	---	15	53	10,49	---	110	50	35,90		
	30	1	23	47,7	---	15	58	20,76	---	111	8	28,85		
November	31	1	25	1,2	---	16	3	30,85	---	111	25	48,14		
	1	1	26	15,9	---	16	8	42,27	---	111	42	27,52		
	3	1	28	49,3	---	16	19	9,33	---	112	14	2,98		
	4	1	30	7,8	---	16	24	24,55	---	112	28	49,94		
	5	1	31	26,7	---	16	29	40,52	---	112	43	2,71		
	10	1	38	17,2	---	16	56	14,91	---	113	44	4,71		
	12	1	41	7,4	---	17	6	58,81	---	114	3	44,97		
	16	1	46	56,3	---	17	28	34,52	---	114	34	31,04		
	17	1	48	21,7	---	17	33	59,99	---	114	40	21,93		
	18	1	49	53,4	---	17	39	25,47	---	114	45	26,96		
	19	1	51	22,6	---	17	44	52,43	---	114	49	48,88		

Apparent Right Ascension and North Polar Distance of VENUS, continued.

1832	Madras Mean Time of Observations			Point Observed.	A. R.			Point Observed.	N. P. D.			REMARKS.	
	h.	m.	s.		h.	m.	s.		°	'	"		
November	23	1	57	21,5	2 L.	18	6	37,81	Centre.	113	59	49,22	
	24	1	58	52,0	---	18	12	4,88	---	115	0	23,97	
1833													
April	5	2	40	41,9	---	3	34	56,47	---	65	32	40,62	
	6	2	39	47,3	---	3	37	28,85	---	65	20	14,56	
May	6	1	0	53,0	---	3	56	34,24	---	64	40	16,87	
	8	0	49	28,0	---	3	53	1,02	---	65	7	36,20	
	23	23	10	59,1	---	3	17	20,99	---	70	33	47,51	
	24	23	5	11,1	---	3	15	28,34	---	70	55	35,03	
	28	23			---				---	72	16	53,49	
	29	22			---				---	72	35	20,34	
	31	22	28	18,6	---	3	6	5,63	---	73	9	30,70	
July	15	20	53	2,8	---	4	27	59,01	---	72	5	50,59	
	23	20	51	26,1	---	4	56	54,80	---	70	54	33,73	
	25	20	51	26,1	---	5	5	47,88	---	70	30	43,62	
	26	20	51	30,5	---	5	9	48,97	---	70	30	43,83	
	28	20	51	43,7	---	5	17	55,18	---	70	16	6,49	
August	29	20	51	53,6	---	5	22	1,80	---	70	9	10,41	
	2	20	52	54,1	---	5	38	48,48	---	69	44	38,79	
	5	20	53	57,7	---	5	51	42,02	---	69	29	51,25	
	7	20	54	50,0	---	6	0	26,75	---	69	21	57,74	
	9	20	55	46,5	---	6	9	17,28	---	69	15	42,75	
	13	20	57	57,7	---	6	27	15,11	---	69	8	29,89	
	14	20	58	33,8	---	6	31	47,94	---	69	7	51,94	
	15	20	59	10,6	---	6	36	20,56	---	69	7	41,23	
September	10	21	18	51,8	---	8	38	35,88	---	72	4	28,98	
	11	21	19	39,8	---	8	43	20,68	---	72	19	22,06	
November	27	22	12	20,6	---	14	39	44,99	---	103	58	46,42	
December	2	22	17	13,8	---	15	4	21,49	---	105	54	50,40	
	9	22	24	53,0	---	15	39	37,99	---	108	19	52,33	
	11	22	27	15,2	---	15	49	53,52	---	106	57	5,48	
	13	22	28	43,1	---	16	0	13,78	---	109	32	12,74	
	17	22	34	51,1	---	16	21	7,64	---				
	18	22	36	7,8	---	16	26	24,07	---	110	50	13,16	
	25	22	45	51,23	---	17	3	44,45	---	112	13	59,69	
	26	22	52	29,0	---	17	9	7,85	---	112	25	22,81	
	30	22	50	24,9	---	17	30	47,82	---	112	56	4,43	

Apparent Right Ascension and North Polar Distance of MARS.

1832	Madras Mean Time of Observations.			Point Observed.	A. R.			Point Observed	N. P. D.			REMARKS.	
	h.	m.	s.		h.	m.	s.		°	'	"		
January	29	21	16	53,0	Centre.	17	44	48,13	Cent e.	113	45	44,33	
February	3	21	13		---				---	113	50	10,94	
	4	21	12	5,0	---	18	9	19,85	---	113	50	25,00	
	5	21	11	18,4	---	18	12	29,18	---	113	50	25,38	
	6	21	10	30,4	---	18	15	38,43	---	113	50	4,07	
	8	21	8	49,0	---	18	21	57,30	---	113	48	52,84	
	22	20	58	3,9	---	19	6	14,40	---	113	13	45,41	

Apparent Right Ascension and North Polar Distance of MARS, continued.

1832	Madras Mean Time of Observations.			Point Observed.	A. R.			Point Observed	N. P. D.			REMARKS.	
	h.	m.	s.		h.	m.	s.		°	'	"		
February	24	20	56	30.6	Centre.	19	12	33,82	Centre.	113	5	1,29	
	27	20	54	11,7		19	23	1,93		112	50	9,87	
	28	20	53	24,6		19	25	11,15		112	44	24,41	
	29	20	52	36,5		19	28	19,85		112	39	8,92	
March	1	20	51	46,6	19	31	28,85	112	33	17,45			
	2	20	50	59,1	19	34	37,48	112	27	15,86			
	3	20	50	11,1	19	37	45,86	112	20	58,68			
	4	20	49	23,2	19	40	54,33	112	14	28,83			
	5	20	48	34,6	19	44	2,45	112	7	41,32			
	6	20	47	46,0	19	47	10,19	112	0	44,30			
	7	20	46	57,0	19	50	17,52	111	53	35,82			
	11	20	43	43,3	20	2	45,78	111	22	41,71			
	12	20	42	50,1	20	5	52,49	111	14	29,56			
	13	20	41	59,9	20	8	58,72	111	6	1,53			
	15	20	40	18,7	20	15	10,20	110	48	33,21			
	19	20	36	52,8	20	27	30,12	110	11	10,93			
	20	20	36	0,3	20	30	34,26	110	1	22,99			
	27	20	29	46,8	20	51	55,48	108	47	2,00			
April	31	20	26	5,8	21	3	59,97	108	0	43,11			
	1	20	25	10,0	21	9	0,29	107	48	43,33			
	2	20	24	13,0	21	10	0,29	107	36	32,83			
	3	20	23	15,9	21	12	59,45	107	24	13,62			
	5	20	21	21,2	21	18	57,50	106	59	7,03			
	6	20	20	23,2	21	21	55,82	106	42	26,18			
	7	20	19	24,6	21	24	53,87	106	33	23,71			
	12	20	14	26,2	21	39	38,65	105	26	33,56			
	13	20	13	25,5	21	42	34,69	105	12	49,08			
	14	20	12	25,0	21	45	30,02	103	58	49,22			
	21	20	5	10,2	22	5	49,00	103	17	51,62			
	30	19	55	23,2	22	31	30,00	101	0	26,25			
	May	1	19	54	16,4	22	34	19,19	100	44	46,00		
		2	19	53	8,6	22	37	8,03	100	28	58,97		
4		19	50	53,0	22	42	45,21	99	57	12,45			
5		19	49	45,7	22	45	33,97	99	41	7,02			
12		19	41	44,8	23	5	2,24	97	47	15,69			
14		19	39	17,5	23	10	33,11	97	14	20,65			
15		19	38	10,6	23	13	18,19	96	57	45,87			
16		19	36	58,1	23	16	2,83	96	41	11,49			
June	31	19	18	28,4	23	56	42,76	92	30	43,90			
	9	19	6	58,5	0	20	38,52	89	59	36,00			
	10	19	5	40,3	0	23	16,93	89	42	58,04			
	11	19	4	21,8	0	25	54,81	89	26	26,22			
	12	19	3	3,3	0	28	32,67	89	9	55,58			
	13	19	1	44,7	0	31	10,54	88	53	23,68			
	14	19	0	26,0	0	33	47,89	88	37	0,50			
	15	18	59	6,7	0	36	25,19	88	20	39,75			
	17	18	56	27,4	0	41	38,18	87	51	6,83			
	22	18	49	42,1	0	54	34,69	86	27	47,22			
	November	9	12	44	40,3	4	0	30,87	68	58	16,56		
15		12	11	47,2	3	51	11,51			
16		12	6	15,7	3	49	35,07	69	8	24,93			
17		12	0	13,7	3	47	28,56	69	10	16,86			
22		11	33	7,3	3	40	0,72	69	21	9,48			
29		10	55	21,1	3	29	41,83	69	37	32,37			
30		10	50	2,1	3	28	21,22	69	39	55,54			

Apparent Right Ascension and North Polar Distance of MARS, continued.

1832	Madras Mean Time of Observations.			Point Observed.	A. R.	Point Observed.	N. P. D.			REMARKS.			
	<i>h.</i>	<i>m.</i>	<i>s.</i>		<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>°</i>	<i>'</i>	<i>"</i>			
December	4	10	29	24,7	Centre.	3	23	26,32	Centre.	69	48	32,52	
	5	10	24	22,1	---	3	22	19,79	---	69	50	32,46	
	6	10	19	25,4	---	3	21	16,19	---	69	52	26,38	
	7	10	14	26,4	---	3	20	15,58	---	69	54	13,71	
	12	9	50	33,2	---	3	16	1,14	---	70	1	29,33	
	13	9	45	56,4	---	3	15	20,07	---	70	2	36,82	
	14	9	41	22,6	---	3	14	42,37	---	
	15	9	36	52,7	---	3	14	8,21	---	70	4	16,50	
	16	9	31	25,8	---	3	13	37,59	---	70	4	53,18	
	17	9	28	2,4	---	3	13	10,02	---	70	5	23,91	
	18	9	23	43,9	---	3	12	46,30	---	70	5	41,99	
	20	9	15	14,0	---	3	0	9,63	---	70	5	51,22	
	21	9	11	4,4	---	3	11	55,05	---	70	5	38,49	
	22	9	6	58,5	---	3	11	44,72	---	70	5	21,76	
	24	8	58	56,0	---	3	11	34,27	---	70	4	12,34	
	25	8	54	59,8	---	3	11	33,84	---	70	3	21,52	
	26	8	51	6,4	---	3	11	36,93	---	
	27	8	47	17,3	---	3	11	43,82	---	70	1	15,47	
	1833												
	January	3	8	21	55,3	---	3	13	53,40	---	69	48	46,01
		4	8	18	29,7	---	3	14	23,77	---	69	46	23,58
		6	8	11	36,2	---	3	15	32,14	---	69	41	14,46
		8	8	5	13,4	---	3	16	51,16	---	69	35	34,79
		9	8	2	0,8	---	3	17	34,74	---	69	32	33,75
		10	7	58	50,8	---	3	18	20,72	---	69	29	22,98
		11	7	55	33,2	---	3	19	9,19	---	69	26	9,69
		14	7	47	22,3	---	3	22	47,53	---	69	15	48,36
15		7	43	36,2	---	3	22	46,07	---	69	12	7,58	
16		7	40	39,5	---	3	23	45,24	---	69	8	21,62	
17		7	37	45,6	---	3	24	47,60	---	69	4	33,05	
18		7	34	55,0	---	3	25	53,42	---	69	0	37,68	
19		7	32	4,7	---	3	26	59,25	---	68	56	37,30	
20		7	29	16,6	---	3	28	7,41	---	68	52	33,07	
21		7	26	31,5	---	3	29	18,26	---	68	48	23,36	
22		7	23	48,0	---	3	31	1,96	---	68	44	9,72	
23		7	21	6,9	---	3	31	45,70	---	68	39	54,05	
24		7	18	27,4	---	3	33	2,06	---	68	35	33,48	
25		7	15	48,6	---	3	34	19,89	---	
27		7	10	38,3	---	3	37	1,80	---	68	22	15,07	
28		7	8	6,2	---	3	38	25,74	---	68	17	45,32	
29	7	5	35,3	---	3	39	50,97	---	68	13	11,20		
30	7	3	6,2	---	3	41	17,88	---	68	8	34,75		
31	7	0	38,4	---	3	42	46,30	---	68	3	57,06		
February	1	6	58	12,0	---	3	44	16,10	---	67	59	17,90	
	2	6	55	48,1	---	3	45	48,49	---	67	54	40,23	
	4	6	51	3,7	---	3	48	56,47	---	67	45	17,33	
	5	6	48	44,3	---	3	50	33,10	---	67	40	36,12	
	6	6	46	25,3	---	3	52	10,38	---	67	35	54,32	
	8	6	41	52,6	---	3	55	28,75	---	67	26	32,32	
	9	6	39	37,6	---	3	57	11,35	---	67	21	52,31	
	10	6	37	24,9	---	3	58	54,05	---	67	17	12,03	
	11	6	34	12,1	---	3	59	37,96	---	67	12	33,89	

Apparent Right Ascension and North Polar Distance of MARS, continued.

1833	Madras Mean Time of Observations.			Point Observed.	A. R.			Point Observed	N. P. D.			REMARKS.	
	h.	m.	s.		h.	m.	s.		°	'	"		
February	12	6	33	1,6	Centre.	4	2	23,97	Centre.	67	7	56,56	
	13	6	30	52,3	---	4	4	10,84	---	67	3	19,87	
	14	6	28	43,6	---	4	5	58,07	---	66	58	42,76	
	15	6	26	36,6	---	4	7	47,23	---	66	54	9,21	
	16	6	24	30,8	---	4	9	37,51	---	66	49	36,72	
	17	6	22	26,0	---	4	11	29,21	---	66	45	6,48	
	18	6	22	22,4	---	4	13	21,71	---	66	40	43,92	
	24	---	---	66	14	56,56	
	25	6	6	27,0	---	4	27	0,06	---	66	10	50,67	
	26	---	---	66	6	47,32	
	27	6	2	37,2	---	4	31	2,83	---	
	March	28	6	0	44,5	---	4	33	6,37	---	65	58	55,67
1		5	58	50,9	---	4	35	8,99	---	65	55	6,03	
2		5	56	58,9	---	4	37	13,45	---	65	51	20,89	
3		5	55	8,0	---	4	39	18,88	---	65	47	40,23	
4		5	53	18,1	---	4	41	24,94	---	65	44	3,69	
6		5	49	40,2	---	4	45	40,64	---	65	37	7,87	
7		5	47	52,5	---	4	47	47,88	---	65	33	46,11	
8		5	46	5,7	---	4	49	57,05	---	65	30	32,11	
9		5	44	19,1	---	4	52	7,11	---	65	27	24,25	
10		5	42	33,5	---	4	54	17,97	---	65	24	18,16	

Observed North Polar Distance of the centre of the Planet Mars and of Stars culminating near to him, together with the Greenwich mean time at which the former passed the Meridian.

1832	NAMES.	Greenwich Mean Time.			N. P. D.			REMARKS.
		h.	m.	s.	°	'	"	
November	9 A ¹ Tauri.....	68	26	24,3	
	♂	69	1	44,9	
	53 Tauri.....	69	19	41,2	
	a Tauri.....	73	53	29,1	
	15 b Tauri.....	69	11	34,5	
	♂	69	8	0,9	
	53 Tauri.....	69	17	38,4	
	a Tauri.....	73	51	24,1	
	16 b Tauri.....	69	11	34,5	
	♂	69	9	49,1	
	a Tauri.....	73	51	24,7	
	17 ♂	69	11	41,8	
	A ¹ Tauri.....	68	24	23,8	
	22 ♂	69	22	24,2	
	b Tauri.....	69	11	32,8	
	A ¹ Tauri.....	68	24	21,4	
	a Tauri.....	73	51	24,8	
	29 ♂	69	38	56,2	
	a Tauri.....	73	51	23,8	
	30 65 Arietis.....	69	49	15,4	
	♂	69	41	17,1	

1832	NAMES.	Greenwich Mean Time.			N. P. D.			REMARKS.
		h.	m.	s.	°	'	"	
December 4	65 Arietis.....				69	52	40,8	
	♂	5	8	15,7	69	53	24,2	
5	a Tauri.....				73	54	52,1	
	♂	5	3	13,1	69	55	24,8	
6	a Tauri.....				73	54	51,9	
	65 Arietis.....				69	52	40,9	
♂		4	58	16,4	69	57	18,7	
	K ¹ Tauri.....				70	55	21,5	
7	a Tauri.....				73	54	53,2	
	65 Arietis.....				69	50	44,4	
♂		4	53	17,4	69	57	9,0	
	K ² Tauri.....				70	53	26,0	
a	Tauri.....				73	52	55,4	
	12	♂	4	29	24,2	70	4	36,9
a	Tauri.....				73	53	8,2	
	13	38 Arietis.....			70	9	12,6	
♂		4	24	47,4	70	5	41,8	
	a Tauri.....				73	53	8,1	
15	38 Arietis.....				70	10	34,1	
	♂	4	15	43,7	70	8	45,4	
a	Tauri.....				73	54	28,9	
	16	38 Arietis.....			70	10	34,0	
♂		4	10	16,8	70	9	22,1	
	a Tauri.....				73	54	29,8	
17	♂	4	6	53,4	70	9	52,4	
	a Tauri.....				73	54	29,3	
18	38 Arietis.....				70	10	36,2	
	♂	4	2	34,9	70	10	11,7	
a	Tauri.....				73	54	30,8	
	20	♂	3	54	5,0	70	13	13,8
65	Arietis.....				69	55	12,8	
	a Tauri.....				73	57	22,1	
21	♂	3	49	55,4	70	13	1,1	
	65 Arietis.....				69	55	12,5	
a	Tauri.....				73	57	22,4	
	22	♂	3	45	49,5	70	12	43,2
65	Arietis.....				69	55	11,7	
	a Tauri.....				73	57	22,2	
24	♂	3	37	47,0	70	11	34,8	
	65 Arietis.....				69	55	13,3	
a	Tauri.....				73	57	22,6	
	25	♂	3	33	50,8	70	10	44,9
65	Arietis.....				69	55	12,3	
	a Tauri.....				73	57	24,3	

The above column *Greenwich Mean Time* is derived from the Madras Mean Time as computed from the observed Transit, by subtracting 5h. 21m. 9s. The column N. P. D. is copied from the Mural Circle Book without any correction whatever having been applied; in the observations it will be noticed that I have not followed the recommendation of Mr. HENDERSON, of observing the first and second limbs on alternate days, but have always bisected the centre of the Planet; my reason for thus deviating from a plan

which as far as it secures uniformity of results is a good one; arises from a conviction, that a perfect contact between the border of the Planet and edge of the wire can never be made to that degree of accuracy which a bisection of the body itself will permit; in support of this opinion I need only refer to the Solar observations made at Greenwich and at Madras, where it will be found, that the irregularity of the differences from the places given in the Nautical Almanac (the errors of observation in fact) are at least three times as large as those which are found in the observations of a fixed Star; added to which on the present occasion, were the limb of the planet observed the Star being observed with reference to the centre of the horizontal wire, and the Planet observed at the edge; we are obliged to know not only the thickness of the wire, but the semi-diameter of the Planet.

Not being possessed of any corresponding observation to the above, I am prevented from applying them to the determination of the parallax of Mars for which purpose it will be understood they have been made.

Apparent Right Ascension and North Polar Distance of JUPITER.

1832	Madras Mean Time of Observations.			Point Observed.	A. R.	Point Observed.	N. P. D.	REMARKS.					
	h.	m.	s.		h.	m.	s.						
May	12	20	9	56,2	Centre.	23	33	18,31	Centre.	94	3	53,73	
	14	20	3	15,3	—	23	34	35,11	—	93	56	3,65	
	15	20	0	2,5	—	23	35	13,84	—	93	52	9,99	
	16	19	56	44,6	—	23	35	52,63	—	93	48	23,26	
	17	19	53	23,4	—	23	36	29,50	—	93	44	39,19	A. R. doubtful
	26	19	23	10,0	—	23	41	40,99	—	93	13	6,42	on account of the clock tripping.
June	31	19	6	9,1	—	23	44	17,82	—	92	57	20,46	
	9	18	34	57,0	—	23	48	31,73	—	92	32	38,09	
	10	18	31	26,1	—	23	48	57,16	—	92	30	10,45	
	11	18	27	54,9	—	23	49	21,92	—	92	27	44,33	
	12	18	24	23,2	—	23	49	46,29	—	92	25	26,93	
	13	18	20	51,0	1 & 2 L.	23	50	9,73	—	92	23	7,36	
	14	18	17	18,1	—	23	50	32,93	—	92	20	57,57	
	15	18	13	44,5	—	23	50	55,44	—	92	18	47,69	
	17	18	6	35,3	—	23	51	37,89	—	92	14	43,87	
September	22	11	29	38,6	—	23	36	11,54	—	94	18	33,43	
	24	11	21	0,1	—	23	35	14,73	—	94	24	43,76	
	25	11	16	34,2	—	23	34	45,20	—	94	27	49,27	
	26	11	12	10,1	—	23	34	16,79	—	94	30	49,93	
	27	11	7	46,2	—	23	33	48,74	—	94	33	49,45	
October	1	1	50	12,5	—	23	31	57,81	—	94	45	24,63	
	8	8	19	41,0	—	23	28	58,51	—	95	3	56,11	
	11	10	6	43,4	—	23	27	47,88	—	95	11	2,35	
	12	10	2	24,6	—	23	27	25,02	—	95	13	18,18	
	13	9	58	6,6	—	23	27	3,71	—	95	15	30,00	
	14	9	53	50,5	—	23	26	42,55	—	95	17	34,81	
	19	9	32	29,6	—	23	25	1,70	—	95	27	16,35	
20	9	24	3,5	—	23	24	26,26	—	95	30	38,12		

Apparent Right Ascension and North Polar Distance of JUPITER, continued.

1832	Madras Mean Time of Observations.			Point Observed.	A. R.	Point Observed.	N. P. D.			REMARKS.				
	<i>h.</i>	<i>m.</i>	<i>s.</i>		<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>°</i>	<i>'</i>	<i>"</i>				
October	22	9	19	51,2	1 & 2 L	23	24	9,64	Centre.	95	32	14 52		
	23	9	15	38,9	---	23	23	53,42	---	95	33	44,23		
	24	9	11	27,2	---	23	23	37,76	---	95	35	10,50		
	25	9	5	17,1	---	23	23	22,88	---	95	36	31,15		
	26	9	3	6,6	---	23	23	8,67	---	95	37	47,32		
	27	8	58	57,2	---	23	22	55,21	---	95	39	1,56		
	28	8	54	48,8	---	23	22	42,98	---	95	40	6,05		
	29	8	50	40,1	---	23	22	30,25	---	95	41	9,42		
	30	8	51	33,1	---	23	22	18,93	---	95	42	8,01		
	31	8	42	27,0	---	23	22	8,66	---	95	42	56,62		
	November	2	8	34	15,6	---	23	21	49,18	---	95	44	30,54	
		4	8	26	8,2	---	23	21	33,43	---	95	45	44,45	
5		8	22	4,7	---	23	21	26,00	---	95	46	11,77		
9		8	6	1,0	---	23	21	5,95	---	95	47	21,28		
10		8	2	1,9	---	23	21	2,57	---	95	47	26,37		
11		7	58	2,8	---	23	20	59,05	---	95	47	22,43		
12		7	54	5,5	---	23	20	58,18	---	95	47	17,40		
15		7	42	17,3	---	23	20	57,57	---	95	46	35,22		
16		7	38	22,8	---	23	20	58,76	---	95	46	10,93		
17		7	34	28,8	---	23	21	1,00	---	95	45	39,84		
18		7	30	35,7	---	23	21	3,76	---	95	45	5,90		
19		7	26	43,4	---	23	21	7,36	---	95	44	26,18		
21		7	19	1,9	---	23	21	17,04	---	95	42	52,44		
22		7	15	10,7	---	23	21	22,59	---	95	41	57,12		
23		7	11	21,7	---	23	21	29,41	---	95	40	59,07		
25	7	3	45,2	---	23	21	44,63	---	95	38	46,98			
29	6	48	43,6	---	23	22	24,56	---	95	33	26,82			
30	6	44	57,4	---	23	22	36,47	---	95	31	57,13			
December	4	6	30	7,9	---	23	23	30,81	---	95	25	1,89		
	6	6	22	49,2	---	23	24	1,80	---	95	21	8,76		
	7	6	19	7,6	---	23	24	18,46	---	95	17	6,02		
	9	6	11	51,2	---	23	24	53,83	---				
	10	6	7	14,3	---	23	25	12,69	---				
	11	6	4	55,0	---	23	25	31,72	---				
	12	6	1	1,5	---	23	25	52,05	---	95	7	46,55		
	13	5	57	26,5	---	23	26	12,62	---	95	5	21,39		
	15	5	50	17,6	---	23	26	56,23	---				
	16	5	46	44,0	---	23	27	18,51	---	94	57	30,16		
	17	5	43	10,9	---	23	27	41,17	---	94	54	48,01		
	19	5	36	6,7	---	23	28	30,21	---	94	49	5,28		
20	5	32	37,1	---	23	28	55,47	---	94	46	8,84			
24	5	18	39,2	---	23	30	41,45	---	94	33	44,57			
1833														
June	29	19	24	53,9	---	1	56	31,07	---	79	21	57,78		
July	8	18	54	35,3	---	2	1	36,25	---	78	56	1,54		
	12	18	40	54,1	---	2	3	39,19	---	78	45	49,80		
October	13	18	37	27,9	---	2	4	8,39	---	78	43	45,08		
	13	12	32	49,5	---	2	1	13,44	---	79	16	34,40		
	14	12	28	23,3	---	2	0	43,38	---	79	18	14,36		
	15	12	23	57,1	---	2	0	13,11	---	79	21	57,69		
	20	12	1	44,4	---	1	57	39,19	---	79	35	44,94		
	22	11	52	51,6	---	1	56	38,07	---	79	41	17,73		
23	11	49	25,0	---	1	56	7,13	---	79	44	4,86			

Apparent Right Ascension and North Polar Distance of JUPITER, continued.

1833	Madras Mean Time of Observations.			Point Observed.	A. R.	Point Observed.	N. P. D.	REMARKS.					
	<i>h.</i>	<i>m.</i>	<i>s.</i>		<i>h.</i>	<i>m.</i>	<i>s.</i>						
November	9	10	33	7,1	1 & 2 L.	1	47	38,66	Centre.	80	28	56,43	
	17	9	58	12,3	---	1	44	9,94	---	80	46	46,91	
	19	9	49	33,5	---	1	43	23,55	---	80	50	44,95	
	20	9	45	14,7	---	1	42	59,88	---	80	52	38,59	
	21	9	40	57,8	---	1	42	37,39	---	80	54	29,33	
	22	8	57	45,0	---	1	42	16,28	---	80	56	15,75	
	23	9	42	20,9	---	1	41	55,81	---	80	58	0,16	
December	2	8	54	21,0	---	1	39	17,11	---	81	10	32,54	
	4	8	46	2,0	---	1	38	49,37	---	81	12	32,33	
	5	8	41	53,0	---	1	38	36,73	---	81	13	26,92	
	6	8	37	45,0	---	1	38	25,00	---	81	14	16,56	
	7	8	33	37,9	---	1	38	13,63	---	81	14	58,46	
	8	8	29	31,3	---	1	38	2,92	---	81	15	44,94	
	10	8	21	20,3	---	1	37	43,55	---	81	16	54,00	
	11	8	17	17,2	---	1	37	36,20	---	81	17	19,46	
	14	8	5	9,5	---	1	37	16,15	---	81	18	17,27	
	18	8	48	59,8	---	1	37	0,14	---	81	18	29,20	
	19	8	45	0,0	---	1	36	58,43	---	81	18	19,59	
	20	7	41	8,2	---	1	36	57,17	---	81	18	7,87	
	22	7	33	23,0	---	1	36	56,91	---	81	17	27,78	
	24	7	25	34,3	---	1	37	0,17	---	81	16	25,27	
	25	7	21	42,6	---	1	37	2,79	---	81	15	54,75	
	26	7	17	48,7	---	1	37	6,33	---	81	15	11,25	
	27	7	13	57,2	---	1	37	10,68	---	81	14	28,13	
	29	7	6	16,8	---	1	37	22,24	---	81	12	44,07	
	30	7	2	27,4	---	1	37	28,69	---	81	11	46,17	
	31	6	58	38,2	---	1	37	35,57	---	81	10	43,87	

Apparent Right Ascension and North Polar Distance of SATURN.

1832	Madras Mean Time of Observations			Point Observed.	A. R.	Point Observed.	N. P. D.	REMARKS.					
	<i>h.</i>	<i>m.</i>	<i>s.</i>		<i>h.</i>	<i>m.</i>	<i>s.</i>						
March	11	11	35	20,2	Centre.	10	52	52,83	Centre.	80	33	27,52	
	13	11	26	48,5	---	10	52	20,29	---	80	29	51,50	
	14	11	22	35,2	---	10	52	3,52	---	80	31	5,77	
	15	11	18	21,9	---	10	51	45,65	---	80	26	21,03	
	16	11	14	8,8	---	10	51	28,43	---	80	24	39,11	
	17	11	9	58,1	---	10	51	19,76	---	80	22	55,55	
	19	11	1	34,4	---	10	50	37,00	---	80	18	36,33	
	22	10	48	56,5	---	10	49	47,96	---	80	14	43,98	
	23	10	44	45,7	---	10	49	32,35	---	80	13	10,25	
	24	10	40	34,2	---	10	49	16,44	---	80	11	36,80	
	25	10	36	20,1	---	10	48	58,05	---	80	13	6,04	
	26	10	32	12,6	---	10	48	46,06	---	80	11	34,29	
	27	10	28	0,9	---	10	48	30,49	---	80	10	7,38	
	28	10	23	50,3	---	10	48	15,61	---	80	5	40,10	
	29	10	19	39,6	---	10	48	1,20	---	80	7	14,92	
	30	10	15	29,7	---	10	47	46,87	---	80	2	51,79	
	31	10	11	19,6	---	10	47	32,65	---	80	4	30,35	
April	1	10	7	9,8	---	10	47	19,07	---	80	3	10,58	

Apparent Right Ascension and North Polar Distance of SATURN, continued.

1832	Madras Mean Time of Observations.			Point Observed.	A. R.			Point Observed.	N. P. D.			REMARKS.		
	h.	m.	s.		h.	m.	s.		°	'	"			
April	2	10	3	0,1	Centre.	10	47	5,63	Centre,	79	58	53,48		
	3	9	58	51,0	—	10	46	52,25	—	79	57	37,66		
	4	9	54	42,0	—	10	46	39,17	—	79	56	23,73		
	5	9	50	33,4	—	10	46	26,11	—	79	59	11,21		
	6	9	46	22,2	—	10	46	14,65	—	79	53	59,27		
	7	9	42	14,0	—	10	46	2,28	—	79	52	49,80		
	10	9	29	57,1	—	10	45	26,57	—	79	49	40,63		
	11	9	25	50,4	—	10	45	15,43	—	79	48	41,22		
	12	9	21	43,2	—	10	45	5,41	—	79	47	46,32		
	13	9	17	36,4	—	10	44	54,62	—	79	46	47,07		
	14	9	13	31,3	—	10	44	44,69	—	79	45	57,15		
	21	8	44	57,9	—	10	43	45,37	—	79	40	59,29		
	22	8	40	55,0	—	10	43	38,28	—	79	40	10,07		
	23	8	36	51,6	—	10	43	30,94	—	79	39	39,71		
	24	8	32	50,3	—	10	43	24,65	—	79	39	7,80		
	26	8	24	45,8	—	10	43	12,87	—	79	41	15,24		
	27	8	20	45,1	—	10	43	8,00	—	79	37	52,42		
	28	8	16	43,8	—	10	43	2,72	—	79	37	30,17		
	29	8	12	43,5	—	10	42	57,93	—	79	40	12,44		
	30	8	8	43,0	—	10	42	53,89	—	79	36	56,94		
	May	3	7	56	45,2	—	10	42	44,18	—	79	36	21,44	
		4	7	52	46,4	—	10	42	40,77	—	79	36	15,84	
		5	7	48	48,3	—	10	42	38,29	—	79	36	10,51	
		6	7	44	51,6	—	10	42	37,21	—	79	36	9,91	
		9	7	32	58,6	—	10	42	32,54	—	79	36	16,21	
		11	7	25	7,7	—	10	42	32,49	—	79	36	37,56	
		12	7	21	11,9	—	10	42	33,02	—	79	36	51,24	
		14	7	13	22,0	—	10	42	35,52	—	79	38	22,64	
		15	7	9	29,0	—	10	42	37,58	—	79	37	44,25	
		16	7	5	34,9	—	10	42	39,66	—	79	38	3,25	
17		7	1	40,9	—	10	42	41,54	—	79	38	30,48		
18		6	57	40,5	—	10	42	44,22	—	79	37	58,29		
19	6	53	59,9	—	10	42	47,63	—	79	37	38,03			
20	6	50	2,5	—	10	42	51,12	—	79	38	23,53			
21	6	46	11,0	—	10	42	55,04	—	79	38	4,08			
1833														
March	13	12	20	21,0	—	11	45	0,35	—	85	41	0,10		
	14	12	16	8,2	—	11	44	43,28	—	85	39	2,11		
	15	12	11	55,5	—	11	44	25,98	—	85	37	7,20		
	16	12	6	36,9	—	11	44	8,40	—	85				
	17	11	46	13,7	—	11	43	51,23	—	85	34	16,29		
	18	11	59	15,0	—	11	43	33,92	—	85	32	22,88		
	19	11	55	1,9	—	11	43	16,12	—	85	29	27,80		
	20	11	50	49,1	—	11	42	59,21	—	85	32	34,91		
	21	11	46	36,0	—	11	42	42,67	—	85	25	42,49		
	22	11	42	22,5	—	11	42	24,82	—	85	23	49,48		
	23	11	38	10,0	—	11	42	7,79	—	85	21	58,76		
	25	11	30	4,3	—	11	41	33,83	—	85	18	28,02		
	26	11	26	1,5	—	11	41	16,84	—	85	16	29,47		
27	11	21	19,1	—	11	41	0,03	—	85	15	2,05			
28	11	17	6,4	—	11	40	48,43	—	85	12	55,52			
29	11	12	53,9	—	11	40	26,87	—	85	12	19,74			
30	11	8	41,3	—	11	40	10,33	—	85	10	24,34			

Apparent Right Ascension and North Polar Distance of SATURN, continued.

1833	Madras Mean Time of Observations.			Point Observed.	A. R.			Point Observed.	N. P. D.			REMARKS.	
	h.	m.	s.		h.	m.	s.		°	'	"		
March	31	11	4	29,3	Centre.	11	39	54,08	Centre.	85	7	42,32	
April	1	11	0	17,4	—	11	39	37,96	—	85	5	59,58	
	2	10	56	5,4	—	11	39	21,77	—	85	4	20,82	
	3	10	51	53,9	—	11	39	6,05	—	85	2	43,02	
	4	10	47	42,8	—	11	38	50,24	—	85	1	3,98	
	5	10	43	0,5	—	11	38	34,94	—	84	59	27,56	
	6	10	39	19,2	—	11	38	19,41	—	84	57	51,95	
	8	10	30	57,7	—	11	37	49,27	—	84	54	48,01	
	13	10	10	5,8	—	11	36	37,73	—	84	47	34,64	
	14	10	5	57,3	—	11	36	24,07	—	84	46	10,71	
	16	9	57	37,5	—	11	35	57,54	—	84	43	35,03	
	17	9	53	29,9	—	84	42	18,90	
	18	9	49	21,8	—	11	35	31,79	—	84	41	4,37	
	19	9	45	13,9	—	11	35	19,44	—	84	39	52,45	
	20	9	41	5,1	—	11	35	7,43	—	84	38	43,25	
	21	9	36	57,3	—	11	34	56,10	—	84	37	35,95	
	22	9	32	50,6	—	11	34	44,24	—	84	36	30,84	
	23	9	28	43,4	—	11	34	33,12	—	84	35	23,50	
	24	9	24	36,7	—	11	34	22,38	—	84	34	24,27	
	25	9	20	30,1	—	11	34	11,54	—	84	33	25,11	
	26	9	16	24,4	—	11	34	1,29	—	84	32	23,30	
	27	9	12	18,5	—	11	33	51,44	—	84	31	34,79	
	30	9	0	2,6	—	11	33	23,56	—	84	29	4,42	
May	2	8	51	53,8	—	11	33	6,41	—	84	27	37,53	
	4	8	43	46,5	—	11	32	50,86	—	84	26	19,12	

Apparent Right Ascension and North Polar Distance of GEORGIAN.

1832	Madras Mean Time of Observations.			Point Observed.	A. R.			Point Observed.	N. P. D.			REMARKS.	
	h.	m.	s.		h.	m.	s.		°	'	"		
August	28	10	46	17,5	Centre.	21	13	59,93	Centre.	
September	11	9	49	21,2	—	21	12	5,52	—	106	58	35,27	
	15	9	33	9,2	—	21	11	36,96	—	106	58	35,27	
	19	9	16	58,9	—	21	11	10,92	—	107	0	30,44	
	22	9	4	52,9	—	21	10	52,09	—	107	1	49,43	
	24	8	56	49,5	—	21	10	40,55	—	107	2	37,12	
	25	8	52	47,60	—	21	10	35,00	—	107	3	58,84	
	27	8	44	45,1	—	21	10	24,67	—	107	4	42,97	
	30	8	32	45,1	—	21	10	9,64	—	107	5	41,68	
October	7	8	4	43,1	—	21	9	42,03	—	107	6	31,43	
	12	7	44	49,4	—	21	9	27,56	—	107	7	27,56	
	14	7	36	53,4	—	21	9	23,13	—	107	7	44,56	
	23	7	1	20,0	—	21	9	12,68	—	107	8	11,40	
	26	6	49	38,2	—	21	9	12,84	—	107	8	2,81	
	27	6	45	47,0	—	21	9	13,50	—	107	7	59,45	
	28	6	41	41,8	—	21	9	14,50	—	107	7	53,53	
	29	6	37	46,6	—	21	9	14,89	—	107	7	54,04	
November	3	6	18	14,7	—	21	9	22,34	—	107	7	8,15	
	5	6	10	27,1	—	21	9	26,78	—	107	6	41,98	
	9	5	54	54,8	—	21	9	28,34	—	107	6	45,54	
	10	5	51	2,0	—	21	9	41,18	—	107	5	28,52	

Apparent Right Ascension and North Polar Distance of GEORGIAN, continued.

1833	Madras Mean Time of Observations.			Point Observed.	A. R.			Point Observed.	N. P. D.			REMARKS.	
	h.	m.	s.		h.	m.	s.		°	'	"		
August	29	10	59	44,0	Centre.	21	30	27,07	Centre.	105	33	18,40	
September	10	10	11	12,0	—	21	29	6,47	—	105	39	19,19	
	11	10	7	15,7	—	21	29	6,64	—	105	39	18,15	
	13	9	59	39,4	—	21	28	22,06	—	105	43	4,55	
	15	9	50	33,1	—	21	28	7,17	—	105	44	15,11	
	17	9	42	26,7	—	21	27	52,50	—	105	45	22,28	
	18	9	38	23,9	—	21	27	45,60	—	105	45	53,94	
	20	9	29	18,1	—	21	26	31,45	—	105	46	56,33	
	21	9	26	16,4	—	21	27	24,73	—	105	47	26,87	
	30	8	50	2,2	—	21	26	31,11	—	105	51	27,09	
October	2	8	41	57,3	—	21	26	21,24	—	105	57	11,77	
	4	8	33	55,9	—	21	26	11,53	—	105	52	54,58	
	6	8	25	54,0	—	21	26	2,71	—	105	53	31,95	
	7	8	21	55,4	—	21	25	58,05	—	105	53	50,75	
	14	7	53	58,9	—	21	25	33,95	—	105	55	32,83	
	15	7	50	0,1	—	21	25	31,09	—	105	55	42,86	
	16	7	46	2,4	—	21	25	29,00	—	105	55	52,95	
	17	7	42	3,6	—	21	25	26,57	—	105	56	4,40	
	22	7	22	6,9	—	21	25	17,70	—	105	56	33,83	
	25	7	10	23,5	—	21	25	14,55	—	105	56	42,10	

Apparent Right Ascension and North Polar Distance of PALLAS.

1832	Madras Mean Time of Observations.			A. R. from Observation.			A. R. from Tables.			Error of Tables.	N. P. D. from Observation.			N. P. D. from Tables.			Error of Tables.	
	h.	m.	s.	h.	m.	s.	m.	s.	s.		°	'	"	'	"	"		
Sept.	24	11	27	39,3	23	41	55,03	41	53,59	—	1,44	95	59	12,82	58	30,9	—	41,9
	25	11	22	58,5	23	41	10,03	41	8,34	—	1,69	96	13	23,18	12	46,4	—	36,8
Oct.	1	10	54	58,7	23	36	44,78	36	43,60	—	1,18	97	37	2,49	36	24,9	—	37,6

Apparent Right Ascension and North Polar Distance of CERES.

1832	Madras Mean Time of Observations.			A. R. from Observation.			A. R. from Tables.			Error of Tables.	N. P. D. from Observation.			N. P. D. from Tables.			Error of Tables.	
	h.	m.	s.	h.	m.	s.	m.	s.	s.		°	'	"	'	"	"		
Oct.	23	12	38	8,6	2	46	56,33	46	56,75	+	0,42	85	40	57,96	40	59,1	+	1,1
	24	12	33	20,5	2	46	4,14	46	4,63	+	0,49	85	43	22,87	43	24,2	+	1,3
	25	12	26	32,8	2	45	11,56	45	11,93	+	0,37	85	45	42,61	45	46,5	+	3,9
	26	12	23	43,2	2	44	18,26	44	18,77	+	0,51	85	48	0,05	48	2,9	+	2,9
	27	12	19	33,8	2	43	24,68	43	25,09	+	0,41	85	50	14,13	50	15,9	+	1,8
	29	12	13	13,1	2	41	36,06	41	36,57	+	0,51	85	54	25,12	54	28,2	+	3,1
	30	12	4	23,1	2	40	41,24	40	41,86	+	0,62	85	56	22,07	56	27,0	+	4,9
	31	11	59	32,6	2	39	46,26	39	46,93	+	0,67	85	58	15,91	58	20,5	+	4,6
Nov.	1	11	54	42,9	2	38	51,43	38	51,78	+	0,35	86	0	4,15	0	8,8	+	4,7
	2	11	49	50,4	2	37	56,16	37	56,54	+	0,38	86	1	46,09	1	51,5	+	5,4
	3	11	44	59,9	2	37	0,78	37	1,26	+	0,48	86	3	24,61	3	28,1	+	3,5

Apparent Right Ascension and North Polar Distance of CERES, continued.

1832	Madras Mean Time of Observations.			A. R. from Observation.			A. R. from Tables.		Error of Tables.	N. P. D. from Observation.			N. P. D. from Tables.		Error of Tables.			
	h.	m.	s.	h.	m.	s.	m.	s.	s.	°	'	"	'	"	"			
Nov.	4	11	40	8,8	2	36	5,51	36	5 99	+	0,48	86	4	55,89	4	58,7	+	2,8
	5	11	35	17,5	2	35	10,31	35	10 82	+	0,51	86	6	19,18	6	22,9	+	4,7
	12	11	0	28,9	2	28	50,76	28	51,26	+	0,50	86	12	55,68	13	1,3	+	5,6

Apparent Right Ascension and North Polar Distance of JUNO.

1833	Madras Mean Time of Observations.			A. R. from Observation.			A. R. from Tables.		Error of Tables.	N. P. D. from Observation.			N. P. D. from Tables.		Error of Tables.			
	h.	m.	s.	h.	m.	s.	m.	s.	s.	°	'	"	'	"	"			
April	27	12	57		15	20		19	59,74		92	47	19,53	48	17,4	-	2,1
	28	12	53	10,5	15	19	16,68	19	13,73	-	2,95	92	42	27,03	42	26,7	-	0,3
	29	12	48	28,1	15	18	29,99	18	27,29	-	2,70	92	36	39,82	36	40,4	+	0,6
May	2	12	34	18,7	15	16	9,87	16	6,17	-	3,70	92	19	52,96	19	49,3	-	3,7
	8	12	5	56,4	15	11	20,33	11	17,24	-	3,09
	9	12	1	12,1	15	10	31,15	10	28,68	-	2,77
	10	11	56	27,6	15	9	43,39	9	40,14	-	3,25	92	39	4,85	38	57,8	-	7,0
	11	11	51	42,9	15	8	54,28	8	51,65	-	2,63	92	34	27,85	34	20,6	-	7,2
	12	11	46	59,8	15	8	6,82	8	3,26	-	3,56
13	11	42	15,8	15	7	18,38	7	15,03	-	3,35	

In consequence of the extreme faintness of JUNO; in making the above observations it was found necessary to exclude all the light from the field, and even then, it was seen with the greatest difficulty; from this circumstance the transits which in general could only be observed at one or two wires are less accurate than the observations of the other Planets.

Apparent Right Ascension and North Polar Distance of VESTA.

1833	Madras Mean Time of Observations.			A. R. from Observation.			A. R. from Tables.		Error of Tables.	N. P. D. from Observation.			N. P. D. from Tables.		Error of Tables.	
	h.	m.	s.	h.	m.	s.	m.	s.	s.	°	'	"	'	"	"	
July	7	11	58		19		0	36,97		113	0	59,64	0	25,0
	8	11	53	42,3	18	59	34,28	59	36,39	+	2,11

The prevalence of clouds and rain prevented further observation of VESTA.

The places with which the observations of the above four Planets are compared are interpolated from the Supplements to the Nautical Almanac which are "deduced from the Berliner Astronomisches Jahrbuch for 1833,

page 109" : not having a copy of this work or indeed any tables of the Planetary Motions, has prevented my offering a similar comparison of the places of the larger Planets.

In the next place we come to the observations of the Moon, before giving which, it will be proper to state the elements which have been employed in the reduction of the observation ; they are as follows.

Ratio of Polar and Equatoreal Axes.....	299 : 300
From which we find the <i>Angle of the Vertical</i>	5' 0"
And the radius of the Earth.....	,999825
Semi-diameter.....	} Computed from the Nautical Almanac.
Parallax.....	

In addition to the above it is necessary I should here state, that the column *mean time* which now follows, is for the instant of the first limb, centre, or second limb, transiting the meridian as the case may be ; at which instant, the Right Ascension of the Moon's *centre* (computed from the observation) is given, and compared with the interpolated place from the Nautical Almanac: Now the observed N.P.D. being necessarily due to the moment of the Moon's *centre* being on the meridian, will correspond to a mean time greater or less than the above according to the circumstance of the first or second limb having been observed ; to obviate the inconvenience which would thus result, I have applied to the reduced North Polar Distance the change of declination due to the interval occupied by the Moon's semi-diameter to pass the Meridian, or in other words the Declination here given is reduced to correspond with the mean time at which the Transit was observed.

Comparison of the observed Right Ascension and North Polar Distance of the Moon with the interpolated place from Nautical Almanac.

1832	Madras Mean Time.			Limb Observed.	Observed A. R. of D's Centre.			A. R. from Nautical Almanac.		Error of Tables.	Limb Observed.	Observed N. P. D. of D's Centre.			N.P.D. from Nautical Almanac.		Error of Tables.					
	h	m	s		°	'	"	'	"	"		°	'	"	'	"	"					
Jan.	13	8	9	54	88	1	54	51	5,4	51	9,7	+	4,3	S.	14	16	36,0	N.	16	28,1	-	7,9
	14	9	7	43	36	1	70	19	44,6	19	47,3	+	2,7	S.	17	33	49,2	N.	33	46,3	-	2,9
	15	10	8	52	03	1	86	38	58,0	38	57,2	-	0,8	S.	19	39	53,6	N.	39	1,4	+	7,8
	17	12	17	27	85	1	120	16	23,4	16	18,9	-	4,5	S.	19	14	9,2	N.	14	9,3	+	0,1
	25	19	0	41	99	2	229	17	7,5	17	5,3	-	2,2	N.	12	47	5,3	S.	46	49,5	-	15,8
Feb.	9	6	2	23	08	1	49	29	11,4	29	1,9	-	9,5	S.	12	49	55,9	N.	50	1,5	+	5,6
	10	6	56	35	32	1	61	4	3,6	4	0,9	-	2,7	S.	16	21	22,7	N.	21	26,2	+	3,5
	11	7	53	58	74	1	79	26	50,9	26	48,2	-	2,7	S.	18	53	26,2	N.	53	34,9	+	8,7

1833	Madras Mean Time.			Limb Observed.	Observed A. R. of D's Centre.			A. R. from Nautical Almanac.			Error of Tables.	Limb Observed.	Observed N. P. D. of D's Centre.			N.P.D. from Nautical Almanac.			Error of Tables.
	h	m	s		o	'	"	o	'	"			+	-	o	'	"	o	
Oct.	25	9	46	55,71	1	0	47	37,2	47	40,3	+ 3,1	S.	5	4	14,1	S.	4	13,3	- 0,8
Nov.	18	5	32	2,87	1	320	34	22,5	34	9,9	- 12,0	S.
	19	6	17	31,84	1	332	57	11,8	57	4,4	- 7,4	S.	15	34	3,3	S.	34	15,1	+ 11,8
	20	7	0	53,70	1	344	48	18,3	48	7,4	- 10,9	S.	11	34	17,8	S.	34	20,3	+ 2,5
	21	7	42	51,08	1	356	18	9,4	18	0,0	- 9,1	S.	7	6	22,1	S.	6	27,5	+ 5,4
	23	8	24	12,04	1	7	39	28,0	39	21,8	- 6,2	S.	2	19	38,9	S.	19	41,7	+ 2,8
Dec.	18	5	37	47,70	1	351	34	1,9	34	2,3	+ 0,4	S.	9	7	42,5	S.	7	41,4	+ 1,1
	19	6	19	4,74	1	2	54	3,7	53	52,9	- 10,8	S.	4	28	41,9	S.	28	39,4	- 2,9
	20	7	0	4,98	1	14	2	42,5	2	53,3	+ 10,8	S.	0	22	42,3	N.	22	49,2	+ 6,9
	22	8	25	20,77	1	37	31	26,6	31	15,8	- 10,3	S.	10	6	3,8	N.	6	15,6	+ 11,8
	23	9	11	39,65	1	50	7	6,2	7	3,6	- 2,6	S.	14	35	12,0	N.	35	17,8	+ 5,8
	24	10	1	30,91	1	63	36	51,6	36	54,1	+ 2,5	S.	18	28	43,5	N.	28	46,5	+ 3,0
	25	10	55	25,29	1	78	6	31,5	6	36,7	+ 5,2	N.	21	27	12,2	N.	27	9,9	- 2,3
	26	11	54	3,13	1	93	31	26,0	31	26,7	+ 0,7	N.	23	10	8,0	N.	10	4,6	- 3,4
	27	12	56	6,54	1	109	31	17,2	31	15,6	- 1,6	S.	23	21	1,0	N.	20	56,6	- 4,4

Observation of the Eclipse of the Moon on the 1st July 1833.

	Shelton's Clock.			Madras Mean Time.			Observed by
	h.	m.	s.	h.	m.	s.	
Beginning of the Eclipse.....	23	3	55	16	25	59,8	
The Shadow covers Mare Ilmoruim.....	} 23	8	36	16	30	40,1	A.
The Shadow touches Grimaldus.....							T.
The Shadow covers Grimaldus.....	} 23	13	28	16	35	31,3	T.
The Shadow touches Tycho.....							A.
The Shadow covers Tycho.....	} 23	14	55	16	36	58,1	T.
The Shadow touches Tycho.....							A.
The Shadow covers Tycho.....	} 23	17	5	16	39	7,7	A.
The Shadow covers Tycho.....							T.
The Shadow covers Tycho.....	} 23	18	25	16	40	27,5	A.
The Shadow covers Tycho.....							T.
The Shadow covers Galileus.....	} 23	22	40	16	44	41,8	T.
The Shadow covers Galileus.....							A.
The Shadow covers Galileus.....	} 23	22	45	16	44	46,8	A.
The Shadow touches Keplerus.....							A.
The Shadow covers Keplerus.....	} 23	27	15	16	49	16,1	A.
The Shadow covers Keplerus.....							A.
The Shadow covers Keplerus.....	} 23	29	55	16	51	55,7	A.
The Shadow touches Copernicus.....							A.
The Shadow covers Copernicus.....	} 23	36	8	16	58	7,6	A.
The Shadow covers Copernicus.....							T.
The Shadow covers Copernicus.....	} 23	36	28	16	58	27,5	T.
The Shadow covers Copernicus.....							A.
The Shadow covers Copernicus.....	} 23	38	10	17	0	9,3	A.
The Shadow covers Copernicus.....							T.
The Shadow covers Aristarchus—somewhat uncertain.....	} 23	38	12	17	0	11,3	T.
The Shadow covers Aristarchus—somewhat uncertain.....							T.
The Shadow covers Aristarchus—somewhat uncertain.....	} 23	43	40	17	5	38,4	T.

Trees prevented further observation,

The Earth's Shadow was not well defined, and the observations altogether were in consequence unsatisfactory particularly towards the latter observations.

The observations marked T were made by myself with Dollond's 5 feet Achromatic with the lowest power (60); those marked A, were made by my Head Assistant with Dollond's 42 Inch Achromatic power 75; a lower power was much wanted for this nature of observation.

Observation of the Eclipse of the Moon on the 27th December 1833.

	Shelton's Clock.			Madras Mean Time.			Observed by
	<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>h.</i>	<i>m.</i>	<i>s.</i>	
Beginning of the Eclipse the Shadow was not well defined.	7	23	50	13	0	11,9	T.
The Shadow covers Aristarchus.....	7	33	28	13	9	48,4	—
The Shadow covers Copernicus.....	7	45	28	13	21	46,4	—
The Shadow touches Plato.....	7	49	23	13	25	40,8	—
The Shadow covers Aristoteles.....	7	58	10	13	34	26,3	—
The Shadow covers Eudoxus.....	7	58	15	13	34	31,3	—
The Shadow touches Tycho.....	7	58	41	13	35	0,2	—
The Shadow covers Tycho.....	7	59	22	13	35	38,1	—
The Shadow covers Plinius.....	8	4	45	13	41	0,2	—
The Shadow covers Meshalæ.....	8	12	2	13	48	16,0	—
The Shadow covers Proclus.....	8	14	35	13	50	46,6	—
The Shadow covers Mare Crisium.....	8	19	34	13	55	46,8	—
The Shadow covers the Moon.....	8	24	28	11	0	40,0	—
End of Total Darkness.....	10	2	55	15	38	51,0	—
The Shadow covers Grimaldus.....	10	5	50	15	41	45,5	—
The Shadow leaves Grimaldus.....	10	6	45	15	42	40,4	—
The Shadow leaves Galileus.....	10	10	14	15	46	8,8	—
The Shadow leaves Aristarchus.....	10	14	16	15	50	10,1	—
The Shadow covers Tycho.....	10	28	13	16	4	4,8	—
The Shadow leaves Tycho.....	10	29	10	16	5	1,8	—
The Shadow leaves Manilius.....	10	41	0	16	16	49,9	—
The Shadow covers Mare Crisium.....	10	58	20	16	34	7,1	—
The Shadow leaves Mare Crisium.....	11	0	0	16	35	46,8	—
End of the Eclipse.....	11	2	50	16	38	36,3	—

The above observations were made by myself with Dollond's 5 feet Achromatic power 60; the air was beautifully clear, and with the exception of the beginning, I have never seen the Earth's Shadow better defined; the observations though few in numbers in consequence of the rapid deposition of dew (which obliged me to stop every five minutes to wipe the object Glass), are nevertheless to be depended upon.

Eclipses of Jupiter's Satellites observed in the years 1832 and 1833.

1832		<i>h. m. s.</i>		<i>h. m. s.</i>	
Sept.	26—Emersion of Jupiter's first Satellite with 5 feet Achromatic power 160 at.....	20	2 45	or	7 40 26,1 M. T.
	Do. with 46 Inches Achromatic power 75 at..	20	2 55	or	7 40 36,1 M. T.
	Air very clear, and good observation.				
Oct.	3—Emersion of Jupiter's first Satellite with 5 feet Achromatic power 130 at.....	22	26 18	or	9 35 31,5 M. T.
	Do. with 46 Inches Achromatic power 75 at..	22	26 18	or	9 35 31,5 M. T.
	A little haze, but observation satisfactory.				
	9—Emersion of Jupiter's second Satellite with 46 Inches Achromatic power 75 at....	23	30 50	or	10 15 35,8 M. T.
	Observation satisfactory.				
Nov.	2—Emersion of Jupiter's first Satellite with 5 feet Achromatic power 120 at.....	2	35 35	or	11 46 41,2 M. T.
	Moon light clear, observation good.				
	3—Emersion of Jupiter's second Satellite with 5 feet Achromatic power 130 at....	22	11 30	or	7 19 23,3 M. T.
	Do. with 46 Inches Achromatic power 75 at..	22	12 10	or	7 20 3,2 M. T.
	Moon light very clear, observation satisfactory.				
	11—Emersion of Jupiter's first Satellite with 46 Inches Achromatic power 75 at.....	23	35 37	or	8 11 34,6 M. T.
	Do. with 5 feet Achromatic power 130 at....	23	35 42	or	8 11 39,6 M. T.
	Observation good.				
	17—Immersion of Jupiter's third Satellite with 46 Inches Achromatic power 75 at ...	0	15 50	or	8 28 32,0 M. T.
	17—Emersion of Jupiter's third Satellite with 46 Inches Achromatic power 75 at....	3	12 30	or	11 24 43,6 M. T.
Dec.	4—Emersion of Jupiter's first Satellite with 46 Inches Achromatic power 75 at.....	1	22 45	or	8 29 39,5 M. T.
	5—Emersion of Jupiter's second Satellite with 46 Inches Achromatic power 75 at....	23	57 30	or	7 0 46,7 M. T.
1833					
Jan.	12—Emersion of Jupiter's first Satellite with 5 feet Achromatic power 130 at.....	2	30 49	or	7 6 58,3 M. T.
	Do. with 46 Inches Achromatic power 75 at..	2	30 55	or	7 7 4,3 M. T.
	13—Emersion of Jupiter's second Satellite with 5 feet Achromatic power 60 at.....	4	47 10	or	9 19 4,6 M. T.
	Planet low, clear, observation good.				
	19—Emersion of Jupiter's first Satellite with 5 feet Achromatic power 110 at.....	4	58 15	or	9 2 34,4 M. T.

		<i>h. m. s.</i>	<i>h. m. s.</i>
Feb.	27—Emersion of Jupiter's first Satellite with 46 Inches Achromatic power 75, at.....	6 7 8	or 7 37 14,7 M. T.
	4° A above the horizon, observation doubtful.		
June	11—Immersion of Jupiter's first Satellite with 5 feet Achromatic power 130, at.....	20 57 46	or 15 37 1,2 M. T.
	Moon light, clear, observation satisfactory.		
	27—Emersion of Jupiter's third Satellite with 5 feet Achromatic power 130, at.....	22 22 25	or 16 0 5,7 M. T.
	Clear, observation good.		
July	4—Immersion of Jupiter's first Satellite with 46 Inches Achromatic power 75, at....	22 31 48	or 15 45 18,8 M. T.
	Thin haze, observation satisfactory.		
	20—Immersion of Jupiter's second Satellite with 46 Inches of Achromatic power 75, at..	0 3 20	or 16 11 8,2 M. T.
	27—Immersion of Jupiter's first Satellite with 46 Inches Achromatic power 75, at....	0 15 45	or 15 56 4,1 M. T.
	Observation satisfactory.		
Aug.	5—Immersion of Jupiter's first Satellite with 46 Inches Achromatic power 75, at....	21 12 26	or 12 17 57,9 M. T.
	9—Immersion of Jupiter's third Satellite with 46 Inches Achromatic power 75, at....	22 53 25	or 13 41 23,8 M. T.
	Thin haze, observation satisfactory.		
	14—Immersion of Jupiter's second Satellite with 5 feet Achromatic power 110, at.....	22 54 34	or 13 23 4,5 M. T.
	Clear, observation good.		
	14—Emersion of Jupiter's second Satellite with 5 feet Achromatic power 110, at.....	1 20 29	or 15 48 34,7 M. T.
Sept.	11—Immersion of Jupiter's first Satellite with 5 feet Achromatic power 110, at.....	3 35 35	or 16 14 22,7 M. T.
	Observation fair.		
	14—Emersion of Jupiter's third Satellite with 46 Inches Achromatic power 75, at....	21 15 45	or 9 43 51,4 M. T.
	Do. with 5 feet Achromatic power 110, at....	21 15 50	or 9 43 56,4 M. T.
	14—Emersion of Jupiter's third Satellite with 5 feet Achromatic power 110, at.....	23 34 35	or 12 2 18,9 M. T.
	21—Immersion of Jupiter's third Satellite with 5 feet Achromatic power 110, at.....	1 45 55	or 13 45 55,0 M. T.
	By reason of haze, this observation can only be considered one of second rate accuracy.		
Oct.	6—Immersion of Jupiter's first Satellite with 5 feet Achromatic power 110, at.....	23 53 36	or 10 55 16,8 M. T.

- h. m. s. h. m. s.
- Oct. 13—Immersion of Jupiter's first Satellite
with 5 feet Achromatic power 180, at..... 2 15 59 or 12 49 51,8 M. T.
Very good observation.
- 15—Immersion of Jupiter's first Satellite
with 46 Inches Achromatic power 75, at.... 20 51 38 or 7 18 34,3 M. T.
Do. with 5 feet Achromatic power 180, at... 20 51 40 or 7 18 36,3 M. T.
- 20—The first appearance of the Emerision of
Jupiter's third Satellite was from be-
hind the body of the Planet at..... 22 8 15 or 8 15 13,5 M. T.
As seen through the 5 feet Achroma-
tic with a power 150.
It appeared in contact with the body of
Jupiter at..... 22 16 20 or 8 23 12,3 M. T.
- 31—Emerision of Jupiter's first Satellite with
5 feet Achromatic power 110, at..... 22 21 20 or 7 44 43,2 M. T.
Do. with 46 Inches Achromatic power 75 at.. 22 21 40 or 7 45 3,1 M. T.
Clear, observation good.
- Nov. 23—Emerision of Jupiter's first Satellite with
5 feet Achromatic power 110, at..... 0 8 58 or 7 59 15,7 M. T.
Do. with 46 Inches Achromatic power 75 at.. 0 9 19 or 7 59 36,6 M. T.
- Dec. 2—Emerision of Jupiter's third Satellite
with 42 Inches Achromatic power 75, at.... 0 56 33 or 8 11 34,1 M. T.
Clear, observation satisfactory.
- 6—Emerision of Jupiter's second Satellite
with 5 feet Achromatic power 180, at..... 2 31 10 or 9 30 14,6 M. T.
- 9—Immersion of Jupiter's third Satellite
with 42 Inches Achromatic power 75, at.... 3 16 35 or 10 3 56,8 M. T.
- 9—Emerision of Jupiter's third Satellite
with 5 feet Achromatic power 110, at..... 5 27 12 or 12 14 2,6 M. T.

Not being possessed of any Greenwich or Cambridge Observations corres-
ponding to these we will now compare them with the times given in the
Nautical Almanac, from which we determine.

The Longitude of the Madras Observatory.

1832	Im. or Em.	I Satellite.	Im. or Em.	II Satellite.	Im. or Em.	III Satellite.	REMARKS.
September 26	E.	5 21 11,1	
October 3	E.	5 20 55,5	
9	E.	5 21 10,8	
November 2	E.	5 20 57,2	
3	E.	5 20 51,3	

The Longitude of the Madras Observatory, continued.

1832	Im. or Em.	I Satellite.	Im. or Em.	II Satellite.	Im. or Em.	III Satellite.	REMARKS.
		<i>h. m. s.</i>		<i>h. m. s.</i>		<i>h. m. s.</i>	
November 11	E.	5 20 57,1	
17	I.	5 23 3,0	
17	E.	5 21 7,6	
December 4	E.	5 21 59,5	
5	E.	5 21 18,7	
1833							
January 12	E.	5 21 31,3	
13	E.	5 22 11,6	
19	E.	5 21 15,4	
27	E.	5 20 36,7	
June 11	I.	5 21 24,2	
27	E.	5 20 58,7	
July 4	I.	5 20 0,0	
20	I.	5 21 36,2	
27	I.	5 21 21,1	
August 5	I.	5 21 10,9	
9	I.	5 24 34,8	
14	I.	5 21 57,5	
14	E.	5 21 58,7	
September 11	I.	5 19 58,7	
14	I.	5 22 42,9	
14	E.	5 22 50,9	
21	I.	5 23 42,0	
October 6	I.	5 20 59,8	
13	I.	5 21 3,8	
15	I.	5 21 6,3	
31	E.	5 20 56,2	
November 23	E.	5 20 50,2	
December 2	E.	5 20 53,1	
6	E.	5 21 19,6	
9	I.	5 22 16,8	
9	E.	5 21 11,6	
Mean.....	..	5 21 0,8	..	5 21 33,0	..	5 22 20,1	

Taking the Mean of the first and second Satellites whose places are much more accurately observed and computed than those of the third Satellite we obtain:

The Longitude of the Madras Observatory.

5h. 21m. 7,7s: East of Greenwich.



Occultations of the Planet SATURN and of Stars, in the years 1832 and 1833.

1832

April 11—Immersion of Saturn behind the Moon's Dark Limb observed with 42 Inches Achromatic power 75.

The first edge of the ring was lost.

Centre of the Body in contact with Moon's Dark Limb at 12h. 4m. 17,0s. by Shelton's Clock or 10h. 41m. 36,7s. Madras Mean Time.

The Second edge of the Limb at 12h. 4m. 39,5s. by Shelton's Clock or 10h. 41m. 59,1s. Madras Mean Time.

Sept. 27—Immersion of a Star behind the Moon's Dark Limb with 42 Inches Achromatic power 75, at 19h. 8m. 22,5s. by Shelton's Clock or 6h. 42m. 20,4. Madras Mean Time.

Clear, observation good.

Nov. 3—Immersion of ψ Aquarii behind the Moon's Dark Limb with 42 Inches Achromatic power 75, at 23h. 10m. 31,0s. by Shelton's Clock or 8h. 18m. 14,6s. Madras Mean Time.

Observation certain to a second.

29—Immersion of β Arietis behind the Moon's Dark Limb with 42 Inches Achromatic power 75, at 2h. 18m. 53,0s. by Shelton's Clock or 9h. 44m. 45,7s. Madras Mean Time.

Clear, observation good.

Dec. 25—Immersion of θ Capricorni behind the Moon's Dark Limb with 42 Inches Achromatic power 75, at 0h. 47m. 28,5s. by Shelton's Clock or 6h. 33m. 30,6. Madras Mean Time.

Observation satisfactory.

27—Immersion of γ Aquarii behind the Moon's Dark Limb with 5 feet Achromatic power 110, at 2h. 8m. 54,5s. by Skelton's Clock or 7h. 47m. 1,0s. Madras Mean Time.

I fancied the Star slightly projected upon the Moon's Disc at Immersion but my Assistant with the 42 Inches Achromatic did not so see it.

Do. —With 42 Inches Achromatic power 75, at 2h. 8m. 54,0s. by Shelton's Clock or 7h. 47m. 1,5s. Madras Mean Time.

1833

March 24—Immersion of μ Ceti behind the Moon's Dark Limb with 5 feet Achromatic power 110, at 7h. 37m. 34,0s. by Shelton's Clock or 7h. 28m. 36,4s. Madras Mean Time.

25—Immersion of a small Star behind the Moon's Dark Limb with 5 feet Achromatic power 110, at 8h. 2m. 55,5s. by Shelton's Clock or 7h. 49m. 57,7s. Madras Mean Time.

Clear, observation good.

26—Immersion of a small Star behind the Moon's Dark Limb with 5 feet Achromatic power 110, at 7h. 9m. 33,0s. by Shelton's Clock or 6h. 52m. 47,1s. Madras Mean Time.

1833

March 26—Immersion of ϵ Tauri behind the Moon's Dark Limb with 5 feet Achromatic power 110, at 7h. 10m. 38,5s. by Shelton's Clock or 6h. 53m. 52,4s. Madras Mean Time.

Clear, observation good.

Emmersion of ϵ Tauri from behind the Moon's Bright Limb with 5 feet Achromatic power 110, at 7h. 46m. 20,0s. by Shelton's Clock or 7h. 29m. 28,1s. Madras Mean Time.

Immersion of a Star in Taurus behind the Moon's Dark Limb with 5 feet Achromatic power 110, at 7h. 53m. 49,5s. by Shelton's Clock or 7h. 36m. 56,4s. Madras Mean Time.

Observed Transits of the Moon and of Stars, culminating near thereto, in the years 1832 and 1833.

1832	NAMES.	Observed Transit.			1832	NAMES.	Observed Transit.		
		h.	m.	s.			h.	m.	s.
Jan. 13	D 1 Limb.....	3	42	12,09	Sept. 5	D 1 Limb.....	19	31	43,97
	48 Tauri.....	4	10	10,03		σ Capricorni.....	20	7	43,06
15	D 1 Limb.....	5	49	24,13		π Capricorni.....	20	15	43,48
	ν Geminorum...	6	1	58,23	Oct. 3	57 Sagittarii.....	19	43	45,10
Feb. 11	D 1 Limb.....	5	18	0,70		D 1 Limb.....	20	6	50,04
	χ^3 Orionis.....	5	54	19,42		ν Capricorni.....	20	31	48,79
12	D 1 Limb.....	6	21	9,84		4 ν Capricorni.....	20	32	1,20
	ζ Geminorum.....	6	54	33,60		D 1 Limb.....	20	58	24,09
23	ϕ Ophiuchi.....	16	22	28,11		δ Capricorni.....	21	39	19,14
	D 2 Limb.....	16	39	3,31		8 P Piscium.....	23	52	9,37
March 13	α^1 Cancri.....	8	49	31,20		S Piscium.....	23	58	49,16
	D 1 Limb.....	9	8	10,03		D 1 Limb.....	0	17	34,42
	ν Leonis.....	9	50	49,63		26 Ceti.....	0	57	15,80
14	ν Leonis.....	9	50	52,15		9 26 Ceti.....	0	57	20,04
	D 1 Limb.....	10	7	48,83		D 1 Limb.....	1	7	34,00
April 13	D 1 Limb.....	12	27	40,59		D 2 Limb.....	1	9	42,22
	k^4 Virginis.....	12	54	20,36		ν Piscium.....	1	34	51,31
14	k^4 Virginis.....	12	54	22,48		30 D 1 Limb.....	19	45	7,18
	D 1 Limb.....	13	18	56,61		σ Capricorni.....	20	10	30,01
	88 Virginis.....	13	38	38,55		ν Capricorni.....	20	31	17,22
15	88 Virginis.....	13	38	14,85		31 σ Capricorni.....	20	10	30,56
	κ Virginis.....	14	2	40,20		ν Capricorni.....	20	31	17,59
	D 2 Limb.....	14	11	23,42		D 1 Limb.....	20	36	49,92
June 9	94 Virginis.....	13	55	58,87		ι Capricorni.....	21	13	42,22
	D 1 Limb.....	14	24	44,27	Nov. 1	ι Capricorni.....	21	13	42,39
	ξ^a Libræ.....	14	46	14,28		D 1 Limb.....	21	27	23,30
12	3 Ophiuchi.....	16	50	25,14		ι Aquarii.....	21	58	11,03
	D 1 Limb.....	16	55	55,09		D Aquarii.....	22	10	49,01
	52 Ophiuchi.....	17	23	35,71		2 ι Aquarii.....	21	58	11,61
Sept. 4	21 Sagittarii.....	18	13	20,36		D Aquarii.....	22	10	49,64
	D 1 Limb.....	18	39	21,07		D 1 Limb.....	22	16	56,43
	138 Sagittarii.....	19	18	55,81		70 Aquarii.....	22	40	29,74
	5,138 Sagittarii.....	19	18	56,79		3 70 Aquarii.....	22	40	30,72

1832		Observed Transit.			1833		Observed Transit.			
NAMES.		h.	m.	s.	NAMES.		h.	m.	s.	
Nov.	3	1	23	5	50,38	May	2	13	28	53,59
	r	Piscium	23	54	12,21		1	13	41	39,84
	4	190	23	40	54,04		2	14	16	30,34
		1	23	54	45,47		3	14	5	59,95
	5	1	0	44	16,75		2	14	16	27,30
	f	Piscium	1	10	17,21		1	14	37	7,11
	v	Piscium	1	33	50,88		2	14	37	7,11
	15	a	10	0	10,53	June	28	15	25	20,65
	p	Leonis	10	24	42,61		7	15	33	50,32
		2	10	36	18,26		1	15	41	27,17
	29	δ	21	37	33,60		χ	16	16	30,42
		1	21	55	46,63		χ	16	16	27,00
	70	Aquarii	22	39	27,82		1	16	35	45,24
	30	σ	22	21	28,72		ρ	17	10	5,84
	70	Aquarii	22	39	23,14		D	17	27	31,37
		1	22	44	9,44		1	17	31	2,03
Dec.	3	26	0	54	41,00	July	1	18	27	50,40
		1	1	8	56,38		1	18	53	38,05
	v	Ceti	1	32	12,23		π	18	58	47,35
	4	v	1	32	8,03		ν ¹	18	42	20,22
		225	1	50	37,10		f	18	52	55,45
		1	1	59	56,12		1	19	2	37,91
	μ	Ceti	2	35	18,50		f	19	34	51,83
	5	μ	2	35	13,90	Aug.	22	21	36	59,69
		1	2	53	52,30		μ	21	43	21,76
	6	1	3	51	29,80		1	22	14	32,37
	179	Tauri	4	35	46,04		2	22	14	32,37
	7	i	4	40	46,09		I	22	33	24,52
		1	4	54	15,52	Sept.	21	18	26	52,70
		2	4	54	15,52		26	18	29	50,28
1833							61	19	8	47,26
Jan.	13	2	14	22	34,92		σ	22	19	51,48
	a ^a	Librae	14	38	9,95	Oct.	20	22	44	16,21
Feb.	4	q	9	10	10,65		f	19	34	26,84
		2	9	17	37,74		1	19	55	33,51
March	2	p	7	19	6,57		20	20	47	56,51
		1	7	35	31,33		1	20	48	25,51
	29	1	7	11	50,45		γ	21	28	41,51
	l	Geminorum	7	47	42,14		δ	21	35	40,75
	31	q	9	11	27,85		γ	21	28	43,06
		1	9	17	56,32		δ	21	35	42,16
April	1	37	10	9	32,10		1	21	38	57,68
		1	10	19	9,54		f	22	19	14,27
	4	θ	13	3	11,66		f	22	19	14,76
		2	13	13	56,45		1	22	27	23,52
	28	1	9	58	25,45		h ¹	22	54	22,05
	p	Leonis	10	26	16,56		χ ³	23	8	11,22
	l	Leonis	10	42	44,27	Nov.	19	21	53	49,40
	29	p	10	26	13,75		1	22	11	17,57
	l	Leonis	10	42	41,42		70	22	40	12,71
		1	10	56	28,90		K	22	45	10,70
	v	Virginis	11	39	29,54		1	22	58	40,90
May	1	γ ¹	12	35	18,56		190	23	40	26,05
		1	12	47	26,24		1	23	44	39,07
	l ³	Virginis	13	28	56,92		60	0	16	26,57
							n	0	21	59,73

1833		NAMES.	Observed Transit.			1833		NAMES.	Observed Transit.		
			<i>h.</i>	<i>m.</i>	<i>s.</i>				<i>h.</i>	<i>m.</i>	<i>s.</i>
Nov.	22	60 Ceti.....	0	16	25,01	Dec.	22	4 Ceti.....	3	1	54,86
		" Ceti.....	0	21	58,17			f Ceti.....	3	21	22,39
		1 Limb.....	0	30	3,09		23	4 Ceti.....	3	1	52,14
Dec.	18	1 Limb.....	23	25	7,09			1 Limb.....	3	29	1,83
		P Piscium.....	23	49	59,57		24	1 Limb.....	4	12	56,52
		S Piscium.....	23	56	39,19			Tauri.....	4	52	45,36
	19	P Piscium.....	23	49	56,80			Tauri.....	4	57	33,86
		S Piscium.....	23	56	36,76		25	Tauri.....	4	52	43,09
		1 Limb.....	0	10	24,54			Tauri.....	4	57	31,44
		m Ceti.....	0	44	18,48			1 Limb.....	5	10	50,35
	20	m Ceti.....	0	44	16,25			Q ² Tauri.....	5	51	12,42
		1 Limb.....	0	55	25,99		26	Q ² Tauri.....	5	51	10,17
		μ Piscium.....	1	21	14,21			γ Geminorum.....	6	4	21,17
		ν Piscium.....	1	32	32,93			1 Limb.....	6	13	35,36
	22	1 Limb.....	2	28	43,38			2 Limb.....			

Selecting from the above those of which corresponding observations have been made at the Greenwich Royal Observatory, we have :

1832	NAMES.	Madras Observations			<i>t</i>	Greenwich Observations			<i>τ</i>	<i>t - τ</i>
		<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>m. s.</i>	<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>m. s.</i>	<i>m. s.</i>
March	13	o ⁺ Cancri.....	8	49	31,20	8	48	35,76
		1 Limb....	9	8	10,03	- 18 38,83	9	20	44,54	- 32 8,68 + 13 29,85
		ν Leonis....	9	50	49,63	+ 42 39,60	9	49	54,97	+ 29 10,43 + 13 29,17
April	13	1 Limb....	12	27	40,59	12	40	50,26
		k ⁺ Virginis...	12	54	20,35	+ 26 39,76	12	56	0 38	+ 15 10,12 + 11 29,66
	14	1 Limb....	13	18	56,61	13	31	53,76
		88 Virginis...	13	38	38,55	+ 19 41,94	13	40	15,96	+ 8 22,20 + 11 19,74
Oct.	8	p Piscium...	23	52	9,47	- 25 24,95	23	50	31,24	- 36 27,76 + 11 2,81
		S Piscium...	23	58	49,23	- 18 45,19	23	56	11,18	- 30 47,82 + 11 2,63
		1 Limb....	0	17	34,42	0	26	59,00
		26 Ceti.....	0	57	15,65	+ 39 41,23	0	55	37,54	+ 28 38,54 + 11 2,69
Nov.	4	190 Aquarii...	23	40	54,10	- 13 51,37	23	40	2,90	- 24 47,02 + 10 55,65
		1 Limb....	23	54	45,47	0	4	49,92
	29	1 Limb....	21	55	46,63	22	7	38,10
		70 Aquarii....	22	39	27,96	+ 43 41,33	22	40	26,90	+ 32 48,80 + 10 52,53

Similarly, for those of which corresponding observations were made at the Observatory at Cambridge, we have :

1832	NAMES.	Madras Observations			<i>t</i>	Cambridge Observations			<i>τ</i>	<i>t - τ</i>
		<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>m. s.</i>	<i>h.</i>	<i>m.</i>	<i>s.</i>	<i>m. s.</i>	<i>m. s.</i>
March	13	1 Limb....	9	8	10,03	9	19	40,20
		ν Leonis....	9	50	49,63	+ 42 39,60	9	48	51,25	+ 29 11,05 + 13 28,55
Sept.	4	1 Limb....	18	39	21,07	18	48	50,34

1832	NAMES.	Madras Observations	t	Cambridge Observations	τ	$t - \tau$
		<i>h. m. s.</i>	<i>m. s.</i>	<i>h. m. s.</i>	<i>m. s.</i>	<i>m. s.</i>
Sept. 4	138 Virginia...	19 18 55,78	+ 39 34,71	19 16 43,87	+ 27 53,53	+ 11 41,18
5	138 Virginia...	19 18 56,79	- 12 47,18	19 16 40,91	- 24 22,91	+ 11 35,73
	D 1 Limb....	19 31 43,97	19 41 3,32
	π Capricorni...	20 15 43,45	+ 43 59,48	20 13 27,46	+ 32 23,64	+ 11 35,84
Oct. 8	P Piscium....	23 52 9,47	- 25 24,95	23 49 12,66	- 36 26,94	+ 11 1,89
	S Piscium....	23 58 49,23	- 18 45,19	23 55 52,54	- 29 46,96	+ 11 1,77
	D 1 Limb....	0 17 34,42	0 25 39,50
Nov. 1	D 1 Limb....	21 27 23,30	21 36 34,29
	i Aquarii....	21 58 11,03	+ 30 47,73	21 56 15,82	+ 19 41,53	+ 11 6,20
4	190 Aquarii...	23 40 54,10	- 13 51,37	23 38 39,55	- 24 46,06	+ 10 54,69
	D 1 Limb....	23 54 45,47	0 3 25,61
1833						
Feb. 4	q Caneri....	9 10 10,65	- 7 27,09	9 9 23,32	- 21 58,12	+ 14 31,03
	D 2 Limb....	9 17 37,74	9 31 21,44
April 1	37 Leonis....	10 9 32,10	- 9 37,44	10 6 53,06	- 22 58,53	+ 13 21,09
	D 1 Limb....	10 19 9,54	10 29 51,59
28	D 1 Limb....	9 58 25,45	10 7 28,38
	ρ Leonis....	10 26 16,58	+ 27 51,13	10 22 12,53	+ 14 44,15	+ 13 6,98
May 3	χ Virginia...	14 5 59,89	- { 30 0,09 32 14,35	14 4 1,02	- { 42 10,40 44 25,16	+ { 12 10,31 12 10,81
	2 Libræ....	14 16 27,26	- { 19 32,72 21 46,98	14 14 28,33	- { 31 43,09 33 57,85	+ { 12 10,37 12 10,87
	D 1 Limb....	14 35 59,98	14 46 11,42
	D 2 Limb....	14 38 14,24	14 48 26,18
June 29	D 1 Limb....	16 35 45,24	16 47 46,49
	D Ophiuchi...	17 32 31,47	+ 56 46,23	17 32 17,83	+ 44 31,34	+ 12 14,89
July 1	D 1 Limb....	18 26 42,62	18 38 55,90
	σ Sagittarii...	18 53 38,14	+ 26 55,52	18 53 27,70	+ 14 31,80	+ 12 23,72
	π Sagittarii...	18 58 47,46	+ 32 4 84	18 58 37,20	+ 19 41,30	+ 12 23,54
Sept. 26	σ Aquarii....	22 19 51,46	- 24 24,75	22 21 1,31	- 34 50,87	+ 10 26,12
	D 1 Limb....	22 44 16,21	22 55 52,18
Oct. 20	f Sagittarii...	19 34 26,86	- 21 6,65	19 35 52,13	- 33 4,28	+ 11 57,63
	D 1 Limb....	19 55 33,51	20 8 56,41
23	f Aquarii....	22 19 14,77	- 8 8,75	22 20 28,23	- 18 40,61	+ 10 31,91
	D 1 Limb....	22 27 23,52	22 39 8,89
	h^1 Aquarii....	22 54 22,03	+ 26 58,51	22 55 35,54	+ 16 26,65	+ 10 31,86
	ψ^2 Aquarii....	23 8 11,18	+ 40 47,66	23 9 24,60	+ 30 15,71	+ 10 31,95

The above observations at Greenwich and Cambridge are extracted from the Monthly reports of the proceedings of the Royal Astronomical Society; but my copy of these not being complete; it is possible that a greater number of corresponding observations may have been made than are now given. In computing the Longitude I have in either case assumed $\alpha = 5h: 21m.$ in preference to computing the horary motion for the middle of the times of passage, and for the other elements of the computation have employed the Nautical Almanac; the values of the Longitude thus deduced are as follows:

1832	Longitude from Greenwich Observations.			Longitude from Cambridge Observations.			REMARKS.
	D. 1 L.	D. 2 L.		D. 1 L.	D. 2 L.		
March 13	h. m. s.	h. m. s.	52,47	h. m. s.	h. m. s.	5 20 27,59	
April 13	5 20	52,97					
14	5 21	4,05					
September 4				5 20	48,62		
5				5 20	20,51		
October 8	5 20	54,37		5 20	28,69		
November 1				5 20	23,01		
4	5 20	52,30		5 20	24,27		
29	5 20	46,60					
1833							
February 4						5 20 33,19	
April 4				5 20	34,31		
28				5 20	50,80		
May 3				5 20	26,25	5 20 34,01	
June 29				5 20	31,14		
July 1				5 20	46,05		
September 26				5 20	22,47		
20				5 20	20,32		
October 23				5 20	23,84		
Mean.....	5 20	55,94		5 20	30,56	5 20 33,60	

The observations of 1831, shew that a correction of about 12 seconds ought to be applied to the Longitude determined from the Observations of the Moon's first limb at Greenwich when compared with the Madras Observations; not having any corresponding observations at Greenwich of the Moon's second limb in 1832 and 1833, it will perhaps be the safer plan to allow these observations to remain as they are until further observations upon the Moon's second limb have been made; the Cambridge Observations offer two results of the longitude from the second limb, which taking the mean and allowing the longitude of Cambridge to be 23,54s. East of Greenwich, we obtain for the longitude of Madras 5h. 20m. 55,62s. a result which is probably from 5 to 10 seconds in defect.

Observation of the Transit of MERCURY, May 5, 1832, observed with Dollond's five feet Achromatic, power 110.

	Mean Time.		
	h.	m.	s.
The Exterior contact at ingress was lost.			
Centre of Planet (by estimation) in contact with the Sun's limb....	2	22	21,5
Interior contact at ingress.....	2	23	37,3
High wind which agitated the telescope prevented further observation.			

SUPPLEMENTARY OBSERVATIONS AND REMARKS.

Hitherto, in the reduction of the Madras Observations I have employed for the aberration, nutation, &c. the tables by Baily, published in the II Volume of the Royal Astronomical Society's Memoirs; the great care which was taken in the computation of these tables, and the superior judgment and skill of the Superintendent renders it unnecessary for me here to offer any explanation as to why I have given to them the preference; in the reduction of the observations of 1831, being obliged to compute the values A, B, C, D, I assumed for the value of the maximum of aberration 20,50s.* instead of 20,36s. which had been recommended by Mr. Baily; but in the reductions for 1832 and 1833, these numbers having been furnished in the Supplements to the Nautical Almanac computed for the latter co-efficient, I have considered the difference too unimportant to render a recomputation necessary. For the refraction, I have employed those very excellent tables by Mr. Henry Atkinson published in the III Volume of the Astronomical Society's Memoirs; independent of the elegance and originality which marks this memoir through every step of the investigation, the actual comparison of observations at low altitudes *below* the Pole at Greenwich, with those *above* the Pole, has given to these tables a preference to those of Ivory, Young, Brinkley, Bradley and Groombridge, with which they have been compared; as this comparison however does not extend to altitudes below 10°, it will at least be *interesting*, if not *useful* to fill up the blank; for this purpose we will select from the Greenwich Catalogue those Stars which are situated at 120° of N. P. D. or 8½° altitude, and downwards to the horizon; for at these altitudes the uncertainty of refraction amounts to several seconds, whereas at the altitudes which Stars thus situated attain to at Madras (from 47° to 39°), the uncertainty is comparatively very unimportant.

The Greenwich Catalogues which I am so fortunate as to possess, are; a Catalogue of 720 Stars observed (I believe) in 1827-29 and reduced to 1830; and the Catalogue published in Part 5, of the observations for 1831; these being reduced to January 1, 1832, and arranged in order of N. P. D. are as follows.

* Deduced from the Greenwich Observations by Mr. W. Richardson.

NAMES.	A. R.	Greenwich N. P. D. for 1832 from Observations.		Concluded N. P. D. Jan. 1, 1832	Bradley's Refraction Bar. 29,6 Ther. 50°.	Presumed Apparent N. P. D.			
		In 1828.	In 1831.						
ξ Sagittarii....	18 51	12	6 37,70	2	6 40,18	120 6 38,05	—	6 7,33	120 0 30,72
α Ault. Piscium	10 19	14	12 50,50	120 12 50,50	—	6 11,70	120 6 38,80
α App. Sculp..	0 50	14	15 58,79	120 15 58,79	—	6 13,85	120 9 44,96
γ ^s Sagittarii...	17 55	2	24 57,80	120 24 57,80	—	6 20,12	120 18 37,68
α Piscis. Aust..	22 48	96	30 36,20	18	30 36,87	120 30 36,31	—	6 24,17	120 24 12,14
ν ^a Eridani....	4 29	6	54 43,20	4	54 42,98	120 54 43,20	—	6 42,64	120 48 0,56
ξ Hyd. & Crat.	11 24	14	55 41,50	2	55 40,89	120 55 41,41	—	6 43,44	120 48 57,97
h Centauri....	13 43	10	5 37,29	121 5 37,29	—	6 51,55	120 58 45,74
θ Piscis. Aust..	21 37	2	40 13,64	121 40 13,64	—	7 22,29	121 32 51,35
k Centauri....	13 42	6	9 24,70	1	9 23,66	122 9 24,55	—	7 51,59	122 1 32,96
* Centauri....	13 36	13	11 26,21	122 11 26,21	—	7 53,90	122 3 32,31
κ ^a Canis Maj..	6 43	14	19 11,50	122 19 11,50	—	8 2,31	122 11 9,19
μ Columbæ....	5 40	2	22 36,21	122 22 36,21	—	8 5,95	122 14 30,26
D Canis Maj..	6 22	10	28 47,14	122 28 47,14	—	8 13,27	122 20 33,87
α Pixed Naut..	8 36	18	35 7,60	122 35 7,60	—	8 20,85	122 26 46,75
4 Piscis. Aust..	21 7	2	52 10,76	122 52 10,76	—	8 41,88	122 43 28,88
β Hydæ.....	11 44	10	58 28,10	122 58 28,10	—	8 49,60	122 49 38,50
χ Lupi.....	15 40	6	6 28,50	2	6 30,77	123 6 29,07	—	9 0,82	122 57 28,25
β Piscis. Aust..	22 22	14	12 17,30	2	12 21,05	123 12 17,77	—	9 8,83	123 3 8,94
λ Canis Maj..	6 16	10	21 28,50	123 21 28,50	—	9 22,05	123 12 6,45
g Centauri....	13 40	8	36 33,29	123 36 33,29	—	9 45,02	123 26 48,27
γ Piscis. Aust..	22 43	6	45 52,75	123 45 52,75	—	10 0,24	123 35 52,51
i Piscis. Aust..	21 35	14	47 17,30	123 47 17,30	—	10 2,56	123 37 14,74
k Scorpii.....	16 54	6	52 42,00	123 52 42,00	—	10 11,60	123 42 30,40
e Scorpii.....	16 39	10	58 49,70	123 58 49,70	—	10 22,33	123 48 27,37
α Columbæ....	5 33	8	10 14,40	6	124 10 14,40	—	10 42,75	123 59 31,65
* Canis Maj..	6 45	124 10 31,59	—	10 43,20	123 59 48,39
41 Eridani....	4 11	4	12 54,90	124 12 54,90	—	10 47,50	124 2 7,40
α Normæ....	16 20	10	19 52,84	124 19 52,84	—	11 1,40	124 8 51,44
α Microscopii..	20 39	6	23 49,30	2	*24 3,33	124 23 49,30	—	11 9,60	124 12 39,70
43 Eridani....	4 17	4	24 48,40	124 24 48,40	—	11 11,70	124 13 36,70
e Sagittarii....	18 12	9	27 22,10	124 27 22,10	—	11 16,90	124 16 5,20
β Pixed Naut..	8 33	2	43 21,05	124 43 21,05	—	11 50,30	124 31 30,75
κ Columbæ....	6 10	8	5 31,40	4	5 35,84	125 5 32,88	—	12 42,40	124 52 50,48
γ Columbæ....	5 51	3	18 31,90	2	+18 46,50	125 18 31,90	—	13 16,30	125 5 15,60
θ Centauri....	13 56	8	32 31,50	125 32 31,50	—	13 53,50	125 18 38,00

* This being only a single observation at each Instrument, I have rejected it and assumed the result of 1828.

† I have rejected this result as erroneous in consequence of the coincidence of the results of 1823, with the R. A. Society's Catalogue, for comparing it with κ Columbæ we have :

R. A. S. Catalogue 1830.	Greenwich Observations of 1828 reduced to 1830.	Greenwich Observations of 1831 reduced to 1830.
. / "	. / "	. / "
125 5 20,20	125 5 31,40	125 5 35,84
125 18 23,30	125 18 31,90	125 18 46,50
-----	-----	-----
Difference..13 3,10	13 0,50	13 10,66
-----	-----	-----

NAMES.	A. R.	Greenwich N. P. D. for 1832 from Observations.		Concluded N. P. D. Jan. 1, 1832.	Bradley's Refraction Bar. 29,6 Ther. 50°.	Presumed Apparent N. P. D.		
		In 1828.	In 1831.					
ϵ Columbæ	5 25	5	36 10,60	2	36 9,75	125 36 10,36	- 14 4,50	125 22 5,86
ϵ Centauri....	13 11	11	49 35,60	125 49 35,60	- 14 46,40	125 34 49,20
β Columbæ....	5 45	2	50 24,40	4	50 29,21	125 50 27,61	- 14 49,20	125 35 38,41
ϕ^2 Lupi.....	15 12	1	15 23,52	126 15 23,52	- 16 13,50	125 59 10,02
θ Lupi.....	15 55	1	20 33,73	126 20 33,73	- 16 33,40	126 4 0,33
β Telescopii...	18 6	6	48 39,70	5	48 30,46	126 48 35,50	- 18 28,70	126 30 6,80
λ Scorpii.....	17 22	1	58 37,30	5	58 38,43	126 58 38,24	- 19 14,20	126 39 24,04
γ Telescopii...	17 38	1	59 25,10	1	58 59,73	126 59 12,41	- 19 16,70	126 39 55,71

It will readily be understood that the above column "Barometer 29,6 Thermometer 50°" is computed from Bradley formula $r = 57",00 \times \tan. (z - 3r)$. If it should so happen that any of the Stars above given were observed under this pressure and at this temperature, then the column "Presumed Apparent N. P. D." will be the N. P. D. as actually observed at Greenwich corrected for aberration, &c. and reduced to the beginning of 1832; but since this can hardly be the case exactly in any one single instance, our "Presumed Apparent N. P. D." will be subject to the error which attaches to Bradley's correction for temperature and pressure; notwithstanding this apparently serious objection which (not being possessed of the Greenwich Observations for 1831) I am unable to remedy, and considering that the numbers for which we are seeking are large in proportion to the *uncertainty* of Bradley's correction for 10° or 15° of temperature or for 3 or 4 tenth of an inch of pressure (the probable extent for which the correction is required) I have ventured as a rough approximation to proceed with these computations.

‡ A single observation at this low altitude cannot much be depended upon; the difference between this Star and the preceding one compared with the Society's Catalogue is as follows:

<i>R. A. S. Catalogue</i> 1830.	<i>Greenwich Observations</i> of 1828, reduced to 1830.	<i>Greenwich Observations</i> of 1831 reduced to 1830.
126 58 3,48	126 58 30,70	126 58 31,83
126 58 41,54	126 59 21,30	126 58 55,93
<hr/> Difference.. 38,06	<hr/> 50,60	<hr/> 24,10

As there appears no reason for attaching an error to one of these rather than to the other, I have attributed the disagreement to error incidental to observations at this altitude, and taken the mean accordingly.

RESULT OF OBSERVATIONS IN 1832 AND 1833.

NAMES.	A. R.	MADRAS.		GREENWICH.		Barometer 29,60 Ther. 50°.			Error of	
		Mean N.P.D. Jan. 1, 1832		Presumed Ap- parent N.P.D. Jan. 1, 1832.		Observed Refraction.	Bradley's Refraction.	Atkinson's Refraction.	Bradley.	Atkin- son.
ζ Sagittarii...	18 51	120 6 44,74		120 0 30,72		6 14,02	6 7,33	6 7,91	- 6,69	- 6,11
α Antl. Pneum.	10 19	120 12 55,91		120 6 38,60		6 17,11	6 11,70	6 12,18	- 5,41	- 4,93
α App. Sculp...	0 50	120 16 2,41		120 9 44,96		6 17,45	6 13,83	6 14,28	- 3,62	- 3,17
γ ^a Sagittarii...	17 55	120 25 2,81		120 18 37,68		6 25,13	6 20,12	6 20,44	- 5,01	- 4,69
α Piscis. Aust...	22 48	120 30 39,01		120 24 12,14		6 26,87	6 24,17	6 24,39	- 2,70	- 2,48
υ ^a Eridani....	4 29	120 54 43,17		120 48 0,56		6 42,61	6 42,64	6 42,73	+ 0,03	+ 0,12
ξ Hyd. & Crat.	11 24	120 55 42,87		120 48 57,97		6 44,90	6 43,44	6 43,47	- 1,46	- 1,43
h Centauri....	13 43	121 5 38,66		120 58 45,74		6 52,92	6 51,55	6 51,05	- 1,37	- 1,87
θ Piscis. Aust...	21 37	121 40 16,01		121 32 51,35		7 24,66	7 22,29	7 21,04	- 2,37	- 3,62
k Centauri....	13 42	122 9 22,74		122 1 32,96		7 49,78	7 51,59	7 49,72	+ 1,81	- 0,06
* Centauri....	13 36	122 11 26,06		122 3 32,31		7 53,75	7 53,90	7 51,92	+ 0,15	- 1,83
κ ^a Canis Maj...	6 43	122 19 9,50		122 11 9,19		8 0,31	8 2,31	7 59,89	+ 2,00	- 0,42
μ Columbæ....	5 40	122 22 30,12		122 14 30,26		7 59,86	8 5,95	8 3,39	+ 6,09	+ 3,53
D Canis Maj...	6 22	122 28 43,76		122 20 33,87		8 9,89	8 13,27	8 10,45	+ 3,38	+ 0,56
α Pixid Naut...	8 36	122 35 3,72		122 26 46,75		8 16,97	8 20,85	8 17,77	+ 3,88	+ 0,80
4 Piscis. Aust...	21 7	122 52 5,40		122 43 28,88		8 36,52	8 41,88	8 38,06	+ 5,36	+ 1,54
β Hydræ.....	11 44	122 58 22,11		122 49 38,50		8 53,61	8 49,60	8 45,48	- 4,01	- 8,13
χ Lupi.....	15 40	123 5 28,93		122 57 28,25		9 0,68	9 0,82	8 56,27	+ 0,14	- 4,41
β Piscis. Aust...	22 22	123 12 14,90		123 3 8,94		9 5,96	9 8,83	9 4,04	+ 2,87	- 1,92
λ Canis Maj...	6 16	123 21 19,79		123 12 6,45		9 13,34	9 22,05	9 16,69	+ 8,71	+ 3,35
g Centauri....	13 46	123 36 28,83		123 26 48,27		9 40,56	9 45,02	9 38,65	+ 4,46	- 1,91
γ Piscis. Aust...	22 43	123 45 47,08		123 35 52,51		9 54,57	10 0,24	9 53,04	+ 5,67	- 1,53
ε Piscis. Aust...	21 35	123 47 14,47		123 37 14,74		9 59,73	10 2,56	9 55,26	+ 2,83	- 4,47
k Scorpii.....	16 54	123 52 38,09		123 42 30,40		10 7,69	10 11,60	10 3,97	+ 3,91	- 3,72
α Scorpii.....	16 39	123 58 45,71		123 48 27,37		10 18,34	10 22,33	10 14,07	+ 3,99	- 4,27
α Columbæ....	5 33	124 10 5,12		123 59 31,65		10 33,47	10 42,75	10 33,68	+ 9,28	+ 0,21
* Canis Maj...	6 45	124 10 25,09		123 59 48,39		10 36,70	10 43,20	10 34,03	+ 6,50	- 2,67
41 Eridani....	4 11	124 12 46,98		124 2 7,40		10 39,58	10 47,50	10 38,59	+ 7,92	- 0,99
α Normæ....	16 20	124 19 49,21		124 8 51,44		10 57,77	11 1,40	10 51,35	+ 3,63	- 6,42
α Microscopii...	20 39	124 23 43,11		124 12 39,70		11 3,41	11 9,60	10 59,26	+ 6,19	- 4,15
43 Eridani....	4 17	124 24 42,65		124 13 36,70		11 5,95	11 11,70	11 0,97	+ 5,75	- 4,98
ε Sagittarii...	18 12	124 27 18,19		124 16 5,20		11 12,99	11 16,90	11 5,85	+ 3,91	- 7,14
β Pixid Naut...	8 33	124 43 1,39		124 31 30,75		11 30,64	11 50,30	11 37,46	+ 19,66	+ 6,82
κ Columbæ....	6 10	125 5 25,85		124 52 50,48		12 35,37	12 42,40	12 28,15	+ 7,03	- 7,22
γ Columbæ....	5 51	125 18 25,92		125 5 15,60		13 10,32	13 16,30	13 0,40	+ 5,98	- 9,92
θ Centauri....	13 56	125 32 24,13		125 18 38,00		13 46,13	13 53,50	13 36,95	+ 7,37	- 9,18
ε Columbæ....	5 25	125 35 56,03		125 22 5,86		13 50,17	14 4,50	13 47,07	+ 14,33	- 3,10
i Centauri....	13 11	125 49 25,99		125 34 49,20		14 36,79	14 46,40	14 29,01	+ 9,61	- 7,78
β Columbæ....	5 45	125 50 11,49		125 35 38,41		14 33,08	14 49,20	14 31,90	+ 16,12	- 1,18
φ ^a Lupi.....	15 12	126 14 56,97		125 59 10,02		15 46,95	16 13,50	15 51,54	+ 26,55	+ 4,59
θ Lupi.....	15 55	126 20 13,64		126 4 0,33		16 13,31	16 33,40	16 10,29	+ 20,09	- 3,02
β Telescopii...	18 6	126 48 8,20		126 30 6,80		18 1,40	18 28,70	18 2,06	+ 27,30	+ 0,66
λ Scorpii.....	17 22	126 58 10,72		126 39 24,04		18 46,68	19 14,20	18 48,20	+ 27,52	+ 1,52
γ Telescopii...	17 38	126 58 43,66		126 39 55,71		18 47,95	19 16,70	18 49,35	+ 28,75	+ 1,40

The above columns will I apprehend require little or no explanation; consulting that headed "Errors of Bradley" we are led at once to the conclusion, that for altitudes below 5°, Bradley's Refraction can in no wise be

* The Greenwich place of this Star is probably 10 or 12 seconds too large.

trusted. For the altitudes above 5° , the errors incidental to one or two observations (which in some cases constitute the result), do not enable us to form an opinion. Consulting the column "Error of Atkinson"; between the limits of $81^\circ 30'$ and $85^\circ 30'$, and between $87^\circ 4'$ and $88^\circ 8'$ of Zenith Distance, we can perceive no error but what can be reasonably supposed to arise from the errors of observations joined to perhaps a small error* in the temperature; an error which must always exist when there is a considerable difference between the "IN" and "OUT" Thermometer; neither of which in this case can be proper to be employed in computing the refraction; *between* these limits, viz. between $85^\circ 30'$ and $87^\circ 4'$ of Zenith Distance we find ten very accordant results, shewing that Atkinson's refractions are too small by about $6''$. To say nothing of the enormous errors which appear in the preceding column (errors of Bradley) which have long since been acknowledged; we cannot but lament that whilst Astronomers have paid such ample and proper attention to the determination of the co-efficient of aberration, nutation, and precession, so much should be left undone in the way of refraction.

PARALLAX OF THE PLANET MARS.

Since Pages 90 and 91 were put to press, I have been favored with Volume VI of the Royal Astronomical Society's Memoirs, containing observations made at the Cape and at St. Helena, of the North Polar Distance of *Mars*,

* Supposing the error of temperature to be only a single degree of Fahrenheit, then the error of the refraction will be :

For 84 Zenith Distance.....	0,8
— 85 — —	1,0
— 86 — —	1,3
— 87 — —	1,7
— 88 — —	2,3
— 89 — —	3,2

Now notwithstanding all the care that is taken at the Greenwich Royal Observatory to equalise the temperature within and without, (and I have good reason for knowing that a very considerable degree of attention is paid to this particular), it is nevertheless not unfrequently found that a difference of 5 or 6 degrees exists between the "IN" and "OUT" Thermometer, it sometimes (most frequently) being warmer in the Observatory than in the open air. and at other times colder.

and of certain Stars, which had been previously pointed out by Mr. Henderson, Astronomer at the Cape, as proper for the determination of the parallax of this Planet; of the observations made at the Cape there have been several corresponding observations made here with which we will now compute the parallax of Mars: from Pages 90 and 91; &c. as above, we obtain as follows.

Difference of Declination between the Centre of the Planet MARS, and of Stars, from observations at

1832.	NAMES.	Madras.				The Cape of Good Hope.				
		Observed Difference.	R.			Observed Difference.	R.	M.		
		o	'	"	"	o	'	"	"	"
Nov. 9	A ¹ Tauri.	0 35 20,6	+ 0,59	+,1399	ρ	= 0 35 16,4	+ 1,79	- 10,30	+,8162	ρ
	53 Tauri.	0 17 56,3	+ 0,30	-,1399	ρ	= 0 17 55,9	+ 0,93	+ 10,30	-,8162	ρ
	a Tauri..	4 51 44,2	+ 4,68	-,1399	ρ	= 4 51 35,3	+ 13,13	+ 10,30	-,8162	ρ
15	b Tauri.	0 3 33,6	+ 0,06	-,1378	ρ^i	= 0 3 29,6	+ 0,12	+ 18,32	-,8152	ρ^i
	53 Tauri.	0 9 37,5	+ 0,16	-,1378	ρ^i	= 0 9 32,5	+ 0,50	+ 18,32	-,8152	ρ^i
	a Tauri.	4 43 23,2	+ 4,42	-,1378	ρ^i	= 4 43 11,5	+ 12,66	+ 18,32	-,8152	ρ^i
16	b Tauri.	0 1 45,4	+ 0,02	-,1370	ρ^{ii}	= 0 1 40,5	+ 0,06	+ 19,18	-,8145	ρ^{ii}
	a Tauri.	4 41 35,6	+ 4,37	-,1370	ρ^{ii}	= 4 41 23,4	+ 12,60	+ 19,18	-,8145	ρ^{ii}
17	A ¹ Tauri.	0 47 18,0	+ 0,78	+,1364	ρ^{iii}	= 0 47 23,1	+ 2,36	- 20,03	+,8142	ρ^{iii}
22	b Tauri..	0 10 51,4	+ 0,18	+,1333	ρ^{iv}	= 0 10 59,3	+ 0,55	- 23,68	+,8123	ρ^{iv}
	A ¹ Tauri.	0 58 2,8	+ 0,97	+,1333	ρ^{iv}	= 0 58 9,8	+ 2,97	- 23,68	+,8123	ρ^{iv}
	a Tauri.	4 29 0,6	+ 4,35	-,1333	ρ^{iv}	= 4 28 43,4	+ 12,01	+ 23,68	-,8123	ρ^{iv}
30	65 Arietis	0 7 58,3	+ 0,13	-,1278	ρ^v	= 0 7 46,1	+ 0,18	+ 23,51	-,8091	ρ^v
Dec. 4	65 Arietis	0 0 43,4	+ 0,01	+,1254	ρ^{vi}	= 0 0 56,3	+ 0,05	+ 20,94	+,8078	ρ^{vi}
	a Tauri.	4 1 27,9	+ 3,83	-,1254	ρ^{vi}	= 4 1 13,3	+ 10,49	+ 20,94	-,8078	ρ^{vi}
5	a Tauri.	3 59 27,1	+ 3,80	-,1248	ρ^{vii}	= 3 59 19,0	+ 10,30	+ 19,74	-,8073	ρ^{vii}
6	65 Arietis	0 4 37,8	+ 0,08	+,1243	ρ^{viii}	= 0 4 44,2	+ 0,22	- 18,74	+,8070	ρ^{viii}
	F ¹ Tauri.	0 58 2,8	+ 0,97	-,1243	ρ^{viii}	= 0 57 58,1	+ 2,82	+ 18,72	-,8070	ρ^{viii}
	a Tauri.	3 57 34,5	+ 3,77	-,1243	ρ^{viii}	= 3 57 24,1	+ 10,29	+ 18,72	-,8070	ρ^{viii}
7	65 Arietis	0 6 24,6	+ 0,11	+,1237	ρ^{ix}	= 0 6 30,5	+ 0,30	- 17,70	+,8067	ρ^{ix}
	F ¹ Tauri.	0 56 17,0	+ 0,94	-,1237	ρ^{ix}	= 0 56 12,4	+ 2,75	+ 17,70	-,8067	ρ^{ix}
	a Tauri.	3 55 46,4	+ 3,77	-,1237	ρ^{ix}	= 3 55 37,9	+ 10,32	+ 15,70	-,8067	ρ^{ix}
12	a Tauri.	3 48 31,3	+ 3,64	-,1217	ρ^x	= 3 48 29,2	+ 9,85	+ 11,32	-,8054	ρ^x
13	38 Arietis	0 3 30,8	+ 0,06	-,1213	ρ^{xi}	= 0 3 31,4	+ 0,16	+ 10,13	-,8052	ρ^{xi}
	a Tauri.	3 47 26,3	+ 3,33	-,1213	ρ^{xi}	= 3 47 27,2	+ 9,85	+ 10,13	-,8052	ρ^{xi}
15	38 Arietis	0 1 48,7	+ 0,03	-,1208	ρ^{xii}	= 0 1 53,3	+ 0,09	+ 7,20	-,8050	ρ^{xii}
22	65 Arietis	0 17 31,5	+ 0,29	+,1205	ρ^{xiii}	= 0 17 16,2	+ 0,82	+ 4,29	+,8048	ρ^{xiii}
	a Tauri.	3 44 39,0	+ 3,50	-,1205	ρ^{xiii}	= 3 44 51,9	+ 9,81	- 4,29	-,8048	ρ^{xiii}
24	65 Arietis	0 16 21,5	+ 0,27	+,1208	ρ^{xiv}	= 0 16 3,3	+ 0,77	+ 7,71	+,8051	ρ^{xiv}
	a Tauri.	3 45 47,8	+ 3,53	-,1208	ρ^{xiv}	= 3 46 3,9	+ 9,91	- 7,71	-,8051	ρ^{xiv}
25	65 Arietis	0 15 32,6	+ 0,26	+,1211	ρ^{xv}	= 0 15 14,4	+ 0,73	+ 9,09	+,8051	ρ^{xv}
	a Tauri.	3 46 39,4	+ 3,55	-,1211	ρ^{xv}	= 3 46 53,1	+ 9,87	- 9,09	-,8051	ρ^{xv}

In the above computation of R, which it will be understood is the difference of the refractions due to the Planet and Star, I have employed Atkinson's Table of Refractions, and have assumed the ratio of the Polar and Equatoreal Axis, 299 : 300; from which we determine :

		For the Observations at	
		Madras.	The Cape.
		/ "	/ "
Angle of the Vertical.....	=	5 0	— 10 38
Logarithm Radius of the Earth.....	=	9,99992	— 9,99958

Not having in my possession any Tables of the Planetary motions, from which I could compute the change of Declination (M.) for the interval between the Planet transiting the meridians of Madras and the Cape of Good Hope, I have been reduced to an interpolation from the observations, on which account errors to the amount of 0',2 in any single measure may be expected; but as these will occur indifferently + or —, the mean result cannot on this account be much affected.

To render these results in a more useful shape, we will now compute P, the Parallax at the time of opposition, when the Planets distance from the Earth was ,50581 whose Logarithm = 9,7040. Employing the Logarithm Distances given in the Supplement to the Nautical Almanac and resolving the above equations.

1832	From the Observations of.	We determine.								
		"		"	"					
Nov.	9 A ¹ Tauri.....	13,30	=	,6763 ρ	or ρ	=	19,666	and P	=	19,697
	53 Tauri.....	10,53	=	,6763 ρ	— ρ	=	15,570	—	=	15,595
	a Tauri.....	9,85	=	,6763 ρ	— ρ	=	14,564	—	=	14,590
	15 b Tauri.....	14,38	=	,6774 ρ ⁱ	— ρ ⁱ	=	21,228	—	=	21,262
	53 Tauri.....	13,56	=	,6774 ρ ⁱ	— ρ ⁱ	=	20,017	—	=	20,050
	α Tauri.....	14,86	=	,6774 ρ ⁱ	— ρ ⁱ	=	21,937	—	=	21,972
	16 b Tauri.....	14,32	=	,6775 ρ ⁱⁱ	— ρ ⁱⁱ	=	21,137	—	=	21,204
	a Tauri.....	15,21	=	,6775 ρ ⁱⁱ	— ρ ⁱⁱ	=	22,450	—	=	22,520
	17 A ¹ Tauri.....	13,45	=	,6778 ρ ⁱⁱⁱ	— ρ ⁱⁱⁱ	=	19,844	—	=	19,944
	22 b Tauri.....	15,41	=	,6790 ρ ^{iv}	— ρ ^{iv}	=	22,695	—	=	23,122
	A ¹ Tauri.....	14,68	=	,6790 ρ ^{iv}	— ρ ^{iv}	=	21,620	—	=	22,027
	a Tauri.....	14,14	=	,6790 ρ ^{iv}	— ρ ^{iv}	=	20,825	—	=	21,216
Dec.	30 65 Arietis.....	11,36	=	,6813 ρ ^v	— ρ ^v	=	16,673	—	=	17,767
	4 65 Arietis.....	8,00	=	,6824 ρ ^{vi}	— ρ ^{vi}	=	11,724	—	=	12,803
	a Tauri.....	13,00	=	,6824 ρ ^{vi}	— ρ ^{vi}	=	19,050	—	=	20,804
	5 α Tauri.....	18,14	=	,6825 ρ ^{vii}	— ρ ^{vii}	=	26,579	—	=	29,247
	6 65 Arietis.....	12,18	=	,6827 ρ ^{viii}	— ρ ^{viii}	=	17,842	—	=	19,784
	F ¹ Tauri.....	15,87	=	,6827 ρ ^{viii}	— ρ ^{viii}	=	18,464	—	=	20,476
	a Tauri.....	14,64	=	,6827 ρ ^{viii}	— ρ ^{viii}	=	21,444	—	=	23,778
	7 65 Arietis.....	11,61	=	,6830 ρ ^{ix}	— ρ ^{ix}	=	16,998	—	=	19,007
	F ¹ Tauri.....	14 91	=	,6830 ρ ^{ix}	— ρ ^{ix}	=	21,830	—	=	24,410
	a Tauri.....	15,75	=	,6830 ρ ^{ix}	— ρ ^{ix}	=	23,060	—	=	25,783
	12 α Tauri.....	15,43	=	,6837 ρ ^x	— ρ ^x	=	22,568	—	=	26,357
	13 38 Arietis.....	10,83	=	,6839 ρ ^{xi}	— ρ ^{xi}	=	15,837	—	=	18,678
	a Tauri.....	17,55	=	,6839 ρ ^{xi}	— ρ ^{xi}	=	25,662	—	=	30,268
	15 38 Arietis.....	11,86	=	,6842 ρ ^{xii}	— ρ ^{xii}	=	17,333	—	=	20,768
	22 65 Arietis.....	10,48	=	,6843 ρ ^{xiii}	— ρ ^{xiii}	=	15,316	—	=	19,748
a Tauri.....	14,92	=	,6843 ρ ^{xiii}	— ρ ^{xiii}	=	21,803	—	=	28,108	

1832	From the Observations of.	We determine.					
		"		"		"	
Dec. 24	65 Arietis.....	9.99 =	,6843	ρ^{xiv} —	ρ^{xiv} =	14,600 and P =	19,224
	a Tauri.....	14.47 =	,6843	ρ^{xiv} —	ρ^{xiv} =	21,145 — =	28,122
25	65 Arietis.....	8.64 =	,6840	ρ^{xv} —	ρ^{xv} =	12,632 — =	16,809
	a Tauri.....	11.03 =	,6840	ρ^{xv} —	ρ^{xv} =	16,126 — =	21,461

The above results it must be confessed are highly unsatisfactory ; a nearer coincidence does however appear to take place between the *individual* determinations of each Star than is found by viewing them collectively, which is better seen by the following arrangement.

Values of P, deduced from observations made at Madras compared with the corresponding observations at the Cape.

1832:	38 Arietis.	65 Arietis.	b. Tauri.	53 Tauri.	A ¹ Tauri.	F ¹ Tauri.	a Tauri.
	"	"	"	"	"	"	"
November 9	15,595	19,697	14,590
15	21,262	20,051	21,972
16	21,204	22,520
17	19,944
22	23,122	22,027	21,216
30	17,707
December 4	12,803	20,804
5	*29,247
6	19,784	20,476	23,778
7	19,007	24,410	25,783
12	26,357
13	18,678	+30,268
15	20,768
22	19,748	28,108
24	19,229	28,122
25	16,809	21,461
Mean.....	19,713	17,869	21,863	17,823	20,556	22,443"	23,155

When we consider that the above values are determined from the *difference of differences* (involving in extreme cases an eight fold amount of error) and that in the course of computation we have multiplied the results by a factor $\frac{100}{68}$ and by another varying from 1 to $\frac{100}{74}$, we are at no loss to account for the irregularities now found in the results of the first six Stars, since on computation it will be found that they do not involve for each single obser-

* The Cape Observation of Mars, appears to be 5" in Error.

+ ————— 2" —————

vation a larger probable error than 1", an amount which might reasonably be expected; but in the case of α Tauri the disagreement assumes so determined a character that we can by no means allow that the errors incidental to observation can account for it: to comprehend this matter more clearly we will again consult the rough observations Pages 90 and 91, Vol. VI of the Royal Astronomical Society's Memoirs, whence we obtain as follows.

Difference of Declination between δ 3 Tauri and α Tauri as observed

<i>at Madras.</i>		<i>at the Cape.</i>		<i>at St. Helena.</i>	
1832	" ' "	1832	" ' "	1832	" ' "
November	9.. 4 33 47,90	October	22.. 4 33 37,40	October	14.. 4 33 45,20
	15.. 4 33 45,70		24.. 4 33 39,10		15.. 4 33 44,80
			25.. 4 33 39,30	November	14.. 4 33 44,30
		November	8.. 4 33 40,30		
			9.. 4 33 39,40		
			10.. 4 33 39,70		
			11.. 4 33 39,00		
			12.. 4 33 41,50		
			13.. 4 33 40,50		
			15.. 4 33 39,00		
Mean.....	4 33 46 80		4 33 39,52		4 33 44,77
Diff. of Refraction.	+ 4,82		+ 12,04		+ 6,39
True Difference...	4 33 51,12		4 33 51,56		4 33 51,16

And further we have the

Difference of Declination between δ 3 Arietis and α Tauri as observed

<i>at Madras.</i>		<i>at the Cape.</i>		<i>at St. Helena.</i>	
1832	" ' "	1832	" ' "	1832	" ' "
December	13.. 3 43 55,50	December	12.. 3 43 55,00	December	18.. 3 43 57,00
	15.. 3 43 54,80		13.. 3 43 55,80		
	16.. 3 43 55,80		22.. 3 43 53,40		
	18.. 3 43 54,60		23.. 3 43 52,30		
			24.. 3 43 52,40		
			25.. 3 43 51,30		
Mean.....	3-43 55,17		3 43 53,37		3-43 57,00
Diff. of Refraction.	+ 3,27		+ 9,69		+ 5,00
True Difference...	3 43 58,44		3 44 3,06		3 44 2,00

Moreover we have the

Difference of Declination between 65 Arietis and a Tauri as observed

<i>at Madras.</i>				<i>at the Cape.</i>				<i>at St. Helena.</i>			
1832	°	'	"	1832	°	'	"	1832	°	'	"
December	4..	4	2 11,30	November 30..	4	2 8,00		November 30..	4	2 14,10	
	6..	4	2 12,30	December 1..	4	2 9,40		December 1..	4	2 13,80	
	7..	4	2 11,00		3..	4	2 6,50		4..	4	2 14,80
	20..	4	2 9,30		4..	4	2 9,60		5..	4	2 15,10
	21..	4	2 9,90		5..	4	2 6,70		6..	4	2 14,00
	22..	4	2 10,55		6..	4	2 8,30		18..	4	2 13 30
	24..	4	2 9,30		7..	4	2 8,40		19..	4	2 11,40
	25..	4	2 12,00		8..	4	2 7,60				
					9..	4	2 6,70				
					10..	4	2 7,60				
Mean.....	4	2	10,70		4	2	7,88		4	2	13,79
Diff. of Refraction.	+		3,84		+		10,56		+		5,47
True Difference...	4	2	14,54		4	2	18,44		4	2	19,26

In each of the above cases, the observations having been made at nearly the same time of the year, the difference of the corrections for aberration, &c. will be very nearly common for the result obtained at each Observatory, and for the present enquiry may be disregarded altogether: Examining the "true difference" the agreement between the results at the Cape and those at St. Helena, affords us good reason for supposing them accurate, and for employing them as standards of comparison. If we accordingly compare the Madras "True difference" with that found from

<i>We determine.</i>	<i>The Cape Observations.</i>	<i>The St. Helena Observations.</i>
$e + e^I$	= - 0,44	= - 0,04
$e + e^{II}$	= - 4,62	= - 3,56
$e + e^{III}$	= - 3,90	= - 4,72

Taking the mean we have:

$$e + \frac{e^I + e^{II} + e^{III}}{3} = - 2,98 \quad \text{---} \quad 2,77$$

In the above, e represents the error of division for α Tauri, e^I that for 53 Tauri, &c. Here (in the case of divisions situated within 3° or 4° of each

other) it appears plain, that if we could obtain a sufficient number of results; the values e^I e^{II} , &c. occurring as they no doubt do with contrary signs, we could determine e to any required degree of accuracy; even with the very limited number we already possess *it appears exceedingly probable that the divisions $73^\circ 50'$ and $73^\circ 55'$ of the Madras Mural Circle are erroneous to the amount of two seconds*, and comparing the mean place of α Tauri from 146 Observations made at Madras, with the Greenwich place from three times that number of observations, I find a difference = $-1''.71$ which so strongly supports the above conjecture that for the present it becomes necessary to suppress altogether the results obtained from α Tauri; were however the circumstances different; the distance of this Star from the Planet Mars ($4^\circ \pm$) would have rendered it questionable if its result ought to be admitted. Taking the mean of the remaining 19 Observations we determine $P = 19''.595$.

Since writing the above — signing as it were the death warrant of the Madras Observations; with a firm conviction that the discordances found in the place of α Tauri did not arise from error of division I have proceeded as follows.

Error of Division of the Madras Mural Circle.

About two years ago I made a set of experiments to ascertain the amount of error in the division of the Madras Mural Circle; having transmitted copy of these observations to England for publication, it is only necessary for me here to remark, that in the course of an examination of every 5th degree; I met with no error which could affect the mean of the four readings to the amount of one second: Now the divisions $69^\circ 50'$, $70^\circ 10'$ and $73^\circ 55'$ on which 56 Arietis, 38 Arietis and α Tauri were respectively observed, not having fallen under this examination, it still becomes necessary, either to admit the remarks at lines 4 — 6; or by a direct appeal to those divisions to ascertain their *actual* amount of error; pursuing the latter course I have adopted a plan on the present occasion similar to that contrived for the above-mentioned examination, which I will now proceed to explain. In the description of the Madras Mural Circle given in the 1st Volume of these results, it is stated, that “the Telescope is attached to the circular ring at each end by appropriate braces, each secured by four strong screws; and is further supported in the middle, by an axis (represented by dotted lines fig. 1) which passes through the axis of the circle, and is secured by a screw C affixed to its smaller end;” from this description it will appear plain that the screws at each end of the Telescope which serve to secure it to the circular ring being loosed; the Telescope is free to turn upon its axis independent of the circle;

by which property we are enabled to measure any angle upon any required divisions, by merely shifting the place of the Telescope upon the ring. To obtain an object which could be distinctly viewed through an Astronomical Telescope, (in which the eye piece is required to be adjusted to the Solar focus,) I availed myself of the well known property of the rays of light, which, diverging from the principal focus of an object glass, after passing through the object glass are transmitted as parallel rays; and hence possess the property of an object placed at an infinite distance; but to be particular; I placed Dolland's 5 feet Achromatic Telescope about 5 feet in front (to the North) of the Mural Circle, with its object glass *vis a vis* to the object glass of the Mural Circle Telescope, and its whole length so disposed that a line passing through the centre of the one Telescope, being continued, would equally pass through the centre of the other: I now rested another Telescope (a 46 Inch Achromatic by Dolland) immediately above the first named Telescope, by means of two pieces of wood A, B, (see fig.) of such dimensions, that the angle subtended by the two Telescopes was nearly that which I required, and such that the Mural Circle Telescope being directed to the said upper Telescope; a line passing through the centre of the one if continued would equally pass through the centre of the other; matters thus arranged I introduced into the principal focus of the upper Telescope a pair of cross lines, which by means of a light placed behind them, were very distinctly seen by the Circle Telescope, and were adjusted to horizontality;—moving the circle and its attached Telescope through the angle subtended by the two Telescopes; a similar pair of lines which had been fitted into the micrometer attached to the lower Telescope now came into view, and the angle formed by the two Telescopes was thus read off from the circle; as this angle did not at first agree with that which was required, the micrometer screw of the lower Telescope enabled me to adjust it to any required degree of accuracy. The Circle was made to read off $70^{\circ} 10' 35''$ (the reading at which 38 Arietis (Page 91) was observed and clamped; the Telescope being released from the circle, was directed to the wires of the *lower Telescope* very nearly, again clamped, and an accurate bisection of them made by the moveable wire of the circle Telescope; the circle was now very carefully read off and the bisection of the cross wires again examined and if necessary improved; unclamping the circle, it was with its now attached Telescope moved to view the cross lines of the upper Telescope, of which an accurate bisection was made and the circle again read off; here the reading was $(73^{\circ} 54' 25'' \pm)$;—as nearly as need be that which was employed in the observations of α Tauri (Page 91), consequently any error arising from ill division which may be attached to the measured difference of declination of these two Stars as found at Page 122, will equally affect the measure of the angle between

the two Telescopes (*collimators*) above described; to ascertain its amount I have measured the above angle between the two collimators on several sets of divisions as follows.

Angular inclination of the two Telescopes on the 2d September 1834, as measured by the Mural Circle, on the divisions employed in the observations of a Tauri and 38 Arietis, together with the measurement of the same upon sundry other divisions.

		A.	B.	C.	D.	Mean.	Angle between the Collimators as measured by					
							Divisions 70° 10' and 73° 54'.			Sundry Divisions.		
°	'	"	"	"	"	"	°	'	"	°	'	"
70	10	36,1	42,1	49,0	40,4	41,90	3	43	52,50	3	43	52,65
73	54	28,5	34,5	42,6	32,0	34,40						
80	11	11,3	20,0	15,7	16,0	15,75	3	43	52,65	3	43	52,28
83	55	0,6	10,7	14,5	7,8	8,40						
80	11	11,9	20,3	16,5	17,0	16,42	3	43	52,28	3	43	51,42
83	55	0,6	11,6	14,0	8,6	8,70						
90	11	9,3	15,3	18,5	13,2	14,08	3	43	51,42	3	43	52,37
93	54	59,3	5,6	12,0	5,1	5,50						
100	11	9,8	15,0	17,2	15,2	14,30	3	43	52,37	3	43	52,42
103	55	0,4	9,5	11,3	5,5	6,67						
70	10	36,8	43,8	49,0	41,5	42,78	3	43	52,42	3	43	52,35
73	54	29,4	36,1	41,3	34,0	35,20						
70	10	31,0	37,2	43,0	36,4	36,90	3	43	52,35	3	43	52,65
73	54	22,2	28,0	35,0	31,8	29,25						
80	11	6,1	14,5	10,3	11,0	10,47	3	43	52,65	3	43	50,35
83	54	55,6	6,4	8,5	2,0	3,12						
90	11	13,9	18,1	22,0	16,3	17,57	3	43	50,35	3	43	51,73
93	55	3,0	8,8	12,0	7,9	7,92						
100	11	9,1	15,1	15,3	13,2	13,17	3	43	51,73	3	43	52,15
103	54	59,8	5,0	8,8	6,0	4,90						
110	11	6,9	12,4	13,3	11,8	11,10	3	43	52,15	3	43	51,75
113	54	57,4	4,0	6,8	4,7	3,25						
120	11	6,2	12,1	6,5	14,0	9,70	3	43	51,75	3	43	52,05
123	54	58,6	1,9	3,8	1,5	1,45						
130	11	10,1	12,1	13,3	11,0	11,62	3	43	52,05	3	43	51,97
133	55	0,3	4,4	7,2	2,8	3,67						
140	11	9,8	13,5	15,3	10,5	12,40	3	43	51,97	3	43	51,02
143	55	1,2	6,0	7,2	3,1	4,37						
150	11	0,0	3,0	6,0	1,2	2,55	3	43	51,02	3	43	51,88
153	54	50,4	53,2	57,0	53,7	53,57						
70	10	36,0	41,1	44,8	40,2	40,52	3	43	51,88	3	43	51,87
73	54	28,0	33,9	37,0	30,7	32,40						
Mean....							3	43	52,29	3	43	51,87

The Telescopes were allowed to remain undisturbed and on the 3d September 1834, the following measures were taken.

		A.	B.	C.	D.	Mean.	Angle between the Collimators as measured by					
							Divisions 70° 10' and 73° 54'.			Sundry Divisions.		
°	'	''	''	''	''	''	°	'	''	°	'	''
70	10	39,9	47,0	47,3	44,3	44,62	3	43	49,90			
73	54	30,0	37,9	34,0	34,2	34,52						
70	10	38,2	44,5	45,3	43,2	42,80	3	43	49,27			
73	54	27,7	33,6	34,5	32,5	32,07						
80	11	14,2	21,2	16,5	18,7	17,65				3	43	49,45
83	55	1,9	9,5	9,0	8,0	7,10				3	43	49,25
90	11	12,0	16,6	15,6	14,7	14,72				3	43	49,00
93	55	0,2	5,1	5,8	4,8	3,97				3	43	49,25
100	11	12,0	17,9	15,7	16,0	15,40				3	43	49,00
103	55	0,0	7,4	5,0	5,2	4,40				3	43	49,00
110	11	12,1	16,4	15,4	16,6	15,12				3	43	49,58
113	55	0,0	8,0	4,5	6,3	4,70				3	43	49,58
120	11	11,9	17,2	15,5	13,2	14,45				3	43	49,77
123	55	1,1	6,6	3,8	5,4	4,22				3	43	49,77
130	11	14,1	16,1	13,2	14,5	14,47				3	43	48,98
133	55	1,3	4,5	3,0	5,0	3,45				3	43	48,98
140	11	12,9	15,5	12,0	14,3	13,68				3	43	49,74
143	55	0,2	7,5	2,0	4,0	3,42				3	43	49,74
150	11	11,6	14,4	13,0	14,6	13,40				3	43	49,95
153	55	0,5	5,4	3,2	4,3	3,35				3	43	49,95
70	11	49,2	54,5	54,0	53,2	52,72	3	43	49,50			
73	54	38,9	44,2	43,3	42,5	42,22						
80	11	12,0	19,2	16,2	16,5	15,97				3	43	49,78
83	55	0,1	9,5	0,6	6,8	5,75				3	43	49,78
90	11	12,4	19,4	16,2	15,8	15,95				3	43	48,20
93	55	0,3	6,5	5,0	4,8	4,15				3	43	48,20
100	11	13,1	16,6	14,6	15,3	14,90				3	43	48,90
103	55	0,2	6,5	3,7	4,8	3,80				3	43	48,90
110	11	12,6	17,8	15,8	17,6	15,95				3	43	48,17
113	55	0,0	6,2	4,3	6,0	4,12				3	43	48,17
120	11	12,2	16,0	15,3	13,8	14,32				3	43	49,43
123	55	0,8	4,5	5,7	4,0	3,75				3	43	49,43
130	11	12,0	13,5	11,6	14,1	12,80				3	43	50,02
133	55	0,7	4,6	2,2	3,8	2,82				3	43	50,02
140	11	12,5	16,5	12,0	13,8	13,95				3	43	50,07
143	55	0,8	7,3	3,7	4,3	4,02				3	43	50,07
153	11	11,8	14,4	13,7	14,8	13,68				3	43	49,94
153	55	0,8	5,2	3,7	4,8	3,62				3	43	49,94
70	11	40,5	46,0	45,0	44,3	43,95	3	43	48,90			
73	54	29,8	35,3	33,6	32,7	32,85						
70	11	39,1	46,0	44,0	43,8	43,22	3	43	48,86			
73	54	29,0	34,1	32,5	32,7	32,08						
Mean....							3	43	49,48	3	43	49,39

The change of 3" in the angle between the observations of September 2d and 3d, no doubt arises from a shrinking of the wooden supports; this however is of no consequence in our present enquiry, since we only require that the angle should remain fixed during the observations of the day, or for the space of an hour which (with one assistant stationed at each microscope) was the time employed on each day.

The angle was now increased to 4° 2' 8" nearly; and its exact quantity measured upon the divisions 69° 50' and 73° 50' (these being the divisions at which 65 Arietis and α Tauri were respectively observed), and upon sundry other divisions as follows.

6th September.

°	'	A.	B.	C.	D.	Mean.	Angle between the Collimators as measured by					
							Divisions 70° 10' and 73° 54'.		Sundry Divisions.			
°	'	"	"	"	"	"	°	'	"			
69	49	14,1	24,8	17,3	23,0	19,80	4	2	7,05			
73	51	23,9	28,1	25,8	29,6	26,85						
79	49	15,8	23,9	18,0	19,5	19,30	4	2	5,98			
83	51	21,2	28,5	24,6	26,8	25,28						
89	49	14,7	22,7	20,3	20,5	19,55	4	2	7,23			
93	51	23,2	29,2	26,1	23,6	26,78						
99	49	15,4	22,5	17,3	20,2	18,85	4	2	6,80			
103	53	23,1	28,6	25,1	25,8	25,65						
109	49	15,8	23,5	18,0	21,4	19,68	4	2	7,07			
113	53	24,4	28,6	24,8	29,2	26,75						
119	49	11,0	17,6	14,6	14,4	14,40	4	2	6,85			
123	53	18,4	23,2	20,7	22,7	21,25						
129	49	13,9	19,4	13,2	18,2	16,18	4	2	7,34			
133	53	21,8	25,2	31,1	26,0	23,52						
139	49	14,1	18,2	13,4	19,0	16,18	4	2	7,14			
143	53	21,6	26,0	20,7	25,0	23,32						
149	49	17,2	22,4	18,2	19,9	19,42	4	2	7,38			
153	53	26,4	28,0	25,0	27,8	26,80						
69	49	15,0	22,6	19,2	21,5	19,58	4	2	7,67			
73	51	26,3	28,4	25,8	28,5	27,25						
Mean....							4	2	7,36	4	2	6,97

7th September.

°	'	A.	B.	C.	D.	Mean.	Angle between the Collimators as measured by		
							Divisions 70° 10' and 73° 54'.		Sundry Divisions.
°	'	"	"	"	"	"	°	'	"
69	49	16,0	25,3	21,4	26,1	22,20	4	2	8,65
73	51	28,0	33,9	29,1	32,4	30,85			
79	49	16,2	25,0	19,1	23,1	20,85	4	2	7,90
83	51	23,3	31,9	28,3	31,5	28,75			
89	49	15,4	24,5	20,9	22,8	20,90	4	2	7,80
93	51	25,0	32,4	27,3	30,1	28,70			
99	49	15,2	23,6	17,0	21,2	19,25	4	2	8,80
103	53	25,0	32,2	26,0	29,0	28,05			
109	49	15,6	24,0	17,1	22,0	19,67	4	2	7,98
113	53	24,0	32,0	24,5	30,1	27,65			
119	49	15,2	22,6	18,6	21,0	19,35	4	2	8,47
123	53	25,4	31,0	26,1	28,8	27,82			

.	A.	B.	C.	D.	Mean.	Angle between the Collimators as measured by					
						Divisions 70° 10' and 73° 54'.			undry Divisions.		
o	i	"	"	"	"	o	i	"	o	i	"
129	49	17,7	24,4	18,6	21,9	20,65	}	4	2	7,15
133	53	26,0	30,8	25,3	29,1	27,80					
139	49	15,0	18,6	13,0	18,0	16,16	}	4	2	8,67
143	53	23,1	29,0	21,2	26,0	24,82					
149	49	14,2	21,5	15,3	15,4	16,60	}	4	2	8,95
153	53	24,2	29,2	23,1	25,7	25,55					
69	49	15,8	24,6	19,1	23,8	20,82	}	4	2	8,93	
73	51	28,7	32,2	28,0	30,1	29,75					
69	49	15,9	25,0	20,2	23,7	21,20	}	4	2	8,05	
73	51	27,7	32,5	27,6	29,1	29,25					
Mean....						4	2	8,54	4	2	8,22

Hence it appears that the angular distance between 38 *Arietis* and *α Tauri* as observed at Madras (Page 122) is erroneous by reason of error of division.

<i>From the observations of</i>	<i>to the amount.</i>
2d September 1834.....	+ 0,42
3d ——— 1834.....	+ 0,09
Mean.....	+ 0,25

It further appears that the angular distance between 65 *Arietis* and *α Tauri* as observed at Madras (Page 123) is erroneous by reason of error of division.

<i>From the observations of</i>	<i>to the amount.</i>
6th September 1834.....	+ 0,39
7th ——— 1834.....	+ 0,32
Mean.....	+ 0,35

The above results, whilst they leave nothing to be desired with regard to the division of the circle, still leave unexplained the discordant results of Page 123; they do not as we have seen arise from error of division in the Madras Mural Circle, and it is *highly improbable* that the Cape or St. Helena Instruments can err from error of division to this amount (4" +). From a comparison of numerous observations of N. P. D. with the mean result, I find that the mean error of a single observation is considerably less than 1", but allowing it to equal this amount, and making a further liberal allowance for

possible errors, we can in no way make up the amount of 4" +. For the present I am compelled to allow this singular and unexpected anomaly to remain unexplained, but venture to hope that in the next Volume of these observations I shall be enabled to offer some sort of conclusion as to its cause.

Error of Observation ; Parallax of α Aquilæ.

With reference to the remarks at Page 129, I had here proposed to give the result of each single observation of the North Polar Distance of one or more of the principal fixed Stars, by way of exhibiting the extent of error committed in this nature of observation; and had commenced for the purpose an examination of the catalogue, to ascertain which Star had been most frequently observed; when the recollection of the reputed annual parallax of α Aquilæ led me to select this Star, and to join to my original enquiry the question of parallax; I must however remark, that the observations which now follow having been made simply for the purpose of determining the Index Error of the Mural Circle and the place of the Star, are not so numerous, or so well disposed for the determination of Parallax as under other circumstances they might have been; if π represent the semiaxis minor of the Earth's orbit (supposed to be a circle) as viewed from α Aquilæ, and λ the Latitude of the Star; we have the semiaxis major or $\rho = \cos. (\text{Long. } \odot - \text{Long. } *) \frac{\pi}{\sin. \lambda}$ nearly: selecting now the observations which are situated near to the positive and negative maximum of Parallax, we have as follows.

1831	N. P. D. from Circle Book.			Refrac. tion.	Aberra- tion, &c.	Index Error.	Annual Preces- sion.	Mean N. P. D. January 1, 1832.										
	°	'	"	"	"	"	"	°	'	"								
February	1	81	35	58.0	+	4.43	+	0.78	-	1 45,77	-	8,67	81	34	8 77	+	,486	ρ
	2	81	35	58.7	+	4.42	+	0.71	-	1 46,47		81	34	8 69	+	,484	ρ
	3	81	36	0.5	+	4.42	+	0.64	-	1 47,29		81	34	9,60	+	,482	ρ
	7	81	36	0.3	+	4.43	+	0.35	-	1 46,58		81	34	9,83	+	,471	ρ
	8	81	36	0.9	+	4.41	+	0.27	-	1 46,58		81	34	10,33	+	,468	ρ
	10	81	36	1.4	+	4.41	+	0.12	-	1 46,80		81	34	10,46	+	,462	ρ
	12	81	36	0.4	+	4.42	-	0.22	-	1 46,96		81	34	8,97	+	,456	ρ
	13	81	35	59.3	+	4.41	-	0.39	-	1 47,14		81	34	7,51	+	,451	ρ
	14	81	36	0.8	+	4.41	-	0.55	-	1 47,14		81	34	8,85	+	,448	ρ
	15	81	36	1.7	+	4.40	-	0.71	-	1 46 50		81	34	10,22	+	,444	ρ
	18	81	36	1.2	+	4.38	-	1,18	-	1 46,57		81	34	9,16	+	,431	ρ
	21	81	36	1.4	+	4.39	-	1,70	-	1 46,95		81	34	8,47	+	,416	ρ
	23	81	36	0.2	+	4.40	-	2,02	-	1 46,89		81	34	7,02	+	,406	ρ
	24	81	36	1.4	+	4.39	-	2,11	-	1 46,89		81	34	8,12	+	,401	ρ
27	81	36	2.8	+	4.40	-	2,43	-	1 48,03		88	34	7,07	+	,386	ρ	

RESULT OF OBSERVATIONS IN 1832 AND 1833.

1831	N. P. D. from Circle Book.			Refrac- tion.	Aberra- tion, &c.	Index Error.	Annual Preces- sion.	Mean N. P. D. January 1, 1832.			
	°	'	"					°	'	"	
March	1	81 36	2,3	+ 4,42	- 2,58	- 1 47,49	- 8,67	81 34	7,98	+ ,375	p
	2	81 36	2,3	+ 4,42	- 2,60	- 1 47,49	81 34	7,90	+ ,369	p
	3	81 36	3,3	+ 4,43	- 2,73	- 1 46,83	81 34	9,50	+ ,363	p
	4	81 36	2,2	+ 4,48	- 2,81	- 1 46,83	81 34	8,37	+ ,357	p
	5	81 36	3,4	+ 4,46	- 2,89	- 1 47,67	81 34	8 63	+ ,351	p
	6	81 36	2,6	+ 4,45	- 2,96	- 1 47,67	81 34	7,75	+ ,345	p
	7	81 36	2,3	+ 4,42	- 3,03	- 1 46,98	81 34	8,04	+ ,338	p
	13	81 36	3,0	+ 4,40	- 3,32	- 1 47,14	81 34	8,27	+ ,305	p
	15	81 36	2,9	+ 4,40	- 3,37	- 1 46,42	81 34	8,84	+ ,291	p
1832											
February	20	81 37	11,1	+ 4,37	- 4,48	- 3 2,03	0,00	81 34	8,96	+ ,423	p
	27	81 37	26,7	+ 4,42	- 4 98	- 3 16,37	81 34	9 77	+ ,387	p
	29	81 37	24,9	+ 4,44	- 5,12	- 3 16 37	81 34	7,85	+ ,375	p
March	1	81 37	25,9	+ 4,45	- 5,19	- 3 16,37	81 34	8,81	+ ,369	p
	2	81 37	24,1	+ 4,45	- 5,26	- 3 16,37	81 34	7,94	+ ,364	p
	3	81 37	25,3	+ 4,44	- 5,33	- 3 16 37	81 34	8,06	+ ,359	p
	6	81 37	25,7	+ 4,43	- 5,52	- 3 15,25	81 34	9,38	+ ,341	p
	8	81 37	26,7	+ 4,42	- 5,65	- 3 15,55	81 34	9,92	+ ,327	p
	9	81 37	25,7	+ 4,44	- 5,71	- 3 15,55	81 34	8,92	+ ,320	p
	11	81 37	38,2	+ 4,44	- 5,77	- 3 27,81	81 34	9 06	+ ,307	p
	12	81 37	38,5	+ 4,44	- 5,81	- 3 28,16	81 34	8,97	+ ,300	p
	13	81 37	37,6	+ 4,45	- 5,85	- 3 28,48	81 34	7 72	+ ,293	p
	15	81 37	38,0	+ 4,45	- 5,93	- 3 28,48	81 34	8,04	+ ,279	p
December	21	81 41	10,6	+ 4,42	+ 11,16	- 7 17,71	81 34	8,47	+ ,433	p
1833.											
January	3	81 36	59,6	+ 4,43	+ 0,30	- 3 5,36	+ 8,67	81 34	7,64	+ ,479	p
	5	81 37	0,9	+ 4,44	- 0,05	- 3 6,05	81 34	7,91	+ ,484	p
	6	81 37	0,4	+ 4,44	- 0,22	- 3 4,56	81 34	8 73	+ ,486	p
	7	81 37	0,7	+ 4,44	- 0,39	- 3 4,56	81 34	8 86	+ ,488	p
March	16	81 35	40,5	+ 4,44	- 8,95	+ 1 36,48	81 34	8,18	+ ,272	p
1831											
July	13	81 37	13,8	+ 4,34	+ 12,41	- 3 10,66	81 34	11,22	- ,495	p
	16	81 37	5,3	+ 4,38	+ 12,97	- 3 4,70	- 8,67	81 34	9,28	- ,497	p
	17	81 37	7,2	+ 4,35	+ 13,14	- 3 4,70	81 34	9,32	- ,498	p
	28	81 37	15,5	+ 4,32	+ 15,02	- 3 15,61	81 34	11,54	- ,497	p
August	10	81 37	12,5	+ 4,33	+ 16,98	- 3 16,38	81 34	9,76	- ,474	p
	11	81 37	12,7	+ 4,33	+ 16,86	- 3 16,38	81 34	9,84	- ,471	p
	22	81 37	11,6	+ 4,34	+ 18,42	- 3 16,43	81 34	10,76	- ,432	p
	23	81 37	11,1	+ 4,33	+ 18,53	- 3 15,51	81 34	9,78	- ,429	p
	26	81 37	11,1	+ 4,36	+ 18,85	- 3 15,55	81 34	10,09	- ,420	p
1833											
August	5	81 35	12,8	+ 4,34	+ 10,55	- 1 26,38	81 34	9,97	- ,486	p
	6	81 35	11,7	+ 4,34	+ 10,69	- 1 26,38	+ 8,67	81 34	9,02	- ,484	p
	7	81 35	12,1	+ 4,33	+ 10,82	- 1 26,38	81 34	9,54	- ,482	p
	16	81 35	10,3	+ 4,34	+ 11,99	- 1 26,38	81 34	8,92	- ,456	p
	29	81 35	9,2	+ 4,36	+ 13,36	- 1 26,84	81 34	8,75	- ,401	p
	30	81 35	8,3	+ 4,36	+ 13,43	- 1 26,66	81 34	8,10	- ,396	p
September	5	81 35	7,8	+ 4,35	+ 13,89	- 1 26,66	81 34	8,05	- ,363	p
	8	81 35	6,2	+ 4,35	+ 14,12	- 1 26,46	*81 34	6,88	- ,346	p

* This is omitted in taking the mean.

1833	N. P. D. from Circle Book.			Refraction.	Aberra- tion, &c.	Index Error.	Annual Preces- sion.	Mean N. P. D. January 1, 1832.										
	°	'	"					°	'	"								
September	10	81	35	9,8	+	4,35	+	14,27	—	1	28,55	81	34	8,54	—	,334	ρ
	11	81	35	9,5	+	4,34	+	14,32	—	1	28,55	81	34	8,28	—	,327	ρ
	12	81	35	9,8	+	4,36	+	14,37	—	1	28,55	81	34	8,65	—	,321	d
	13	81	35	9,3	+	4,36	+	14,42	—	1	28,55	81	34	8,20	—	,314	ρ
	15	81	35	10,6	+	4,34	+	14,51	—	1	28,55	81	34	9,57	—	,300	ρ

Taking the mean we have:

		Mean N.P.D. January 1, 1832:		
		°	'	"
From 24 Observations in the Winter of 1830-1831	—	81	34	8,682 + ,408 ρ
From 13 ———— 1831-1832	—	81	34	8,723 + ,342 ρ
From 6 ———— 1832 1833	—	81	34	8,298 + ,440 ρ
From 9 ———— Summer of 1831	—	81	34	10,177 — ,468 ρ
From 12 ———— 1833	—	81	34	8,900 — ,389 ρ
$\therefore \rho = 0'',978$				

or the angle under which the Earth's orbit is seen at α Aquilæ* = $1'',96$. Considering the disagreement which is found to exist between the numerous results of the Greenwich and Dublin Instruments when applied to the determination of the Parallax of α Aquilæ, it would appear that the above result as far as the determination of parallax is concerned, is entitled to very little if any credit; one circumstance however will be found to affect *these* observations which goes far to diminish the weight of this objection. It must be recollected that in the Latitude of Greenwich the meridian altitude of α Aquilæ is about 47° and that it arrives at the positive and negative maximum of parallax in the middle of Winter and in the height of Summer respectively, whereby a considerable uncertainty exists as to the amount of Refraction; in the case of the Madras Observations however, the meridian altitude being 85° and the variation of temperature at the times of the Star arriving at the + and — maximum amounting to little or nothing, no such uncertainty exists. With regard to error of observation the above speak so well for themselves that it is unnecessary for me to offer any further remarks.

* From a few very accordant Transit Observations the Parallax of α Aquilæ comes out $0'',49$, or the diameter of the Earth's orbit viewed from α Aquilæ = $0'',98$: these it will be as well to reserve for a future opportunity when a greater number of observations shall have been made.

PLACES OF THE FIXED STARS.

At the commencement of my Superintendance of the Madras Observatory in 1830, I selected for observation a Catalogue of about 1200 of the brightest Stars, from the Catalogue of 2881 given in the 2d Volume of the Memoirs of the Royal Astronomical Society, and set to work, intending to make at least five Observations of each Star: towards the end of 1831, finding that the greater part of this Catalogue was then already completed, I determined to extend my observations to the remaining Stars of the Society's Catalogue, and to devote the Instruments during the years 1832 and 1833 solely to this purpose; the result of the three years observation are given in the pages which follow reduced to January 1, 1832. It may at first sight appear superfluous that the result of the observations for 1831 which have already been given in Vol. I. should again appear in the present work; but several of the Stars observed in 1831 having been again observed in 1832 and 1833, it became necessary to state the former results in order to obtain the mean of *all* the observations; added to which, the peculiar circumstances to which the Transit Instrument has been subject (by reason of the very rapid and unequal wear of the pivots, and the meridian marks having undergone a change of position), renders it desirable that the nature of the agreement between the observations of one year and another should be distinctly pointed out.

It will be noticed that I have retained *all the names*, and consequently the same *numbers*, as given in the Society's Catalogue, notwithstanding that from twenty to thirty Stars (from being situated near the South Pole) are invisible at Madras, and that about double of that number have not been observed at either the Transit or the Mural Circle; my reason for so doing was for the sake of uniformity and facility of reference, and to allow me to fill up the blanks with a pen from the observations of 1834 and 1835.

The magnitudes and Annual Precessions* are copied from the Society's Catalogue: the *Greenwich place* is derived from the Catalogue of 720 Stars for 1830; in which the place of the equinox is assumed Dr. Maskelyne + 0".20, and it is from this point which the places in this Catalogue are likewise

† Where an asterisk is attached to the Annual Precession it denotes, that the proper motion exceeds 0".5s. of space (according to M. Piazzi) and that it is included with the precession.

reckoned. Under the head "No. 1831", "No. 1832", &c. is found the number of observations made in each year, and the corresponding mean result on the supposition that the pivots remained unaltered during the three years; to make the requisite correction, we must have recourse to the table at Page 8 and proceed as follows—thus for α Cassiopeæ N. P. D. $45^{\circ} 34'$ which was observed at the commencement of 1832 and 1833, we find :

From Observa- tions in	Mean Place Ja- nuary 1, 1832.			Correc- tion.	No. of Obs.		
	<i>h.</i>	<i>m.</i>	<i>s.</i>		<i>s.</i>		<i>s.</i>
1831.....	0	31	1,42	+ 0 00	= 1,42	× 12	= 17,04
1832.....	0	0	1,24	+ 0 04	= 1,28	× 20	= 25 60
1833.....	0	0	1,09	+ 0 09	= 1,18	× 28	= 33,04
					Sum	60	75,68

$$\frac{\text{Sum}}{\text{No. of Observ.}} = \frac{75\ 68}{60} = 1',26$$

The place thus deduced is set down in column "Mean", and compared with the Greenwich, and the Astronomical Society's Catalogue.

The column "Mean N. P. D." is derived from the three preceding columns in the usual way with reference to the number of observations; the Greenwich N. P. D. is derived from the Catalogue of 720 Stars for 1830, increased by two years precession, and reduced by the table Vol. I. Pages 62 and 147, in order to render the results which were computed by Bradley's table of Refractions, in terms of Atkinson's: this Catalogue rests upon the supposition that the Latitude of the Greenwich Royal Observatory deduced from Bradley's table of Refraction = $51^{\circ} 28' 39",00$.

A

GENERAL CATALOGUE

OF THE

PRINCIPAL FIXED STARS

FROM

OBSERVATIONS AT THE MADRAS OBSERVATORY

IN THE YEARS 1831, 1832 AND 1833,

COMPARED WITH

THE GREENWICH, AND ASTRONOMICAL SOCIETY'S CATALOGUE.

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion			
			No.	1831	No.				1832	No.		1833	Green.	A. S.
			s.	s.	s.				h.	m.		s.	s.	s.
1	7	Ceti	2	6,82	5	6,79	—	0 0 6,80	15,71	6,45	+0,35	+3,068		
2	2.3	11 Cassiop. β	6	15,55	—	—	0 0 15,55	—	14,73	-0,16	+0,82	3,124*		
3	6	87 Pegasi μ	—	—	5	23,10	—	0 0 23,10	—	22,46	+0,64	3,069		
4	6	AppSculp κ^1	1	46,56	2	46,68	4	46,68	—	46,07	+0,57	3,066		
5	4	Phœnicis ϵ	6	51,81	—	—	0 0 51,81	—	51,51	+0,30	+0,30	3,063		
6	7	Piscium	—	—	5	19,05	0 1 19,05	—	18,63	+0,42	+0,42	3,068		
7	6	34 Piscium E^1	—	—	6	24,70	0 1 24,70	—	23,93	+0,77	+0,77	3,069		
8	5	22 Androm B	5	37,04	—	—	0 1 37,04	—	36,98	+0,06	+0,06	3,077		
9	6.7	Ceti	—	—	5	42,92	0 1 42,92	—	42,54	+0,38	+0,38	3,067		
10	5	Octantis γ^3	—	—	—	—	Invisible	—	10,01	—	—	2,967		
11	6	6 Ceti f	—	—	4	42,72	0 2 42,72	—	42,05	+0,67	+0,67	3,064		
12	5.6	AppSculp κ^2	6	1,77	—	—	0 3 1,77	—	1,65	+0,12	+0,12	3,059		
13	2.3	88 Pegasi γ	12	35,66	24	35,69	16	35,74	35,71	35,51	-0,02	+0,18	3,075	
14	6	89 Pegasi χ	2	55,09	3	55,37	—	0 5 55,26	—	54,95	+0,31	3,080		
15	5.6	7 Ceti h	—	—	5	6,26	—	0 6 6,26	—	5,86	+0,40	3,055		
16	6	35 Piscium B	—	—	5	20,02	—	0 6 20,02	—	19,60	+0,42	3,073		
17	6.7	36 Piscium	—	—	5	56,61	—	0 7 56,61	—	55,85	+0,76	3,074		
18	5	24 Androm θ	6	20,26	—	—	0 8 20,26	—	19,48	+0,78	+0,78	3,105		
19	6.7	33 Piscium	6	10,31	—	—	0 9 10,31	—	9,89	+0,42	+0,42	3,069		
20	4	8 Ceti i	6	52,14	5	52,16	2	52,22	52,15	51,75	+0,01	+0,41	3,057	
21	5	Tucanæ ζ	6	15,80	—	—	0 11 15,80	—	12,83	+2,97	+2,97	2,923		
22	6	40 Piscium	2	16,03	5	15,92	—	0 11 15,95	—	15,22	+0,73	3,086		
23	5.6	41 Piscium d	4	57,47	7	57,66	5	57,58	57,60	57,82	-0,01	-0,23	3,077	
24	6	Ap. Sculp i	—	—	6	4,26	—	0 13 4,25	—	4,00	+0,25	3,025		
25	6	9 Ceti	6	15,10	—	—	0 14 15,10	—	15,22	-0,12	-0,12	3,049		
26	6.7	Ceti	1	54,79	5	54,88	2	55,07	—	54,67	+0,24	3,063		
27	3	Hydri β	—	—	—	—	Invisible	—	37,97	—	—	2,606		
28	6	44 Piscium t	7	47,63	—	—	0 16 47,63	—	47,19	+0,44	+0,44	3,070		
29	6	45 Piscium	—	—	6	2,68	—	0 17 2,68	—	2,01	+0,67	3,080		
30	5	Phœnicis κ	5	55,61	—	—	0 17 55,61	—	54,29	+1,32	+1,32	2,966		
31	2	Phœnicis α	6	57,86	11	57,84	—	0 17 57,84	—	57,21	+0,63	2,970		
32	6	10 Ceti	—	—	5	0,78	4	0,75	—	0,40	+0,37	3,066		
33	6	47 Piscium	—	—	3	18,05	3	18,26	—	17,76	+0,39	3,102		
34	6	48 Piscium	6	29,83	—	—	0 19 29,83	—	28,96	+0,87	+0,87	3,099		
35	6	28 Androm	3	16,42	2	16,60	—	0 21 16,51	—	15,95	+0,56	3,136		
36	6	Ceti	—	—	6	21,04	—	0 21 21,04	—	20,23	+0,81	3,033		
37	6	12 Ceti π	—	—	5	28,17	2	28,20	—	27,77	+0,41	3,057		
38	6	Ceti	—	—	6	58,44	—	0 21 58,44	—	58,39	+0,05	3,009		
39	5	14 Cassiopeæ λ	6	32,86	—	—	2	32,67	—	32,00	+0,83	3,245		
40	5	Phœnicis λ^1	6	17,64	—	—	0 23 17,64	—	16,68	+0,96	+0,96	2,909		
41	4	15 Cassiopeæ κ	5	30,89	—	—	0 23 30,89	30,79	29,80	+0,10	+1,09	3,324		
42	6.7	51 Piscium	—	—	5	44,17	—	0 23 44,17	—	43,80	+0,37	3,083		
43	6	52 Piscium	—	—	5	47,94	—	0 23 47,94	—	47,36	+0,58	3,116		
44	4	Tucanæ β^1	6	48,36	—	—	0 23 48,36	—	48,41	-0,05	-0,05	2,786		
45	4	Tucanæ β^2	—	—	3	49,35	—	0 23 49,22	—	48,77	+0,45	2,786		

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
	"	"	"				"	"	
1	—	—	4 9 25,31	93 9 25,31	—	9 25,79	—	0,48	-20,043
2	5 46 37,06	4 46 36,52	—	31 46 36,82	46 37,91	46 39,46	-1,09	2,64	20,043
3	—	—	4 43 16,97	72 43 16,97	—	43 19,12	—	2,15	20,043
4	—	—	5 55 20,45	118 55 20,45	—	55 23,01	—	2,56	20,043
5	5 40 24,30	—	—	136 40 24,30	—	40 25,62	—	1,32	20,042
6	—	—	5 29 43,17	93 29 43,17	—	29 45,10	—	1,93	20,042
7	—	—	5 47 25,89	79 47 25,89	—	47 19,22	+	6,67	20,042
8	5 51 50,12	—	—	44 51 50,12	—	51 46,46	+	3,66	20,042
9	2 10 55,69	4 10 55,94	—	96 10 55,86	—	10 55,20	+	0,66	20,042
10	—	—	—	173 —	—	9 29,62	—	—	20,042
11	4 23 23,42	1 23 24,39	—	106 23 23,61	—	23 27,96	—	4,35	20,041
12	—	3 44 4,62	3 44 4,39	118 44 4,51	—	44 8,62	—	4,11	20,041
13	22 45 3,67	8 45 2,98	—	75 45 3,49	45 2,93	45 1,81	+0,56	1,68	20,039
14	5 43 40,51	—	—	70 43 40,51	—	43 35,11	+	5,40	20,036
15	2 51 51,97	4 51 53,75	—	109 51 53,16	—	51 43,51	+	4,65	20,036
16	—	5 7 45,44	—	82 6 45,44	—	6 41,81	+	3,63	20,035
17	3 41 33,71	2 41 33,44	—	82 41 33,60	—	41 32,31	+	1,29	20,031
18	4 15 5,59	9 15 5,67	—	52 15 5,63	—	15 5,23	+	0,40	20,030
19	5 14 36,35	—	—	89 14 36,35	—	14 44,52	—	8,17	20,027
20	5 45 23,44	6 45 24,96	—	99 45 24,27	45 20,55	45 17,08	+3,72	7,19	20,021
21	5 51 45,54	5 51 46,99	—	155 51 46,27	—	53 47,78	+	121,51	20,019
22	5 40 57,20	—	—	74 40 57,20	—	40 52,59	+	4,61	20,019
23	6 44 33,38	1 44 32,92	—	82 44 33,31	44 36,50	44 33,93	-3,19	0,62	20,016
24	—	5 55 42,09	—	119 54 42,09	—	54 38,51	+	3,58	20,011
25	3 8 36,53	2 8 38,37	—	103 8 37,27	—	8 38,53	—	1,26	20,004
26	5 8 52,53	1 8 51,07	—	93 8 52,29	—	8 56,48	—	4,19	19,995
27	—	—	—	168 —	—	12 1,02	—	—	19,990
28	—	5 59 26,90	—	88 59 26,90	—	59 25,11	+	1,79	19,990
29	—	5 14 16,88	—	83 14 16,88	—	14 15,08	+	1,80	19,988
30	5 36 46,74	5 36 48,41	—	134 36 47,57	—	36 40,38	+	7,19	19,982
31	8 13 5,03	5 13 5,57	—	133 13 5,30	—	12 56,19	+	9,11	19,982
32	5 58 50,28	—	—	90 58 50,29	—	58 47,37	+	2,02	19,981
33	8 2 12,96	3 2 14,83	—	73 2 13,47	—	2 14,08	—	0,61	19,972
34	—	5 29 5,07	—	74 29 5,07	—	29 1,94	+	3,13	19,971
35	—	5 10 30,50	—	61 10 30,50	—	10 32,62	—	2,12	19,957
36	—	2 47 32,21	3 47 32,22	105 47 32,62	—	47 32,67	—	0,05	19,957
37	—	—	4 53 15,65	94 53 15,65	—	53 10,85	+	4,80	19,956
38	—	2 43 7,23	3 43 6,52	114 43 6,80	—	43 4,93	+	1,47	19,951
39	5 24 21,52	5 24 22,17	—	36 24 21,85	—	24 11,86	+	9,99	19,947
40	5 44 1,60	5 44 2,45	—	139 44 2,02	—	43 6,42	+	55,60	19,940
41	5 59 44,70	1 59 45,63	6 59 43,40	27 59 44,12	59 48,44	59 49,77	-4,32	5,65	19,938
42	4 58 20,66	—	—	83 58 20,66	—	58 19,91	+	6,75	19,936
43	—	2 37 55,06	3 37 55,32	70 37 55,22	—	37 50,85	+	4,37	19,936
44	—	—	—	153 —	—	53 18,49	—	—	19,935
45	—	—	—	153 —	—	53 40,49	—	—	19,935

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832			Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	h.	m.	s.			Green.	A. S.	
46	5	Tucanae β^3	—	—	5 2,80	0	25	2,65	—	2,44	+0,21	+2,771	
47	7	Piscium	—	5 28,99	—	0	25	28,99	—	28,47	+0,52	3,092	
48	7	Ceti	—	6 55,43	—	0	25	55,43	—	55,12	+0,31	3,054	
49	5.7	Piscium	—	5 13,29	—	0	26	13,30	—	13,04	+0,26	3,102	
50	6	13 Ceti	—	6 36,35	—	0	26	36,35	—	35,42	+0,93	3,056	
51	6.7	Piscium (120)	—	3 55,65	1 55,36	0	26	55,58	—	55,63	-0,05	3,064	
52	4	17 Cassiopeae ζ	7 39,27	—	—	0	27	39,27	39,18	38,86	+0,09	+0,41	3,280
53	4.5	29 Androm π	6 55,69	—	—	0	27	55,69	55,71	55,77	-0,02	-0,08	3,172
54	6	53 Piscium	5 2,86	—	—	0	28	2,86	—	2,16	+0,70	3,109	
55	6	Ceti	—	4 42,63	—	0	28	42,62	—	39,05	+3,57	2,988	
56	7	Piscium	—	5 51,86	—	0	28	51,86	—	51,19	+0,67	3,074	
57	7	15 Ceti	—	6 29,44	—	0	29	29,44	—	28,76	+0,68	3,064	
58	4	30 Androm ϵ	6 41,73	—	—	0	29	41,73	41,81	41,24	-0,08	+0,49	3,161
59	3	31 Androm δ	6 21,74	—	—	0	30	21,74	21,81	21,32	-0,07	+0,42	3,169
60	3	18 Cassiopeae α	12 1,42	20 1,24	28 1,09	0	31	1,21	1,26	0,76	-0,05	+0,45	3,330
61	6	55 Piscium	—	5 5,79	—	0	31	5,79	—	5,40	+0,39	3,135	
62	6.7	Ceti	5 9,38	—	—	0	32	9,38	—	8,89	+0,49	3,051	
63	5	Phœnicis μ	6 22,30	—	—	0	32	22,30	—	22,26	+0,02	2,861	
64	6	Ceti	1 45,76	4 45,55	—	0	33	45,59	33,45,34	33,45,34	+0,25	3,024	
65	5	20 Cassiopeae π	6 12,38	—	—	0	34	12,38	—	11,58	+0,80	3,274	
66	6	Ceti	—	5 18,70	—	0	34	18,70	—	18,46	+0,24	2,991	
67	2.3	16 Ceti β	7 9,18	8 9,42	8 9,26	0	35	9,29	9,27	9,23	+0,02	+0,06	2,998
68	5	17 Ceti ϕ^1	5 42,82	—	—	0	35	42,82	—	42,63	+0,19	3,026	
69	5	Phœnicis η	4 46,86	—	—	0	35	46,86	—	45,80	+1,06	2,731	
70	6	Ceti	—	5 25,50	—	0	36	25,50	—	25,32	+0,18	2,979	
71	6	Ceti	—	5 51,34	—	0	36	51,34	—	50,92	+0,42	3,047	
72	6	18 Ceti	—	7 2,50	—	0	37	2,50	—	1,95	+0,55	3,015	
73	6.7	57 Piscium	—	5 45,99	—	0	37	45,99	—	46,06	-0,07	3,125	
74	6	58 Piscium	—	5 16,17	—	0	38	16,17	—	15,29	+0,88	3,111	
75	6	59 Piscium	—	5 21,78	—	0	38	21,78	—	21,28	+0,50	3,143	
76	4	34 Androm ζ	6 26,94	—	—	0	38	26,94	27,01	27,01	-0,07	-0,07	3,164
77	6	60 Piscium	—	4 42,79	—	0	38	42,79	—	41,87	+0,92	3,091	
78	4	24 Cassiopeae ν	4 59,32	2 59,19	—	0	38	59,31	59,16	58,27	+0,15	+1,04	3,533*
79	6	Piscium	—	5 34,84	—	0	39	34,84	—	32,81	+2,03	3,086	
80	6	62 Piscium	—	2 35,11	5 35,00	0	39	35,04	—	34,36	+0,68	3,094	
81	5	63 Piscium δ	6 58,51	—	3 58,65	0	39	58,56	—	58,30	+0,26	3,095	
82	5.6	64 Piscium γ	—	5 9,73	—	0	40	9,74	—	9,49	+0,25	3,135	
83	4	35 Androm ν	5 34,79	1 35,08	4 34,63	0	40	34,78	34,84	34,25	-0,06	+0,53	3,266
84	6	65 Piscium i	5 52,82	—	—	0	40	52,82	—	52,14	+0,68	3,187	
85	6	19 Ceti ϕ^2	—	7 42,94	—	0	41	42,94	—	42,81	+0,13	3,019	
86	5	20 Ceti m	7 25,64	8 25,73	2 25,74	0	44	25,69	25,68	25,72	+0,01	-0,03	3,059
87	6	66 Piscium	—	6 42,44	—	0	45	42,45	—	41,80	+0,65	3,155	
88	6	36 Androm	6 59,24	1 59,33	—	0	45	59,26	—	58,94	+0,32	3,179	
89	3	27 Cassiopeae γ	5 37,85	12 37,65	8 37,61	0	46	37,67	37,83	36,79	-0,16	+0,88	3,531
90	6	67 Piscium k	2 57,31	2 57,35	—	0	46	57,33	—	56,97	+0,36	3,202	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue:	A. S. Catalogue.	Difference from:		Annual Precession.			
	No.	1831	No.				1832	No.		1833	Green.	A. S. C
	"	"	"				"	"		"	"	"
46	4 57	22,87	5 57	24,16	—	153 57	23,65	57 27,57	—	3,92	19,924	
47	—	—	2 37	18,60	3 37	18,69	80 37	18,64	37 11,70	+	6,94	19,920
48	—	—	4 28	29,46	2 28	31,23	95 28	30,07	28 26,54	+	3,53	19,916
49	3 33	9,42	5 33	10,35	—	—	77 33	10,90	33 11,52	—	1,52	19,913
50	—	—	1 31	10,22	5 3	8,57	94 31	8,84	31 5,37	+	3,47	19,909
51	—	—	5 25	44,95	—	—	91 25	44,95	25 43,05	+	1,90	19,905
52	5 1	45,03	6 1	44,54	4 1	44,11	37 1	44,57	1 43,57	+1,00	+ 1,62	19,898
53	6 12	24,53	—	—	—	—	57 12	24,53	12 23,44	+1,09	+ 2,73	19,895
54	5 41	37,56	—	—	—	—	75 41	37,56	41 35,06	+	2,50	19,894
55	—	—	1 41	35,15	5 41	34,67	115 41	34,75	41 33,43	+	1,32	19,887
56	—	—	—	—	5 47	14,80	87 47	14,80	47 10,98	+	3,82	19,885
57	—	—	4 25	40,87	—	—	91 25	40,87	25 40,38	+	0,49	19,878
58	4 36	5,22	—	—	3 36	1,00	61 36	3,41	36 4,10	—0,69	— 1,70	19,876
59	5 3	37,56	6 3	38,47	—	—	60 3	38,06	3 32,69	+5,37	+ 6,53	19,868
60	16 23	8,74	15 23	8,77	—	—	34 23	8,76	23 7,54	+1,22	+ 1,27	19,861
61	5 29	3,63	—	—	—	—	60 29	3,63	29 2,08	+	1,55	19,860
62	4 16	31,72	1 16	32,32	—	—	95 16	31,34	16 27,58	+	4,26	19,847
63	5 0	28,80	—	—	—	—	137 0	28,80	0 18,30	+	10,50	19,832
64	4 43	30,63	1 43	32,11	—	—	102 43	30,92	43 33,41	—	2,49	19,827
65	5 53	44,03	4 53	42,52	—	—	43 53	43,36	53 46,03	—	2,67	19,821
66	—	—	4 6	55,34	—	—	111 6	55,34	6 55,43	—	0,09	19,820
67	5 54	33,23	5 54	34,09	—	—	108 54	33,66	54 33,21	—1,27	— 1,55	19,809
68	4 31	30,42	2 31	30,69	—	—	101 31	30,48	31 38,57	—	3,09	19,801
69	5 23	3,42	—	—	—	—	118 23	3,42	23 3,54	—	0,12	19,800
70	3 55	46,30	2 55	51,90	—	—	112 55	48,54	55 51,83	—	3,29	19,791
71	—	—	5 33	6,38	—	—	95 33	6,38	33 4,43	+	1,95	19,786
72	—	—	4 47	27,17	—	—	103 47	27,17	47 32,26	—	5,09	19,783
73	—	—	4 26	32,52	—	—	75 26	32,52	26 27,39	+	5,13	19,773
74	—	—	—	—	5 56	35,05	78 56	35,05	56 33,65	+	1,40	19,766
75	1 20	26,00	—	—	4 20	26,33	71 20	26,26	20 22,95	+	3,31	19,764
76	5 38	50,85	—	—	7 38	52,13	66 38	51,59	38 52,76	—1,17	— 1,24	19,763
77	4 10	38,32	—	—	—	—	84 10	38,32	10 33,68	+	4,64	19,759
78	5 4	39,71	—	—	—	—	33 4	39,71	4 42,23	—2,52	— 0,98	19,755*
79	—	—	5 35	8,20	—	—	85 35	8,20	34 27,41	—	—	19,747
80	—	—	—	—	5 37	4,41	83 37	4,41	37 7,54	—	3,13	19,746
81	5 19	47,91	5 19	46,83	—	—	83 19	47,37	19 46,61	+	0,76	19,740
82	—	—	5 58	1,34	—	—	73 58	1,34	58 1,48	—	0,14	19,737
83	5 50	19,58	5 50	18,25	—	—	49 50	18,91	50 17,29	+3,86	+ 1,62	19,731
84	5 12	19,88	—	—	—	—	63 12	19,98	12 22,67	—	2,79	19,726
85	—	—	5 33	1,33	—	—	101 33	1,33	33 1,64	—	0,31	19,713
86	5 3	23,93	5 3	26,03	—	—	92 3	27,48	3 29,13	—1,65	— 1,39	19,669
87	5 43	25,11	—	—	—	—	71 43	25,11	43 27,36	—	2,25	19,647
88	5 16	56,54	—	—	—	—	67 16	56,54	16 57,56	—	1,02	19,642
89	9 11	45,60	5 11	44,85	9 11	46,42	30 11	45,76	11 37,18	+3,28	+ 8,28	19,632
90	—	—	4 42	6,73	—	—	63 42	6,73	42 7,39	—	0,66	19,625

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832			Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	h.	m.	s.			Green.	A. S.	
91	5	Cephei	4 10,56	—	—	0 47 10,56	—	—	4,94	—	+5,62	+3,996	
92	6	Ceti	—	5 13,26	—	0 47 13,26	—	—	12,81	—	+0,45	3,028	
93	7	Piscium	—	5 20,89	—	0 47 20,90	—	—	20,51	—	+0,39	3,131	
94	4	37 Androm	5 27,42	—	3 26,93	0 47 27,26	—	27,48	27,17	-0,22	+0,09	3,359*	
95	6	22 Ceti	—	6 36,21	1 36,00	0 47 36,18	—	—	35,71	—	+0,47	3,009	
96	5	38 Androm	6 15,20	1 15,14	—	0 48 15,19	—	—	14,35	—	+0,84	3,183	
97	6	68 Piscium	—	4 45,88	—	0 48 45,90	—	—	45,11	—	+0,79	3,218	
98	6.7	Piscium	—	5 6,25	—	0 49 6,26	—	—	5,88	—	+0,38	3,132	
99	6	23 Ceti	1 19,22	5 19,23	—	0 50 19,22	—	—	18,95	—	+0,27	3,005	
100	5	App Sculp.	6 30,31	—	—	0 50 30,31	—	—	30,36	—	-0,05	2,898	
101	6.7	Piscium	—	5 4,62	—	0 51 4,62	—	—	7,03	—	-2,41	3,097	
102	7	Piscium	3 45,77	—	—	0 53 45,77	—	—	45,21	—	+0,56	3,111	
103	4	71 Piscium	6 13,96	8 13,86	4 13,85	0 54 13,90	—	13,96	14,00	-0,06	-0,10	3,106	
104	6	25 Ceti	—	5 32,96	—	0 54 32,96	—	—	32,50	—	+0,46	3,036	
105	6.7	26 Ceti	—	5 10,71	—	0 55 10,71	—	—	10,11	—	+0,60	3,071	
106	6.7	73 Piscium	—	5 10,83	—	0 56 10,83	—	—	10,64	—	+0,19	3,095	
107	6	72 Piscium	—	3 13,99	3 14,12	0 56 14,08	—	—	13,30	—	+0,78	3,149	
108	5.6	74 Piscium	2 41,41	1 41,86	—	0 56 41,57	—	—	41,31	—	+0,26	3,191	
109	6	27 Ceti	4 12,06	—	—	0 57 12,06	—	—	11,77	—	+0,29	3,005	
110	6	28 Ceti	8 39,57	—	—	0 57 39,57	—	—	39,49	—	+0,08	3,005	
111	6.7	75 Piscium	—	5 43,99	—	0 57 43,97	—	—	43,74	—	+0,23	3,139	
112	3.4	Phœnicis	6 34,66	—	—	0 58 34,66	—	—	34,50	—	+0,16	2,698	
113	6	79 Piscium	—	—	3 57,39	0 58 57,42	—	—	56,88	—	+0,54	3,190	
114	6	30 Ceti	—	—	3 19,31	0 59 19,30	—	—	19,12	—	+0,18	3,004	
115	2.3	1 Ursae Min.	—	—	—	1 0 2,19	—	3,14	1,81	-0,95	-0,38	15,430*	
116	5	80 Piscium	2 43,54	—	2 43,30	0 59 43,42	—	4 3,45	43,51	-0,03	-0,09	3,097	
117	5	42 Androm	—	—	2 47,08	0 59 47,18	—	—	46,96	—	+0,22	3,428	
118	3.4	31 Ceti	—	—	3 8,37	1 0 8,37	—	8,46	8,63	-0,69	-0,26	3,000	
119	2	43 Androm	—	—	3 21,16	1 0 21,23	—	20,96	20,28	+0,27	+0,95	3,309	
120	6	81 Piscium	—	—	3 51,26	1 0 51,29	—	—	50,75	—	+0,54	3,187	
121	4.5	33 Cassiopeae	—	—	3 55,11	1 1 55,24	—	55,19	55,20	+0,05	+0,04	3,555	
122	6	Piscium	—	—	2 17,75	1 1 17,77	—	—	16,98	—	+0,79	3,161	
123	5	Phœnicis	—	—	2 18,29	1 1 18,18	—	—	18,08	—	+0,10	2,542	
124	6	32 Ceti	—	—	2 46,21	1 1 46,20	—	—	46,20	—	0,00	3,007	
125	6	33 Ceti	—	1 55,44	2 55,23	1 1 55,30	—	—	54,79	—	+0,51	3,078	
126	6	83 Piscium	—	—	3 25,98	1 2 26,03	—	—	24,93	—	+1,10	3,268	
127	5	84 Piscium	—	—	3 26,31	1 2 26,34	—	—	26,54	—	-0,20	3,200	
128	7	Piscium	—	—	3 41,17	1 2 41,19	—	—	41,36	—	-0,17	3 128	
129	6.7	34 Ceti	—	—	5 11,09	1 3 11,09	—	—	10,68	—	+0,41	3,048	
130	6.7	35 Ceti	2 54,16	3 54,19	—	1 3 54,18	—	—	53,87	—	+0,31	3,078	
131	6	85 Piscium	—	5 38,52	—	1 4 38,54	—	—	38,41	—	-0,13	3,331	
132	6	86 Piscium	—	5 57,82	—	1 4 57,82	—	57,79	57,76	+0,03	+0,06	3 112	
133	6.7	87 Piscium	—	4 12,98	—	1 5 12,99	—	—	12,13	—	+0,86	3,170	
134	6	37 Ceti	—	—	3 56,53	1 5 56,53	—	—	53,92	—	+0,61	3,009	
135	6.7	88 Piscium	6 58,57	—	—	1 5 58,57	—	—	57,98	—	+0,59	3,108	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession	
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.		
	" "	" "	" "				" "	" "		
91	5 38 58,35	—	—	4 38 58,35	—	39 0,28	—	1,93	19,625	
92	—	4 15 22,84	—	98 15 22,84	—	15 23,80	—	0,96	19,621	
93	—	4 57 34,60	1 57 33,07	76 57 34,47	—	57 33,65	+	0,82	19,618	
94	5 24 47,07	5 24 46,98	5 24 49,89	52 24 48,28	24 46,29	24 49,12	+1,99	—	0,84	19,617
95	—	5 11 40,81	—	102 11 40,81	—	10 39,72	+	1,09	19,614	
96	5 29 23,91	5 29 22,06	—	67 29 22,98	—	29 25,67	—	2,69	19,602	
97	4 55 2,58	—	—	61 55 2,58	—	55 3,90	—	1,32	19,593	
98	—	5 12 49,41	—	77 12 49,41	—	12 49,17	+	0,24	19,586	
99	—	5 17 17,50	—	102 17 17,50	—	17 17,87	—	0,37	19,563	
100	5 16 2,70	5 16 2,12	—	120 16 2,41	—	16 0,13	+	2,28	19,560	
101	2 25 30,55	3 25 30,34	—	81 25 30,44	—	25 27,97	+	2,47	19,548	
102	5 5 1,44	—	—	82 5 1,44	—	4 58,62	+	2,82	19,496	
103	7 0 55,88	6 0 55,27	5 0 54,95	83 0 55,42	0 57,81	0 57,83	-2,39	—	2,41	19,486
104	2 44 18,03	3 44 12,26	—	95 44 12,57	—	44 8,66	+	3,91	19,480	
105	—	5 32 10,96	—	89 32 10,96	—	32 7,92	+	3,04	19,467	
106	—	5 14 48,99	2 14 47,68	85 14 48,62	—	14 42,79	+	5,83	19,446	
107	—	5 57 32,92	—	75 57 32,92	—	57 31,39	+	1,53	19,445	
108	5 25 41,23	—	—	69 25 41,23	—	25 41,23	—	0,00	19,435	
109	—	5 51 47,54	—	100 51 47,54	—	52 43,68	+	3,86	19,424	
110	—	5 44 29,64	—	100 44 29,64	—	44 24,81	+	4,83	19,414	
111	—	1 56 46,03	4 56 43,60	77 56 44,08	—	56 46,01	—	1,93	19,412	
112	5 37 11,75	5 37 9,75	—	137 37 10,75	—	37 2,81	+	7,94	19,394	
113	—	4 9 24,62	—	70 9 24,62	—	9 19,22	+	5,40	19,386	
114	—	3 41 9,89	—	100 41 9,89	—	41 11,51	—	1,62	19,377	
115	10 35 12,17	11 35 12,57	3 35 12,94	1 35 12,44	35 12,60	35 12,20	-0,16	+	0,24	19,375
116	5 14 30,74	—	3 14 31,69	65 14 31,10	14 28,36	14 28,02	+2,74	+	3,08	19,368
117	5 39 19,22	—	—	43 39 19,22	—	39 22,03	—	2,81	19,367	
118	5 4 27,40	—	—	101 4 27,40	4 29,30	4 25,08	-1,90	+	2,32	19,359
119	5 16 22,94	2 16 22,30	—	55 16 22,76	16 19,41	16 17,61	+3,35	+	5,15	19,355
120	—	—	6 14 23,10	71 14 23,10	—	14 22,55	+	0,55	19,343	
121	4 44 47,01	1 44 47,33	—	35 44 47,07	44 46,75	44 48,41	+0,32	—	1,34	19,341
122	—	5 13 23,85	—	75 13 23,85	—	13 18,06	+	5,79	19,333	
123	—	—	4 8 43,27	146 8 43,27	—	8 31,30	+	11,97	19,332	
124	—	5 43 6,81	—	99 43 6,81	—	47 56,65	—	—	—	19,321
125	—	5 26 59,38	—	88 26 59,38	—	27 0,61	—	1,23	19,318	
126	—	5 48 17,42	—	60 48 17,42	—	48 13,60	+	3,82	19,306	
127	6 51 41,98	—	—	60 51 41,98	—	51 38,18	+	3,80	19,306	
128	—	4 36 13,89	—	80 36 13,89	—	36 10,81	+	3,08	19,300	
129	3 8 38,31	—	2 8 40,17	93 8 39,06	—	8 42,26	—	3,20	19,288	
130	—	4 25 1,65	—	88 25 1,65	—	24 58,94	+	2,71	19,271	
131	—	—	5 18 26,71	66 18 26,71	—	18 28,67	—	1,96	19,253	
132	1 18 53,12	—	5 18 53,73	83 18 53,63	18 54,82	18 53,38	-1,19	+	0,25	19,245
133	—	—	5 45 30,31	74 45 30,31	—	45 28,26	+	2,05	19,240	
134	5 49 35,96	—	—	98 49 35,96	—	49 36,27	—	0,31	19,221	
135	—	5 53 42,15	—	83 53 42,15	—	53 39,62	+	2,58	19,221	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832		Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833					Green.	A. S.	
			s.	s.	s.	h. m. s.	s.					
136	6	38 Ceti	7 14,92	—	—	1 6 14,92	—	14,70	—	+0,22	+3,056	
137	6	39 Ceti	1 4,78	—	4 4,87	1 8 4,85	—	4,70	—	+0,15	3,045	
138	6	40 Ceti	—	3 23,24	2 23,64	1 8 23,40	—	23,29	—	+0,11	3,046	
139	6	89 Piscium	f	5 8,51	—	1 9 8,51	—	8,16	—	+0,35	3,087	
140	5.6	90 Piscium	v	5 14,89	—	1 10 14,89	—	14,82	—	+0,07	3,268	
141	6	42 Ceti	z	1 13,44	5 13,35	—	1 11 13,37	—	13,18	—	+0,19	3,058
142	6	91 Piscium	i	—	5 51,05	—	1 11 51,07	—	51,17	—	-0,10	3,285
143	5	46 Androm	z	6 29,10	1 29,04	—	1 12 29,09	—	28,54	—	+0,55	3,478
144	7	Ceti	—	5 58,44	—	1 13 58,44	—	58,32	—	+0,12	3,074	
145	6.7	43 Ceti	z	—	5 59,78	—	1 13 59,78	—	59,41	—	+0,37	3,058
146	4.5	36 Cassiopea	↓	6 10,30	2 10,44	1 10,29	1 14 10,39	10,45	9,43	-0,06	+0,96	4,079
147	3	37 Cassiopea	δ	2 53,88	2 53,50	5 53,54	1 14 53,69	—	55,11	—	-1,42	4,833
148	6	44 Ceti	—	—	5 35,76	—	1 15 35,76	—	35,46	—	+0,30	3,000
149	3.	45 Ceti	θ	—	6 37,84	—	1 15 37,84	—	37,55	—	+0,29	2,999
150	5.6	93 Piscium	ρ	—	6 12,74	—	1 17 12,76	—	12,40	—	+0,36	3,214
151	5	Phoenicia	—	6 14,34	—	—	1 17 14,34	—	14,29	—	+0,05	2,665
152	5	46 Ceti	c	6 21,62	—	—	1 17 21,62	—	21,40	—	+0,22	2,946
153	6.7	94 Piscium	ρ	—	—	5 38,16	1 17 38,18	—	37,87	—	+0,31	3,215
154	6.7	Ceti	—	—	5 51,88	—	1 17 51,88	—	51,51	—	+0,37	3,058
155	6	47 Ceti	—	—	5 34,39	—	1 17 34,38	—	34,97	—	-0,59	2,957
156	7	95 Piscium	—	—	5 57,04	—	1 18 57,04	—	56,87	—	+0,17	3,103
157	7	Piscium	—	—	—	5 22,97	1 18 22,99	—	22,32	—	+0,67	3,200
158	7	Piscium	—	—	—	5 35,08	1 19 35,09	—	34,86	—	+0,23	3,124
159	6.7	96 Piscium	—	—	—	5 17,49	1 20 17,50	—	16,91	—	+0,59	3,120
160	6.7	97 Piscium	—	—	6 49,64	—	1 20 49,66	—	49,01	—	+0,65	3,213
161	3	Phoenicia	γ	6 3,86	—	—	1 21 3,86	—	3,20	—	+0,66	2,619
162	5	98 Piscium	μ	5 23,52	1 23,32	7 23,43	1 21 23,46	23,32	23,09	+0,24	+0,37	3,111
163	6	48 Ceti	—	—	5 32,63	—	1 21 32,62	—	32,55	—	+0,07	2,875
164	6	Ceti	—	—	5 46,73	—	1 21 46,71	—	—	—	—	2,836
165	6	Ap. Scalp	—	—	5 27,84	—	1 22 27,83	—	27,02	—	+0,80	2,828
166	4	99 Piscium	π	6 30,46	6 30,40	—	1 22 30,44	30,35	30,27	+0,09	+0,17	3,189
167	4	Phoenicia	δ	5 15,00	—	—	1 24 15,00	—	14,69	—	+0,31	2,497
168	7	Piscium	—	—	6 30,35	—	1 24 30,35	—	30,11	—	+0,24	3,130
169	6	Piscium	—	—	5 44,65	—	1 25 44,67	—	43,86	—	+0,81	3,223
170	7	100 Piscium	—	—	5 56,91	—	1 25 56,92	—	56,41	—	+0,51	3,169
171	5.8	49 Ceti	—	—	5 25,06	—	1 26 25,64	—	24,72	—	+0,92	2,922
172	6	101 Piscium	—	—	5 48,19	—	1 26 48,21	—	47,45	—	+0,76	3,189
173	6	Piscium	—	—	3 50,36	3 50,51	1 26 50,45	—	49,82	—	+0,63	3,215
174	5	50 Androm	v	6 58,17	1 57,55	—	1 26 58,08	—	57,50	—	+0,58	3,491
175	6.7	Piscium	—	—	5 15,99	—	1 27 16,09	—	15,76	—	+0,24	3,127
176	3.4	51 Androm	R	6 43,35	—	—	1 27 43,35	43,28	42,93	+0,07	+0,42	3,617
177	6	50 Ceti	—	—	—	6 47,50	1 27 47,49	—	46,82	—	+0,67	2,922
178	6	102 Piscium	π	4 12,19	—	6 12,17	1 28 12,19	12,18	11,90	+0,01	+0,29	3,168
179	6	Ceti	—	—	5 14,20	—	1 29 14,19	—	13,08	—	+1,11	2,976
180	6.7	104 Piscium	—	—	—	5 16,05	1 30 16,06	—	16,30	—	-0,24	3,190

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	No. 1831	No. 1832	No. 1833				Green.	A. S. C	
136		2 52 29,33	3 52 30,25	91 52 29,88		52 27,00	+ 2,88	+ 19,214	
137	4 23 8,47	1 23 9,09		93 23 8,59		23 8,41	+ 0,18	19,167	
138		5 9 38,60		93 9 38,60		9 38 64	+ 0,04	19,159	
139	4 16 19,55	1 16 17,85		87 16 19,21		16 14,40	+ 4,81	19,140	
140		5 37 14,43		63 37 14,43		37 12,26	+ 2,17	19,111	
141		5 23 35,87		91 23 35,87		23 31,64	+ 4,23	19,085	
142	3 8 30,33	1 8 30,27		62 8 30,32		8 31,16	- 0,84	19,069	
143	7 21 18,45	3 21,18 43		45 21 18,45		21 11,24	+ 7,21	19,052	
144	5 9 8,75			89 9 8,75		9 9,39	- 0,64	19,011	
145		5 19 48,96		91 19 48,96		19 47,47	+ 1,49	19,010	
146	6 44 58,73	3 44 59,26		22 44 58,91	45 1,67	45 2,29	- 2,76	19,006	
147	6 38 29,52	5 38 28,74		30 38 29,16		38 25,79	+ 3,37	18,986	
148	1 52 55,81	4 52 56,69		98 52 56,51		52 53,92	+ 2,59	18,965	
149	5 3 6,39		2 3 8,80	99 3 7,04		3 7,25	- 0,17	18,964	
150		7 42 15,69		71 42 15,69		42 12,35	+ 3,34	18,919	
151	5 22 5,77		1 22 8,83	132 22 6,23		22 9,01	- 2,73	18,918	
152	5 28 32,62	3 28 32,00		105 28 32,39		28 26,62	+ 5,77	18,915	
153		5 37 56,66		71 37 56,66		37 55,11	+ 1,55	18,907	
154	3 16 25,24		2 16 25,07	91 16 25,17				18,900	
155			5 55 54,73	103 55 54,73		55 56,71	- 1,98	18,879	
156			5 30 58,50	85 30 58,50		30 53,20	+ 5,30	18,868	
157		5 47 30,58		73 47 30,58		47 31,57	- 0,99	18,856	
158			5 54 40,69	82 54 40,69		54 42,07	- 1,38	18,850	
159		5 34 29 53		83 34 29,53		34 29,06	+ 0,47	18,829	
160	5 30 51,82			72 30 51,82		30 54,23	- 2,41	18,813	
161	5 10 49,87	1 10 50,89		134 10 50,04		10 45,91	+ 4 13	18,805	
162	4 43 32,04		6 43 32,09	84 43 32,07	43 23,74	43 33,89	+ 8,33	18,795	
163		4 29 59,68	1 30 0,46	112 29 59,83		30 0,62	- 0,79	18,790	
164	1 29 20,02	3 29 19,88		116 29 19,91		29 18,88	+ 1 03	18,785	
165		1 4 40,11	4 4 39,84	117 4 39,89		4 40,02	- 0,13	18,762	
166	6 31 23,26	5 31 25,36	4 31 24,06	75 31 24,17	31 22,61	31 20,08	+ 1,56	+ 4,09 18,761	
167	5 56 56,23	3 56 56,78		139 56 56,44		56 40,27	+ 16,17	18,706	
168	5 39 17,65			82 39 17,65		39 18,96	- 1,31	18,699	
169	5 23 57,69			72 23 57,69		23 53,57	+ 4,12	18,660	
170		5 18 13,19		78 18 13,19		18 12,05	+ 1,14	18,653	
171		5 32 22,60		106 32 22,60		32 22,58	+ 0,02	18,638	
172		5 12 0,14		76 12 0,14		11 57,60	+ 2,54	18,626	
173		2 25 41,35	3 25 42,29	73 25 41,91		25 44,59	- 2,68	18,625	
174	5 26 18,21			49 26 18,21		26 15,89	+ 2,32	18,621	
175			5 13 0,17	83 13 0,17		13 1,36	- 1,19	18,611	
176	5 13 35,35		3 13 36,15	42 13 35,65	13 34,61	13 34,35	+ 1,04	+ 1,30 18,596	
177		5 15 43,04		106 15 43,04		15 38,10	+ 4,94	18,594	
178	5 43 9,33			78 43 9,33	43 14,98	43 11,35	- 5,65	- 2,02 18,580	
179		5 16 3,82		100 16 3,82		16 3,24	+ 0,58	18,546	
180		4 34 8,71	1 34 7,39	76 34 8,45		34 9,14	- 0,69	18,511	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833				Green.	A. S.	
			s	s	s				s.	s.	
181	6	105 Piscium	—	4 37,79	5 37,78	1 30 37,81	37,71	37,67	+0,10	+0,14	+3,211
182	1	Eridani α	5 27,19	3 27,31	3 27,39	1 31 27,23	—	26,51	—	+0,72	2,235
183	7	Piscium	—	5 44,73	—	1 31 44,74	—	43,94	—	+0,80	3,140
184	5	106 Piscium	6 41,57	6 41,90	6 41,90	1 32 41,79	41,86	41,60	-0,07	+0,19	3,111
185	5	54 Androm	5 10,53	—	—	1 33 10,53	—	10,36	—	+0,17	3,693
186	5.6	107 Piscium	4 23,42	5 23,56	—	1 33 23,51	—	23,29	—	+0,31	3,255
187	6.7	109 Piscium	—	6 46,12	—	1 35 46,14	—	45,72	—	+0,42	3,257
188	3.4	52 Ceti τ	5 15,98	7 16,06	—	1 36 16,01	16,01	15,64	0,00	+0,37	2,779*
189	5	110 Piscium	6 31,85	4 31,81	3 31,93	1 36 31,86	31,82	31,70	+0,04	+0,16	3,148
190	6	Ceti	—	5 33,71	—	1 37 33,71	—	33,27	—	+0,44	3,004
191	5	App Sculp ϵ	6 46,78	1 46,78	—	1 37 46,78	—	46,01	—	+0,77	2,800
192	6.7	4 Arietis	—	5 4,99	—	1 39 5,01	—	4,81	—	+0,20	3,230
193	6	Arietis	—	5 52,60	—	1 40 52,62	—	52,13	—	+0,49	3,291
194	5	53 Ceti α^2	5 20,30	1 20,35	—	1 41 20,31	—	19,40	—	+0,91	2,952
195	6	54 Ceti	1 57,53	5 57,39	—	1 41 57,85	—	56,76	—	+1,09	3,171
196	3.4	45 Cassiopeae ϵ	6 24,11	—	—	1 42 24,11	23,49	23,47	+0,62	+0,64	4,191
197	3	55 Ceti ζ	6 10,25	1 10,46	—	1 43 10,28	10,35	9,82	-0,07	+0,46	2,953
198	3.4	2 Trianguli α	6 31,59	1 31,48	5 31,50	1 43 31,56	31,52	31,23	+0,04	+0,33	3,388
199	4.5	5 Arietis γ^1	5 19,56	3 19,56	4 19,65	1 44 19,60	19,58	19,51	+0,02	+0,09	3,264
200	5.6	111 Piscium ξ	—	5 52,00	—	1 44 52,00	—	51,79	—	+0,21	3,092
201	3	6 Arietis β	—	12 22,54	16 22,52	1 45 22,55	22,55	21,77	0,00	+0,78	3,233
202	6	7 Arietis	—	5 29,93	—	1 46 29,95	—	29,45	—	+0,50	3,319
203	7	Piscium	—	5 13,24	—	1 47 13,24	—	12,92	—	+0,32	3,079
204	5	Phoenicis ϕ	6 23,78	—	—	1 47 23,78	—	23,77	—	+0,01	2,499
205	6	8 Arietis ι	—	6 11,23	4 11,25	1 48 11,26	11,24	11,16	+0,02	+0,10	3,253
206	5	48 Cassiopeae	4 18,15	—	—	1 48 18,15	—	18,17	—	-0,02	4,744
207	5.6	9 Arietis λ	—	5 35,05	1 35,01	1 48 35,06	34,98	35,17	+0,08	-0,11	3,324
208	6	56 Ceti ν^1	—	—	5 48,23	1 48 48,21	—	47,30	—	+0,91	2,804
209	4.5	50 Cassiopeae	4 15,21	2 14,79	—	1 49 15,13	15,46	14,23	-0,33	+0,90	4,903
210	4	Eridani χ	6 25,12	—	—	1 49 25,12	—	23,86	—	+1,26	2,270
211	6	Arietis	—	—	5 17,14	1 50 17,16	—	16,48	—	+0,68	3,296
212	7	Piscium	—	—	5 26,91	1 50 26,93	—	26,50	—	+0,43	3,194
213	4.5	Hydri η^2	2 41,32	2 41,17	—	1 50 41,19	—	40,73	—	+0,46	1,495
214	7	Ceti	—	4 10,93	—	1 51 10,93	—	10,45	—	+0,48	3,125
215	6	112 Piscium	—	3 25,44	2 25,71	1 51 25,55	—	25,20	—	+0,35	3,093
216	6	57 Ceti ι	—	4 52,26	—	1 51 52,24	—	51,92	—	+0,32	2,819
217	4.5	59 Ceti ν^2	6 5,50	—	—	1 52 5,30	5,36	4,99	+0,14	+0,51	2,816
218	5	113 Piscium α	6 21,68	1 21,80	2 21,70	1 53 21,70	21,65	21,73	+0,05	-0,03	3,090
219	3	Hydri α	3 28,77	2 28,65	—	1 53 28,69	—	28,87	—	-0,16	1,854
220	3.4	57 Androm γ	—	7 37,11	2 37,06	1 53 37,12	37,10	36,71	+0,02	+0,41	3,630
221	7	Arietis	—	—	6 0,78	1 54 0,80	—	0,31	—	+0,49	3,183
222	6	Arietis Λ	—	2 30,66	3 30,93	1 54 30,84	—	30,33	—	+0,51	3,269
223	6	60 Ceti	—	1 35,29	4 35,14	1 54 35,17	—	35,29	—	-0,12	3,060
224	5	Phoenicis χ	6 57,98	—	—	1 54 57,98	—	58,34	—	-0,36	2,414
225	6	12 Arietis κ	1 10,36	4 10,95	—	1 57 10,96	—	10,43	—	+0,53	3,330

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N P D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831		No. 1832		No. 1833					Green.	A. S. C.	
	' "	' "	' "	' "	' "	' "						
181	—	—	5 26 58,70	—	—	74 26 58,70	26 58,50	26 55,71	+0,20	+ 2,99	+18,499	
182	7 5 30,61	6 5 30,27	—	3 5 30,75	148 5 30,49	148 5 30,49	—	5 35,18	—	— 4,69	18,471	
183	3 5 37,89	—	—	—	82 5 37,89	82 5 37,89	—	5 38,18	—	— 0,29	18,462	
184	5 21 59,47	4 21 58,03	—	5 21 58,14	85 21 58,58	85 21 58,58	21 55,67	21 55,29	+2,91	+ 3,29	18,429	
185	5 9 44,47	—	—	—	40 9 44,47	40 9 44,47	—	9 40,43	—	+ 4,04	18,413	
186	4 33 4,46	1 33 6,31	—	—	70 33 4,83	70 33 4,83	—	32 58,88	—	+ 5,95	17,835*	
187	3 45 31,92	2 45 33,16	—	—	70 45 32,42	70 45 32,42	—	45 29,66	—	+ 2,76	18,322	
188	5 49 27,29	5 49 27,23	—	—	106 49 27,26	106 49 27,26	49 29,75	49 24,77	-2,49	+ 2,49	19,144*	
189	5 41 26,30	5 41 25,94	—	7 41 25,71	81 41 25,95	81 41 25,95	41 26,67	41 25,66	-0,72	+ 0,29	18,295	
190	—	6 34 34,31	—	—	96 34 34,31	96 34 34,31	—	34 34,60	—	— 0,29	18,258	
191	5 53 40,46	1 53 41,78	—	—	115 53 40,68	115 53 40,68	—	53 41,21	—	— 0,53	18,250	
192	4 52 59,63	—	—	1 53 1,73	73 53 0,07	73 53 0,07	—	53 0,30	—	— 0,23	18,203	
193	1 33 38,70	4 33 40,02	—	—	68 33 39,76	68 33 39,76	—	33 43,08	—	— 3,32	18,137	
194	5 31 13,53	5 31 11,94	—	—	101 31 12,73	101 31 12,73	—	31 13,87	—	— 1,14	18,119	
195	5 47 33,14	—	—	—	79 47 33,14	79 47 33,14	—	47 30,92	—	+ 2,22	18,096	
196	6 9 42,79	—	—	5 9 48,59	27 9 43,15	27 9 43,15	9 43,98	9 44,25	-0,78	+ 1,10	18,081	
197	5 10 4,07	—	—	5 10 5,31	101 10 4,69	101 10 4,69	10 6,40	10 9,49	-1,71	+ 4,80	18,050	
198	5 14 35,14	6 14 34,89	—	2 14 33,72	61 14 34,81	61 14 34,81	14 33,90	14 36,82	+0,91	+ 2,01	18,037	
199	2 31 58,67	3 31 58,71	—	1 31 59,56	71 31 58,73	71 31 58,73	32 0,83	31 55,20	-2,10	+ 3,53	18,006	
200	3 38 33,32	2 38 39,52	—	—	87 38 33,80	87 38 33,80	—	38 44,50	—	— 5,70	17,985	
201	5 1 2,84	6 1 3,29	—	11 1 3,48	70 1 3,28	70 1 3,28	0 59,91	0 56,81	+3,57	+ 6,47	17,966	
202	2 14 57,21	4 14 57,90	—	—	67 14 57,67	67 14 57,67	—	14 57,09	—	+ 0,58	17,922	
203	—	5 59 10,45	—	—	88 59 10,45	88 59 10,45	—	59 14,57	—	— 4,12	17,893	
204	5 19 21,76	2 19 21,48	—	2 19 23,86	133 19 22,16	133 19 22,16	—	19 25,90	—	+ 3,74	17,885	
205	—	1 0 19,80	—	7 0 18,94	73 0 18,99	73 0 18,99	0 21,66	0 21,64	-2,67	+ 2,65	17,855	
206	4 54 51,19	—	—	—	19 54 51,19	19 54 51,19	—	54 47,87	—	+ 3,82	17,852	
207	3 13 37,10	1 13 33,82	—	5 13 36,02	67 13 36,14	67 13 36,14	13 37,67	13 33,27	-1,53	+ 2,87	17,839	
208	—	2 20 57,52	—	3 20 58,33	113 20 58,01	113 20 58,01	—	21 1,82	—	— 3,81	17,830	
209	5 23 48,34	1 23 47,19	—	—	18 23 48,14	18 23 48,14	23 49,41	23 50,50	-1,27	+ 2,36	17,815	
210	5 26 47,82	2 26 47,03	—	—	142 26 47,85	142 26 47,85	—	26 58,47	—	— 10,62	17,805	
211	1 45 41,26	3 45 41,08	—	—	69 45 41,12	69 45 41,12	—	45 37,56	—	+ 3,56	17,771	
212	—	—	—	5 31 22,49	78 31 22,49	78 31 22,49	—	31 21,03	—	+ 1,46	17,764	
213	5 28 24,88	—	—	—	158 28 24,88	158 28 24,88	—	28 34,33	—	— 10,05	17,753	
214	—	—	—	6 47 1,54	84 47 1,54	84 47 1,54	—	46 57,00	—	+ 4 54	17,735	
215	—	5 42 34,23	—	—	87 42 34,23	87 42 34,23	—	42 35,77	—	— 1,54	17,735	
216	2 38 34,54	3 38 36,63	—	—	111 38 35,82	111 38 35,82	—	38 37,51	—	— 1,69	17,706	
217	4 53 41,09	3 53 40,63	—	—	111 53 40,89	111 53 40,89	53 51,98	53 38,09	-11,09	+ 2,80	17,697	
218	5 3 0,91	3 3 1,63	—	2 3 0,60	88 3 1,06	88 3 1,06	3 4,32	3 0,44	-3,26	+ 0,62	17,645	
219	4 23 18,88	—	—	—	152 23 18,88	152 23 18,88	—	23 9,83	—	+ 9,05	17,638	
220	5 28 47,23	—	—	—	48 28 47,23	48 28 47,23	28 49,93	28 50,74	-2,70	+ 3,51	17,635	
221	—	5 47 45,54	—	—	79 47 45,54	79 47 45,54	—	47 39,19	—	+ 6,35	17,618	
222	—	2 33 24,21	—	4 33 24,96	72 33 24,71	72 33 24,71	—	33 26,63	—	+ 1,92	17,597	
223	—	—	—	5 41 8,27	90 41 8,27	90 41 8,27	—	41 2,12	—	+ 6,15	17,593	
224	3 31 29,77	1 31 30,53	—	—	185 31 29,96	185 31 29,96	—	31 39,06	—	— 10,00	17,576	
225	—	5 9 18,30	—	—	68 9 18,30	68 9 18,30	—	9 18,44	—	+ 0,14	17,484	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^b January 1, 1832			A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831			No. 1832				Green.	A. S.	
			s.	s.	s.	h.	m.	s.				
226	6	Arietis	—	5 12,27	—	1 57 12,29	—	12,11	—	+0,18	+3,373	
227	3	13 Arietis	α 10 43,13	5 43,18	46 43,22	1 57 43,22	43,17	42,80	+0,05	+0,42	3,342	
228	4	4 Trianguli	β 6 34,35	5 34,40	—	1 59 34,39	34,34	34,49	+0,05	-0,10	3,520	
229	5.6	14 Arietis	—	—	—	1 59 52,70	—	52,47	—	+0,23	3,381	
230	6.7	62 Ceti	—	2 55,53	—	2 0 55,35	—	55,07	—	+0,28	3,108	
231	6	15 Arietis	—	4 19,66	2 19,61	2 1 19,67	—	19,09	—	+0,57	3,296	
232	6.7	64 Ceti	—	5 29,56	—	2 2 29,56	—	29,42	—	+0,14	3,161	
233	5.6	6 Trianguli	—	5 38,74	1 38,98	2 2 38,81	—	38,53	—	+0,28	3,453	
234	6	63 Ceti	—	6 4,50	—	2 3 4,50	—	3,71	—	+0,79	3,037	
235	6	17 Arietis	γ 3 24,71	2 24,61	—	2 3 24,68	—	24,11	—	+0,57	3,323	
236	7	19 Arietis	—	7 54,30	—	2 3 54,34	54,24	53,62	+0,10	+0,72	3,245	
237	5	65 Ceti	ζ ¹ 7 6,40	6 6,43	7 6,43	2 4 6,44	6,47	5,84	-0,03	+0,60	3,165	
238	6	67 Ceti	—	5 36,62	—	2 8 36,62	—	36,18	—	+0,44	2,978	
239	6	22 Arietis	θ ¹ 2 47,62	4 47,83	3 47,84	2 8 47,82	47,80	47,39	+0,02	+0,43	3,315	
240	6	Ceti	—	6 18,10	—	2 9 18,10	—	17,41	—	+0,69	3,080	
241	4	Eridani	φ 6 30,55	3 30,30	—	2 10 30,16	—	30,11	—	+0,35	2,136	
242	5	9 Persei	—	6 42,23	3 42,23	2 10 42,27	—	41,92	—	+0,35	4,100	
243	Var.	68 Ceti	—	—	6 52,10	2 10 52,10	—	51,80	—	+0,30	3,021	
244	6	69 Ceti	—	5 20,70	—	2 13 20,70	—	19,76	—	+0,94	3,063	
245	6	70 Ceti	—	1 39,26	4 39,37	—	2 13 39,35	—	38,82	+0,53	3,047	
246	6	Fornacia	—	5 51,50	—	2 14 51,48	—	50,42	—	+1,06	2,729	
247	4.5	Cassiopeae	—	4 20,50	5 20,38	10 19,69	2 15 20,13	20,22	20,48	-0,09	-0,35	4,788
248	6.7	Ceti	—	2 32,57	4 32,49	—	2 15 32,51	—	32,35	—	+0,16	3,185
249	6	24 Arietis	ζ ¹ —	5 49,38	—	2 15 49,38	—	49,13	—	+0,25	3,197	
250	6	71 Ceti	—	5 29,44	—	2 16 29,44	—	29,28	—	+0,16	3,022	
251	6	Arietis	—	2 45,59	2 45,75	2 45,62	2 17 45,66	—	45,44	—	+0,22	3,198
252	5	72 Ceti	ρ 6 50,33	—	—	2 17 50,33	—	50,13	—	+0,20	2,893	
253	6	12 Trianguli	—	1 20,44	5 20,39	2 18 20,43	—	20,15	—	+0,28	3,487	
254	4	Hydri	δ 4 46,56	2 47,45	—	2 18 47,06	—	46,02	—	+1,04	1,041	
255	5	73 Ceti	ξ ² —	12 14,29	12 14,29	2 19 14,29	14,25	14,11	+0,04	+0,18	3,171	
256	4.5	Eridani	—	4 49,63	4 49,52	—	2 20 49,56	—	47,60	—	+1,96	2,199
257	6.7	Arietis	—	—	5 53,53	—	2 20 53,55	—	53,00	—	+0,55	3,419
258	6.7	26 Arietis	—	—	5 14,14	—	2 21 14,16	—	13,54	—	+0,62	3,335
259	6	27 Arietis	—	—	—	6 36,09	2 21 36,12	—	35,78	—	+0,34	3,304
260	6	Fornacia	—	1 53,16	4 53,38	—	2 22 53,32	—	53,13	—	+0,19	2,732
261	5.6	75 Ceti	—	—	5 36,94	—	2 23 36,94	—	37,17	—	-0,23	3,044
262	6.7	29 Arietis	—	—	—	4 42,98	2 23 43,00	—	42,59	—	+0,41	3,267
263	5	76 Ceti	—	6 7,81	1 7,53	—	2 24 7,77	—	7,54	—	+0,23	2,843
264	6.7	Arietis	—	—	—	11 13,94	2 24 13,96	—	13,58	—	+0,38	3,325
265	6.7	Ceti	—	—	5 11,07	—	2 26 11,07	—	10,83	—	+0,24	3,162
266	6	77 Ceti	—	—	5 25,78	—	2 26 25,78	—	25,42	—	+0,36	2,948
267	6	Fornacia	—	—	5 29,47	—	2 26 29,45	—	29,36	—	+0,09	2,627
268	6.7	Ceti	—	3 52,63	2 52,67	—	2 26 52,72	—	48,59	—	+4,13	3,153
269	4.5	78 Ceti	—	6 4,00	5 3,92	7 4,01	2 27 3,98	4,02	3,53	-0,04	+0,45	3,136
270	6	30 Arietis	—	—	2 17,58	4 17,58	2 27 17,61	—	16,55	—	+1,06	3,423

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from.		Annual Precession.
	1831		1832		1833					Green.	A. S. C	
	No.	"	No.	"	No.	"				"	"	
226	3 58	33,35	2 58	33,65	—	64 58	33,47	—	—	—	-17,488	
227	30 20	7,65	11 20	7,82	50 20	8,22	67 20	7,98	20 8,67	20 6,62	-0,69 + 1,36	17,461
228	6 18	41,49	5 48	41,79	—	—	55 48	41,63	48 41,93	48 37,14	-0,30 + 4,49	17,581
229	—	—	5 51	35,69	—	—	64 51	35,69	—	51 31,04	+ 4,65	17,368
230	—	—	—	—	6 34	0,23	86 34	0,23	—	33 58,34	+ 1,89	17,322
231	—	—	3 17	44,35	2 17	44,00	71 17	44,21	—	17 44,14	+ 0,07	17,304
232	4 13	13,02	1 13	14,28	—	—	82 13	13,27	—	13 12,78	+ 0,49	17,252
233	—	—	—	—	4 29	18,03	60 29	18,03	—	29 13,87	+ 4,16	17,246
234	—	—	3 37	6,74	—	—	93 37	6,74	—	37 4,79	+ 1,95	17,226
235	—	—	—	—	5 34	59,64	69 34	59,64	—	34 53,44	+ 6,20	17,212
236	—	—	5 30	42,75	—	—	75 30	42,75	30 41,21	30 39,57	+1,54 + 3,18	17,190
237	5 56	42,78	5 56	41,38	—	—	81 56	42,08	56 41,57	56 41,00	+0,51 + 1,08	17,180
238	5 12	0,01	—	—	—	—	97 12	0,01	—	11 59,24	+ 0,77	16,974
239	4 52	47,19	2 52	47,29	—	—	70 52	47,22	52 47,23	52 46,37	-0,01 + 0,85	16,966
240	—	—	5 2	23,91	—	—	89 2	23,91	—	2 36,85	- 12,94	16,942
241	5 17	33,24	5 17	33,14	—	—	142 17	33,19	—	17 33,83	- 0,64	16,883
242	4 55	48,97	5 55	49,99	—	—	34 55	49,54	—	55 47,79	+ 1,75	16,877
243	—	—	—	—	—	—	93 44	—	—	44 35,16	—	16,868
244	5 22	35,49	—	—	—	—	90 22	35,49	—	22 32,30	+ 3,19	16,750
245	5 39	10,89	—	—	—	—	91 39	10,89	—	39 11,69	- 0,80	16,735
246	—	—	5 34	58,99	—	—	114 34	58,99	—	34 57,91	+ 1,08	16,677
247	5 21	32,95	5 21	30,86	3 21	31,25	23 21	31,75	21 34,08	21 36,73	-2,33 - 4,98	16,656
248	—	—	5 3	0,30	—	—	81 3	0,30	—	2 57,20	+ 3,10	16,644
249	—	—	4 9	19,38	—	—	80 9	19,38	—	9 14,75	+ 4,63	16,630
250	6 32	37,29	—	—	—	—	93 32	37,29	—	32 37,97	- 0,68	16,597
251	4 11	45,28	—	—	—	—	80 11	45,28	—	11 41,06	+ 4,22	16,535
252	5 3	3,96	5 3	4,18	—	—	103 3	4,07	—	3 2,80	+ 1,27	16,531
253	—	—	5 5	7,08	—	—	61 5	7,08	—	5 2,98	+ 4,10	16,307
254	5 25	35,45	—	—	—	—	159 25	35,45	—	25 36,10	- 0,65	16,481
255	6 17	49,27	—	—	5 17	49,56	82 17	49,41	17 48,68	17 50,54	+0,73 - 1,13	16,461
256	5 27	36,78	5 27	35,28	—	—	138 27	36,03	—	27 43,84	- 7,81	16,382
257	—	—	5 30	50,09	—	—	65 30	50,09	—	30 48,04	+ 2,05	16,379
258	5 53	41,69	—	—	—	—	70 53	41,69	—	53 37,39	+ 4,30	16,362
259	5 2	34,14	—	—	—	—	73 2	34,14	—	2 32,32	+ 1,82	16,343
260	2 17	34,48	3 17	35,67	—	—	113 17	35,19	—	17 35,08	+ 0,11	16,277
261	—	—	5 46	49,86	—	—	91 46	49,86	—	46 50,41	- 0,55	16,240
262	—	—	1 42	51,07	4 42	49,67	75 42	49,95	—	42 47,63	+ 2,32	16,235
263	5 59	9,24	5 59	8,84	—	—	105 59	9,04	—	59 1,81	+ 7,23	16,213
264	—	—	5 51	54,51	—	—	71 51	54,51	—	51 52,87	+ 1,64	16,209
265	—	—	5 15	49,63	—	—	83 15	49,63	—	15 45,80	+ 3,83	16,107
266	5 35	45,14	—	—	—	—	98 35	45,44	—	35 45,39	+ 0,05	16,094
267	5 53	21,09	—	—	—	—	118 53	24,09	—	53 25,05	- 0,96	16,090
268	—	—	1 55	5,86	4 55	9,39	83 55	8,68	—	55 54,64	- 45,96	16,075
269	5 8	43,96	1 8	43,00	5 8	42,97	85 8	43,42	8 38,36	8 36,24	+5,06 + 7,18	16,062
270	—	—	—	—	10 5	17,99	66 5	17,99	—	5 19,00	- 1,01	16,052

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.		Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832		Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
				No. 1831	No. 1832	No. 1833	h.	m. s.			Green.	A. S.	
				s	s	s					s.	s.	
271	6.7	Arietis			2 20,33	3 20,50	2 27 20,42				+0,44	+3,423	
272	6	31 Arietis	<i>v</i>	2 28,92		6 28,99	2 27 28,98		28,72		+0,26	3,234	
273	6.7	Ceti		5 41,68			2 27 41,68		41,52		+0,16	3,166	
274	6	80 Ceti	<i>e</i> ²			4 44,30	2 2 44,30		44,01		+0,29	2,947	
275	5.6	81 Ceti	<i>d</i> ³		4 14,30		2 29 14,30		14,41		-0,11	3,010	
276	5.6	32 Arietis	<i>v</i>			6 17,62	2 29 17,64		17,19		+0,45	3,382	
277	6	33 Arietis				4 53,13	2 30 53,16		52,59		+0,57	3,472	
278	4	82 Ceti	<i>δ</i>	6 52,81	6 52,77		2 30 52,79	52,81	52,77	-0,02	+0,02	3,062	
279	7	Ceti			3 25,01	1 25,30	2 31 25,23		24,20		+1,03	3,145	
280	4.5	83 Ceti	<i>ε</i>	6 26,47	5 26,57		2 31 26,52	26,59	26,41	-0,07	+0,11	2,885	
281	6	84 Ceti				5 38,17	2 3 38,17		37,37		+0,80	3,048	
282	4	13 Persei	<i>θ</i>		3 46,00	1 46,09	2 32 46,07	45,86	45,64	-0,21	+0,43	4,046*	
283	6	34 Arietis	<i>μ</i>			5 54,69	2 32 54,71		53,61		+1,10	3,357	
284	7	Arietis				5 58,69	2 32 58,71		57,55		+1,16	3,211	
285	5	Eridani			5 23,78		2 33 23,75		23,78		-0,03	2,278	
286	6	85 Ceti				3 27,00	2 33 27,01		26,59		+0,42	3,214	
287	4	35 Arietis	<i>α</i>		5 36,79		2 33 36,81	36,75	36,11	+0,06	+0,70	3,490	
288	4.5	Eridani	<i>ι</i>		5 2,48		2 34 2,46		1,57		+0,89	2,356	
289	3	86 Ceti	<i>γ</i>	6 36,11			2 34 36,11	36,26	35,98	-0,15	+0,13	3,105	
290	7	36 Arietis				2 57,54	2 34 57,56		56,96		+0,60	3,324	
291	6.7	37 Arietis	<i>ο</i>			3 18,39	2 35 18,11		17,77		+0,64	3,286	
292	5.6	38 Arietis		1 48,99		7 49,14	2 35 49,13		48,41		+0,72	3,242	
293	4	87 Ceti	<i>μ</i>	6 52,30	5 52,21		2 35 52,27	52,23	51,66	+0,04	+0,61	3,207	
294	4	89 Ceti	<i>π</i>	6 7,86	2 7,95		2 36 7,69	7,79	7,46	+0,10	+0,33	2,849	
295	5	Hydri	<i>ε</i>				2 36		54,60			0,868	
296	5	Hydri	<i>ξ</i>	1 2,03	3 2,55	6 2,39	2 37 2,29		0,65		+1,64	0,866	
297	5.6	1 Eridani	<i>τ</i> ¹			4 16,30	2 37 16,28		15,65		+0,63	2,772	
298	4	39 Arietis	<i>β</i>		6 55,41		2 37 55,44		55,26		+0,18	3,530	
299	5	Persei			4 20,51	2 29,76	2 38 29,65		29,55		+0,10	4,292	
300	6.7	16 Trianguli			2 0,99	2 0,86	2 39 0,86		0,19		+0,77	3,457	
301	6	40 Arietis				7 7,90	2 39 7,62		7,31		+0,61	3,339	
302	5	42 Arietis	<i>π</i>		4 55,85	2 55,73	2 39 55,62		55,03		+0,79	3,326	
303	4.5	16 Persei	<i>p</i> ¹		6 0,38		2 40 0,42	0,31	0,47	+0,11	-0,05	3,729	
304	3	41 Arietis	<i>c</i>	3 6,77	4 6,81		2 40 6,80	6,76	6,63	+0,04	+0,17	3,497	
306	5	Fornacis	<i>v</i>	5 55,51			2 41 55,61		55,74		-0,23	2,388	
306	5	Fornacis	<i>β</i>		5 3,74		2 42 3,72		3,36		+0,36	2,502	
307	6	43 Arietis	<i>σ</i>	1 14,00	1 13,79	2 14,11	2 42 14,01		13,01		+1,00	3,291	
308	5	18 Persei	<i>τ</i>	6 23,77	1 23,53	1 23,61	2 42 23,74	23,54	22,99	+0,20	+0,75	4,182	
309	6	Fornacis	<i>γ</i> ¹			5 24,63	2 42 24,60		24,45		+0,15	2,658	
310	5	Hydri	<i>ζ</i>		1 59,13	4 59,21	2 42 59,07		58,63		+0,44	0,874	
311	4.5	2 Eridani	<i>τ</i> ²	1 25,27	6 25,24	7 25,33	2 43 25,27	25,41	25,02	-0,14	+0,25	2,720	
312	7	Arietis		1 51,26	4 51,25		2 43 51,26		50,77		+0,49	3,316	
313	6	45 Arietis	<i>p</i> ²	1 23,03	4 22,84		2 46 22,90		22,35		+0,55	3,350	
314	6	46 Arietis	<i>p</i> ³	1 58,12	2 57,84	5 57,95	2 46 57,96	57,92	57,86	+0,04	+0,10	3,346	
315	6.7	Arietis		1 15,63	2 15,49	3 15,33	2 47 15,34		15,24		+0,10	3,188	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in					Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession	
	No. 1831		No. 1832		No. 1833				Green.	A. S. C.		
	'	"	'	"	'				"	"		"
271	—	—	—	—	—	66	—	5 20,40	—	—	-16,048	
272	—	—	—	—	5 17 3,81	78 17 3,81	—	17 2,98	+	0,83	16,040	
273	—	—	—	—	5 0 18,04	83 0 18,04	—	0 18,81	-	0,77	16,028	
274	—	—	—	—	6 33 57,28	98 33 57,28	—	33 52,84	+	4,44	16,026	
275	—	—	5 7 36,91	—	—	94 7 36,91	—	7 35,82	+	1,09	15,947	
276	—	—	5 46 13,53	—	—	68 46 13,53	—	46 11,91	+	1,62	15,945	
277	—	—	5 39 51,92	—	—	63 39 51,92	—	39 51,52	+	0,40	15,860	
278	5 24 4,86	—	3 24 3,96	—	1 24 5,79	90 24 4,47	24 2,00	24 3,52	+ 2,47	+	0,95	15,859
279	3 36 57,15	—	2 36 58,86	—	—	84 36 57,83	—	36 56,40	+	1,43	15,831	
280	5 35 20,08	—	4 35 18,86	—	—	102 35 19,54	35 11,73	35 19,43	+ 7,81	+	0,11	15,829
281	—	—	3 24 53,23	—	2 24 51,96	91 24 52,73	—	24 48,49	+	4,24	15,766	
282	4 29 15,63	—	—	—	5 29 17,63	41 29 16,74	29 16,71	29 16,09	+ 0,03	+	0,65	15,760
283	—	—	—	—	5 42 31,23	70 42 31,23	—	42 29,06	+	2,17	15,752	
284	—	—	—	—	5 10 42,50	80 10 42,50	—	10 34,76	+	7,74	15,747	
285	5 36 53,83	—	—	—	—	133 36 53,83	—	36 53,92	-	2,09	15,723	
286	—	—	—	—	7 58 50,43	79 58 50,43	—	58 38,07	+	12,36	15,722	
287	4 0 42,14	—	5 0 44,01	—	—	63 0 43,17	0 46,40	0 45,32	- 3,23	-	2,15	15,713
288	4 34 40,94	—	—	—	—	130 34 40,94	—	34 38,47	-	2,47	15,688	
289	5 28 34,63	—	—	—	—	87 28 34,63	28 35,90	28 36,38	- 1,27	-	1,75	15,658
290	—	—	5 57 1,71	—	—	72 57 1,71	—	57 5,05	-	3,34	15,640	
291	—	—	—	—	5 24 20,15	75 24 20,15	—	24 11,62	-	8,53	15,621	
292	3 15 52,51	—	—	—	2 15 55,75	78 15 53,81	—	15 53,70	+	0,11	15,593	
293	5 36 2,08	—	5 36 0,49	—	5 36 1,52	80 36 1,36	35 58,82	35 54,02	+ 2,54	+	7,34	15,590
294	5 31 30,30	—	5 34 28,73	—	—	104 34 29,51	34 28,58	34 17,34	+ 0,93	+	12,17	15,574
295	—	—	—	—	—	158	—	58 49,90	—	—	—	15,528
296	—	—	—	—	5 59 20,50	158 59 20,50	—	59 17,78	+	2,72	15,522	
297	—	—	5 17 16,07	—	—	109 17 16,07	—	17 15,79	+	0,28	15,511	
298	5 27 22,64	—	4 27 20,49	—	4 27 20,82	61 27 21,42	27 21,39	27 20,26	+ 0,03	+	1,16	15,476
299	6 43 32,77	—	—	—	—	34 48 32,77	—	48 30,91	+	1,86	15,446	
300	—	—	5 31 6,28	—	—	65 31 6,28	—	31 3,60	+	2,68	15,416	
301	—	—	2 25 10,43	—	—	72 25 10,43	—	25 8,12	+	2,31	15,409	
302	5 14 18,16	—	5 14 19,43	—	—	73 14 18,60	—	14 18,69	-	0,09	15,364	
303	—	—	5 22 44,17	—	—	52 22 44,17	22 42,40	22 40,43	+ 1,77	+	3,74	15,360
304	—	—	1 26 11,10	—	9 26 11,60	63 26 11,55	26 12,40	26 12,05	- 0,85	-	0,50	15,354
305	—	—	—	—	5 6 23,31	128 6 23,31	—	6 26,48	-	3,17	15,249	
306	3 6 52,17	—	1 6 53,84	—	—	123 6 52,59	—	7 2,16	-	9,57	15,242	
307	3 36 51,03	—	3 36 56,37	—	—	75 36 55,20	—	36 51,89	+	3,31	15,234	
308	5 55 52,31	—	—	—	12 55 51,75	37 55 51,86	55 54,30	55 57,78	- 2,44	-	5,92	15,227
309	5 15 19,65	—	—	—	—	115 15 19,65	—	15 13,42	+	6,23	15,222	
310	5 19 22,34	—	—	—	—	158 19 22,34	—	19 15,49	+	6,85	15,187	
311	5 42 2,77	—	5 42 0,69	—	—	111 42 1,73	41 59,00	41 56,62	+ 2,73	+	5,11	15,165
312	5 12 24,83	—	—	—	—	74 12 24,83	—	12 22,52	+	2,31	15,142	
313	3 21 14,04	—	2 21 14,41	—	—	72 21 14,19	—	21 9,12	+	5,07	14,996	
314	5 39 6,05	—	—	—	—	72 39 6,05	39 5,30	39 5,78	+ 0,75	+	0,27	14,961
315	5 17 55,56	—	—	—	—	82 17 55,56	—	17 54,51	+	1,05	14,944	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^b January 1, 1832			A. S. Catal.	Difference from		Annual Precession	
			No. 1831	No. 1832	No. 1833	1832				Green.	A. S.		
			s.	s.	s.	h.	m.	s.		s.	s.		
316	3	3 Eridani	6 13,45	4 13,60	2 13,50	2	48	13,51	13,57	13,61	-0,06	-0,10	+2,917
317	6	47 Arietis	—	—	5 29,25	2	48	29,27	—	27,87	—	+1,40	3,394
318	7	Arietis	—	3 16,53	3 16,18	2	49	16,37	—	15,61	—	+0,76	3,412
319	5	48 Arietis	6 37,30	6 37,30	5 37,27	2	49	37,30	37,27	36,98	+0,08	+0,32	3,408
320	5.6	4 Eridani	—	—	6 55,75	2	49	55,73	—	55,56	—	+0,17	2,656
321	5.6	6 Eridani	—	—	4 37,53	2	50	37,51	—	37,41	—	+0,10	2,660
322	5.6	91 Ceti	—	—	3 43,36	2	50	43,36	—	42,56	—	+0,80	3,199
323	6	5 Eridani	—	—	3 12,62	2	51	12,62	—	12,57	—	+0,05	3,018
324	5	Horologii	—	—	—	2	51	—	—	18,98	—	—	1,222
325	4.5	Eridani	—	5 53,78	—	2	51	53,68	—	53,86	—	-0,18	2,277
326	6	49 Arietis	—	—	3 1,30	2	52	1,33	—	1,07	—	+0,26	3,508
327	6	Fornacis	—	—	3 12,37	2	52	12,35	—	12,17	—	+0,18	2,624
328	7	51 Arietis	—	—	2 29,57	2	52	29,60	—	28,48	—	+1,12	3,512
329	4	23 Persei	6 40,78	1 40,89	—	2	52	40,80	40,60	40,28	+0,20	+0,52	4,273
330	5.6	8 Eridani	1 54,81	—	5 54,88	2	52	54,86	—	54,64	—	+0,22	2,934
331	5	Persei	—	5 58,33	—	2	52	58,39	—	58,41	—	-0,02	4,426
332	2.3	92 Ceti	9 30,35	—	14 30,36	2	53	30,36	30,32	30,15	+0,04	+0,21	3,123
333	6	Fornacis	—	—	4 23,27	2	54	23,24	—	22,36	—	+0,88	2,563
334	4	25 Persei	5 26,21	2 26,28	—	2	54	26,26	26,27	25,77	-0,01	+0,49	3,792
335	5	9 Eridani	3 27,73	6 27,86	—	2	54	27,82	—	27,60	—	+0,22	2,933
336	4	11 Eridani	—	3 59,33	3 59,24	2	54	59,26	—	59,15	—	+0,11	2,651
337	6	Fornacis	—	—	—	2	55	—	—	0,29	—	—	2,663
338	6.7	52 Arietis	—	3 36,59	1 36,72	2	55	36,65	—	35,61	—	+1,04	3,492
339	5	10 Eridani	—	3 1,80	2 1,81	2	56	1,80	—	1,43	—	+0,37	2,933
340	4	Persei	—	5 59,42	1 59,28	2	56	59,41	58,76	54,63	+0,65	+4,78	4,138
341	2.3	26 Persei	5 16,03	3 15,87	1 16,10	2	57	16,00	15,97	15,43	+0,03	+0,57	3,659
342	6	53 Arietis	—	—	5 58,81	2	57	58,83	—	58,61	—	+0,22	3,358
343	5	27 Persei	6 11,99	4 12,09	—	2	58	12,01	—	11,53	—	+0,48	3,979
344	6.7	54 Arietis	—	—	5 50,71	2	58	50,73	—	49,85	—	+0,88	3,376
345	7	Arietis	—	—	6 43,01	2	59	43,03	—	42,59	—	+0,44	3,413
346	6.7	Arietis	—	1 29,77	4 29,51	3	0	29,58	—	25,30	—	+4,28	3,535
347	6	Fornacis	—	4 40,39	—	3	0	40,37	—	39,82	—	+0,55	2,554
348	4	57 Arietis	6 2,16	6 2,18	6 2,16	3	2	2,18	2,09	2,03	+0,09	+0,15	3,398
349	5	Hydri	—	5 58,68	—	3	1	58,57	—	57,45	—	+1,12	0,034
350	6.7	Ceti	—	—	6 9,12	3	2	9,14	—	8,69	—	+0,25	3,278
351	6	56 Arietis	—	2 14,82	3 14,94	3	2	14,91	—	13,93	—	+0,98	3,546
352	5.6	94 Ceti	—	6 12,36	—	3	4	12,36	—	12,08	—	+0,28	3,037
353	3.4	12 Eridani	6 56,20	2 56,40	2 56,32	3	4	56,26	56,41	56,59	-0,15	-0,33	2,561
354	5	58 Arietis	2 15,50	12 15,51	9 15,60	3	5	15,56	15,55	15,01	+0,01	+0,55	3,427
355	4	13 Eridani	5 40,67	7 40,68	3 40,59	3	7	40,66	40,67	40,48	-0,01	+0,18	2,906
356	6	14 Eridani	—	—	5 27,89	3	8	27,89	—	27,05	—	+0,84	2,899
357	5.6	95 Ceti	—	5 47,13	—	3	9	47,13	—	47,17	—	-0,04	3,041
358	6.7	59 Arietis	—	—	5 54,84	3	9	54,86	—	54,25	—	+0,61	3,568
359	5.6	Tauri	—	4 11,58	1 11,59	3	10	11,60	—	10,71	—	+0,89	3,601
360	6	96 Ceti	—	—	6 33,61	3	10	33,61	—	33,13	—	+0,48	3,115

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.	
	1831		1832		1833					Green.	A. S. C.		
	No.	"	No.	"	No.	"				"	"		
316	4	34 19,53	5	34 17,49	—	99	34 18,40	34	10,68	+2,24	+ 7,72	14,887	
317	—	—	5	0 41,40	—	70	0 41,40	0	30,76	—	+ 10,64	14,874	
318	—	—	4	3 32,35	—	69	3 32,35	—	—	—	—	14,827	
319	5	20 11,08	5	20 13,26	—	69	20 12,62	20	9,02	-0,47	+ 3,60	14,806	
320	—	—	5	32 28,15	—	114	32 28,15	32	20,66	—	+ 7,49	14,786	
321	5	17 11,18	—	—	—	114	17 11,18	17	7,08	—	+ 4,10	14,745	
322	2	45 59,08	—	—	—	81	45 59,08	46	0,73	—	- 1,65	14,741	
323	—	—	—	—	5	8 16,66	98	8 16,66	8	16,04	+ 0,62	—	14,711
324	—	—	—	—	—	158	—	48	0,39	—	—	14,701	
325	5	58 53,13	—	—	—	130	58 53,13	58	54,04	—	- 0,91	14,669	
326	—	—	3	12 28,53	—	64	12 28,53	12	28,36	—	+ 0,17	14,664	
327	—	—	—	—	5	57 6,41	115	57 6,41	57	7,46	—	- 1,05	14,651
328	—	—	—	—	5	3 3,00	64	3 3,00	3	6,14	—	- 3,14	14,637
329	5	9 32,30	5	9 31,23	—	37	9 31,76	9	31,64	+0,69	+ 0,12	14,627	
330	2	19 46,07	5	19 45,47	3	19 45,90	98	19 45,72	19	44,09	—	+ 1,68	14,610
331	5	57 39,88	—	—	—	33	57 39,88	57	43,70	—	- 3,82	14,609	
332	15	34 25,68	9	34 27,48	4	34 26,91	86	34 26,43	34	25,80	-1,27	+ 0,63	14,575
333	—	—	—	—	6	44 17,10	118	44 17,10	44	7,64	—	+ 9,46	14,521
334	4	49 2,19	—	—	—	51	49 2,19	49	0,11	+2,08	+ 2,59	14,520	
335	3	21 1,54	2	20 58,70	1	20 59,86	98	21 0,31	21	4,58	—	- 4,27	14,517
336	2	17 14,19	—	—	3	17 13,57	114	17 13,82	17	14,71	-0,89	+ 3,77	14,484
337	—	—	—	—	2	38 21,58	113	38 21,58	38	46,58	—	- 25,00	14,483
338	—	—	5	24 16,71	—	65	24 16,71	24	14,28	—	+ 2,43	14,449	
339	1	15 44,91	2	15 42,95	—	98	15 43,60	15	44,87	—	- 1,27	14,422	
340	5	2 7,51	5	2 8,34	—	41	2 7,92	2	8,21	-0,29	- 0,31	14,370	
341	12	41 55,98	5	41 55,43	—	49	41 55,79	41	52,93	+2,86	+ 14,95	14,349	
342	—	—	4	46 25,80	—	72	46 25,86	46	26,30	—	- 0,44	14,304	
343	5	47 8,51	3	47 8,78	—	45	47 8,61	47	9,76	—	- 1,15	14,292	
344	—	—	—	—	5	51 19,66	71	51 19,66	51	16,84	—	+ 2,82	14,251
345	2	53 13,11	—	—	4	53 16,13	69	53 15,12	53	12,68	—	+ 2,44	14,197
346	—	—	—	—	5	45 3,81	63	45 3,81	—	—	—	—	14,153
347	4	28 42,21	—	—	—	118	28 42,21	28	48,92	—	- 6,71	14,136	
348	4	54 52,34	6	54 53,70	6	54 52,29	70	54 52,84	54	51,33	+1,51	+ 2,91	14,053
349	—	—	—	—	1	33 29,13	162	33 29,13	33	19,60	—	+ 9,53	14,051
350	2	35 37,37	3	35 37,10	—	77	35 37,21	35	39,93	—	- 2,77	14,046	
351	—	—	1	22 53,39	4	22 51,09	63	22 53,95	22	54,13	—	- 0,18	14,041
352	5	49 48,04	1	49 44,34	—	91	49 47,42	49	46,48	—	+ 0,94	13,916	
353	5	39 15,73	4	39 14,95	—	119	39 15,38	39	8,29	+2,58	+ 7,09	14,689	
354	5	35 2,61	9	35 3,53	5	35 4,49	69	35 3,54	35	1,70	+1,84	+ 4,43	13,851
355	5	26 58,06	5	26 55,20	—	99	26 56,63	26	55,61	+1,02	+ 3,46	13,696	
356	5	46 55,14	—	—	—	99	46 55,14	46	51,90	—	+ 3,24	13,646	
357	5	32 49,39	—	—	—	91	32 49,39	32	51,69	—	- 2,30	13,561	
358	2	33 28,00	4	33 29,19	—	63	33 28,79	32	29,33	—	- 0,54	13,554	
359	—	—	2	33 58,83	3	33 59,64	61	33 59,32	33	57,56	—	+ 1,76	13,537
360	—	—	—	—	5	15 6,41	87	15 6,41	15	7,94	—	- 1,53	13,512

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831 No. 1832 No. 1833						Green.	A. S.	
			s	s	s						
361	5.6	15 Eridani	—	—	5 56,66	3 10 56,64	—	56,62	+0,02	+2,646	
362	6	61 Arietis	—	—	3 32,39	3 11 32,41	—	32,42	-0,01	3,439	
363	3.4	16 Eridani	1 2,81	6 2,80	1 2,70	3 12 2,78	2,95	2,51	-0,17	+0,27	2,659
364	6	62 Arietis	—	—	5 7,29	3 12 7,81	—	7,16	+0,65	3,574	
365	2.3	33 Persei	8 22,05	5 22,11	2 21,87	3 12 22,05	22,14	21,77	-0,09	+0,28	4,221
366	6	97 Ceti	—	—	5 20,00	3 12 20,00	—	19,97	+0,03	3,121	
367	7	63 Arietis	—	—	5 6,09	3 13 6,11	5,90	5,24	+0,21	+0,87	3,433
368	4	Eridani	6 13,19	4 13,31	2 13,72	3 13 13,30	—	4,55	—	2,114	
369	5.6	64 Arietis	—	—	5 24,09	3 14 24,11	—	24,04	+0,07	3,517	
370	6	65 Arietis	—	14 45,66	—	3 14 45,68	—	44,87	+0,81	3,437	
371	4	Camelopard	6 31,97	—	—	3 15 31,97	31,50	31,53	+0,47	+0,44	4,765
372	4.5	1 Tauri	2 47,02	5 46,72	—	3 15 46,80	47,00	46,66	-0,20	+0,14	3,217
373	4.5	Camelopard	—	4 33,55	—	3 16 33,55	33,51	33,23	+0,04	+0,32	4,702
374	7	Tauri	—	—	4 28,59	3 17 28,55	—	28,40	+0,15	3,400	
375	4	2 Tauri	—	6 4,49	10 4,53	3 18 4,52	4,51	4,15	+0,01	+0,37	3,291
376	6.7	66 Arietis	—	—	5 33,29	3 18 33,31	—	33,13	+0,18	3,484	
377	5	35 Persei	6 45,99	—	—	3 18 45,99	—	46,60	-0,61	4,178	
378	6	4 Tauri	—	5 14,15	—	3 21 14,16	—	13,91	+0,25	3,263	
379	5.6	5 Tauri	—	—	15 36,69	3 21 36,71	36,66	36,18	+0,05	+0,53	3,293
380	4.5	17 Eridani	6 17,23	9 17,28	2 17,35	3 22 17,27	17,31	17,18	-0,04	+0,09	2,966
381	6.7	6 Tauri	—	—	5 31,18	3 23 31,20	—	30,71	+0,49	3,228	
382	5	Eridani	3 14,40	5 14,56	—	3 24 14,48	—	14,42	+0,06	2,134	
383	6	7 Tauri	—	3 30,41	3 30,45	3 24 30,45	—	29,60	+0,85	3,529	
384	5	37 Persei	5 35,35	6 35,37	—	3 24 35,39	—	34,59	+0,80	4,208	
385	4	18 Eridani	—	7 1,39	6 1,27	3 25 1,32	1,49	1,22	-0,17	+0,10	2,884
386	4	19 Eridani	6 22,16	2 22,28	4 22,41	3 26 22,26	22,26	21,97	0,00	+0,29	2,641
387	6	9 Tauri	—	5 6,10	—	3 27 6,11	—	6,08	+0,03	3,506	
388	5	10 Tauri	4 18,51	7 18,34	—	3 28 18,40	—	18,42	-0,02	3,065	
389	6	20 Eridani	—	—	6 38,36	3 28 38,34	—	38,08	+0,26	2,725	
390	7	Tauri	—	4 56,25	1 56,37	3 29 56,28	—	55,86	+0,42	3,371	
391	6	21 Eridani	—	3 43,96	3 43,34	3 30 43,90	—	43,35	+0,55	2,953	
392	6	11 Tauri	—	5 45,06	—	3 30 45,08	—	44,78	+0,30	3,558	
393	3.4	39 Persei	6 59,87	6 59,93	—	3 30 59,92	59,84	59,76	+0,08	+0,16	4,217
394	5	Eridani	5 4,14	1 4,22	—	3 31 4,15	—	4,21	-0,06	2,149	
395	6	12 Tauri	—	—	4 6,72	3 31 6,77	—	6,50	+0,27	3,114	
396	6	Fornacis	—	—	5 48,63	3 31 48,61	—	48,62	-0,01	2,489	
397	5.6	22 Eridani	—	—	5 19,71	3 32 19,71	—	19,34	+0,37	2,960	
398	6.7	13 Tauri	—	3 38,25	1 38,34	3 32 38,29	—	38,01	+0,28	3,439	
399	4.5	41 Persei	5 48,50	4 48,57	—	3 33 48,54	—	47,49	+1,05	4,035	
400	4	Persei	—	5 48,04	—	3 33 48,06	48,43	48,13	-0,37	-0,07	3,732
401	7	14 Tauri	—	—	3 5 19	3 34 5,22	—	4,58	+0,64	3,440	
402	5.6	16 (Pleiadum g)	—	—	4 49,38	3 34 49,90	50,11	49,48	-0,21	+0,42	3,542
403	4.5	17 (Pleiadum b)	3 35,01	6 54,79	—	3 34 54,87	54,75	54,59	+0,12	+0,28	3,538
404	7	18 (Pleiadum m)	—	—	5 9,21	3 35 9,24	—	8,60	+0,64	3,555	
405	5	19 (Pleiadum c)	6 13,42	1 13,71	—	3 35 13,47	13,38	12,87	+0,09	+0,60	3,546

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession	
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.		
	"	"	"				"	"		
361	—	—	5 7 45,57	113 7 45,57	—	7 44,74	+	0,83	13,485	
362	—	—	5 27 52,71	69 27 52,71	—	27 49,71	+	3,00	13,448	
363	5 22 24,07	5 22 24,04	—	112 22 24,05	22 27,35	22 27,58	-3,30	-	3,53	13,414
364	—	—	5 0 5,10	63 0 5,10	—	0 4,27	+	0,83	13,411	
365	29 44 41,29	6 44 40,08	2 44 40,76	40 44 41,08	44 39,84	44 36,33	+1,27	+	4,75	13,397
366	5 56 5,20	—	—	86 56 5,20	—	56 3,53	+	1,67	13,396	
367	—	1 51 52,68	4 51 55,57	69 51 54,99	51 52,61	51 54,14	+2,38	+	0,85	13,348
368	4 42 57,22	5 42 57,83	—	133 42 57,56	—	43 25,88	-	28,32	13,346	
369	—	—	5 52 37,11	65 52 37,11	—	52 30,66	+	6,45	13,262	
370	3 47 56,69	8 47 57,53	—	69 47 57,30	—	47 50,13	+	7,17	13,239	
371	5 39 16,27	5 39 15,23	—	39 39 15,75	39 14,58	39 15,52	+1,17	+	0,23	13,191
372	—	6 34 3,80	5 34 3,75	81 34 3,78	34 4,28	34 2,02	-0,50	+	1,76	13,171
373	5 42 44,86	5 42 43,78	—	31 42 44,32	42 45,88	42 45,71	-1,56	-	1,39	13,123
374	—	1 50 14,59	4 50 13,84	71 50 13,99	—	50 15,09	-	1,10	13,059	
375	5 51 31,83	5 51 31,93	—	80 51 31,88	51 29,75	51 28,47	+2,13	+	3,41	13,019
376	5 46 48,68	—	—	67 46 48,68	—	46 48,85	-	0,17	12,982	
377	5 35 29,89	2 35 28,41	—	42 35 29,47	—	35 33,92	-	4,45	12,974	
378	3 14 46,39	2 14 45,06	—	79 14 45,86	—	14 39,07	+	6,79	12,808	
379	—	2 38 39,31	6 38 38,74	77 38 38,88	38 40,25	38 39,50	-1,37	-	0,62	12,783
380	4 39 25,93	6 39 21,22	—	95 39 24,90	39 23,86	39 22,98	+1,04	+	1,92	12,736
381	5 11 59,93	—	—	81 11 59,93	—	11 58,42	+	1,51	12,654	
382	4 56 25,83	3 56 27,69	—	131 56 26,63	—	56 16,93	+	0,70	12,602	
383	5 6 18,15	—	—	66 6 18,15	—	6 18,40	-	0,31	12,588	
384	5 22 25,27	5 22 25,32	—	42 22 25,30	—	22 30,52	-	5,22	12,584	
385	5 1 58,93	5 1 58,21	—	100 1 58,57	1 55,34	1 55,55	+3,23	+	3,02	12,550
386	5 12 5,07	3 12 2,72	—	112 12 4,19	12 3,74	12 3,07	+0,45	+	1,12	12,458
387	6 21 2,72	—	—	67 21 2,72	—	20 59,48	+	3,24	12,409	
388	8 8 15,91	5 8 16,40	—	90 8 16,19	—	8 8,18	+	7,92	12,325	
389	—	—	—	108 —	—	1 33,81	—	—	12,302	
390	1 0 53,16	4 0 56,50	—	74 0 55,83	—	0 54,88	+	0,95	12,214	
391	—	5 10 12,55	—	96 10 12,55	—	10 8,27	+	4,28	12,158	
392	—	1 13 15,29	4 13 16,26	65 13 16,07	—	13 12,74	+	3,33	12,158	
393	5 45 26,20	5 45 25,59	12 45 25,76	42 45 25,82	45 27,03	45 29,39	-1,21	-	3,57	12,142
394	5 49 48,86	5 49 48,25	—	130 49 48,55	—	49 47,64	+	0,91	12,132	
395	—	—	5 29 41,79	87 29 41,79	—	29 38,41	+	3,38	12,131	
396	—	1 29 43,36	4 29 44,45	118 29 44,23	—	29 47,08	-	2,85	12,081	
397	—	1 45 32,49	4 45 33,15	95 45 33,02	—	45 30,67	+	2,35	12,046	
398	1 50 35,93	3 50 38,14	—	70 50 37,89	—	50 36,18	+	1,71	12,026	
399	5 57 31,42	4 57 29,45	—	47 57 30,55	57 35,23	57 33,68	-4,67	-	3,13	11,946
400	5 15 0,79	5 15 1,64	—	58 15 1,22	—	15 2,49	-	1,27	11,944	
401	—	1 53 18,63	4 52 20,64	70 52 20,24	—	52 16,90	+	3,34	11,924	
402	—	—	3 14 43,54	66 14 43,54	14 44,55	14 42,42	-1,01	+	1,12	11,872
403	4 25 15,65	7 25 16,02	—	66 25 15,89	25 16,45	25 11,26	-0,56	+	4,63	11,866
404	—	—	7 41 42,10	65 41 42,10	—	41 38,41	+	3,69	11,850	
405	4 3 56,95	—	5 3 57,68	66 3 57,36	4 0,15	3 54,45	-3,79	+	2,91	11,845

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^h January 1, 1832		A. S. Catal.	Difference from		Annual Preces- sion	
			No. 1831	No. 1832	No. 1833	s.	h. m. s.		s.	s.		s.
			s.	s.	s.							
406	3.4	28 Eridani δ		5 12,42		3 35 12,41	12,26	12,45	+0,15	-0,04	+2,871	
407	6	Eridani			5 32,78	3 35 32,77		32,48		+0,29	2,858	
408	5	Fornacis δ		5 34,22		3 35 34,19		34,37		-0,18	2,861	
409	5	20 (Pleiadum c)		2 50,71	2 50,46	3 35 50,59		50,25		+0,34	3,545	
410	5	23 (Pleiadum d)		5 22,05		3 36 22,07	22,05	22,06	+0,02	+0,01	3,537	
411	5	Eridani h			5 36,72	3 36 36,74		35,93		+0,81	2,227	
412	6	29 Tauri u^1			5 45,45	3 36 45,45		44,93		+0,52	3,173	
413	7	(Pleiadum)			3 30,38	3 37 30,40		30,17		+0,23	3,546	
414	3	25 Tauri u	3 30,76	1 30,66	1 30,65	3 37 30,73	30,65	30,27	+0,08	+0,46	3,542	
415	5	26 Eridani π		6 12,35		3 38 12,34		12,17		+0,17	2,824	
416	7	Tauri			5 24,79	3 38 24,82		24,73		+0,09	3,523	
417	6	30 Tauri e		5 4,13		3 39 4,15		3,11		+1,04	3,273	
418	5	27 (Pleiadum f)	3 10,98	6 11,17		3 39 11,12		10,72		+0,40	3,543	
419	5.6	28 (Pleiadum h)			5 12,28	3 39 12,30		11,65		+0,65	3,545	
420	6	Fornacis e			5 36,87	3 39 36,83		36,62		+0,21	2,440	
421	5	27 Eridani m^1	6 37,44	3 37,40		3 39 37,42		37,31		+0,11	2,587	
422	7	Tauri		4 3,20	1 3,55	3 40 3,31		3,06		+0,25	3,504	
423	6.7	(Pleiadum)			5 14,08	3 40 14,11		13,34		+0,77	3,580	
424	5	28 Eridani m^2		5 26,50		3 40 26,47		26,96		-0,49	2,571	
425	4	Reticuli β		4 7,20		3 42 7,16		6,56		+0,60	0,668	
426	5	Eridani		6 24,07		3 42 24,06		23,71		+0,35	2,202	
427	6	31 Tauri z^2		5 3,24		3 43 3,24		2,52		+0,72	3,184	
428	5	Eridani g	5 9,94			3 43 9,94		10,50		-0,56	2,244	
429	7	Tauri		5 34,25		3 43 34,25		33,40		+0,85	3,402	
430	3.4	44 Persei ζ	6 35,36	2 35,33	13 35,36	3 43 35,37	35,35	35,07	+0,02	+0,30	3,742	
431	6	30 Eridani			5 24,18	3 44 24,18		23,70		+0,48	2,954	
432	5	32 Eridani		5 51,69		3 45 51,69		51,65		+0,04	3,001	
433	3.4	45 Persei ϵ		12 36,19	5 36,03	3 46 36,19	36,18	36,03	+0,01	+0,16	3,988	
434	5.6	33 Eridani λ			5 34,31	3 47 34,28		33,68		+0,60	2,545	
435	6	32 Tauri			5 57,39	3 47 57,41		56,77		+0,64	3,519	
436	6.7	33 Tauri		5 6,88		3 47 6,90		6,39		+0,51	3,535	
437	5	Eridani i	5 14,82	4 15,00		3 47 14,89		15,03		-0,14	2,278	
438	5	46 Persei ξ	6 5,15	4 5,08		3 48 5,13		4,37		+0,76	3,861	
439	3	Hydri γ		4 57,00	1 57,41	3 49 56,94		55,50		+1,44	-1,068	
440	2.3	34 Eridani γ^1	1 11,89	6 11,71	12 11,76	3 50 11,74	11,73	11,53	+0,01	+0,21	+2,787	
441	6.7	Tauri		5 8,62		3 51 8,63		8,26		+0,37	3,429	
442	7	34 Tauri		4 20,47	1 20,65	3 51 20,51		18,96		+1,55	3,473	
443	4	35 Tauri λ	6 22,97	2 22,95	7 22,82	3 51 22,90	22,88	22,36	+0,02	+0,54	3,309	
444	5	36 Eridani k	4 46,08	1 45,83		3 52 46,03		45,68		+0,35	2,551	
445	5	35 Eridani		6 1,68		3 53 1,68		1,58		+0,10	3,028	
446	5	38 Tauri ν		6 13,64		3 54 13,64		13,60		+0,04	3,178	
447	6.7	36 Tauri		5 19,54		3 54 19,55		19,32		+0,23	3,567	
448	5	37 Tauri A^1	6 46,42	7 46,58	5 46,50	3 54 46,51	46,48	46,12	+0,03	+0,39	3,520	
449	6.7	39 Tauri A^2		2 24,24	3 24,42	3 55 24,37		23,99		+0,38	3,519	
450	5	Reticuli δ		5 6,41		3 56 6,31		5,87		+0,44	0,925	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.			
	1831		1833				Green.	A. S. C.				
	No.	"	No.				"	"		"		
406	5 20	17,16	5 20	18,50	100 20	17,83	20 19,43	20 13,19	-1,60	+ 4,64	-11,243	
407	—	—	—	—	101 1	23,25	—	1 21,53	+	1,72	11,820	
408	5 28	44,61	5 28	45,78	122 28	45,19	—	28 44,09	+	1,10	11,817	
409	1 9	46,07	—	—	4 9	49,13	—	66 9 48,72	+	2,02	11,801	
410	—	—	—	—	7 34	52,10	66 34 52,10	34 54,15	34 42,51	-2,05	+ 9,59	11,763
411	—	—	5 50	49,97	127 50	49,97	—	50 51,80	—	1,83	11,744	
412	—	—	1 28	58,45	4 28	55,30	—	28 56,18	—	0,25	11,735	
413	—	—	4 14	13,97	66 14	13,97	—	14 12,24	+	1,73	11,683	
414	5 25	12,74	4 25	13,43	66 25	13,05	25-15,35	25 14,50	-2,30	- 1,45	11,682	
415	4 37	59,78	—	—	1 37	59,01	—	38 0,89	—	1,26	11,631	
416	—	—	—	—	5 6	6,06	—	6 2,32	+	3,74	11,618	
417	2	—	—	—	3 22	47,91	—	22 44,89	+	3,02	11,572	
418	4 27	59,19	5 27	59,24	—	—	—	28 0,02	—	0,80	11,563	
419	—	—	5 22	58,05	—	—	—	22 59,47	—	1,42	11,562	
420	—	—	—	—	5 51	56,75	119 51 56,75	51 54,52	+	2,23	11,530	
421	5 44	59,81	—	—	113 44	59,81	—	45 3,52	—	3,71	10,939*	
422	—	—	—	—	5 16	23,32	—	16 22,08	+	1,24	11,501	
423	1 56	7,92	—	—	4 56	8,02	—	55 59,42	+	8,58	11,489	
424	5 24	1,21	—	—	114 24	1,21	—	23 57,14	+	4,07	11,470	
425	—	—	5 20	13,78	—	—	—	20 20,21	—	6,43	11,346	
426	—	—	5 8	18,42	—	—	—	8 13,25	+	5,17	11,329	
427	—	—	5 58	30,49	—	—	—	58 32,01	—	1,52	11,285	
428	5 42	45,84	5 42	48,38	—	—	—	42 45,02	+	2,09	11,273	
429	—	—	—	—	7 10	45,05	—	73 10 45,05	—	0,14	11,248	
430	5 37	17,80	7 37	18,32	5 37	17,91	58 37 18,04	37 22,78	-1,32	- 4,74	11,247	
431	—	—	5 52	7,20	—	—	—	52 3,52	+	3,68	11,186	
432	5 27	23,63	5 27	24,47	—	—	—	27 24,37	—	0,32	11,080	
433	5 29	2,46	5 29	2,59	5 29	3,08	50 29 2,70	29 0,65	+2,05	+ 3,16	11,028	
434	—	—	5 6	55,56	—	—	—	6 51,98	+	3,58	11,027	
435	—	—	4 0	42,95	1 0	42,60	68 0 42,88	0 42,00	+	0,88	11,002	
436	—	—	—	—	5 19	7,75	67 19 7,75	19 7,75	—	0,00	10,990	
437	5 14	4,04	5 14	4,88	—	—	—	13 59,48	+	4,98	10,976	
438	5 42	2,65	5 41	59,35	—	—	—	41 58,44	+	2,56	10,920	
439	—	—	—	—	1 45	40,02	164 45 40,02	45 11,90	+	28,12	10,772	
440	7 59	28,25	5 59	28,25	5 59	28,98	103 59 28,49	59 30,53	-2,04	+ 0,11	10,762	
441	5 17	9,35	5 17	9,99	—	—	—	17 8,89	+	1,08	10,693	
442	—	—	5 16	44,03	—	—	—	16 40,30	+	3,73	10,680	
443	5 59	21,69	6 59	23,29	6 59	22,52	77 59 22,54	59 25,63	-3,09	+ 2,73	10,676	
444	4 29	51,08	5 29	53,40	—	—	—	29 47,30	+	5,07	10,571	
445	5 1	36,04	5 1	35,88	—	—	—	1 35,21	+	0,75	10,552	
446	5 28	57,68	5 29	0,99	—	—	—	28 54,85	+	4,48	10,463	
447	—	—	4 21	46,34	—	—	—	21 46,73	—	0,30	10,457	
448	5 22	58,15	8 22	59,07	3 22	59,64	68 22 58,89	23 2,68	-3,79	+ 5,15	10,424	
449	5 27	0,20	2 27	2,13	—	—	—	27 1,46	—	0,71	10,376	
450	5 52	34,78	5 52	35,37	—	—	—	52 18,39	+	16,68	10,318	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833				Green.	A. S.	
451	6	41 Tauri			5 18,99	3 56 19,01		18,96		+0,05	3,655
452	5	48 Persei	c	4 29,80	2 29,99	3 56 29,88		29,18		+0,70	4,308
453	5.6	42 Tauri	ψ			5 38,16	3 56 38,18	37,92		+0,26	3,692
454	6	Tauri			5 22,75	1 22,83	3 58 22,78	22,46		+0,32	3,418
455	5	Reticuli	γ	5 29,41	3 29,49		3 58 29,43	27,32		+2,11	0,841
456	6	Eridani	A		4 42,36	2 41,97	3 58 42,22	41,59		+0,63	2,452
457	6	43 Tauri	ω ¹			5 23,55	3 59 23,57	23,29		+0,28	3,469
458	6.7	Tauri				6 39,39	3 59 39,41	38,83		+0,58	3,334
459	6.7	44 Tauri	P			6 37,02	4 0 37,04	36,49		+0,55	3,634
460	5.6	37 Eridani			6 11,20		4 2 11,19	10,89		+0,30	2,918
461	6	45 Tauri			5 24,22		4 2 24,23	23,82		+0,41	3,171
462	4.5	51 Persei	μ	5 35,48	5 35,49	5 35,15	4 2 35,41	35,34	-0,04	+0,07	4,360
463	7	Tauri			4 54,27	2 54,52	4 2 54,36	54,11		+0,25	3,538
464	4.5	38 Eridani	ο	6 40,16	6 40,18	6 40,32	4 3 40,22	39,60	+0,03	+0,62	2,919
465	6	46 Tauri				5 30,86	3 4 30,86	30,36		+0,50	3,217
466	5.6	47 Tauri			1 48,60	4 48,81	4 4 48,77	48,42		+0,35	3,250
467	5	Persei	δ	4 38,47	4 38,54		4 5 38,49	38,02		+0,47	4,459
468	6	48 Tauri		1 14,63	5 14,52		4 6 14,55	14,14		+0,41	3,382
469	5	49 Tauri	μ	1 25,03	5 25,09		4 6 25,09	24,88		+0,21	3,243
470	5	39 Eridani	A		5 24,45		4 6 24,45	24,27		+0,18	2,846
471	5.6	50 Tauri	ω ²		5 25,62		4 7 25,63	25,23		+0,40	3,500
472	5	40 Eridani	d		6 32,51		4 7 32,51	32,18		+0,33	2,757*
473	7	51 Tauri			2 27,40	3 27,48	4 8 27,47	26,98		+0,49	3,525
474	5	Horologii	α	5 26,32	3 26,66		4 8 26,52	25,98		+0,54	1,978
475	6.7	53 Tauri			5 32,54		4 9 32,56	32,03		+0,53	3,516
476	6.7	56 Tauri				5 40,75	4 9 40,77	40,29		+0,48	3,531
477	6	52 Tauri	φ			5 2,08	4 10 2,10	2,08	+0,02	-0,02	3,670
478	3.4	54 Tauri	γ	6 14,61	11 14,50		4 10 14,55	14,42	+0,13	+0,32	3,390
479	6	57 Tauri	η ¹			5 30,69	4 10 30,71	30,60		+0,11	3,355
480	6	58 Tauri	δ ²		2 5,31	3 5,28	4 11 5,31	4,42		+0,89	3,379
481	6.7	Tauri				5 26,27	4 11 26,29	26,14		+0,15	3,352
482	3.4	41 Eridani	X	5 32,38	6 32,55		4 11 32,47	32,42	+0,05	+0,84	2,259
483	4	Doradus	γ	2 38,17	2 38,14		4 11 38,14	37,10		+1,04	1,550
484	6	59 Tauri	κ			5 22,15	4 12 22,17	22,05	+0,12	+0,36	3,629
485	3.4	Reticuli	κ		5 17,04		4 12 16,96	16,50		+0,46	0,741
486	7	Tauri				5 30,53	4 12 30,55	30,10		+0,45	3,511
487	7	60 Tauri	η ³			3 36,00	4 12 36,03	35,37		+0,66	3,358
488	4	61 Tauri	δ ¹	5 15,30	5 15,28	6 15,51	4 13 15,38	15,31	+0,07	+0,50	3,436
489	5	Reticuli	ε		4 35,94		4 13 35,90	35,83		+0,07	1,023
490	6	63 Tauri			4 47,24	1 47,15	4 13 47,22	46,95		+0,27	3,419
491	7	62 Tauri				3 52,62	4 13 52,64	52,71		-0,07	3,598
492	4.5	64 Tauri	δ ²		3 25,21		4 14 25,23	25,15	+0,08	+0,12	3,435
493	6	Eridani	ο ²			3 33,17	4 14 33,13	32,97		+0,16	2,481
494	5.6	66 Tauri	r			3 42,82	4 14 42,85	41,71		+1,14	3,259
495	6	42 Eridani	ξ			5 19,35	4 15 19,35	18,86		+0,49	2,981

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession	
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.		
	"	"	"				"	"		
451	5 51 33,26	4 51 34,15	—	62 51 33,66	—	51 34,49	—	0,83	10,308	
452	5 44 39,34	5 44 37,56	—	42 44 38,45	—	44 43,60	—	5,15	10,297	
453	—	5 27 36,04	—	61 27 36,04	—	27 30,89	+	5,15	10,285	
454	—	1 6 56,47	4 6 55,29	73 6 55,53	—	6 56,16	—	0,63	10,153	
455	5 37 46,32	5 37 44,07	—	152 37 45,20	—	37 43,82	+	1,38	10,140	
456	—	—	5 6 57,00	118 6 57,00	—	7 2,85	—	5,65	10,126	
457	—	—	5 50 31,44	70 50 31,44	—	50 21,34	+	10,10	10,076	
458	3 3 10,56	1 3 11,53	—	77 3 10,80	—	3 9,69	+	1,11	10,056	
459	5 57 50,87	—	—	63 57 50,87	—	57 55,58	—	4,71	9,984	
460	5 22 7,81	6 22 6,74	—	97 22 7,24	—	22 6,27	+	0,97	9,863	
461	—	5 55 19,34	—	84 55 19,34	—	55 14,22	+	5,12	9,847	
462	5 1 36,44	5 1 35,84	5 1 36,19	42 1 36,16	1 34,82	1 38,39	+1,34	—	2,23	9,836
463	1 2 27,21	4 2 29,23	—	68 2 28,83	—	1 30,55	—	1,72	9,810	
464	5 16 54,00	5 16 52,35	—	97 16 53,17	16 54,52	16 52,82	-1,35	+	0,35	9,750
465	—	—	5 43 9,56	82 43 9,56	—	43 8,29	+	1,27	9,686	
466	—	5 10 8,17	—	81 10 8,17	—	10 5,03	+	3,14	9,663	
467	5 7 40,57	5 7 41,94	—	40 7 41,25	—	7 36,12	+	5,13	9,603	
468	—	5 1 38,54	—	75 1 38,54	—	1 26,93	+	11,61	9,554	
469	4 32 5,77	4 32 5,22	—	81 32 5,50	—	32 4,14	+	1,36	9,540	
470	5 40 46,58	5 40 46,97	—	100 40 46,77	—	40 42,05	+	4,72	9,540	
471	5 50 33,53	5 50 34,37	—	69 50 33,95	—	50 27,22	+	6,73	9,463	
472	5 55 7,48	5 55 8,34	—	97 55 7,91	—	55 6,70	+	1,21	5,852*	
473	5 50 16,05	—	—	68 50 16,05	—	50 16,58	—	0,53	9,384	
474	5 42 41,21	5 42 44,10	—	132 42 42,65	—	42 38,14	+	4,51	9,381	
475	1 16 14,17	4 16 15,69	—	69 16 15,39	—	16 15,40	—	0,01	9,300	
476	—	3 38 18,35	3 38 18,57	68 38 18,46	—	—	—	—	9,289	
477	—	—	5 3 28,41	63 3 28,41	3 30,92	3 30,53	-2,51	—	2,12	9,261
478	5 47 7,80	11 47 8,57	3 47 9,15	74 47 8,54	47 4,78	47 4,92	+3,76	+	3,62	9,245
479	—	—	5 22 36,55	76 22 36,55	—	22 27,82	+	8,73	9,224	
480	—	7 18 53,33	—	75 18 53,33	—	18 45,85	+	7,48	9,180	
481	3 32 36,71	2 32 39,62	—	76 32 37,87	—	32 34,43	+	3,44	9,152	
482	5 12 46,93	5 12 47,03	—	124 12 46,98	12 47,90	12 46,64	-0,92	+	0,34	9,142
483	5 54 49,95	5 54 49,31	—	141 54 49,63	—	55 9,83	—	20,20	9,133	
484	—	4 46 31,29	1 46 30,96	64 46 31,22	46 27,95	46 26,68	+3,27	+	4,54	9,080
485	4 53 42,89	5 53 42,76	—	152 53 42,82	—	53 28,30	+	14,52	9,080	
486	—	—	5 35 3,55	69 35 3,55	—	34 58,32	+	5,23	9,069	
487	—	—	5 19 35,63	76 19 35,63	—	19 31,27	+	4,36	9,062	
488	5 51 28,24	5 51 28,07	—	72 51 28,15	51 30,36	51 24,03	-2,21	+	4,12	9,011
489	5 42 30,66	5 42 31,93	—	149 42 31,30	—	42 30,11	+	1,19	8,977	
490	1 37 16,99	—	6 37 16,85	73 37 16,44	—	37 12,61	+	3,83	8,969	
491	—	—	5 5 50,57	66 5 50,57	—	5 50,42	+	0,15	8,962	
492	5 57 4,21	5 57 6,25	—	72 57 5,23	57 7,96	57 3,70	-2,73	+	1,53	8,919
493	—	—	5 7 43,66	116 7 43,66	—	7 40,52	+	3,14	8,906	
494	—	1 56 15,58	4 56 13,40	80 56 13,84	—	56 10,95	+	2,89	8,897	
495	—	—	5 8 23,83	94 8 23,83	—	8 21,31	+	2,52	8,848	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names:		Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^h January 1, 1832		A. S. Catal.	Difference from		Annual Preces- sion
				No. 1831	No. 1832	No. 1833	Green.			A. S.		
				s.	s.	s.	h. m.	s.		s.	s.	
496	5.6	65 Tauri	α^1	—	1 21,78	—	4 15 21,80	21,99	21,51	-0,19	+0,29	+3,550
497	6.7	Tauri		—	—	5 25,38	4 15 25,41	—	25,10	—	+0,31	3,790
498	6.7	67 Tauri	α^2	—	4 25,21	5 25,18	4 15 25,21	25,22	25,01	-0,01	+0,20	3,548
499	5	68 Tauri	β^3	7 46,92	—	5 46,69	4 15 46,84	46,78	46,09	+0,06	+0,75	3,447
500	5	Reticuli	θ	—	4 48,78	—	4 15 48,72	—	46,89	—	+1,88	0,643
501	7	70 Tauri		—	—	5 2,61	4 16 2,63	—	2,02	—	+0,61	3,402
502	5	69 Tauri	ν^1	—	5 15,88	—	4 16 15,89	15,66	15,27	—	+0,62	3,564
503	5.6	71 Tauri		—	—	4 47,04	4 16 47,04	—	46,02	—	+1,02	3,395
504	5	73 Tauri	π	6 7,35	6 7,53	1 7,15	4 17 7,40	—	7,19	—	+0,21	3,375
505	6	72 Tauri	ν^2	—	—	5 15,29	4 17 15,32	—	14,93	—	+0,39	3,569
506	4.5	43 Eridani		—	3 43,82	—	4 17 43,80	—	43,54	—	+0,26	2,242
507	4	74 Tauri	ϵ	3 48,93	12 49,02	6 48,80	4 18 48,93	48,93	48,63	0,00	+0,30	3,479
508	6	75 Tauri		—	5 50,65	—	4 18 50,65	—	50,69	—	-0,04	3,414
509	7	76 Tauri		—	1 52,94	4 52,93	4 18 52,95	—	52,08	—	+0,87	3,377
510	5	77 Tauri	θ^1	6 59,14	—	—	4 18 59,14	59,13	58,68	+0,01	+0,46	3,405
511	5.6	78 Tauri	θ^2	2 4,76	—	6 4,67	4 19 4,71	4,76	4,67	-0,05	+0,04	3,403
512	6	79 Tauri	b	—	—	4 25,98	4 19 25,99	—	25,59	—	+0,40	3,340
513	5.6	44 Eridani	k^1	—	—	5 51,47	4 19 51,47	—	51,09	—	+0,38	3,089
514	5	Reticuli	η	—	4 5,94	—	4 20 5,88	—	5,48	—	+0,40	0,608
515	6	80 Tauri		—	—	11 34,38	4 20 34,40	—	34,01	—	+0,39	3,309
516	5.6	Tauri		—	—	3 57,23	4 20 57,25	—	56,89	—	+0,36	3,412
517	5.6	81 Tauri		—	—	2 4,49	4 21 4,51	—	4,18	—	+0,33	3,400
518	6	83 Tauri		—	6 10,49	—	4 21 10,49	—	10,01	—	+0,48	3,356
519	7	84 Tauri		—	—	5 35,68	4 21 35,70	—	35,27	—	+0,43	3,387
520	6	85 Tauri		—	3 16,64	2 16,50	4 22 16,59	—	16,03	—	+0,56	3,405
521	6	45 Eridani	k^2	—	4 17,28	1 17,35	4 23 17,29	—	16,92	—	+0,37	3,059
522	7	Tauri		—	—	6 8,00	4 24 8,03	—	7,50	—	+0,53	3,734
523	5	86 Tauri	p	6 19,64	6 19,31	—	4 24 19,48	—	19,05	—	+0,43	3,383
524	6	46 Eridani		—	—	6 43,88	4 25 43,88	—	43,47	—	+0,41	2,915
525	5	Cæli Scalp	δ	6 41,39	6 41,78	—	4 25 41,57	—	41,33	—	+0,24	1,830
526	6	Eridani		—	—	5 4,04	4 26 4,04	—	3,73	—	+0,31	2,913
527	5	47 Eridani		6 6,37	6 6,51	—	4 26 6,54	—	6,24	—	+0,30	2,883
528	1	87 Tauri	α	12 17,35	23 17,28	49 17,32	4 26 17,31	17,36	16,94	-0,05	+0,37	3,423
529	5	88 Tauri	d	4 25,76	—	—	4 26 25,76	—	25,47	—	+0,29	3,280
530	4	48 Eridani	ν^2	2 56,00	7 55,80	—	4 27 55,85	55,71	55,49	+0,14	+0,36	2,958
531	7	89 Tauri		—	—	5 33,07	4 28 33,09	—	32,70	—	+0,39	3,414
532	6	49 Eridani	k^3	—	—	4 34,60	4 28 34,60	—	34,21	—	+0,39	3,082
533	5	90 Tauri	c^1	2 46,58	3 46,63	—	4 28 46,62	—	46,35	—	+0,27	3,333
534	3	52 Eridani	ν^2	—	6 1,48	—	4 29 1,46	1,47	1,42	-0,01	+0,04	2,330
535	5.6	51 Eridani	c	—	—	4 9,13	4 29 9,13	—	8,86	—	+0,27	3,007
536	5.6	91 Tauri	σ^1	—	—	4 34,13	4 29 34,15	—	33,98	—	+0,17	3,409
537	5.6	92 Tauri	σ^2	—	—	6 40,36	4 29 40,38	—	40,54	—	-0,16	3,412
538	3	Doradus	α	—	3 22,78	3 22,92	4 30 22,78	—	22,04	—	+0,74	1,278
539	4	53 Eridani		—	6 29,43	—	4 30 29,42	29,53	29,16	-0,11	+0,26	2,745
540	5	93 Tauri	c^2	6 42,62	6 42,71	—	4 30 42,67	—	42,31	—	+0,36	3,327

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	1831		1832				Green.	A. S. C.	
	No.	"	No.						
496	—	—	5 5 51,08	68 5 51,08	5 51,78	5 51,28	-0,70	- 0,20	8,846
497	—	—	56 53,10	58 56 53,10	—	56 49,12	+	3,98	8,842
498	1 11 23,85	—	5 11 26,84	68 11 26,34	11 28,98	11 27,55	-2,64	- 1,21	8,841
499	4 27 46,92	5 27 46,25	1 27 46,42	72 27 46,54	27 48,44	27 46,14	-1,90	+ 0,40	8,813
500	5 39 46,96	5 39 45,89	1 —	153 39 46,93	—	39 47,53	—	0,60	8,805
501	—	5 27 1,94	—	74 27 1,94	—	26 54,75	+	7,19	8,792
502	5 34 25,64	4 34 24,17	—	67 34 24,88	—	34 28,34	—	3,46	8,775
503	—	—	5 46 15,33	74 46 15,33	—	46 10,02	+	5,31	8,735
504	5 40 25,26	5 40 24,95	—	75 40 25,10	—	40 20,37	+	4,78	8,707
505	—	—	5 23 21,67	67 23 21,67	—	23 22,29	—	0,62	8,697
506	5 24 42,84	5 24 42,46	—	124 24 42,65	24 40,45	24 42,16	+ 2,20	+ 0,49	8,656
507	5 11 56,71	8 11 57,79	—	71 11 57,37	11 57,42	11 55,70	-0,05	+ 1,67	8,574
508	—	6 1 21,56	—	74 1 21,56	—	1 21,90	—	0,34	8,571
509	—	—	5 38 25,83	75 38 25,83	—	38 24,49	+	1,34	8,569
510	5 25 4,00	5 25 5,09	—	74 25 4,54	25 4,08	24 58,57	+0,46	+ 5,97	8,560
511	—	2 30 33,40	5 30 35,12	74 30 34,63	30 32,08	30 24,21	+2,55	+ 10,42	8,552
512	5 19 53,62	—	—	77 19 53,62	—	19 53,53	+	0,09	8,525
513	—	—	5 59 51,36	88 59 51,36	—	59 47,39	+	3,97	8,490
514	5 47 6,22	5 47 5,98	—	153 47 6,10	—	47 9,25	—	3,15	8,465
515	—	—	—	74 —	—	44 5,31	—	—	8,434
516	2 10 42,17	3 10 41,89	—	74 10 42,00	—	10 38,91	+	3,09	8,404
517	—	3 40 51,48	—	74 40 51,48	—	40 46,03	+	5,45	8,394
518	—	—	5 38 51,72	76 38 51,72	—	38 49,12	+	2,60	8,387
519	—	—	5 15 55,74	75 15 55,74	—	15 48,88	+	6,86	8,353
520	5 31 0,98	4 31 0,13	—	74 31 0,60	—	30 52,41	+	8,19	8,299
521	—	—	5 24 42,94	90 24 42,94	—	24 39,30	+	3,64	8,218
522	—	—	5 23 52,69	61 23 52,69	—	23 49,69	+	3,00	8,152
523	5 30 58,80	5 30 58,87	—	75 30 58,83	—	30 49,59	+	9,24	8,136
524	5 5 51,25	5 5 51,88	—	97 5 51,56	—	5 50,42	+	1,14	8,022
525	5 19 7,80	4 19 7,07	—	135 19 7,48	—	19 10,19	—	2,71	8,022
526	—	—	3 11 36,92	97 11 36,92	—	11 31,84	+	5,08	7,995
527	5 35 15,67	5 35 18,14	—	98 35 16,90	—	35 18,47	—	1,57	7,991
528	44 50 5,41	37 50 6,39	65 50 6,34	73 50 6,08	50 7,77	50 3,73	-1,69	+ 2,35	7,979
529	5 11 25,17	5 11 28,38	—	80 11 26,78	—	11 28,06	—	1,28	7,967
530	—	5 42 4,43	—	93 42 4,43	42 8,59	42 6,61	-4,16	- 2,18	7,845
531	—	—	5 18 38,64	74 18 38,64	—	18 35,19	+	3,45	7,797
532	5 20 54,10	—	—	89 20 54,10	—	20 48,32	+	5,78	7,794
533	5 49 58,74	5 49 59,20	—	77 49 58,97	—	49 58,79	+	0,18	7,778
534	7 54 41,04	5 54 44,69	—	120 54 42,56	54 43,40	54 41,38	-0,84	+ 1,18	7,755
535	—	—	4 48 54,84	92 48 54,84	—	48 52,83	+	2,01	7,747
536	—	—	5 32 20,14	74 32 20,14	—	32 16,15	+	3,99	7,714
537	—	—	5 25 20,99	74 25 20,99	—	25 17,83	+	3,16	7,705
538	5 23 46,59	5 23 45,42	—	145 23 46,01	—	23 33,75	+	12,26	7,644
539	5 38 18,18	4 38 17,17	—	104 38 17,73	38 16,66	38 12,21	+1,07	+ 5,52	7,638
540	5 8 15,55	5 8 16,13	—	78 8 15,84	—	8 20,74	—	4,90	7,622

xxvi *Comparison of the Observed Places of the Principal Fixed Stars*

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833				Green.	A. S.	
			s	s	s				s.	s.	
541	6.7	Tauri	—	4 49,45	2 49,70	4 30 49,55	—	49,14	—	+0,41	+3,733
542	6	Eridani	—	—	5 36,45	4 31 36,43	—	36,11	—	+0,32	2,743
543	5	Tauri	1 10,49	6 10,28	—	4 32 10,33	10,40	10,08	-0,07	+0,25	3,584
544	7	95 Tauri	—	—	4 4,11	4 33 4,14	—	3,51	—	+0,63	3,614
545	4	54 Eridani	6 5,82	6 5,86	6 5,86	4 33 5,84	5,86	5,60	-0,02	+0,24	2,616
546	6	Eridani	—	—	5 7,97	4 33 7,94	—	7,76	—	+0,18	2,494
547	6	Tauri	—	5 7,64	—	4 35 7,65	—	7,05	—	+0,60	3,306
548	4.5	Cœli Scalp α	6 9,09	6 9,26	—	4 35 9,15	—	9,54	—	-0,39	1,939
549	5	Cœli Scalp β	5 7,37	4 7,46	—	4 36 7,40	—	7,09	—	+0,31	2,111
550	6	Tauri	—	6 28,74	—	4 36 28,76	—	28,31	—	+0,45	3,484
551	5	57 Eridani	4 6,65	6 6,42	—	4 37 6,51	—	6,05	—	+0,46	2,990
552	4.5	Camelopardi	—	3 24,49	6 24,29	4 37 24,47	24,17	24,02	+0,30	+0,45	5,881
553	6	Eridani	—	5 43,52	—	4 39 43,50	—	43,32	—	+0,18	2,390
554	6	58 Eridani	—	—	6 3,99	4 40 3,96	—	3,57	—	+0,39	2,678
555	6	96 Tauri	—	5 7,86	—	4 40 7,85	—	7,26	—	+0,59	3,419
556	4	1 Orionis	6 43,56	6 43,54	8 43,63	4 40 43,58	43,66	43,79	-0,08	-0,21	3,251*
557	6	59 Eridani	—	—	5 59,35	4 40 59,33	—	59,51	—	-0,18	2,692
558	5	2 Orionis	6 27,75	6 27,82	3 27,82	4 41 27,80	27,73	27,59	+0,07	+0,21	3,258
559	5.6	97 Tauri	—	2 33,21	5 33,21	4 41 33,23	—	32,69	—	+0,54	3,490
560	4	3 Orionis	5 15,68	6 15,96	—	4 42 15,83	15,97	16,12	-0,14	-0,29	3,185
561	7	Aurigæ	—	—	5 18,02	4 42 18,05	—	17,42	—	+0,63	3,727
562	6	60 Eridani	—	—	8 37,81	4 42 37,79	—	36,89	—	+0,90	2,694
563	5	4 Orionis	5 2,34	6 2,16	—	4 43 2,24	—	1,62	—	+0,62	3,382
564	5	7 Camelop.	—	5 50,51	—	4 43 50,55	—	50,60	—	-0,05	4,773
565	6	5 Orionis	—	1 37,17	2 37,65	4 44 37,49	—	37,01	—	+0,48	3,117
566	5	61 Eridani	—	6 38,77	—	4 44 38,77	—	38,50	—	+0,27	2,941
567	6	6 Orionis	—	—	5 28,36	4 45 28,38	—	27,49	—	+0,89	3,317
568	4.5	8 Orionis	1 30,31	4 30,33	3 30,44	4 45 30,37	30,39	29,54	-0,02	+0,83	3,116
569	5.6	7 Orionis	—	—	5 39,22	4 45 39,23	—	38,51	—	+0,72	3,288
570	4	3 Aurigæ	4 3,81	5 3,89	5 3,85	4 46 3,86	3,88	0,61	-0,02	+3,25	3,887
571	5	9 Orionis	6 56,05	6 56,02	—	4 46 56,04	—	55,46	—	+0,58	3,367
572	6.7	Tauri	—	—	5 37,61	4 47 37,63	—	37,34	—	+0,29	3,625
573	6.7	Tauri	—	—	5 40,50	4 47 40,51	—	40,66	—	-0,15	3,453
574	5	4 Aurigæ	6 51,73	6 51,93	—	4 47 51,84	—	51,63	—	+0,21	4,047
575	6	98 Tauri	—	—	5 53,00	4 47 53,02	—	52,12	—	+0,90	3,654
576	6	62 Eridani	—	—	5 8,13	4 48 8,13	—	7,73	—	+0,40	2,947
577	4.5	10 Camelop.	2 30,34	1 30,11	3 30,36	4 48 30,37	30,45	30,49	-0,08	-0,12	5,286
578	7	Tauri	—	5 27,66	—	4 49 27,67	—	27,85	—	-0,18	3,392
579	5.6	10 Orionis	—	—	5 50,85	4 49 50,85	—	50,63	—	+0,22	3,100
580	4	7 Aurigæ	4 55,67	5 55,70	4 55,68	4 49 55,72	55,76	55,32	-0,04	+0,40	4,280
581	7	101 Tauri	—	1 6,20	4 6 14	4 50 6,16	—	5,96	—	+0,20	3,425
582	4	8 Aurigæ	—	6 44,97	3 44,89	4 50 44,98	45,04	44,84	-0,06	+0,14	4,170
583	5	63 Eridani	1 53,79	6 53,68	—	4 51 53,70	—	53,75	—	-0,05	2,831
584	6	64 Eridani	—	—	5 7,81	4 52 7,80	—	6,94	—	+0,86	2,778
585	4.5	102 Tauri	6 3,77	7 3,76	7 3,76	4 53 3,78	3,76	3,80	+0,02	+0,48	3,568

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession	
	No. 1831		No. 1832				No. 1833	Green.		A. S. C.
	"	"	"							
541	—	—	5 43 6,16	61 43 6,16		43 4,58	+	1,58	7,614	
542	4 41 29,02	—	—	104 41 29,02		41 22,14	+	6,88	7,548	
543	5 22 20,10	4 22 20,93	5 22 20,85	67 22 20,43	22 21,26	22 20,13	-0,83	+	0,30	7,504
544	—	5 14 13,17	—	66 14 13,17		14 10,52	+	2,65	7,482	
545	4 0 2,09	4 0 4,27	—	110 0 3,18	59 58,97	59 54,49	+4,21	+	8,69	7,426
546	5 49 2,36	1 49 1,56	—	114 49 2,23		48 55,62	+	6,61	7,423	
547	1 10 24,40	5 10 25,76	—	79 10 25,53		10 25,53	+	0,00	7,264	
548	5 11 19,58	5 11 21,27	3 11 18,59	132 11 20,00		11 16,60	+	3,40	7,257	
549	5 28 34,61	5 28 35,70	—	127 28 35,16		28 43,47	-	8,31	7,179	
550	—	5 34 35,54	—	71 34 35,54		34 31,24	+	4,30	7,154	
551	5 34 4,04	5 34 5,10	—	93 34 4,57		34 4,46	+	0,11	7,101	
552	5 57 18,09	5 57 18,67	—	23 57 18,38	57 22,09	57 20,69	-3,71	-	2,31	7,084
553	5 23 45,18	5 23 44,65	—	118 23 44,91		23 44,71	+	0,20	6,884	
554	—	5 14 51,81	—	107 14 51,81		14 48,70	+	3,11	6,857	
555	—	2 23 48,45	2 23 48,33	74 23 48,39		23 45,54	+	2,85	6,854	
556	5 20 17,62	5 20 20,08	6 20 19,37	83 20 19,04	20 20,81	20 21,38	-1,77	-	2,34	6,804
557	—	—	5 37 58,12	106 37 58,12		37 57,36	+	0,76	6,781	
558	5 23 42,48	4 23 41,75	5 23 42,85	81 23 42,41	23 42,53	23 41,40	-0,12	+	1,01	6,744
559	—	—	5 27 11,79	71 27 11,79		27 8,09	+	3,70	6,737	
560	5 41 23,60	3 41 25,52	5 41 23,73	84 41 24,10	41 20,43	41 17,57	+3,67	+	6,53	6,677
561	—	—	5 23 29,36	62 23 29,36		23 27,33	+	2,03	6,677	
562	—	1 30 51,18	4 30 53,97	106 30 53,41		30 51,43	+	1,98	6,647	
563	5 2 11,92	5 2 11,04	—	76 2 11,48		2 10,44	+	1,04	6,615	
564	5 31 39,06	5 31 37,44	—	36 31 38,25		31 40,22	-	1,97	6,551	
565	—	4 46 32,82	1 46 32,63	87 46 32,72		46 24,90	+	7,82	6,483	
566	4 44 27,39	5 44 24,84	—	95 44 26,11		44 19,82	+	6,29	6,480	
567	1 51 17,31	1 51 16,33	4 51 17,93	78 51 17,56		51 19,18	-	1,62	6,414	
568	5 50 26,39	3 50 27,06	—	87 50 26,64	50 29,32	50 25,07	-2,68	+	1,57	6,410
569	—	—	5 7 29,05	80 7 29,05		7 26,39	+	2,66	6,398	
570	5 6 29,91	5 6 32,14	—	57 6 31,02		6 26,77	+	4,25	6,369	
571	5 45 28,65	5 45 30,92	—	76 45 29,78		45 25,33	+	4,45	6,292	
572	—	—	6 19 16,52	66 19 16,52		19 16,41	+	0,11	6,235	
573	—	—	5 7 0,82	73 7 0,82		7 1,07	-	0,25	6,230	
574	5 22 20,35	5 22 21,07	—	52 22 20,71		22 18,70	+	2,01	6,216	
575	—	—	5 13 3,26	65 13 3,26		13 1,20	+	2,06	6,215	
576	—	5 26 40,73	—	95 26 40,73		26 34,47	+	6,26	6,191	
577	5 48 57,25	5 48 58,62	—	29 48 57,93	48 56,36	48 54,17	+1,57	+	3,76	6,166
578	1 43 15,32	—	—	75 43 15,32		43 14,88	+	0,44	6,081	
579	—	—	5 33 0,33	88 33 0,33		33 0,82	-	0,49	6,049	
580	5 26 4,82	6 26 3,83	5 26 3,89	46 26 4,16	26 5,50	26 4,35	-1,34	-	0,19	6,045
581	—	1 20 37,91	4 20 39,22	74 20 38,96		20 32,36	+	6,60	6,028	
582	5 10 43,38	6 10 42,64	6 10 42,90	49 10 42,95	10 44,13	10 43,45	-1,18	-	0,50	5,976
583	5 30 55,27	4 30 56,69	—	100 30 55,90		30 51,36	+	1,54	5,877	
584	—	2 47 24,26	3 47 26,19	102 47 25,42		47 24,03	+	1,39	5,858	
585	5 39 24,25	5 39 23,97	4 39 23,58	68 39 23,96	39 27,48	39 25,94	-3,52	-	1,98	5,782

xxviii *Comparison of the Observed Places of the Principal Fixed Stars*

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833				Green.	A. S.	
			s.	s.	s.				s.	s.	
586	5	65 Eridani ↓	3 17,81	6 17,76	—	4 58 17,78	—	17,61	—	+ 0,17	+ 2,901
587	7	Tauri	—	4 21,29	2 21,35	4 54 21,32	—	21,10	—	+ 0,22	3,561
588	4	10 Aurigæ η	5 44,71	5 44,84	5 44,72	4 54 44,79	44,75	44,47	+ 0,04	+ 0,32	4,182
589	5	11 Orionis γ ¹	3 58,44	3 58,42	—	4 54 58,43	—	58,16	—	+ 0,27	3,416
590	5.6	Leporis	—	—	5 20,13	4 55 20,10	—	19,76	—	+ 0,34	2,428
591	6	1 Leporis	—	—	5 39,94	4 55 39,92	—	39,56	—	+ 0,36	2,522
592	5	104 Tauri m	1 31,95	7 31,80	—	4 57 31,82	31,66	31,77	+ 0,16	+ 0,05	3,542*
593	5.6	106 Tauri n	—	5 52,24	1 52,12	4 57 52,24	—	51,86	—	+ 0,38	3,541
594	6	Tauri	—	—	5 52,73	4 57 52,76	—	52,66	—	+ 0,10	3,642
595	6	105 Tauri	—	—	5 53,13	4 57 53,15	53,01	53,47	+ 0,14	- 0,32	3,574
596	7	Tauri	—	—	5 2,60	4 58 2,53	—	2,22	—	+ 0,41	3,754
597	4	2 Leporis ε	5 21,14	6 21,16	—	4 58 21,15	21,27	20,40	- 0,12	+ 0,75	2,532
598	5	Cæli Scalp γ ¹	1 22,20	6 22,12	—	4 58 22,11	—	21,72	—	+ 0,39	2,142
599	6	66 Eridani	—	—	5 27,57	4 58 27,57	—	27,43	—	+ 0,14	2,958
600	6	Leporis	—	—	5 27,35	4 58 27,32	—	26,81	—	+ 0,51	2,429
601	6	14 Orionis i	—	—	4 44,37	4 58 44,38	—	44,41	—	- 0,03	3,255
602	7	107 Tauri l ²	—	—	2 56,07	4 58 56,09	—	55,81	—	+ 0,28	3,528
603	3	67 Eridani β ²	6 35,69	6 35,73	1 35,88	4 59 35,72	36,01	35,58	- 0,29	+ 0,14	2,948
604	5	15 Orionis γ ²	2 5,49	6 5,45	—	5 0 5,47	—	5,22	—	+ 0,25	3,423
605	6	16 Orionis h	—	3 5,24	2 5,56	5 0 5,38	—	5,17	—	+ 0,21	3,286
606	6	68 Eridani	—	—	5 24,42	5 0 24,42	—	24,12	—	+ 0,30	2,962
607	4	69 Eridani λ	1 6,32	5 6,64	—	5 1 6,58	6,66	6,41	- 0,08	+ 0,17	2,864
608	5	11 Aurigæ μ	—	5 56,39	—	5 1 56,41	—	56,34	—	+ 0,07	4,088
609	6.7	Orionis γ ³	—	—	4 3,18	5 2 3,20	—	2,58	—	+ 0,62	3,435
610	5	Doradus ζ	1 38,24	6 38,52	—	5 2 38,45	—	39,71	—	- 1,26	1,021
611	1	13 Aurigæ α	17 17,46	19 17,45	33 17,16	5 4 17,36	17,43	17,11	- 0,07	+ 0,25	4,402
612	5	14 Aurigæ α	—	3 28,09	—	5 4 28,11	—	28,78	—	- 0,67	3,894
613	4.5	3 Leporis i	—	6 27,83	—	5 4 27,82	27,81	27,50	+ 0,01	+ 0,32	2,791
614	5	17 Orionis ρ ¹	4 30,79	1 30,72	—	5 4 30,78	—	30,46	—	+ 0,32	3,128
615	7	108 Tauri	—	6 22,14	—	5 5 22,15	—	21,53	—	+ 0,62	3,595
616	5	5 Leporis μ	6 23,26	4 23,28	—	5 5 23,27	—	22,94	—	+ 0,33	2,686
617	4	Orionis	—	—	6 29,10	5 5 29,09	28,94	28,57	+ 0,15	+ 0,52	2,878
618	5	4 Leporis x	—	—	6 28,50	5 5 28,49	—	28,36	—	+ 0,13	2,765
619	1	19 Orionis β	10 28,07	7 28,09	11 28,08	5 6 28,08	28,05	27,68	+ 0,03	+ 0,20	2,876
620	6	18 Orionis	—	—	5 44,45	5 6 44,47	—	41,17	—	+ 0,30	3,324
621	5	15 Aurigæ λ	—	4 19,90	—	5 7 19,92	—	19,16	—	+ 0,76	4,157
622	6	Columbæ	—	—	5 40,42	5 8 40,39	—	40,10	—	+ 0,29	2,400
623	5.6	109 Tauri n	—	—	4 11,24	5 9 11,27	—	10,48	—	+ 0,79	3,592
624	7	Tauri	—	—	6 18,77	5 9 18,79	—	18,70	—	+ 0,69	3,541
625	4	20 Orionis τ	7 27,10	6 27,21	—	5 9 27,15	27,12	27,21	+ 0,03	- 0,06	2,907
626	6	Leporis	—	—	5 57,52	5 9 57,50	—	—	—	—	2,750
627	7	Tauri	—	6 24,31	—	5 10 24,32	—	23,87	—	+ 0,45	3,527
628	6	21 Orionis	—	6 25,60	—	5 10 25,60	—	25,61	—	- 0,01	3,123
629	6.7	Aurigæ	—	2 31,90	4 31,84	5 10 31,88	—	31,76	—	+ 0,12	3,803
630	5	Columbæ o	6 25,68	5 25,75	—	5 11 25,70	—	25,32	—	+ 0,38	2,151

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in					Mean N. P. D. January 1, 1833.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.		
	No.	1831	No.	1832	No.				1833	Green.		A. S. C	
586	5 25	34,23	5 25	36,10		—	97 25	35,17		25 33,52	+ 1,65	— 5,760	
587			5 57	53,10		—	68 57	53,10		57 55,21	— 2,11	5,673	
588	3 0	6,78	5 0	5,95	5 0	6,22	49 0	6,24	0 7,83	0 7,54	— 1,59	— 1,30	5,642
589	5 50	17,18	5 50	18,43		—	74 50	17,80		50 11,48	+ 6,32	5,621	
590	3 31	5,05	4 31	4,66		—	116 31	4,83		31 2,65	+ 2,18	5,588	
591			4 2	25,33	1 2	24,62	113 2	25,19		2 31,02	— 5,88	5,560	
592	5 35	16,40	5 35	17,06	7 35	15,66	71 35	16,29	35 16,93	35 13,95	— 0,64	+ 2,34	5,406
593			5 48	41,10		—	69 48	41,10		48 38,05	+ 3,05	5,377	
594			4 57	50,91	1 57	51,77	65 57	51,08		57 53,14	— 2,06	5,377	
595					5 31	26,22	68 31	26,22	31 31,38	31 28,50	— 5,16	— 2,28	5,375
596					5 57	18,99	61 57	18,99		57 21,08	— 2,09	5,364	
597	5 36	5,93	5 36	5,32		—	112 36	5,62	36 10,05	36 2,33	— 4,43	+ 3,29	5,335
598	5 43	4,96	5 43	8,31		—	125 43	6,63		43 1,23	+ 5,40	5,331	
599					5 53	16,59	94 53	16,59		53 10,01	+ 6,58	5,326	
600					5 23	4,09	116 23	4,09		23 0,64	+ 3,45	5,325	
601					5 43	39,58	81 43	39,58		43 36,41	+ 3,17	5,303	
602					5 22	0,51	70 22	0,51		21 56,02	+ 4,49	5,288	
603	5 18	38,69	5 18	40,08		—	95 18	39,39	18 36,52	18 33,96	+ 2,87	+ 5,43	5,230
604	5 37	31,37	6 37	31,30		—	74 37	31,33		37 22,61	+ 8,72	5,190	
605			1 23	43,56	4 23	45,75	80 23	45,31		23 33,91	+ 11,40	5,189	
606	3 40	54,30		—	2 40	56,49	94 40	55,17		40 48,04	+ 7,13	5,162	
607	5 58	32,39	5 58	31,63		—	98 58	31,96	58 32,10	58 31,85	— 0,14	+ 0,11	5,102
608	5 43	26,15	5 43	24,95		—	51 43	25,55		43 29,97	— 4,42	5,085	
609			2 10	9,86	4 10	12,22	74 10	11,47		10 7,83	+ 3,64	5,024	
610	2 42	13,59	5 42	15,70		—	147 42	14,31		41 33,18	+ 41,63	4,965	
611	40 10	57,39	25 10	57,57	47 10	57,80	44 10	57,61	10 56,14	10 58,03	+ 1,47	— 0,42	4,837
612						—	57	—		30 54,16		—	4,819
613	4 4	38,27	5 4	40,30		—	102 4	39,24	4 38,93	4 33,58	+ 0,31	+ 5,66	4,818
614			5 20	44,52		—	87 20	44,52		20 41,09	+ 3,43	4,815	
615					5 54	52,25	67 54	52,25		54 54,34	— 2,09	4,744	
616	5 24	35,74	1 24	37,07		—	106 24	35,96		24 34,50	+ 1,46	4,739	
617	5 21	5,47		—		—	98 21	5,47	21 7,68	—	— 2,21	4,732	
618			5 8	44,12		—	103 8	44,12		8 45,54	— 1,42	4,732	
619	29 24	5,06	14 24	4,51	23 24	5,43	98 24	5,07	24 7,78	24 5,40	— 2,71	— 0,33	4,647
620			5 51	17,14		—	78 51	17,14		51 13,91	+ 3,23	4,626	
621	5 3	39,48	5 3	39,48		—	50 3	39,48		3 32,20	+ 7,28	4,578	
622					5 8	13,99	117 8	13,99		8 10,78	+ 3,21	4,459	
623					5 5	4,75	68 5	4,75		5 2,75	+ 2,00	4,419	
624					5 3	1,77	70 3	1,77		2 56,71	+ 5,06	4,407	
625	5 1	55,65	4 1	56,38		—	97 1	55,97	1 57,62	1 56,11	— 1,65	— 0,14	4,398
626	2 42	13,70		—	3 42	14,56	103 42	14,22		42 15,79	— 1,57	4,352	
627					5 36	8,66	70 36	8,66		36 6,18	+ 2,48	4,314	
628					5 35	5,36	87 35	5,36		35 3,22	+ 2,14	4,310	
629					5 36	35,09	60 36	35,09		36 30,85	+ 4,24	4,304	
630	5 3	55,89	5 3	58,53		—	125 3	57,21		3 36,48	+ 20,73	4,223	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.		Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832			Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
				No. 1831	No. 1832	No. 1833						Green.	A. S.	
				s	s	s	h.	m.	s.			s.	s.	
631	4.5	6 Leporis	λ	5 50,16	5 50,32	—	5	11	50,23	50,34	49,90	-0,11	+0,33	+2,758
632	5.6	7 Leporis	ν	1 11,75	6 11,57	—	5	12	11,61	—	11,27	—	+0,34	2,778
633	6	Columbæ		—	—	5 42,35	5	12	42,32	—	41,53	—	+0,79	2,386
634	7	22 Aurigæ		—	—	5 44,68	5	12	44,71	—	44,49	—	+0,22	3,786
635	5.6	22 Orionis	o	—	—	—	5	13	—	—	10,74	—	—	3,055
636	7	Aurigæ		—	—	5 49,09	5	13	49,12	—	48,92	—	+0,20	3,856
637	7	Aurigæ		—	—	4 49,32	5	13	49,35	—	49,28	—	+0,07	3,854
638	7	110 Tauri		—	—	6 55,83	5	13	55,85	—	55,47	—	+0,38	3,457
639	5	23 Orionis	m	5 0,33	6 0,54	—	5	14	0,44	—	0,37	—	+0,07	3,145
640	6	111 Tauri		—	—	6 37,55	5	14	37,57	—	36,78	—	+0,79	3,474
641	6	Eridani		—	4 52,70	2 53,18	5	14	52,94	—	52,51	—	+0,43	2,459
642	2	112 Tauri	β	12 40,69	15 40,61	—	5	15	40,66	40,70	40,53	-0,04	+0,13	3,779
643	6	8 Leporis	ξ	—	—	3 49,12	5	15	49,10	—	49,00	—	+0,10	2,739
644	5.6	29 Orionis	ε	—	—	4 51,64	5	15	51,63	—	51,31	—	+0,32	2,884
645	5.6	27 Orionis	p	—	—	5 56,69	5	15	56,69	—	56,55	—	+0,14	3,044
646	5.6	25 Orionis	↓ ¹	—	—	5 1,93	5	16	1,93	—	1,27	—	+0,66	3,107
647	4.5	28 Orionis	η	6 2,00	6 2,03	—	5	16	2,01	2,03	1,86	-0,02	+0,15	3,009
648	2	24 Orionis	γ	1 7,36	8 7,49	—	5	16	7,48	7,40	7,42	+0,08	+0,06	3,210
649	6	113 Tauri		—	—	5 23,58	5	16	23,60	—	23,02	—	+0,58	3,458
650	5	24 Aurigæ	φ	1 31,08	5 31,09	—	5	16	31,10	—	31,72	—	-0,62	3,964
651	5.6	115 Tauri		—	1 22,53	3 22,46	5	17	22,49	—	22,15	—	+0,34	3,490
652	5	114 Tauri	o	3 33,15	6 32,95	—	5	17	33,02	32,96	32,78	+0,06	+0,24	3,593
653	5	30 Orionis	↓ ²	6 2,37	6 2,42	—	5	18	2,40	2,34	2,08	+0,06	+0,32	3,136
654	6	116 Tauri		—	—	5 6,65	5	18	6,66	—	6,35	—	+0,31	3,438
655	6	117 Tauri		—	—	5 16,87	5	18	16,88	—	16,49	—	+0,39	3,472
656	7	Tauri		—	—	5 28,41	5	18	28,42	—	28,32	—	+0,10	3,452
657	7	118 Tauri		—	—	6 56,14	5	18	56,16	—	56,27	—	-0,11	3,681
658	6	Leporis		—	3 15,32	2 15,62	5	19	15,42	—	14,07	—	+1,35	2,787
659	4	9 Leporis	β	6 2,96	6 2,94	5 3,03	5	21	2,97	3,00	2,96	-0,03	+0,01	2,565
660	5	31 Orionis	t	4 12,26	6 12,28	—	5	21	12,27	—	11,43	—	+0,84	3,040
661	5	25 Aurigæ	χ	6 47,99	6 47,94	—	5	21	47,96	—	39,75	—	—	3,893
662	5	32 Orionis	A	—	4 47,90	—	5	21	47,90	—	47,51	—	+0,39	3,202
663	5.6	119 Tauri		—	3 22,03	2 22,08	5	22	22,07	—	21,65	—	+0,42	3,508
664	6	33 Orionis	n	—	—	5 25,95	5	22	25,95	—	25,53	—	+0,42	3,141
665	2	34 Orionis	δ	6 25,54	16 25,63	1 25,72	5	23	25,63	25,69	25,55	-0,06	+0,08	3,058
666	6.7	Tauri		—	—	6 39,81	5	23	39,83	—	39,90	—	-0,07	3,557
667	6	120 Tauri		—	—	5 41,33	5	23	41,35	—	40,75	—	+0,60	3,507
668	5	36 Orionis	υ	1 48,49	4 48,54	—	5	23	48,53	—	48,85	—	-0,32	2,896
669	6	10 Leporis	ο	—	—	5 56,68	5	23	56,66	—	56,12	—	+0,54	2,562
670	7	35 Orionis	υ	—	—	5 21,69	5	24	21,71	—	21,02	—	+0,69	3,402
671	6	121 Tauri		—	—	4 11,82	5	25	11,84	—	11,48	—	+0,36	3,654
672	4	Columbæ	ε	1 14,94	5 15,03	—	5	25	15,01	15,00	14,80	+0,01	+0,21	2,122
673	3.4	11 Leporis	α	1 19,19	5 19,41	—	5	25	19,36	19,45	19,21	-0,09	+0,15	2,640
674	6.7	Aurigæ		—	—	5 23,26	5	25	23,28	—	23,15	—	+0,13	3,757
675	6	38 Orionis		—	—	5 26,61	5	25	26,61	—	26,17	—	+0,44	3,152

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
631	5 21 21,79	5 21 20,79	—	103 21 21,29	21 23,18	21 21,05	-1,89	+ 0,24	- 4,189
632	1 29 33,95	6 29 37,24	—	102 29 36,77	—	29 37,02	—	0,25	4,159
633	—	1 32 45,87	5 32 46,13	117 32 46,09	—	32 46,79	—	0,70	4,115
634	—	—	5 13 55,04	61 13 55,04	—	13 53,71	—	1,33	4,114
635	—	—	5 33 19,66	90 33 19,66	—	33 10,05	—	9,61	4,075
636	—	—	5 56 27,95	58 56 27,95	—	56 27,09	—	0,86	4,023
637	—	—	5 1 17,95	59 1 17,95	—	1 19,21	—	1,26	4,022
638	—	—	5 27 58,29	73 27 58,29	—	27 59,87	—	1,58	4,012
639	5 37 27,38	5 37 27,09	—	86 37 27,23	—	37 26,69	—	0,54	4,004
640	—	—	5 46 46,80	72 46 46,80	—	46 48,75	—	1,95	3,953
641	—	—	5 56 34,07	114 56 34,07	—	56 26,43	—	7,64	3,928
642	32 32 32,59	25 32 33,83	14 32 33,74	61 32 33,22	32 33,29	32 34,31	-0,07	- 1,09	3,863
643	—	—	4 5 27,05	104 5 27,05	—	5 25,57	—	1,48	3,848
644	—	—	5 58 5,50	97 58 5,50	—	58 5,69	—	0,19	3,845
645	—	—	5 3 34,02	91 3 34,02	—	3 31,01	—	3,01	3,838
646	—	—	3 18 50,62	88 18 50,62	—	18 50,40	—	0,22	3,831
647	5 33 28,48	5 33 28,81	—	92 33 28,65	33 30,84	33 29,95	-2,19	- 1,30	3,830
648	6 18 29,80	5 48 31,36	5 18 32,59	83 48 31,16	48 34,21	48 35,25	-3,05	- 4,09	3,823
649	—	—	1 27 23,83	73 27 23,83	—	27 24,82	—	0,99	3,801
650	5 40 36,68	5 40 37,12	—	55 40 36,90	—	40 37,79	—	0,89	3,790
651	—	—	5 11 24,06	72 11 24,06	—	11 24,90	—	0,84	3,717
652	5 12 50,08	5 12 49,13	—	68 12 49,60	12 51,38	12 47,88	-1,78	+ 1,72	3,702
653	5 3 23,41	3 3 24,58	—	87 3 23,90	3 25,40	3 22,39	-1,50	+ 1,51	3,658
654	—	1 16 31,89	4 16 32,45	74 16 32,34	—	16 26,24	—	6,10	3,653
655	—	—	5 54 28,21	72 54 28,21	—	54 26,51	—	1,70	3,639
656	—	—	5 42 23,86	73 42 23,86	—	—	—	—	3,622
657	—	—	5 59 41,76	64 59 41,76	—	59 39,12	—	2,64	3,582
658	2 2 55,28	2 2 54,79	—	102 2 55,03	—	2 52,46	—	2,57	3,554
659	5 53 58,45	5 53 57,48	—	110 53 57,96	53 56,80	53 54,03	+1,16	+ 3,93	3,397
660	5 13 53,43	5 13 53,98	—	91 13 53,71	—	13 50,93	—	2,78	3,387
661	5 56 24,55	5 56 24,37	—	57 56 24,46	—	56 28,97	—	4,51	3,337
662	5 11 13,67	5 11 14,43	—	84 11 14,05	—	11 9,87	—	4,18	3,335
663	1 32 15,50	3 32 17,42	7 32 17,41	71 32 17,22	—	32 17,41	—	0,19	3,287
664	—	—	6 50 33,19	86 50 33,19	—	50 35,88	—	2,69	3,280
665	5 25 52,80	14 25 51,95	—	90 25 52,17	25 50,10	25 49,95	+2,07	+ 2,22	3,194
666	—	—	5 39 13,32	69 39 13,32	—	39 9,31	—	4,01	3,175
667	—	—	—	71 35 —	—	35 10,19	—	—	3,173
668	2 25 53,45	2 25 52,06	—	97 25 52,75	—	25 55,52	—	2,77	3,160
669	1 59 37,99	4 59 38,15	—	110 59 38,12	—	59 52,38	—	14,26	3,148
670	—	—	6 49 13,05	75 49 13,05	—	49 8,83	—	4,22	3,115
671	—	—	5 4 48,26	66 4 48,26	—	4 50,21	—	1,95	3,043
672	5 35 56,03	—	—	125 35 56,03	—	35 50,54	—	5,49	3,034
673	6 56 51,61	5 56 53,11	—	107 56 52,29	56 54,00	56 54,19	-1,71	- 1,90	3,029
674	—	—	5 27 17,86	62 27 17,86	—	27 15,42	—	2,44	3,026
675	—	—	5 21 17,96	86 21 17,96	—	21 16,26	—	1,70	3,020

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^b January 1, Catal.			A. S. Catal.		Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	1832					Green.	A. S.	
			s.	s.	s.	h.	m.	s.	s.	s.	s.	s.	
676	4.5	37 Orionis ϕ^1	236,19	436,09	—	5 25 36,12	36,04	35,67	+0,08	+0,45	+3,286		
677	4	39 Orionis λ	—	653,33	—	5 25 53,33	53,41	53,00	-0,08	+0,33	3,297		
678	7	Tauri	—	139,66	539,51	5 26 39,56	—	39,19	—	+0,37	3,736		
679	6	41 Orionis θ^1	—	—	—	5 27 —	—	1,52	—	—	2,941		
680	5	42 Orionis c^1	5 6,10	6 6,09	—	5 27 6,09	—	5,53	—	+0,56	2,953		
681	6	43 Orionis θ^2	—	—	12 8,08	5 27 8,08	—	7,72	—	+0,36	2,940		
682	3.4	44 Orionis ι	213,00	313,11	—	5 27 13,07	13,15	13,48	-0,08	-0,41	2,928		
683	6	122 Tauri	—	—	419,16	5 27 19,18	—	18,66	—	+0,52	3,471		
684	3.4	123 Tauri ζ	636,56	636,53	—	5 27 36,55	36,51	35,81	+0,04	+0,74	3,577		
685	5	40 Orionis ϕ^2	—	640,80	—	5 27 40,80	—	40,28	—	+0,52	3,282		
686	2.3	46 Orionis ϵ	241,50	1041,50	—	5 27 41,50	41,50	41,35	0,00	+0,15	3,038		
687	5	26 Aurigæ l	—	—	650,15	5 27 50,17	—	50,85	—	-0,68	3,844		
688	6	125 Tauri	—	—	519,69	5 29 19,71	19,73	19,03	-0,02	+0,68	3,708		
689	6	Columbæ	—	—	336,28	5 29 36,25	—	36,07	—	+0,18	2,339		
690	4	48 Orionis σ	1118,86	618,92	—	5 30 18,88	18,95	18,57	-0,07	+0,31	3,005		
691	6	47 Orionis ω	—	—	619,24	5 30 19,24	—	24,20	—	—	3,161		
692	6	Columbæ ν^1	—	—	538,71	5 30 38,68	—	38,06	—	+0,62	2,364		
693	5	49 Orionis d	145,19	645,55	—	5 30 45,50	—	45,68	—	-0,18	2,898		
694	6	Orionis	—	—	5 9,16	5 31 9,16	—	8,85	—	+0,31	2,983		
695	6	Columbæ ν^2	—	611,39	—	5 31 11,38	—	11,25	—	+0,13	2,339		
696	5.6	126 Tauri	—	635,32	—	5 31 35,33	—	34,66	—	+0,67	3,459		
697	4	Doradus β	—	610,70	—	5 32 10,64	—	10,18	—	+0,46	3,509		
698	3	50 Orionis ζ	616,92	1717,09	217,12	5 32 17,02	17,09	17,12	-0,07	-0,10	3,021		
699	2	Columbæ α	734,14	634,11	1234,26	5 33 34,15	34,17	33,81	-0,02	+0,34	2,167		
700	6	51 Orionis δ	—	447,56	247,52	5 33 47,55	—	47,35	—	+0,20	3,100		
701	7	Tauri	—	—	626,06	5 34 26,08	—	26,11	—	-0,03	3,401		
702	6	12 Leporis	110,20	5 9,99	—	5 35 10,02	—	9,08	—	+0,94	2,519		
703	6	128 Tauri M	—	612,65	—	5 35 12,66	—	11,75	—	+0,91	3,449		
704	6	129 Tauri	—	1 5,92	5 6,00	5 37 6,01	—	5,52	—	+0,49	3,443		
705	4	13 Leporis γ	1527,72	627,76	—	5 37 27,73	27,79	27,32	-0,06	+0,41	2,517		
706	6	130 Tauri N	—	—	538,74	5 37 38,76	—	38,49	—	+0,27	3,491		
707	6	131 Tauri O	—	—	639,12	5 37 39,14	—	39,30	—	-0,16	3,410		
708	6	133 Tauri	—	—	511,40	5 38 11,42	—	11,20	—	+0,22	3,396		
709	5	132 Tauri B	542,55	642,56	142,47	5 38 42,56	42,70	41,87	-0,14	+0,69	3,674		
710	6	52 Orionis	—	—	458,89	5 38 58,89	—	58,50	—	+0,39	3,217		
711	4.5	14 Leporis ζ	520,69	620,82	220,67	5 39 20,75	20,84	20,24	-0,09	+0,51	2,714		
712	5	Columbæ μ	—	445,64	—	5 39 45,62	—	45,24	—	+0,38	2,224		
713	3	53 Orionis κ	247,38	647,44	—	5 39 47,43	47,40	47,20	+0,03	+0,23	2,840		
714	5	32 Aurigæ ν	—	650,91	—	5 39 50,93	—	50,63	—	+0,30	4,149		
715	5	31 Camelopard.	—	655,50	—	5 39 55,54	—	54,77	—	+0,77	5,358		
716	5.6	134 Tauri P	—	—	5 6,86	5 40 6,88	—	6,14	—	+0,74	3,365		
717	7	Tauri	—	—	523,10	5 40 23,13	—	22,87	—	+0,26	3,773		
718	5	30 Aurigæ ξ	146,25	546,34	—	5 40 46,36	—	46,28	—	+0,08	5,017		
719	6	135 Tauri	—	455,68	255,69	5 40 55,69	—	54,84	—	+0,85	3,406		
720	7	Tauri	—	—	514,84	5 41 14,86	—	15,09	—	-0,23	3,410		

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	1831		1832		1833					Green.	A. S. O	
	No.	"	No.	"	No.	"						
676	5 37	56,66	5 37	55,84	—	80 37	56,25	37 53,08	+	3,17	— 3,007	
677	5 11	12,99	5 11	12,20	5 11	12,44	80 11	12,54	11 8,42	11 4,05	+ 4,12 + 8,49 2,982	
678	—	—	—	—	5 11	18,16	63 11	18,16	11 19,34	—	1,18 2,917	
679	1 30	25,04	—	—	4 30	27,93	95 30	26,63	30 22,32	+	4,31 2,882	
680	5 57	24,45	5 57	26,26	—	—	94 57	25,35	57 16,49	+	8,86 2,877	
681	—	—	2 32	1,85	—	—	95 32	1,85	31 55,59	+	6,26 2,873	
682	4 1	34,26	4 1	35,12	—	—	96 1	34,69	1 36,98	1 29,85	-2,29 + 4,84 2,865	
683	—	—	—	—	5 4	14,63	73 4	14,63	4 16,55	—	1,92 2,859	
684	6 58	0,92	4 58	0,62	6 58	2,27	68 58	1,36	58 3,69	58 1,84	-2,33 - 0,48 2,835	
685	3 48	27,31	5 48	30,42	—	—	80 48	29,25	48 27,68	+	1,57 2,827	
686	10 18	57,47	9 18	57,76	—	—	91 18	57,61	18 58,10	18 57,02	-0,49 + 0,59 2,825	
687	5 37	0,86	—	—	—	—	59 37	0,86	36 49,72	+	11,14 2,814	
688	1 12	18,16	—	—	4 12	20,33	64 12	19,90	12 19,31	12 16,00	+ 0,56 + 3,90 2,686	
689	—	—	—	—	5 49	2,88	118 49	2,88	49 5,84	—	2,96 2,657	
690	5 42	11,14	5 42	13,62	—	—	92 42	12,39	42 13,76	42 9,10	-1,37 + 3,29 2,598	
691	—	—	6 58	53,96	—	—	85 58	53,96	58 50,04	+	3,92 2,590	
692	—	—	5 58	25,75	—	—	117 58	25,75	58 25,29	+	0,46 2,568	
693	5 18	47,11	4 18	46,12	—	—	97 18	46,67	18 44,97	+	2,40 2,558	
694	—	—	—	—	5 39	52,08	93 39	52,08	39 53,05	—	0,97 2,525	
695	—	—	—	—	5 47	42,96	118 47	42,96	47 46,54	—	3,58 2,520	
696	—	—	—	—	6 33	37,09	73 33	37,09	33 35,29	+	1,80 2,489	
697	4 36	1,70	5 36	1,62	—	—	152 36	1,61	36 0,50	+	1,14 2,429	
698	4 2	15,61	16 2	18,36	—	—	92 2	17,81	2 17,94	2 16,64	-0,13 + 1,17 2,427	
699	40 10	5,11	—	—	3 10	5,12	124 10	5,12	10 7,40	10 5,56	-2,28 - 0,44 2,313	
700	—	—	5 36	47,78	—	—	88 36	47,78	36 46,82	+	0,96 2,296	
701	—	—	—	—	4 54	35,48	75 54	35,48	—	—	— 2,241	
702	—	—	1 27	39,42	—	—	112 27	39,42	27 38,09	+	1,33 2,176	
703	—	—	5 59	35,54	—	—	73 59	35,54	59 34,25	+	1,29 2,175	
704	—	—	5 15	2,06	—	—	74 15	2,06	14 57,31	+	4,75 2,010	
705	5 30	29,28	5 30	29,05	3 30	28,51	112 30	29,01	30 31,05	30 30,59	-2,04 - 1,58 1,976	
706	—	—	7 20	29,10	—	—	72 20	29,10	20 25,66	+	3,44 1,962	
707	—	—	—	—	6 34	55,42	75 34	55,42	34 52,86	+	2,56 1,961	
708	—	—	—	—	4 10	9,32	76 10	9,32	10 3,26	+	6,06 1,915	
709	4 29	50,04	4 29	49,68	6 29	50,08	65 29	49,95	29 50,75	29 46,92	-0,80 + 3,09 1,871	
710	—	—	—	—	6 36	41,92	83 36	41,92	36 42,66	—	0,74 1,845	
711	5 53	29,35	5 53	30,94	—	—	104 53	30,14	53 27,10	53 23,45	+ 3,04 + 6,69 1,812	
712	5 22	30,39	5 22	29,86	—	—	122 22	30,12	22 29,25	+	0,87 1,775	
713	5 44	11,24	4 44	11,59	—	—	90 44	11,40	44 8,61	44 4,28	+ 2,79 + 7,12 1,774	
714	5 54	37,51	2 54	36,75	—	—	50 54	37,29	54 37,90	—	0,61 1,772	
715	5 9	47,31	—	—	—	—	30 9	47,31	9 42,39	+	4,92 1,770	
716	—	—	3 24	31,61	2 24	32,79	77 24	32,14	24 27,00	+	5,14 1,748	
717	—	—	—	—	5 5	29,57	62 5	29,57	5 28,14	+	1,43 1,725	
718	4 20	40,59	—	—	—	—	34 20	40,59	20 35,61	+	4,98 1,694	
719	—	—	—	—	6 45	5,43	75 45	5,43	45 55,31	+	10,12 1,677	
720	—	—	—	—	5 36	48,91	75 36	48,91	36 44,13	+	4,78 1,648	

xxiv Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832			Green ^h Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833						Green.	A. S.	
			s.	s.	s.	h.	m.	s.			s.	s.	
721	7	Tauri			4 19,06	5 41 19,08			17,66		+1,42	+3,400	
722	4.5	136 Tauri C	13 46,29	5 46,29		5 42 46,29	46,31	46,01	-0,02	+0,28	3,763		
723	6	137 Tauri D			5 49,98	5 42 50,00		49,69		+0,31	3,403		
724	6	Leporis			5 52,90	5 42 52,87		52,36		+0,51	2,502		
725	6	55 Orionis			5 15,54	5 43 15,53		15,22		+0,31	2,891		
726	5.6	56 Orionis			5 43,37	5 43 43,37		43,04		+0,33	3,110		
727	6.7	Aurigæ			7 4,73	5 44 4,76		4,87		-0,11	3,890		
728	5	15 Leporis δ	4 5,89	5 5,92		5 44 5,90		5,67		+0,23	2,559		
729	5	54 Orionis χ ¹	5 26,34	6 26,08		5 44 26,20	26,27	25,82	-0,07	+0,38	3,559		
730	5	Doradus δ		6 29,22		5 44 29,15		28,16		+0,99	0,102		
731	6	57 Orionis χ ²			4 0,11	5 44 0,13	0,03	0,30	+0,10	-0,17	3,546		
732	3	Columbæ β	3 2,60	6 2,49		5 45 2,51		2,32		+0,19	2,105		
733	3.4	33 Aurigæ δ		4 41,92		5 45 41,96	41,82	41,24	+0,14	+0,72	4,921		
734	1	58 Orionis α	7 4,67	19 4,71	54 4,75	5 46 4,74	4,79	4,66	-0,05	+0,08	3,241		
735	2	34 Aurigæ β	5 12,36	1 12,44		5 47 12,38	12,40	12,25	-0,02	+0,13	4,398		
736	5	35 Aurigæ α		6 28,21		5 47 28,23		27,92		+0,31	4,445		
737	5.6	139 Tauri			5 34,35	5 47 34,38		34,12		+0,26	3,717		
738	4	37 Aurigæ β	14 15,95	6 16,05		5 48 15,99	15,92	15,63	+0,07	+0,36	4,081		
739	4	16 Leporis η	6 45,56	6 45,38		5 48 45,47	45,50	45,06	-0,03	+0,41	2,730		
740	6	59 Orionis		6 40,98		5 49 40,98		40,54		+0,44	3,110		
741	5	Doradus ε		5 4,35		5 50 4,28		2,74		+1,54	-0,069		
742	6	60 Orionis B		6 11,50		5 50 11,50		11,01		+0,49	+3,080		
743	7	Aurigæ			5 27,12	5 50 27,15		26,90		+0,25	3,765		
744	5.6	2 Monocer A ²			6 5,90	5 51 5,89		5,58		+0,31	2,843		
745	6	141 Tauri Q ²			8 33,00	5 51 33,08		32,63		+0,40	3,618		
746	4	Columbæ γ	9 34,86	6 34,99	6 35,05	5 51 34,94	34,86	34,80	+0,08	+0,14	2,122		
747	5	61 Orionis μ	6 8,55	6 8,49	6 8,65	5 53 8,57	8,59	7,85	-0,02	+0,72	3,295		
748	5.6	64 Orionis χ ⁴		6 30,74		5 53 30,76		30,94		-0,18	3,546		
749	5	1 Geminor. H	6 54,69	6 54,74		5 53 54,73	54,63	54,32	+0,10	+0,41	3,042		
750	5	62 Orionis χ ³	3 56,55	6 56,65		5 53 56,63		56,67		-0,04	3,558		
751	5.6	3 Monocerotis		6 56,36		5 53 56,35		55,92		+0,43	2,818		
752	6	66 Orionis C		6 5,89		5 56 5,89		5,53		+0,36	3,165		
753	5.6	Leporis			6 29,77	5 56 29,74		29,19		+0,55	2,408		
754	6.7	2 Geminorum			6 34,17	5 56 34,20		33,98		+0,22	3,653		
755	7	Orionis			5 3,17	5 57 3,19		2,87		+0,32	3,440		
756	5.6	17 Leporis p		5 29,54	2 29,59	5 57 29,55		29,12		+0,43	2,673		
757	4.5	67 Orionis v	19 58,92	4 58,93	8 58,87	5 57 58,91	58,85	58,45	+0,06	+0,46	3,421		
758	4.5	18 Leporis θ	1 33,10	7 33,28	6 33,28	5 58 33,26	33,24	33,28	+0,02	-0,02	2,712		
759	6.7	Tauri		4 24,73	2 24,71	5 59 24,74		24,73		+0,01	3,614		
760	6	3 Geminorum		8 31,89		5 59 31,90		31,54		+0,36	3,639		
761	5	Camelopardi		6 19,13		6 0 19,21		19,01		+0,20	6,616		
762	7	4 Geminorum		3 18,50		6 0 18,51		17,94		+0,57	3,636		
763	6	19 Leporis τ			6 23,26	6 0 23,24		22,77		+0,47	2,604		
764	5	40 Camelopardi	1 34,37	5 34,76		6 0 34,76		34,43		+0,33	5,385		
765	6	4 Monocer D ²			2 33,35	6 0 33,34					2,505		

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
	" "	" "	" "				" "	" "	
721	—	5 11 1,42	—	70 11 1,42	—	—	—	—	1,644
722	5 26 7,12	4 26 7,85	7 26 7,69	62 26 7,56	26 8,60	—	-1,04	—	1,517
723	—	6 52 43,78	—	75 52 43,78	—	52 42,39	+	1,39	1,510
724	—	—	4 1 39,36	113 1 39,36	—	1 43,84	-	4,48	1,504
725	—	—	5 33 27,69	97 33 27,69	—	34 9,99	—	—	1,472
726	—	—	5 11 33,57	88 11 33,57	—	11 33,51	+	0,06	1,432
727	—	—	5 19 55,91	58 19 55,91	—	19 52,58	+	3,33	1,402*
728	5 53 56,66	5 53 56,26	—	110 53 56,46	—	53 56,09	+	0,37	0,777
729	5 45 48,04	6 45 48,79	5 45 48,43	69 45 48,43	45 45,67	45 45,24	+2,76	+ 3,19	1,371
730	6 47 55,78	—	—	155 47 55,78	—	47 54,13	+	1,65	1,357
731	—	5 17 29,80	—	70 17 29,80	17 26,90	17 22,49	+2,90	+ 7,31	1,321
732	5 50 12,17	4 50 10,63	—	125 50 11,49	—	50 20,58	-	9,09	1,314
733	5 44 24,71	—	—	35 44 24,71	44 21,46	44 23,42	+3,25	+ 1,29	1,265
734	45 37 52,16	24 37 52,54	49 37 52,79	82 37 52,50	37 53,70	37 53,42	-1,20	- 0,92	1,226
735	6 4 50,55	5 4 51,02	—	45 4 50,77	4 45,67	4 46,38	+5,10	+ 4,39	1,181
736	5 5 16,61	—	—	44 5 16,61	—	5 17,61	-	1,00	1,109
737	—	—	—	64	—	4 31,13	—	—	1,097
738	5 48 27,83	5 48 27,73	2 48 27,60	52 48 27,75	48 29,38	48 26,94	-1 63	+ 0,81	1,038
739	5 12 13,50	5 12 15,01	—	104 12 14,26	12 17,10	12 10,23	-2,64	+ 4,03	0,991
740	—	—	5 11 13,88	88 11 13,88	—	11 9,25	+	4,63	0,912
741	5 56 35,25	4 56 32,70	—	156 56 34,12	—	56 52,04	-	17,92	0,870
742	—	5 28 12,33	—	89 28 12,33	—	28 9,52	+	2,81	0,867
743	—	4 26 38,72	1 26 39,71	62 26 38,92	—	26 40,66	-	1,74	0,846
744	—	—	5 34 36,44	99 34 36,44	—	34 30,94	+	5,50	0,787
745	—	—	5 36 41,12	67 36 41,12	—	36 38,49	+	2,63	0,750
746	5 18 24,55	5 18 27,30	—	125 18 25,92	—	18 21,86	+	4,06	0,742
747	5 21 40,54	5 21 40,27	5 21 40,02	80 21 40,28	21 35,33	21 37,32	+4,95	+ 2,96	0,610
748	—	5 18 54,90	—	70 18 54,90	—	18 48,24	+	6,66	0,577
749	5 44 7,60	6 44 7,59	5 44 7,61	66 44 7,60	44 8,77	44 6,63	-1,17	+ 0,97	0,543
750	5 52 0,45	4 51 57,74	—	69 51 59,25	—	51 56,05	+	3,20	0,540
751	4 36 24,61	—	—	100 36 24,61	—	36 23,32	+	1,29	0,539
752	—	6 50 20,40	—	85 50 20,40	—	50 19,86	+	0,54	0,351
753	—	4 17 21,52	—	116 17 21,52	—	17 21,36	+	0,16	0,314
754	—	4 21 15,28	—	66 21 15,28	—	21 6,84	+	6,44	0,311
755	—	5 27 44,41	—	74 27 44,41	—	26 40,66	+	3,75	0,268
756	—	—	5 28 44,63	106 28 44,63	—	28 48,06	-	3,43	0,228
757	11 13 12,25	5 13 13,56	11 13 12,87	75 13 12,75	13 7,21	13 2,83	+5,54	+ 9,92	0,187
758	7 55 37,34	4 55 30,42	—	104 55 38,10	55 35,48	55 39,58	+2,62	- 1,48	0,134
759	—	5 47 24,68	—	67 47 24,68	—	47 27,85	—	3,17	0,062
760	—	5 52 1,35	—	66 52 1,35	—	52 1,73	-	0,38	0,052
761	5 38 12,01	5 38 12,08	—	20 38 12,94	—	38 6,44	+	5,60	+ 0,009
762	—	—	5 58 46,12	66 58 46,12	—	58 45,15	+	0,97	0,016
763	—	—	5 9 7,12	109 9 7,12	—	9 6,13	+	0,90	0,026
764	4 58 7,02	5 58 5,58	—	29 58 6,22	—	58 1,07	+	5,15	0,034
765	—	—	4 7 38,77	101 7 38,77	—	7 38,35	+	0,42	0,038

xxvi *Comparison of the Observed Places of the Principal Fixed Stars*

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^b January 1, Catal. 1832			A. S. Catal.	Difference from		Annual Preces- sion	
			No. 1831	No. 1832	No. 1833	h.	m.	s.	s.	s.	s.		
			s.	s.	s.								Green.
766	7	5 Geminorum			6 13,96	6	1	13,98		13,90		+0,08	+3,676
767	5	Columbæ δ	5 46,24	6 46,15		6	1	46,18		45,68		+0,50	2,053
768	6	68 Orionis E ¹			4 4,42	6	2	4,44		3,78		+0,66	3,550
769	6.7	6 Geminorum			5 7,76	6	2	7,78		7,24		+0,54	3,634
770	5	1 Lyncis α	4 24,85	2 25,04		6	2	24,94		24,69		+0,25	5,535
771	6	69 Orionis f ¹			5 22,12	6	2	22,14		21,51		+0,63	3,455
772	5	70 Orionis ζ	8 23,28	6 23,27		6	2	23,28		23,13		+0,15	3,407
773	6	Canis Maj.		6 53,74		6	3	53,72		53,25		+0,47	2,384
774	4	44 Aurigæ η	5 40,46	1 40,28		6	4	40,43	40,34	39,66	+0,09	+0,77	3,825
775	4.5	7 Geminor. ν	2 44,18	5 44,16	7 44,20	6	4	44,20	44,24	43,82	-0,04	+0,38	3,623
776	4.5	2 Lyncis b		4 47,81	2 47,50	6	4	47,78	47,51	47,31	+0,27	+0,47	5,297
777	5.6	71 Orionis E ²	1 57,76	5 57,72		6	4	57,74		58,02		-0,28	3,533
778	6	72 Orionis f ²		5 44,03		6	5	44,04		43,56		+0,48	3,456
779	7	8 Geminor. k ¹			5 2,96	6	6	2,99		2,72		+0,27	3,663
780	6	73 Orionis k ¹			6 18,79	6	6	18,81		18,44		+0,37	3,367
781	4.5	5 Monocer α	18 39,74	6 39,80	14 39,80	6	6	39,77	39,79	39,49	-0,02	+0,28	2,922
782	7	9 Geminorum		5 43,77		6	6	43,78		43,26		+0,52	3,657
783	5.6	74 Orionis k ²			5 0,65	6	7	0,67		0,32		+0,05	3,360
784	7	Aurigæ		4 49,05	2 48,91	6	7	49,01		48,90		+0,11	3,756
785	6	75 Orionis l			5 51,51	6	7	51,51		51,25		+0,26	3,303
786	7	11 Geminorum		5 5,69	3 5,61	6	9	5,67		5,14		+0,53	3,649
787	4.5	Columbæ η	16 34,56	6 34,61		6	10	34,57	34,74	34,56	-0,17	+0,01	2,130
788	6	7 Monocerotis		5 37,32		6	11	37,32		37,00		-0,28	2,886
789	5	46 Aurigæ d	3 57,31	5 57,27		6	11	57,30		57,00		+0,30	4,623
790	3	13 Geminor μ	8 47,66	7 47,82	7 47,71	6	12	47,75	47,83	47,44	-0,08	+0,31	3,623
791	3	1 Canis Maj. ζ	6 51,94	6 51,95	6 51,96	6	13	51,94	52,01	51,95	-0,07	-0,01	2,298
792	7	Geminorum		6 22,78		6	14	22,79		22,60		+0,19	3,694
793	6	Monocerotis		6 27,27		6	14	27,27		27,09		+0,18	3,158
794	5.6	8 Monocer b		6 52,00		6	14	52,00		51,82		+0,18	3,177
795	7	Geminorum		5 19,30		6	15	19,31		18,82		+0,49	3,648
796	7	Geminorum			6 20,08	6	15	20,11		19,87		+0,24	3,645
797	2.3	2 Canis Maj. β	7 18,10	6 18,15	20 18,25	6	15	18,19	18,21	17,94	-0,02	+0,25	2,638
798	4	3 Canis Maj. λ	2 58,63	6 58,52		6	15	58,54	58,47	58,29	+0,07	+0,25	2,191
799	6	15 Geminorum			5 45,75	6	17	45,77		45,37		+0,40	3,576
800	6	48 Aurigæ z			4 46,16	6	17	46,19		45,96		+0,23	3,856
801	6	16 Geminorum			5 57,20	6	17	57,22		56,81		+0,41	3,569
802	6	77 Orionis D ¹			5 36,23	6	18	36,23		36,33		-0,10	3,077
803	6	78 Orionis D ²			6 40,61	6	18	40,61		40,40		+0,21	3,064
804	5	15 Geminorum υ	18 59,26	7 59,24	7 59,08	6	18	59,21	59,17	58,76	+0,04	+0,45	3,561
805	7	17 Geminorum		5 45,69		6	17	45,70		9,12			3,588
806	6	10 Monocerotis		6 39,98		6	19	39,98		39,84		+0,14	2,059
807	1	Argus α	12 13,22		12 13,42	6	20	13,23		13,23		+0,05	1,327
808	7	Geminorum		2 29,29	5 29,32	6	21	29,34		29,04		+0,30	3,918
809	6.7	19 Geminorum			7 57,75	6	21	57,77		57,36		+0,41	3,450
810	5	Canis Maj. D ¹	5 56,98	6 56,70		6	21	56,82		56,67		+0,15	2,221

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.					
	No.	1831	No.				1832	No.		1833	Green.	A. S. O		
	"	"	"				"	"		"	"	"		
766	—	—	5 33	5,06	65 33	5,06	33	2,17	+	2,89	+0,097			
767	4 14	1,76	5 14	3,08	127 14	2,49	14	2,90	—	0,41	0,148			
768	—	—	1 10	52,67	4 10	52,89	70 10	52,85	+	8,84	0,170			
769	—	—	5 3	41,49	67 3	41,49	3	39,94	+	1,55	0,175			
770	4 26	32,57	5 26	33,01	—	—	26	38,73	—	5,91	0,195			
771	—	—	6 50	19,95	—	—	73 50	19,95	+	3,07	0,196			
772	4 45	43,73	5 45	40,14	—	—	75 45	41,74	+	1,00	0,199			
773	—	—	5 7	20,80	—	—	117 7	20,80	+	1,30	0,333			
774	5 26	56,85	5 26	56,09	5 26	57,21	60 26	56,72	26 53,58	26 57,63	+3,14	—	0,91	0,396
775	10 27	5,65	4 27	6,47	5 27	6,63	67 27	6,08	27 9,47	27 7,90	—3,39	—	1,82	0,403
776	7 56	27,19	5 56	27,62	5 56	26,66	30 56	27,16	56 26,68	56 26,15	+0,48	+	1,01	0,403
777	—	—	—	—	5 47	41,77	70 47	41,77	—	47 32,65	—	+	9,12	0,424
778	—	—	—	—	5 48	44,34	73 48	44,34	—	48 39,13	—	+	5,21	0,491
779	—	—	5 59	0,79	—	—	65 59	0,79	—	58 59,86	—	+	0,93	0,518
780	—	—	—	—	5 24	12,72	77 24	12,72	—	24 15,62	—	—	2,90	0,542
781	5 13	48,75	9 13	47,85	—	—	96 13	48,17	13 48,38	13 43,75	—0,21	+	4,42	0,574
782	—	—	—	—	5 12	36,69	66 12	36,69	—	12 39,37	—	—	2,68	0,577
783	5 41	17,61	—	—	5 41	18,29	77 41	17,95	—	41 15,88	—	+	2,07	0,603
784	—	—	5 43	57,05	—	—	62 43	57,05	—	43 56,15	—	+	0,90	0,672
785	—	—	5 0	19,48	—	—	80 0	19,48	—	0 13,23	—	+	6,25	0,677
786	—	—	5 28	19,93	—	—	66 28	19,93	—	28 18,89	—	+	1,04	0,784
787	5 5	26,03	5 5	25,68	—	—	125 5	25,85	—	5 22,00	—	+	3,85	0,918
788	—	—	5 45	30,52	—	—	97 45	30,52	—	45 29,17	—	+	1,35	1,008
789	5 38	15,33	5 38	14,11	—	—	40 38	14,72	—	38 16,08	—	—	1,36	1,008
790	20 24	27,86	5 24	26,69	18 24	27,56	67 24	27,59	24 29,67	24 27,46	—2,08	+	0,13	1,107
791	5 59	42,83	5 59	43,71	—	—	119 59	43,27	59 37,70	59 44,18	+5,57	—	0,91	1,205
792	—	—	6 52	19,95	—	—	64 52	19,95	—	52 16,25	—	+	3,70	1,246
793	3 9	28,02	2 9	29,67	—	—	86 9	28,68	—	9 26,56	—	+	2,12	1,254
794	—	—	1 19	47,25	4 19	47,58	85 19	47,52	—	19 36,16	—	+	11,36	1,290
795	—	—	7 28	24,84	—	—	66 28	24,84	—	28 26,70	—	—	1,86	1,328
796	—	—	3 35	17,21	—	—	66 35	17,21	—	35 17,35	—	—	0,14	1,329
797	27 52	40,11	5 52	39,92	5 52	40,29	107 52	40,08	52 41,98	52 48,36	—1,90	—	8,28	1,329
798	5 21	20,55	5 21	19,03	—	—	123 21	19,79	21 22,50	21 26,02	—2,71	—	6,23	1,389
799	—	—	5 6	54,99	—	—	69 6	54,99	—	6 55,01	—	—	0,02	1,541
800	—	—	5 24	44,47	—	—	59 24	44,47	—	24 39,22	—	+	5,25	1,541
801	—	—	5 24	40,35	—	—	69 24	40,35	—	24 36,28	—	+	4,07	1,558
802	—	—	—	—	6 36	28,13	89 36	28,13	—	36 24,30	—	+	3,83	1,616
803	—	—	—	—	4 11	0,40	90 11	0,40	—	10 53,91	—	+	6,49	1,622
804	5 41	24,20	6 41	22,38	5 41	24,32	69 41	23,68	41 21,39	41 18,67	+2,29	+	5,01	1,648
805	—	—	5 7	1,90	—	—	69 7	1,90	—	7 3,86	—	—	1,96	1,663
806	—	—	5 39	57,98	—	—	94 39	57,98	—	39 56,51	—	+	1,47	1,709
807	59 36	22,19	15 36	23,38	13 36	21,78	142 36	22,33	—	36 31,02	—	—	8,69	1,762
808	—	—	5 26	2,67	—	—	57 26	2,67	—	26 1,62	—	+	1,05	1,865
809	—	—	5 59	10,15	—	—	73 59	10,15	—	59 7,24	—	+	2,91	1,907
810	6 28	44,84	5 28	42,47	—	—	122 28	43,76	—	28 47,17	—	—	3,41	1,910

xxxviii Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag.	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832		Green ^h Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831		No. 1832	No. 1833	Green.			A. S.		
			s.	s.	s.	h. m. s.						
811	7	21 Geminorum	—	—	5 30,10	6 22 30,12	—	30,32	—	-0,20	+3,497	
812	6	12 Monocer <i>e</i>	—	6 24,19	—	6 23 24,19	—	24,22	—	-0,03	3,184	
813	5	13 Monocer <i>f</i>	6 49,18	5 49,12	—	6 23 49,15	—	49,04	—	+0,11	3,242	
814	6.7	Geminorum	—	6 3,84	—	6 24 3,86	—	3,50	—	+0,36	3,406	
815	6	CanisMaj. <i>U</i>	—	5 7,37	—	6 24 7,35	—	7,61	—	-0,26	2,372	
816	6	Canis Maj.	—	1 32,15	4 32,09	6 24 32,08	—	32,01	—	+0,07	2,638	
817	6	49 Aurigæ <i>c</i>	—	—	5 36,91	6 24 36,94	—	37,45	—	-0,51	3,779	
818	5.6	4 CanisMaj <i>ξ</i> ¹	—	—	4 51,58	6 24 51,55	—	50,95	—	+0,60	2,495	
819	6	14 Monocer <i>g</i>	—	—	5 40,53	6 25 40,54	—	41,23	—	+0,31	3,248	
820	3	24 Geminor. <i>γ</i>	3 0,06	4 0,34	2 0,19	6 28 0,23	—	59,93	-0,29	+0,30	3,462	
821	5	5 CanisMaj <i>ξ</i> ²	4 1,20	2 1,01	—	6 28 1,17	—	0,74	—	+0,43	2,510	
822	6	54 Aurigæ	—	6 57,33	—	6 28 57,35	—	57,03	—	+0,32	3,785	
823	5	7 CanisMaj <i>ν</i> ²	4 21,39	6 21,48	—	6 29 21,44	—	21,32	—	+0,12	2,609	
824	5.6	8 CanisMaj <i>μ</i> ³	—	6 30,20	—	6 30 30,19	—	29,32	—	+0,27	2,635	
825	7	25 Geminorum	—	3 45,50	2 45,31	6 30 45,45	—	44,84	—	+0,61	3,782	
826	5	55 Aurigæ	2 50,99	4 50,83	—	6 30 50,91	—	50,86	—	+0,05	4,377	
827	6	15 Monocer <i>k</i>	—	4 43,49	2 43,60	6 31 43,54	—	43,40	—	+0,14	3,302	
828	5.6	26 Geminor <i>π</i>	—	—	7 37,18	6 32 37,20	—	37,13	37,25	-0,05	3,493	
829	3	Argus <i>ν</i>	4 37,24	6 37,39	—	6 32 37,32	—	37,32	—	0,00	1,832	
830	5	42 Camelopardi	5 23,39	—	—	6 33 23,39	—	23,12	—	+0,27	6,209	
831	3	27 Geminor <i>ε</i>	5 35,69	4 35,62	8 35,52	6 33 35,62	35,63	35,28	—	0 01	+0,34	3,693
832	6	28 Geminorum	—	4 6,61	2 6,48	6 34 6,59	6,33	6,05	—	+0,26	+0,54	3,305
833	5.6	30 Geminor <i>ξ</i> ¹	—	6 30,92	—	6 34 30,93	—	29,95	—	+0,38	3,383	
834	5	Camelopardi	—	5 26,75	—	6 35 26,88	—	26,36	—	+0,22	3,868	
835	5	43 Camelop. <i>g</i>	—	6 32,76	—	6 35 32,84	—	34,81	—	—	6,520	
836	4	31 Geminor <i>ξ</i> ²	6 51,75	5 51,54	—	6 35 51,66	51,53	51,57	—	+0,13	+0,09	3,375
837	6	16 Monocerotis	—	6 22,66	—	6 37 22,66	—	22,02	—	+0,64	3,271	
838	1	9 Canis Maj <i>α</i>	21 44,62	10 44,59	77 44,67	6 37 44,63	44,55	44,56	—	+0,08	+0,07	2,643*
839	5	17 Monocer <i>i</i>	4 12,47	4 12,43	—	6 38 12,45	—	12,03	—	+0,42	3,258	
840	5	18 Monocer <i>k</i>	1 5,77	5 6,09	—	6 39 6,05	—	5,81	—	+0,24	3,138	
841	6	11 CanisMaj <i>λ</i> ²	—	6 11,33	—	6 39 11,32	—	10,78	—	+0,54	2,734	
842	6	33 Geminor <i>G</i>	—	6 9,52	—	6 40 9,53	—	8,56	—	+0,97	3,455	
843	6	35 Geminorum	—	—	6 56,40	6 40 56,42	—	56,22	—	+0,20	3,386	
844	6.7	36 Geminor <i>d</i>	—	6 28,89	—	6 41 28,90	—	28,62	—	+0,28	3,508	
845	5	Arginup <i>x</i>	7 36,40	6 36,38	—	6 41 36,38	—	36,43	—	-0,05	2,051	
846	5	34 Geminor <i>β</i>	3 42,62	5 42,43	—	6 41 42,52	—	42,14	—	+0,38	3,960	
847	5	15 Lyncis <i>e</i>	4 42,39	1 42,44	—	6 42 42,34	—	41,78	—	+0,56	5,222	
848	4	13 CanisMaj <i>μ</i> ²	6 34,06	6 34,13	7 34,11	6 43 34,09	34,13	33,91	—	-0,04	+0,18	2,238
849	5	Canis Maj	7 45,69	5 45,83	—	6 44 45,74	—	45,70	—	+0,04	2,178	
850	6	37 Geminorum	—	5 58,54	2 58,47	6 44 58,53	—	58,01	—	+0,52	3,695	
851	5.6	38 Geminor <i>el</i> ¹	—	—	5 9,78	6 45 9,89	—	9,65	—	+0,15	3,389	
852	4	Argus <i>σ</i>	—	6 46,24	1 45,93	6 45 46,16	—	45,85	—	+0,31	1,484	
853	5.6	15 CanisMaj <i>π</i> ¹	—	—	6 17,06	6 46 17,06	—	16,84	—	+0,22	2,591	
854	5	14 Canis Maj <i>θ</i>	—	6 23,37	—	6 46 23,36	—	23,01	—	+0,35	2,794	
855	7	Geminorum	—	6 30,24	—	6 46 30,25	—	29,90	—	+0,35	3,492	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N P D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
							Green.	A. S. C.	
	No. 1831	No. 1832	No. 1833						
811	—	5 6 19,49	—	72 6 19,49	—	6 17,87	+	1,62	+1,955
812	—	4 1 57,15	1 1 56,04	85 1 56,93	—	1 47,73	+	9,20	2,034
813	5 33 4,08	5 33 0,58	—	82 33 2,33	—	33 6,48	—	4,15	2,070
814	—	—	5 43 26,80	75 43 26,80	—	43 20,57	+	6,23	2,090
815	—	5 39 25,37	—	117 39 25,37	—	39 27,03	—	1,66	2,099
816	—	4 56 46,34	—	107 56 46,34	—	56 49,57	—	3,23	2,134
817	—	5 51 18,68	—	61 51 18,68	—	51 20,12	—	1,44	2,138
818	—	5 17 9,11	—	113 17 9,11	—	18 10,44	—	1,33	2,162
819	—	5 18 15,65	—	82 18 15,65	—	18 16,32	—	0,67	2,231
820	10 27 50,90	5 27 51,74	16 27 51,43	73 27 51,31	27 53,66	27 51,35	-2,35	0,04	2,432
821	5 50 11,08	5 50 10,88	—	112 50 10,98	—	50 7,83	+	3,15	2,436
822	—	8 35 48,62	—	61 35 48,62	—	35 50,69	—	2,07	2,514
823	5 7 4,24	5 7 3,68	—	109 7 3,96	—	7 3,72	+	0,24	2,553
824	—	5 5 50,72	—	108 5 50,72	—	5 49,18	+	1,54	2,652
825	—	2 39 24,37	—	61 39 24,37	—	39 22,45	+	1,92	2,670
826	5 19 27,37	5 19 28,14	—	45 19 27,75	—	19 24,34	+	3,41	2,677
827	5 57 27,21	3 57 27,45	—	79 57 27,33	—	57 22,25	+	5,08	2,756
828	—	5 11 53,32	—	72 11 53,32	11 51,84	11 51,21	+1,48	2,11	2,833
829	5 3 8,02	5 3 5,48	—	133 3 6,75	—	3 10,64	—	3,89	2,838
830	5 15 26,34	5 15 24,94	—	22 15 25,64	—	15 26,56	—	0,92	2,891
831	16 42 40,30	6 42 42,83	23 42 41,68	64 42 41,33	42 39,04	42 36,57	+2,29	4,76	2,916
832	—	5 52 4,35	—	60 52 4,35	52 2,39	52 0,70	+1,96	3,65	2,960
833	—	1 36 34,87	4 36 31,02	76 36 32,51	—	36 34,91	—	2,40	2,996
834	5 49 39,46	5 49 40,82	—	12 49 40,14	—	49 40,70	—	0,56	3,061
835	5 55 51,48	5 55 51,95	—	20 55 51,71	—	55 49,66	+	2,05	3,080
836	5 55 49,01	5 55 48,57	1 55 48,65	76 55 48,77	55 49,24	55 45,24	-0,47	3,53	3,113
837	—	5 14 32,79	—	81 14 32,79	—	14 29,35	+	3,44	3,244
838	52 29 30,40	24 29 30,08	54 29 30,74	106 29 30,46	29 31,08	29 27,58	-0,62	2,88	4,418*
839	5 47 18,42	5 47 18,22	—	81 47 18,32	—	47 17,85	+	0,47	3,316
840	5 24 37,24	5 24 37,78	—	87 24 37,51	—	24 31,34	+	6,17	3,394
841	—	5 15 6,70	—	104 15 6,70	—	15 0,70	+	6,00	3,402
842	—	5 36 47,15	—	73 36 47,15	—	36 45,75	+	1,40	3,433
843	—	5 24 4,46	—	76 24 4,46	—	24 1,48	+	2,98	3,551
844	—	5 2 51,89	—	68 2 51,59	—	2 51,28	+	0,61	3,597
845	5 44 53,06	5 44 51,86	—	127 44 52,91	—	44 56,30	—	3,39	3,613
846	5 50 42,31	5 50 44,33	—	55 50 43,35	—	50 43,43	—	0,06	3,615
847	5 22 9,30	5 22 6,90	—	31 22 8,10	—	22 11,50	—	3,40	3,697
848	5 19 9,35	4 19 9,69	—	122 19 9,50	19 9,20	19 8,20	+0,30	1,50	3,781
849	5 10 24,22	5 10 25,96	—	124 10 25,09	—	10 27,88	—	2,79	3,883
850	—	5 25 21,22	—	64 25 21,22	—	25 17,72	+	3,50	3,897
851	5 36 55,81	—	1 36 57,74	76 36 56,13	—	36 55,14	+	0,99	3,914
852	5 25 4,84	5 25 4,86	—	140 25 4,85	—	24 50,70	+	5,15	3,971
853	5 1 23,22	2 1 24,89	—	110 1 23,70	—	1 20,12	+	3,58	4,013
854	—	—	—	101 —	—	50 3,72	—	—	4,021
855	—	5 3 10,94	—	72 3 10,94	—	3 8,28	+	2,66	4,029

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832			Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	1832					Green.	A. S.	
			s.	s.	s.	h	m.	s.			s.	s.	
856	4	Equul Pict α	2 28,05	4 28,04	—	6 46	28,01	—	—	27,74	+0,27	+0,631	
857	4	16 Canis Maj. α^1	5 9,84	6 9,91	—	6 47	9,87	9,92	9,65	-0,05	+0,22	2,486	
858	6	17 Canis Maj. α^2	—	2 47,64	4 47,64	6 47	47,62	—	47,47	—	+0,15	2,587	
859	7	Geminorum	—	—	4 55,72	6 47	55,74	—	55,48	—	+0,26	3,496	
860	5.6	19 Canis Maj. α^3	—	—	5 20,62	6 48	20,60	—	20,03	—	+0,57	2,594	
861	6.7	39 Geminor. γ^1	—	—	11 25,76	6 48	25,79	—	25,07	—	+0,72	3,714	
862	5.6	18 Canis Maj. μ	—	—	6 24,91	6 48	24,90	—	24,28	—	+0,62	2,716	
863	4.5	20 Canis Maj. ν	10 38,81	5 38,78	3 38,90	6 48	38,81	38,76	38,45	+0,05	+0,36	2,673	
864	6.7	40 Geminor. γ^2	—	—	5 5,27	6 49	5,30	—	4,95	—	+0,35	3,709	
865	7	Geminorum	—	2 11,61	2 11,41	6 50	11,53	—	11,19	—	+0,34	3,446	
866	6.7	41 Geminorum	—	6 36,40	—	6 50	36,41	—	35,98	—	+0,43	3,450	
867	6	Canis Maj.	—	5 38,08	—	6 50	38,07	—	37,79	—	+0,28	2,476	
868	6	Canis Maj.	—	6 43,03	—	6 51	43,02	—	42,72	—	+0,30	2,455	
869	2.3	21 Canis Maj. ϵ	11 1,65	10 1,53	6 1,59	6 52	1,58	1,59	1,29	-0,01	+0,29	2,354	
870	6	42 Geminor. ω^1	—	—	6 10,16	6 52	10,18	—	10,05	—	+0,13	3,660	
871	6.7	Geminorum	—	1 49,58	4 49,22	6 52	49,30	—	48,78	—	+0,52	3,808	
872	4	43 Geminorum ζ	7 8,47	4 8,39	12 8,45	6 54	8,45	8,42	8,33	+0,03	+0,12	3,562	
873	5.6	19 Monocer δ	—	6 34,38	—	6 54	34,38	—	33,78	—	+0,60	2,977	
874	4.5	Camelopardi	4 14,18	5 15,36	—	6 55	14,99	15,27	14,68	-0,28	+0,31	13,217	
875	3.4	22 Canis Maj. σ	6 1,72	5 1,83	—	6 55	1,76	1,81	1,53	-0,05	+0,23	2,387	
876	6.7	44 Geminor. ω^2	—	6 11,31	1 10,99	6 55	11,28	—	10,76	—	+0,52	3,616	
877	4	24 Canis Maj. ω^2	4 0,64	6 0,76	—	6 56	0,70	0,79	0,38	-0,09	+0,32	2,502	
878	4	23 Canis Maj. γ	7 9,50	9 9,53	8 9,54	6 56	9,51	9,46	9,21	+0,05	+0,30	2,711	
879	6	45 Geminor. θ	—	6 43,74	—	6 58	43,75	—	43,56	—	+0,19	3,444	
880	5	63 Aurigæ	6 5,36	5 5,32	1 5,31	7 0	5,32	—	5,58	—	-0,26	4,135	
881	5	46 Geminor. τ	5 26,21	5 26,33	—	7 0	26,29	—	26,02	—	+0,27	3,829	
882	6	47 Geminorum	—	6 57,58	—	7 0	57,60	—	57,25	—	+0,35	3,729	
883	3.4	25 Canis Maj. δ	5 33,50	7 33,76	6 33,78	7 1	33,66	33,69	33,52	-0,03	+0,14	2,436	
884	5.6	29 Monocerotis	—	6 53,04	—	7 1	53,04	—	52,96	—	+0,08	2,979	
885	6	48 Geminor. m	—	6 13,73	2 13,45	7 2	13,69	—	13,22	—	+0,47	3,652	
886	5.6	Canis Maj.	—	4 47,85	2 47,81	7 2	47,82	—	47,74	—	+0,08	2,469	
887	4.5	22 Monocer. m	7 17,00	5 17,08	2 17,16	7 3	17,08	17,08	17,12	0,00	-0,04	3,063	
888	5	51 Geminorum	5 43,26	1 43,15	6 43,14	7 3	43,20	43,29	43,15	-0,09	+0,05	3,447	
889	7	52 Geminor. n	—	—	7 24,93	7 4	24,95	—	24,81	—	+0,14	3,671	
890	6	26 Canis Maj.	—	6 19,91	—	7 5	19,89	—	19,94	—	-0,05	2,452	
891	6	53 Geminor. α	—	—	6 27,17	7 5	27,20	—	26,73	—	+0,47	3,755	
892	5	64 Aurigæ	5 20,55	4 20,58	—	7 6	20,58	—	20,16	—	+0,42	4,188	
893	7	Geminorum	—	6 6,09	—	7 7	6,11	—	5,94	—	+0,17	3,446	
894	4.5	27 Canis Maj. ϵ^1	9 24,48	6 24,55	—	7 7	24,49	—	24,33	—	+0,16	2,443	
895	5	Arg. in pup 1	5 46,33	6 46,33	—	7 7	46,31	—	46,36	—	-0,05	1,722	
896	6	28 Canis Maj. ω^3	—	5 59,73	—	7 7	59,70	—	59,12	—	+0,58	2,431	
897	5	Arg in pup L1	4 12,17	4 11,85	4 11,92	7 8	11,90	—	11,41	—	+0,49	1,795	
898	4.5	54 Geminor. λ	4 26,20	6 26,18	8 26,03	7 8	26,12	25,98	26,13	+0,14	-0,01	3,455	
899	6	Canis Maj.	—	6 51,15	—	7 9	51,14	—	50,95	—	+0,19	2,402	
900	3.4	55 Geminor. δ	6 5,05	2 5,16	12 4,97	7 10	5,02	5,11	4,82	-0,09	+0,20	3,590	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	1831		1832		1833					Green.	A. S. C	
	No.	"	No.	"	No.	"						
856	5 45	44,37	1 45	44,39	—	—	151 45	44,37	45 47,25	—	2,88	+ 4,034
857	5 58	43,43	5 58	41,52	—	—	113 58	42,48	58 43,51	-4,05	1,03	4,088
858	—	—	4 11	50,38	—	—	110 11	50,38	11 45,57	—	4,81	4,142
859	—	—	—	—	5 53	3,06	71 53	3,06	53 1,09	—	1,97	4,151
860	—	—	4 55	45,26	—	—	109 55	45,26	55 37,57	—	7,69	4,188
861	—	—	—	—	3 42	19,73	63 42	19,73	—	—	—	4,192
862	—	—	5 49	54,38	—	—	103 49	54,38	49 54,21	—	0,17	4,194
863	6 50	31,38	5 50	33,53	—	—	106 50	32,36	50 28,11	-0,14	4,25	4,214
864	—	—	1 51	54,23	—	—	63 51	54,60	51 54,13	—	0,47	4,249
865	—	—	—	—	3 50	9,21	73 50	9,21	50 7,82	—	1,39	4,344
866	—	—	5 41	45,46	1 41	45,78	73 41	45,51	41 41,89	—	3,62	4,379
867	—	—	2 25	3,23	2 25	0,43	114 25	1,83	25 1,24	—	0,59	4,385
868	—	—	4 11	36,27	—	—	115 11	36,27	11 30,20	—	6,07	4,477
869	35 44	51,48	6 44	50,00	17 44	50,29	118 44	50,98	44 51,30	-3,39	0,82	4,504
870	—	—	5 33	11,42	—	—	65 33	11,42	33 7,31	—	4,11	4,512
871	—	—	5 23	23,61	—	—	60 23	23,61	22 58,32	—	25,29	4,567
872	5 11	26,04	8 11	25,23	25 11	26,76	69 11	26,35	11 27,39	-1,04	2,25	4,681
873	—	—	5 0	6,64	—	—	94 0	6,64	0 0,45	—	6,19	4,718
874	5 17	33,08	5 17	33,12	—	—	7 17	33,10	17 32,10	—	1,61	4,747
875	6 41	56,18	5 41	57,00	—	—	117 41	56,55	41 59,66	-0,25	3,11	4,759
876	—	—	5 7	8,38	—	—	67 7	8,38	7 6,43	—	1,95	4,769
877	5 35	31,46	5 35	31,11	—	—	113 35	31,29	35 36,12	-6,57	4,83	4,842
878	5 23	29,60	5 23	29,91	—	—	105 23	29,75	23 23,78	—	5,97	4,854
879	—	—	6 48	27,84	—	—	73 48	27,84	48 26,82	—	1,02	5,070
880	5 24	55,02	5 24	55,09	—	—	50 24	55,05	24 48,43	—	6,62	5,184
881	4 29	19,18	5 29	18,60	—	—	59 29	18,86	29 12,34	—	6,52	5,213
882	4 52	27,77	—	—	—	—	62 52	27,77	52 30,51	—	2,73	5,257
883	5 7	53,46	5 7	51,59	—	—	116 7	52,52	7 52,59	-1,70	0,07	5,312
884	5 58	50,06	—	—	—	—	93 58	50,06	58 47,44	—	2,62	5,338
885	3 35	53,27	—	—	—	—	65 35	53,27	35 52,25	—	1,02	5,364
886	—	—	5 57	56,99	—	—	114 57	56,99	57 51,77	—	5,22	5,416
887	5 13	19,33	4 13	18,68	—	—	90 13	19,04	13 16,55	—	3,96	5,456
888	5 33	44,45	5 33	44,13	10 33	44,83	73 33	44,56	33 44,79	-1,50	0,23	5,491
889	—	—	5 49	55,81	—	—	64 49	55,81	49 51,26	—	4,55	5,519
890	3 40	1,25	—	—	—	—	115 40	1,25	39 53,05	—	8,20	5,630
891	—	—	—	—	5 49	5,64	61 49	5,64	49 2,90	—	2,74	5,635
892	5 49	34,01	5 49	33,93	—	—	48 49	33,97	49 38,42	—	4,45	5,709
893	—	—	4 33	53,34	3 33	52,93	73 33	53,17	33 54,85	—	1,68	5,775
894	5 4	4,16	5 4	3,71	—	—	116 4	3,93	4 1,68	-1,40	2,25	5,803
895	5 28	55,43	6 28	54,45	—	—	136 28	54,90	28 58,59	—	3,69	5,836
896	5 29	9,14	—	—	5 29	9,42	116 29	9,28	29 7,08	—	2,20	5,852
897	5 53	44,39	5 53	43,43	—	—	134 53	43,91	53 33,75	—	10,16	5,871
898	5 9	44,69	5 9	48,32	10 9	46,30	73 9	46,40	9 47,46	-3,16	1,06	5,887
899	—	—	5 35	20,45	—	—	117 35	20,45	35 21,76	—	1,31	6,003
900	5 42	51,86	10 12	51,26	8 12	55,71	67 42	51,89	42 56,32	-2,89	1,43	6,024

No.	Mag.	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Precession
			No. 1831	No. 1832	No. 1833				Green.	A. S.	
901	5	Piscis Vol γ	2 8,77	4 8,82	—	7 10 8,72	—	—	—	—	—
902	5	65 Aurigæ	5 48,38	4 48,49	—	7 10 48,44	—	—	—	—	—
903	3.4	Argus π	2 12,39	6 12,57	4 12,95	7 11 12,63	—	—	—	—	—
904	6	29 Canis Maj.	—	6 40,80	—	7 11 40,78	—	—	—	—	—
905	6	30 Canis Maj. d	—	—	5 44,64	7 11 44,61	—	—	—	—	—
906	5.6	56 Geminor. q	—	7 1,64	—	7 12 1,65	1,77	1,76	-0,12	-0,11	3,550
907	6	57 Geminor. A	—	4 13,60	1 13 80	7 13 13,66	—	—	—	—	—
908	7	58 Geminorum	1 22,07	6 22,15	1 21,96	7 13 22,13	—	—	—	—	—
909	6.7	59 Geminorum	—	6 5,75	—	7 14 5,76	—	—	—	—	—
910	6	Canis Maj.	—	6 10,51	—	7 14 10,50	—	—	—	—	—
911	4	60 Geminor i	12 17,12	7 17,18	13 16,95	7 15 17,05	17,04	16,94	+0,01	+0,11	3,744
912	6	1 Canis Min.	—	—	6 37,87	7 15 37,88	—	—	—	—	—
913	6	2 Canis Min. ϵ	—	6 27,76	—	7 16 27,76	—	—	—	—	—
914	5	Piscis Vol δ	5 53,32	1 52,61	—	7 16 53,36	—	—	—	—	—
915	3	31 Canis Maj. η	6 27,15	6 27,16	3 27,27	7 17 27,18	27,22	26,37	-0,01	+0,81	2,370
916	6	63 Geminor p	—	6 45,70	3 45,73	7 17 45,72	45,67	45,61	+0,05	+0,11	3,572
917	3	3 Canis Min. β	6 2,40	6 2,18	21 2,29	7 18 2,29	2,21	1,73	+0,08	+0,56	3,259
918	5	62 Geminor r	4 17,87	3 18,02	—	7 18 17,94	—	—	—	—	—
919	3.6	64 Geminor b^1	—	2 51,84	4 51,68	7 18 51,75	—	—	—	—	—
920	6	5 Canis Min. η	—	—	6 59,73	7 18 59,73	—	—	—	—	—
921	5.6	4 Canis Min. γ	—	6 0,79	—	7 19 0,79	—	—	—	—	—
922	5.6	65 Geminor b^2	—	—	5 21,18	7 19 21,20	—	—	—	—	—
923	5.6	6 Canis Min. σ	—	6 26,56	—	7 20 26,57	—	—	—	—	—
924	6	Argus	—	8 19,22	—	7 21 19,21	—	—	—	—	—
925	6	7 Canis Min. δ^1	—	5 22,19	—	7 23 22,19	—	—	—	—	—
926	7	67 Geminorum	—	—	6 49,42	7 23 49,43	—	—	—	—	—
927	3	66 Geminor α	30 51,96	15 52,13	60 52,11	7 23 52,10	52,18	52,00	-0,08	+0,10	3,856
928	4	Argus π	5 51,21	5 54,10	—	7 23 54,14	—	—	—	—	—
929	5	68 Geminor k	5 1,13	5 0,99	—	7 24 1,07	—	—	—	—	—
930	5.6	8 Canis Min. δ^2	—	9 23,17	—	7 24 23,17	—	—	—	—	—
931	7	Geminorum	—	—	6 27,49	7 24 27,53	—	—	—	—	—
932	6	9 Canis Min δ^3	—	1 26,85	2 26,90	7 25 26,86	—	—	—	—	—
933	5	69 Geminor v	—	4 33,76	6 33,72	7 25 33,76	33,76	33,74	0,00	+0,02	3,709
934	7	Geminorum	—	7 12,45	1 12,44	7 27 12,46	—	—	—	—	—
935	6	Arg. in pup n^1	—	6 12,74	—	7 27 12,73	—	—	—	—	—
936	6	Arg. in pup n^2	—	—	5 13,58	7 27 13,56	—	—	—	—	—
937	5.6	Arg. in pup p	—	—	5 38,28	7 28 38,26	—	—	—	—	—
938	6	25 Monocerotis	—	5 55,51	—	7 28 55,51	—	—	—	—	—
939	7	Geminorum	—	2 8,89	3 8,61	7 29 8,75	—	—	—	—	—
940	6	74 Geminor f	—	—	5 46,12	7 29 46,15	46,20	46,16	-0,05	-0,01	3,471
941	1.2	10 Canis Min. α	41 30,27	16 30 29	48 30,29	7 30 30,29	30,35	30,14	-0,06	+0,15	3,143*
942	6	Arg. in pup m	—	6 18,60	—	7 31 18,59	—	—	—	—	—
943	6	75 Geminor σ	—	5 48,13	—	7 32 48,14	—	—	—	—	—
944	4.5	26 Monocer n	5 13,11	6 13,35	—	7 33 13,24	13,30	13,02	-0,06	+0,22	2,870
945	7	Geminorum	—	6 21,35	2 21,39	7 33 21,35	—	—	—	—	—

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession	
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.		
	"	"	"				"	"		
901	8 13 32,65	5 13 32,41	—	160 13 32,50	—	13 31,32	+	1,18	+ 6,040	
902	5 55 53,20	5 55 51,94	—	52 55 52,57	—	55 50,40	+	2,17	6,082	
903	5 48 2,49	5 47 59,41	—	126 48 0,95	—	48 1,26	—	0,31	6,121	
904	5 15 25,37	5 15 26,78	—	114 15 26,08	—	15 25,87	+	0,21	6,159	
905	5 39 12,78	—	—	114 39 12,78	—	39 10,51	+	2,27	6,165	
906	2 14 45,90	—	5 14 48,21	69 14 47,55	14 48,09	14 46,38	-0,54	+	1,17	6,186
907	—	5 38 4,17	—	61 28 4,17	—	38 1,16	+	3,01	6,285	
908	—	—	4 44 19,86	66 44 19,86	—	44 20,79	—	0,93	6,296	
909	—	10 2 43,28	—	62 2 43,28	—	2 43,66	—	0,38	6,357	
910	—	5 34 56,83	—	115 34 56,83	—	34 53,91	+	2,92	6,367	
911	5 52 34,93	10 52 33,97	9 52 32,98	61 52 33,81	52 32,91	52 28,63	+0,90	+	5,16	6,456
912	—	—	4 0 28,71	78 0 28,71	—	0 30,05	—	1,34	6,486	
913	—	5 24 0,91	—	80 24 0,91	—	23 57,53	+	3,38	6,551	
914	4 38 54,03	5 38 51,98	—	157 38 52,89	—	38 51,80	+	1,09	6,599	
915	13 58 46,76	5 58 45,42	—	118 58 46,38	58 49,36	58 48,85	-3,00	—	2,47	6,638
916	—	—	5 13 5,16	68 13 5,16	13 5,88	13 4,75	-0,72	+	0,41	6,661
917	8 22 44,25	9 22 44,29	—	81 22 44,27	22 44,24	22 40,03	+0,03	+	4,24	6,684
918	5 53 21,35	3 53 18,69	—	57 53 20,35	—	53 20,59	—	0,24	6,703	
919	7 32 35,42	—	—	61 32 35,42	—	32 35,62	—	0,20	6,751	
920	—	5 43 20,42	—	82 43 20,42	—	43 14,77	+	5,65	6,764	
921	1 44 30,66	4 44 30,45	—	80 44 30,51	—	44 28,04	+	2,47	6,765	
922	—	4 44 41,78	—	61 44 41,79	—	44 36,92	+	4,87	6,791	
923	—	5 39 9,40	—	77 39 9,40	—	39 6,59	+	2,81	6,881	
924	—	6 49 4,34	—	118 49 4,34	—	49 5,66	—	1,32	6,957	
925	—	5 44 5,80	—	87 44 5,80	—	44 3,00	+	2,80	7,122	
926	—	6 0 26,62	—	74 0 26,62	—	0 21,81	+	4,81	7,159	
927	33 45 3,91	23 45 3,61	77 45 3,65	57 45 3,71	15 4,22	45 3,45	-0,51	+	0,26	7,161
928	5 57 53,50	5 57 49,24	—	132 57 51,37	—	57 53,76	—	7,39	7,169	
929	5 49 5,48	5 49 4,57	—	73 49 5,02	—	49 2,76	+	2,26	7,174	
930	6 21 29,32	—	—	86 21 29,32	—	21 27,44	+	1,88	7,205	
931	—	5 40 54,18	—	58 40 54,18	—	40 54,73	—	0,55	7,210	
932	6 16 13,47	—	—	86 16 13,47	—	16 10,30	+	3,17	7,291	
933	5 44 16,03	5 44 15,93	—	62 44 15,98	44 17,52	44 13,36	-1,54	+	2,62	7,300
934	—	5 28 19,21	—	69 28 19,21	—	28 13,55	+	5,06	7,434	
935	—	5 6 46,47	—	113 6 46,47	—	6 44,26	+	2,21	7,137	
936	—	5 6 44,89	—	113 6 44,89	—	6 47,81	—	2,52	7,439	
937	—	2 0 7,98	4 0 7,62	118 0 7,31	—	0 6,93	+	0,38	7,553	
938	—	5 44 26,37	—	93 44 26,37	—	44 32,55	—	6,18	7,574	
939	—	5 36 46,18	—	57 36 46,18	—	36 42,85	+	3,33	7,591	
940	—	6 57 1,22	—	71 57 1,22	56 59,14	56 57,33	+2,08	+	3,89	7,642
941	60 21 3,78	33 21 3,78	75 21 3,05	84 21 3,85	21 2,91	20 57,47	+0,94	+	6,38	8,682*
942	—	4 59 24,41	—	114 59 24,44	—	59 18,10	+	6,31	7,768	
943	—	5 43 5,54	—	60 43 5,54	—	43 0,59	+	5,15	7,885	
944	4 9 51,82	—	—	99 9 51,82	9 51,80	9 51,10	-2,98	+	0,72	7,921
945	—	5 12 39,39	—	67 12 39,39	—	12 41,33	—	1,94	7,930	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Greenb January 1, Catal. 1832			A. S. Catal.	Difference from		Annual Preces- sion	
			No. 1831	No. 1832	No. 1833	h.	m.	s.		s.	s.		s.
			s.	s.	s.								
946	6	76 Geminor. c		3 51,68	3 51,24	7 33	51,48		51,29		+0,19	+3,671	
947	4	77 Geminor. x	5 18,01	5 17,91		7 34	17,97	17,81	17,05	+0,16	+0,92	3,634	
948	2	78 Geminor. β	24 1,62	17 1,57	63 1,50	7 35	1,56	1,60	1,14	-0,04	+0,42	3,682*	
949	7	79 Geminorum		6 17,38		7 35	17,39		17,38		+0,01	3,530	
950	6	1 Argus			5 52,09	7 35	52,07		51,82		+0,25	2,474	
951	6	81 Geminor. g			8 23,42	7 36	23,44	23,61	23,08	-0,17	+0,36	3,486	
952	5.6	Argus		6 45,56		7 36	45,54		45,04		+0,50	2,420	
953	6	11 Canis Min π		6 1,07		7 37	1,08		0,80		+0,28	3,309	
954	5	3 Argus	4 3,91	6 3,91		7 37	3,91		3,67		+0,24	2,405	
955	5.6	4 Argus		4 12,73		7 38	12,72		12,27		+0,45	2,761	
956	7	82 Geminor. B		6 30,49	2 30,43	7 38	30,49		29,79		+0,70	3,598	
957	4	Arg in pup c	6 16,32	4 16,27		7 39	16,29		16,42		-0,13	2,135	
958	5.6	Arg in pup o		6 6,27		7 41	6,26		5,87		+0,39	2,491	
959	5.6	6 Argus		6 5,95		7 42	5,94		6,78		-0,84	2,704	
960	7.	Geminorum		5 10,08	1 9,86	7 42	10,05		9,94		+0,11	3,501	
961	4	7 Argus ξ	7 13,99	6 13,93		7 42	13,95	13,90	13,73	+0,05	+0,22	2,520	
962	5.6	13 Canis Min ζ		6 59,13		7 42	59,13		58,78		+0,35	3,114	
963	5	83 Geminor. Φ	7 12,35	6 12,29		7 43	12,34	12,37	11,88	-0,03	+0,46	3,686	
964	5	9 Argus	5 59,50	1 60,00	5 59,71	7 43	59,64		59,21		+0,43	2,781	
965	4.5	Arg in pup P	4 7,32		2 7,53	7 44	7,38		7,36		+0,02	1,827	
966	6	10 Argus		5 35,10	2 34,93	7 44	35,04		34,54		+0,50	2,760	
967	6.7	85 Geminor. l		5 51,17	4 51,11	7 45	51,16		51,08		+0,08	3,511	
968	6	Canis Min		6 23,54		7 46	23,54		23,57		-0,03	3,264	
969	5	Arg in pup b	6 42,04	3 42,07		7 46	42,04		41,65		+0,39	2,121	
970	6	1 Cancri		5 26,83		7 47	26,84		26,36		+0,48	3,415	
971	5	Arg in pup R	6 21,95	6 21,82		7 48	21,86		22,08		-0,22	1,762	
972	7	Cancri		6 56,01		7 48	56,02		55,72		+0,30	3,431	
973	6	14 Canis Min		6 37,89		7 49	37,89		37,67		+0,22	3,123	
974	5.6	11 Argus		6 38,43		7 49	38,41		38,39		+0,02	2,578	
975	7	Cancri		6 11,97		7 50	11,98		11,70		+0,28	3,357	
976	6	2 Cancri α ¹			5 45,19	7 50	45,22		44,81		+0,41	3,641	
977	6	Argus		3 58,55	3 58,50	7 50	58,50		57,73		+0,77	2,388	
978	6	3 Cancri	1 9,20	5 9,35	1 9,33	7 51	9,33	9,33	8,94	0,00	+0,39	3,447	
979	6.7	4 Cancri α ²			6 35,21	7 51	35,23		34,89		+0,34	3,633	
980	6	12 Argus			6 53,61	7 51	53,59		52,70		+0,89	2,571	
981	6	5 Cancri r		6 55,35		7 51	55,37		54,80		+0,57	3,427	
982	3	Argus x	5 30,11	3 30,24		7 52	30,14		30,34		-0,20	1,530	
983	5.6	28 Monocer p		5 40,55		7 52	40,55		39,79		+0,76	3,049	
984	5.6	6 Cancri		6 11,34		7 53	11,36		10,89		+0,47	3,701	
985	5	13 Argus	8 31,43	5 31,49		7 53	31,45		31,31		+0,14	3,125	
986	6	8 Cancri		6 42,65		7 55	42,67		42,44		+0,23	3,351	
987	5	27 Lync.s k	7 47,27	3 46,99		7 55	47,18		46,99		+0,19	4,564	
988	5	55 Camelopardi	5 59,12	1 59,30		7 55	59,18		58,19		+0,99	6,107	
989	6	9 Cancri μ ¹		6 20,57		7 56	20,58	20,57	20,32	+0,01	+0,26	3,567	
990	3	Argus ζ	8 40,92	9 40,90	2 41,15	7 57	40,93		40,74		+0,19	2,108	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.	
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.		
	" " "	" " "	" " "				" " "	" " "		
946	—	5 49 22,44	—	63 49 22,44	—	49 16,87	+	3,57	+7,970	
947	5 12 25,06	5 12 25,56	—	65 12 25,31	12 22,54	12 22,70	+2,77	+	2,61	8,005
948	33 34 29,70	29 34 30,08	69 34 29,98	61 34 29,93	34 30,39	34 30,86	-0,46	-	0,93	8,064
949	—	5 17 16,11	—	69 17 16,11	—	17 15,10	+	1,01	8,086	
950	—	5 57 27,40	—	115 57 27,40	—	57 23,05	+	4,35	8,135	
951	—	4 5 14,12	—	71 5 14,12	5 13,03	5 10,69	+1,09	+	3,43	8,174
952	—	5 0 55,02	—	118 0 55,02	—	0 55,65	-	0,63	8,206	
953	—	5 49 42,62	—	78 49 42,62	—	49 33,04	+	4,58	8,224	
954	5 33 24,86	5 33 23,38	—	118 33 24,12	—	33 26,95	-	2,33	8,230	
955	—	5 9 39,44	—	104 9 39,44	—	9 36,20	+	3,24	8,320	
956	1 26 58,98	5 26 57,88	—	66 26 58,06	—	27 0,49	-	2,43	8,341	
957	5 33 52,36	5 33 49,63	—	127 33 51,00	—	33 54,77	-	3,77	8,407	
958	9 31 32,20	—	—	115 31 32,20	—	31 27,76	+	4,44	8,551	
959	5 48 20,07	—	—	106 48 20,07	—	48 17,80	+	2,27	8,630	
960	—	5 15 9,37	—	70 15 9,37	—	15 3,48	+	5,39	8,632	
961	5 26 37,02	5 26 35,08	—	114 26 36,05	26 35,83	26 33,47	+0,22	+	2,58	8,640
962	—	5 48 36,56	—	87 48 36,56	—	48 38,92	—	2,36	8,698	
963	5 48 21,53	5 48 21,61	16 48 22,10	62 48 21,90	48 23,92	48 24,92	-2,02	-	3,02	8,713
964	5 27 24,61	5 27 23,74	—	103 27 24,17	—	27 28,44	-	4,27	8,778	
965	5 57 12,83	5 57 13,52	—	135 57 13,17	—	57 12,25	+	0,92	8,791	
966	—	5 25 13,94	—	104 25 13,94	—	25 6,34	+	7,60	8,824	
967	1 40 45,47	4 40 46,95	—	69 40 46,65	—	40 42,73	+	3,92	8,922	
968	6 41 52,84	—	—	80 41 52,84	—	41 48,39	+	4,45	8,965	
969	5 25 51,23	—	—	128 25 51,23	—	25 52,51	-	1,28	8,992	
970	7 46 0,62	—	—	73 46 0,62	—	45 58,67	+	1,95	9,047	
971	5 40 1,25	5 40 0,57	—	137 40 0,91	—	40 7,91	-	7,00	9,123	
972	2 2 4,96	5 2 3,38	—	73 2 4,17	—	2 7,22	-	3,05	9,163	
973	—	4 19 59,46	—	87 19 59,46	—	20 1,71	-	2,25	9,218	
974	—	5 26 11,96	—	112 26 11,96	—	26 8,31	+	3,65	9,220	
975	—	5 18 26,41	—	76 18 26,41	—	18 24,82	+	1,59	9,261	
976	—	—	4 9 15,08	64 9 15,08	—	9 14,63	+	0,45	9,303	
977	—	5 53 19,38	—	119 53 19,38	—	53 13,97	+	5,41	9,323	
978	4 14 14,98	3 14 15,64	—	72 14 15,26	14 13,64	14 13,13	+1,62	+	2,13	9,335
979	—	5 27 20,75	—	64 27 20,75	—	27 16,58	+	4,17	9,368	
980	—	5 51 29,39	—	112 51 29,39	—	51 24,07	+	5,32	9,393	
981	—	2 5 14,53	—	73 5 14,53	—	5 13,78	+	0,75	9,394	
982	15 32 3,87	—	—	142 32 3,87	—	32 9,53	-	5,66	9,445	
983	—	5 55 54,78	—	90 55 54,78	—	55 54,31	+	0,47	9,453	
984	1 14 29 02	1 44 28,94	6 44 30,70	61 44 30,27	—	44 30,05	+	0,22	9,491	
985	5 12 36,43	5 12 35,39	—	87 12 35,91	—	12 31,48	+	4,43	9,519	
986	—	5 24 31,09	—	76 24 31,09	—	24 27,33	+	6,76	9,636	
987	5 1 2,46	5 1 2,41	—	38 1 2,45	—	1 10,38	-	7,93	9,639	
988	5 2 34,34	5 2 34,84	—	21 2 34,59	—	—	—	—	9,699	
989	—	5 55 29,08	—	66 55 29,08	53 28,57	53 26,06	+0,51	—	9,734	
990	12 32 4,42	9 32 3,59	20 32 4,07	129 32 4,06	—	32 0,49	+	3,57	9,840	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833				Green.	A. S.	
			s.	s.	s.				s.	s.	
991	6.7	10 Cancri μ^2	—	6 52,28	2 52,14	7 57 52,28	—	51,84	+0,39	+3,540	
992	7	11 Cancri	—	—	2 32,25	7 58 32,28	—	32,51	-0,23	3,685	
993	6	12 Cancri σ	1 18,66	—	2 18,67	7 59 18,68	—	18,61	+0,07	3,360	
994	5.6	29 Monocerotis	—	5 8,91	—	8 0 8,91	—	8,73	+0,18	3,018	
995	3.4	15 Argus	6 23,33	5 23,45	—	8 0 23,38	23,55	22,94	-0,17	+0,44	2,558
996	5.6	16 Argus	—	6 31,77	—	8 1 31,76	—	30,38	+1,38	2,677	
997	7	Cancri	—	—	—	8 2 —	—	6,32	—	3,278	
998	6	16 Cancri	—	—	1 34,29	8 2 34,31	—	33,94	+0,37	3,445	
999	6	15 Cancri ζ	—	—	2 43,33	8 2 43,36	—	42,53	+0,83	3,696*	
1000	6	18 Argus $\downarrow s$	—	—	2 52,62	8 2 52,60	—	52,74	-0,14	2,796	
1001	6	19 Argus	—	6 23,35	—	8 3 23,35	—	22,82	+0,53	2,815	
1002	5	Argus γ^1	14 18,87	4 18,91	—	8 4 18,87	—	18,89	-0,02	1,847	
1003	2	Argus γ^2	16 21,19	6 21,48	4 21,47	8 4 21,29	—	21,50	-0,21	1,848	
1004	7	Cancri	1 33,78	3 33,70	—	8 4 33,72	—	33,46	+0,26	3,444	
1005	5	20 Argus	5 36,94	6 36,77	—	8 5 36,84	—	36,43	+0,41	2,756	
1006	5	Argus	5 45,83	1 45,78	5 45,81	8 5 45,79	—	45,06	+0,73	2,024	
1007	5	Arg in pup r	6 9,14	—	7 9,43	8 7 9,29	—	9,04	+0,25	2,261	
1008	4	17 Cancri β	7 24,04	6 24,10	6 24,02	8 7 24,05	24,00	23,56	+0,05	+0,49	3,262
1009	5	Piscis Vol s	—	6 21,58	—	8 7 21,52	—	21,12	+0,40	0,242	
1010	6	21 Argus	—	5 41,05	—	8 9 41,04	—	40,75	+0,29	2,750	
1011	6	18 Cancri α	—	6 50,95	—	8 9 50,97	—	50,37	+0,60	3,662	
1012	6	19 Cancri λ	—	6 32,15	—	8 10 32,17	—	31,73	+0,44	3,542	
1013	6.7	Cancri	1 32,92	5 32,61	2 32,50	8 10 32,63	—	31,90	+0,73	3,506	
1014	5	31 Lyncis m	9 18,48	1 18,22	—	8 11 18,45	—	17,99	+0,46	4,142	
1015	4.5	Arg in pup q	6 16,43	5 16,33	—	8 12 16,37	—	16,56	-0,19	2,250	
1016	6	20 Cancri d^1	—	7 44,36	1 44,24	8 13 44,35	—	44,44	-0,09	3,449	
1017	7	21 Cancri f	—	6 43,65	—	8 14 43,66	—	43,15	-0,51	3,288	
1018	6	22 Argus	—	5 53,13	—	8 14 53,13	—	52,96	+0,17	2,821	
1019	6	Argus	—	6 44,14	—	8 15 44,13	—	44,01	+0,12	2,532	
1020	4.5	1 Ursæ Maj e	5 14,30	2 14,27	—	8 16 14,29	14,38	13,47	-0,09	+0,32	5,089
1021	6	1 Hydræ	—	5 12,65	—	8 16 12,65	—	11,86	+0,79	3,006	
1022	6.7	22 Cancri Φ^1	—	4 13,76	2 13,75	8 16 13,84	—	13,35	+0,49	3,668	
1023	6	25 Cancri d^2	1 18,72	—	5 18,81	8 16 18,80	—	18,34	+0,46	3,419	
1024	6	23 Cancri Φ^2	—	—	—	8 16 —	—	36,45	—	3,643	
1025	7	24 Cancri v^1	—	—	6 39,55	8 16 39,57	—	39,63	-0,06	3,586	
1026	6	Cancri	—	—	6 53,88	8 16 53,88	—	53,82	+0,06	3,226	
1027	5.6	30 Monocer q	—	—	6 15,99	8 17 15,99	—	15,60	+0,39	3,003	
1028	6.7	27 Cancri e	—	—	6 26,20	8 17 26,20	—	25,81	+0,39	3,327	
1029	6	Argus	—	6 48,33	—	8 17 48,37	—	48,02	+0,35	2,589	
1030	6	2 Hydræ	—	—	3 3,49	8 18 3,49	—	3,25	+0,24	3,002	
1031	6.7	28 Cancri v^2	—	8 33,59	1 33,47	8 18 33,58	—	37,86	+0,72	3,574	
1032	2	Argus ϵ	12 3,65	1 3,39	10 3,56	8 19 3,55	—	3,63	-0,08	1,243	
1033	6	29 Cancri	—	6 14,49	—	8 19 14,50	—	14,09	+0,41	3,357	
1034	6.7	30 Cancri ω^3	—	4 33,89	—	8 21 33,91	—	33,56	+0,35	3,568	
1035	5.6	31 Cancri θ	1 0,52	6 0,56	1 0,58	8 22 0,57	0,39	59,84	+0,18	+0,73	3,436

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.			
	1831		1832				Green.	A. S. C.				
	No.	"	No.									
991	2 56	9,21	2 56	9,67	—	67 56 9,44	56 10,48	—	1,04	+ 9,850		
992	—	—	—	—	3 2	16,08	62 2 16,08	2 14,19	+	1,89	9,901	
993	—	—	—	—	2 52	37,00	75 52 37,00	52 31,54	+	5,46	9,960	
994	3 29	59,07	5 29	57,97	—	—	92 29 58,38	29 59,26	—	0,88	10,025	
995	5 49	26,26	5 49	23,32	—	—	113 49 24,79	49 24,77	+ 4 45	+	0,02	10,044
996	5 45	27,06	3 45	27,00	—	—	108 45 27,04	45 30,42	—	3,38	10,128	
997	—	—	—	—	2 41	15,10	79 41 15,10	41 9,47	+	5,63	10,172	
998	—	—	—	—	2 51	8,28	71 51 8,28	51 5,12	+	3,16	10,206	
999	—	—	—	—	—	—	59 —	50 48,65	—	—	10,216	
1000	—	—	—	—	1 18	34,90	103 18 34,90	18 31,32	+	3,58	10,232	
1001	2 25	59,81	3 26	2,49	—	—	102 26 1,42	26 0,10	+	1,32	10,269	
1002	5 51	10,81	—	—	—	—	136 51 10,81	51 12,66	—	1,85	10,342	
1003	—	—	5 50	38,05	—	—	136 50 38,05	50 43,67	—	5,62	10,345	
1004	—	—	4 49	26,82	—	—	71 49 26,82	49 25,67	+	1,15	10,356	
1005	5 17	17,73	5 17	17,42	—	—	105 17 17,57	17 8,72	+	8,85	10,436	
1006	5 29	22,56	5 29	21,78	—	—	132 29 22,17	29 22,25	—	0,08	10,449	
1007	5 23	50,93	5 23	47,99	—	—	125 23 49,16	23 44,43	+	5,03	10,552	
1008	8 18	12,41	6 18	11,46	14 18	13,06	80 18 12,53	18 7,07	+ 3 23	+	5,46	10,568
1009	—	—	—	—	—	—	158 —	7 29,56	—	—	10,572	
1010	1 46	10,95	4 46	12,35	—	—	105 46 12,07	46 12,74	—	0,67	10,738	
1011	—	—	4 14	40,63	—	—	62 14 40,63	14 39,98	+	0,65	10,748	
1012	—	—	5 27	17,78	—	—	65 27 17,78	27 18,11	—	0,33	10,799	
1013	—	—	5 43	40,28	—	—	68 43 40,28	43 40,67	—	0,39	10,799	
1014	5 16	49,39	5 16	48,62	—	—	46 16 49,00	16 48,00	+	1,00	10,854	
1015	5 8	35,39	5 8	31,99	—	—	126 8 33,69	8 34,07	—	0,38	10,931	
1016	5 8	4,64	5 8	1,57	—	—	71 8 3,10	8 4,46	+	1,36	11,035	
1017	2 49	53,52	5 49	54,53	—	—	78 49 54,24	49 51,89	+	2,35	11,107	
1018	2 31	7,72	3 31	8,54	—	—	102 31 8,21	31 9,78	—	1,57	11,120	
1019	—	—	5 48	55,23	—	—	115 48 55,23	48 50,87	+	4,36	11,182	
1020	5 43	43,10	6 43	43,83	12 43	43,18	26 43 43,34	43 47,26	— 1 40	—	3,92	11,212
1021	—	—	5 12	40,91	—	—	93 12 40,91	12 40,18	+	0,73	11,215	
1022	—	—	5 33	33,23	—	—	61 33 33,23	33 34,19	—	0,96	11,215	
1023	—	—	5 24	20,83	—	—	72 24 20,83	24 22,03	—	1,20	11,222	
1024	—	—	—	—	—	—	62 —	31 22,41	—	—	11,243	
1025	—	—	—	—	7 55	16,53	64 55 16,53	55 13,43	+	3,10	11,247	
1026	—	—	4 53	39,18	—	—	81 53 39,18	53 37,25	+	1,93	11,265	
1027	—	—	2 21	44,68	2 21	47,01	93 21 45,84	21 45,13	+	0,71	11,292	
1028	—	—	5 47	49,75	—	—	76 47 49,75	47 48,11	+	1,64	11,303	
1029	—	—	5 30	18,88	—	—	113 30 18,88	30 20,63	—	1,75	11,332	
1030	—	—	3 26	24,91	—	—	93 26 24,91	26 23,41	+	1,50	11,349	
1031	—	—	5 18	14,79	—	—	65 18 14,79	18 13,24	+	1,55	11,389	
1032	—	—	—	—	—	—	148 —	58 22,79	—	—	11,426	
1033	—	—	5 13	24,52	—	—	75 13 24,52	14 15,13	+	9,39	11,438	
1034	—	—	4 21	33,72	—	—	65 21 33,72	21 28,61	+	5,11	11,599	
1035	3 20	34,63	5 20	36,02	—	—	71 20 35,50	20 34,94	— 1 43	+	0,56	11,630

xlvi*i* Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833				Green.	A. S.	
			s.	s.	s.				s.	s.	
1036	5	Chamæl <i>α</i>	—	—	—	8 22 invisible	43,88	—	—	-1,411	
1037	6	33 Cancrī <i>η</i>	—	7 59,06	—	8 22 59,07	58,98	58,77	+0,09	+0,30	+3,485
1038	6.7	34 Cancrī <i>h</i>	—	6 31,25	—	8 23 31,26	—	30,57	—	+0,69	+3,271
1039	5	Piscis Vol <i>η</i>	7 30,88	—	—	8 23 30,88	—	30,28	—	+0,60	-0,441
1040	5	Piscis Vol <i>β</i>	6 53,46	2 53,38	—	8 23 53,43	—	44,93	—	+8,50	+0,686
1041	6	Monocerotis	—	5 58,17	—	8 23 58,16	—	57,48	—	+0,68	+2,696
1042	5	4 Ursæ Maj <i>π²</i>	7 26,08	2 26,26	—	8 25 26,13	—	26,08	—	+0,05	+5,368
1043	5	Chamæl <i>θ</i>	—	—	—	8 25 invisible	—	31,02	—	—	-1,506
1044	6	Hydræ	—	6 54,85	—	8 26 54,85	—	54,63	—	+0,22	+3,204
1045	7	36 Cancrī <i>κ¹</i>	—	5 58,67	—	8 27 58,67	—	58,53	—	+0,14	3,261
1046	4	4 Hydræ <i>δ</i>	11 45,32	6 45,60	9 45,55	8 28 45,46	45,48	45,06	-0,02	+0,46	3,185
1047	5	5 Hydræ <i>σ</i>	6 58,66	3 58,61	5 58,57	8 29 58,61	—	58,06	—	+0,55	3,141
1048	7	38 Cancrī <i>ο</i>	—	6 2,92	—	8 30 2,93	—	2,77	—	+0,16	3,462
1049	7	Cancrī	—	4 11,97	1 11,86	8 30 11,96	—	11,41	—	+0,55	3,457
1050	6	39 Cancrī	—	6 26,36	2 26,06	8 30 26,28	—	25,63	—	+0,65	3,466
1051	6	40 Cancrī	—	2 31,30	2 31,25	8 30 31,29	—	31,17	—	+0,12	3,465
1052	7	Cancrī	—	4 43,06	—	8 30 43,07	—	42,64	—	+0,43	3,459
1053	6	Pixid Naut <i>η</i>	—	5 41,83	—	8 30 41,82	—	41,50	—	+0,32	2,560
1054	6.7	41 Cancrī <i>ε</i>	—	—	2 48,09	8 30 48,11	—	—	—	—	3,456
1055	6.7	Cancrī	—	3 48,40	6 48,22	8 30 48,30	—	48,24	—	+0,06	3,456
1056	5	Arg in Vel <i>κ¹</i>	8 44,82	6 44,51	—	8 31 44,40	—	44,87	—	+0,03	2,106
1057	5.6	6 Hydræ	—	—	6 4,05	8 32 4,05	—	3,52	—	+0,53	2,847
1058	7	Cancrī	—	—	7 10,49	8 32 10,50	—	10,14	—	+0,36	3,464
1059	6	Pixid Naut <i>ζ</i>	—	—	5 44,38	8 32 44,37	—	44,03	—	+0,34	2,487
1060	5	43 Cancrī <i>γ</i>	8 33,08	7 33,26	6 33,15	8 33 33,17	83,20	32,83	-0,03	+0,34	3,493
1061	5	Pixid Naut <i>β</i>	8 31,78	—	6 31,95	8 33 31,84	—	31,14	—	+0,70	2,342
1062	6.7	45 Cancrī <i>λ¹</i>	—	6 56,40	—	8 33 56,41	—	56,05	—	+0,36	3,315
1063	6	9 Hydræ	—	6 55,73	—	8 33 55,72	—	55,39	—	+0,33	2,781
1064	5	7 Hydræ <i>η</i>	8 26,53	1 26,49	—	8 34 26,53	—	26,16	—	+0,37	3,141
1065	5	Arg. in Vel <i>b</i>	6 3,32	6 3,27	—	8 35 3,28	—	3,28	—	0,00	1,987
1066	4.5	47 Cancrī <i>δ</i>	5 7,75	2 7,64	6 7,81	8 35 7,78	7,78	7,35	0,00	+0,43	3,422
1067	4	Argus <i>ο</i>	4 28,89	2 28,87	—	8 35 28,87	—	29,16	—	-0,29	1,721
1068	6.7	49 Cancrī <i>b</i>	—	5 37,62	—	8 35 37,63	—	36,93	—	+0,70	3,264
1069	5.6	48 Cancrī <i>i</i>	—	5 31,01	—	8 36 31,02	—	30,69	—	+0,33	3,652
1070	4.5	Pixid Naut <i>α</i>	6 50,85	—	5 50,85	8 36 50,84	50,92	50,65	-0,08	+0,19	2,406
1071	5	Arg in Car <i>d</i>	—	4 54,05	—	8 36 54,01	—	54,05	—	-0,04	1,884
1072	6	50 Cancrī <i>λ²</i>	1 42,89	6 43,23	—	8 37 43,19	—	42,50	—	+0,69	3,301
1073	4	11 Hydræ <i>ε</i>	4 52,56	5 52,39	5 52,54	8 37 52,50	52,50	52,25	-0,06	+0,25	3,195
1074	6	12 Hydræ	—	6 26,52	—	8 38 26,52	—	25,88	—	+0,64	2,832
1075	6	Hydræ	—	6 43,70	—	8 38 43,70	—	44,02	—	-0,32	3,045
1076	5	13 Hydræ <i>ρ</i>	8 31,62	4 32,03	—	8 39 31,78	—	31,50	—	+0,28	3,184
1077	3	Argus <i>δ</i>	5 3,46	3 3,96	3 4,29	8 40 3,92	—	3,91	—	-0,01	1,655
1078	5	Arg in Vel <i>a</i>	9 20,01	—	6 20,31	8 40 20,13	—	20,06	—	+0,07	2,030
1079	5.6	14 Hydræ <i>B</i>	—	5 55,15	—	8 40 55,15	—	54,43	—	+0,72	3,018
1080	7	Cancrī	—	6 9,00	—	8 41 9,01	—	8,53	—	+0,48	3,412

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	1831		1832		1833					Green.	A. S. C	
	No.	"	No.	"	No.	"				"	"	
1036	—	—	—	—	—	—	166 Invisible	23 16,59	—	—	+11,694	
1037	—	—	5 59 36,03	—	—	—	68 59 36,03	59 38,42	59 38,53	-2,39	- 2,50	11,700
1038	—	—	5 22 14,24	—	—	—	79 22 14,24	—	22 9,73	—	+ 4,51	11,738
1039	5 51 21,43	—	—	—	—	—	162 51 21,43	—	51 31,99	—	- 10,56	11,747
1040	6 34 37,42	—	5 34 34,91	—	—	—	155 34 36,28	—	34 28,14	—	+ 8,14	11,761
1041	—	—	5 0 53,91	—	—	—	109 0 53,91	—	0 52,28	—	+ 1,63	11,771
1042	3 5 45,19	—	4 5 41,25	—	—	—	25 5 41,65	—	5 42,01	—	+ 2,64	11,869
1043	—	—	—	—	—	—	166 Invisible	—	56 26,18	—	—	11,891
1044	1 47 54,77	—	5 47 51,87	—	—	—	82 47 51,85	—	47 50,95	—	+ 3,96	11,978
1045	—	—	4 15 55,28	—	—	—	79 45 55,28	—	15 58,16	—	- 2,88	12,058
1046	5 42 55,61	—	7 42 51,46	—	12 42 54,31	—	83 42 54,92	42 57,02	42 52,30	2,10	+ 2,62	12,107
1047	5 4 26,02	—	3 4 28,59	—	—	—	86 4 26,98	—	4 25,13	—	+ 1,85	12,192
1048	—	—	5 38 10,12	—	—	—	69 38 10,12	—	38 9,13	—	+ 0,96	12,196
1049	—	—	5 52 4,43	—	—	—	69 52 4,43	—	52 26,43	—	- 16,06	12,206
1050	—	—	5 24 19,98	—	—	—	69 24 19,98	—	24 15,56	—	+ 4,42	12,223
1051	—	—	5 26 31,13	—	—	—	69 26 31,13	—	26 25,34	—	+ 5,79	12,229
1052	—	—	5 14 33,48	—	—	—	69 14 33,48	—	44 33,46	—	+ 0,02	12,242
1053	—	—	—	—	6 40 17,54	—	115 40 17,54	—	40 16,18	—	+ 2,36	12,243
1054	—	—	—	—	7 52 4,63	—	69 52 4,63	—	52 1,88	—	+ 3,25	12,247
1055	—	—	—	—	—	—	69	—	51 59,80	—	—	12,240
1056	5 24 17,38	—	4 21 16,78	—	—	—	132 24 17,06	—	24 16,57	—	+ 0,49	12,317
1057	—	—	5 53 11,82	—	—	—	101 53 11,82	—	53 10,92	—	+ 0,90	12,337
1058	—	—	5 32 0,30	—	—	—	60 32 0,30	—	31 58,53	—	+ 1,77	12,343
1059	—	—	5 57 59,31	—	—	—	118 57 59,31	—	57 58,64	—	+ 0,67	12,384
1060	6 55 55,15	—	—	—	7 55 53,89	—	67 55 56,24	55 58,08	55 55,22	-1,81	+ 1,02	12,438
1061	5 43 2,49	—	6 43 0,47	—	—	—	124 43 1,39	—	42 57,86	—	+ 3,53	12,439
1062	—	—	5 43 19,16	—	—	—	76 43 19,16	—	43 18,48	—	+ 0,98	12,465
1063	—	—	5 20 42,85	—	—	—	105 20 42,85	—	20 33,14	—	+ 7,71	12,465
1064	5 0 13,51	—	5 0 12,45	—	10 0 12,43	—	86 0 12,71	—	0 12,36	—	+ 0,35	12,499
1065	6 3 16,47	—	5 3 17,97	—	—	—	136 3 17,15	—	3 13,63	—	+ 3,52	12,544
1066	5 14 0,06	—	—	—	4 14 1,06	—	71 14 0,51	14 0,73	14 0,60	-0,22	+ 0,51	12,546
1067	3 19 39,21	—	5 19 43,03	—	—	—	142 19 42,07	—	19 42,51	—	- 0,44	12,574
1068	—	—	5 18 57,19	—	—	—	79 18 57,19	—	18 51,57	—	+ 2,62	12,580
1069	—	—	5 37 51,67	—	—	—	60 37 54,67	—	37 51,97	—	+ 2,70	12,640
1070	5 35 4,82	—	4 35 2,63	—	—	—	122 35 3,85	35 4,83	35 0,29	-0,98	- 2,44	12,665
1071	—	—	4 9 45,47	—	—	—	149 9 45,47	—	9 53,30	—	- 7,83	12,671
1072	—	—	5 16 43,22	—	—	—	77 16 43,22	—	16 38,59	—	+ 4,63	12,722
1073	—	—	6 58 9,48	—	—	—	82 58 9,48	58 12,61	58 13,57	-3,13	- 4,09	12,733
1074	3 55 13,03	—	2 56 13,98	—	—	—	102 56 13,41	—	56 17,33	—	- 3,92	12,771
1075	—	—	5 17 9,64	—	—	—	91 17 9,64	—	17 10,61	—	- 0,97	12,791
1076	6 32 47,09	—	5 32 45,38	—	—	—	83 32 46,31	—	32 46,93	—	- 0,62	12,844
1077	6 5 44,49	—	4 5 41,25	—	—	—	144 5 43,05	—	5 51,99	—	- 8,94	12,884
1078	4 25 51,51	—	5 25 51,01	—	—	—	135 25 51,24	—	25 48,64	—	+ 2,60	12,901
1079	—	—	5 19 26,01	—	—	—	92 49 26,01	—	49 23,68	—	+ 2,93	12,937
1080	—	—	5 22 38,00	—	—	—	71 22 38,00	—	22 38,28	—	- 0,28	12,952

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in				Mean A. R. January 1, 1832	Green ^h Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831		No. 1832					Green.	A. S.	
			s.	s.	s.	s.						
1081	7	Canceri	—	4 10,90	1 10,67	8 41 10,88	—	10,89	—	-0,01	+3,428	
1082	6.7	54 Canceri	1 39,60	—	5 39,78	8 41 39,76	—	38,98	—	+0,78	3,359	
1083	6	55 Canceri ρ^2	—	5 34,57	—	8 42 34,58	—	34,21	—	+0,37	3,630	
1084	6	Pixid Naut	—	4 0,04	—	8 43 0,63	—	59,33	—	+0,70	2,511	
1085	6	Pixid Naut γ	—	5 24,32	—	8 43 24,31	—	24,42	—	-0,11	2,551	
1086	7	Canceri	1 43,37	6 43,39	—	8 43 43,40	—	43,28	—	+0,12	3,396	
1087	6	58 Canceri ρ^4	—	6 34,88	—	8 45 34,89	—	34,50	—	+0,39	3,614	
1088	7	Canceri	1 54,44	5 54,64	2 54,63	8 45 54,63	—	54,33	—	+0,30	3,591	
1089	4	16 Hydræ ζ	18 30,80	6 30,50	5 30,54	8 46 30,69	30,52	30,98	+0,17	-0,29	3,183	
1090	6	60 Canceri α^1	—	6 44,66	—	8 46 44,67	44,74	44,50	-0,07	+0,17	3,255	
1091	5	Chamœl η	—	—	—	8 46 invisible	—	50,17	—	—	-1,767	
1092	3 4	9 Ursæ Maj ι	7 40,05	2 40,09	3 40,02	8 47 40,09	40,07	39,55	+0,02	+0,54	+4,131*	
1093	6	62 Canceri ρ^1	—	4 52,35	—	8 47 52,36	—	51,74	—	+0,62	3,352	
1094	6	63 Canceri ρ^3	—	5 11,96	—	8 48 11,97	—	11,07	—	+0,90	3,357	
1095	6	Pixid Naut δ	—	4 19,42	—	8 48 19,40	—	18,63	—	+0,77	+2,562	
1096	5	Chamœl	—	—	—	8 48 invisible	—	43,79	—	—	-1,775	
1097	5	65 Canceri α^3	7 17,56	1 17,54	6 17,60	8 49 17,58	17,50	17,24	+0,08	+0,34	+3,287	
1098	7	Canceri	—	5 40,93	1 40,52	8 49 40,87	—	40,63	—	+0,24	3,404	
1099	4.5	Ursæ Maj. κ	6 7,23	2 7,05	5 6,99	8 52 7,14	7,19	7,29	-0,05	-0,15	4,147	
1100	6	69 Canceri ν	4 54,35	4 54,35	—	8 52 54,36	—	53,80	—	+0,56	3,525	
1101	5	Arg. in car b^1	9 51,83	3 51,83	—	8 52 51,82	—	51,67	—	+0,15	1,474	
1102	5	Arg. in car b^2	8 16,63	—	6 16,59	8 55 16,70	—	17,10	—	-0,40	1,498	
1103	6	Lyncis	7 49,22	—	5 48,96	8 55 49,14	—	48,88	—	+0,26	3,851	
1104	6	18 Hydræ ω	2 7,51	6 7,72	—	8 57 7,67	—	7,31	—	+0,36	3,165	
1105	6	Arg. in Vel c	14 22,03	6 21,87	—	8 58 21,98	—	22,26	—	-0,28	2,068	
1106	5.6	76 Canceri κ	—	6 38 72	—	8 58 38,73	38,37	39,22	+0,36	-0,49	3,259	
1107	6.7	75 Canceri	—	5 53,21	—	8 58 53,22	—	52,89	—	+0,33	3,559	
1108	7	78 Canceri	1 36,61	6 36,77	3 36,63	8 59 36,73	—	36,03	—	+0,70	3,379	
1109	5.6	77 Canceri ζ	2 41,36	5 41,36	—	8 59 41,36	41,26	41,14	+0,10	+0,22	3,465	
1110	5	Piscis Vol α	7 46,71	—	—	8 59 46,71	—	46,25	—	+0,46	0,966	
1111	6	79 Canceri	—	6 41,31	—	8 59 41,32	—	40,81	—	+0,51	3,462	
1112	5.6	Pixid Naut	—	6 40,45	—	9 0 40,44	—	40,14	—	+0,36	2,625	
1113	6	20 Hydræ λ^2	—	4 22,91	—	9 1 22,91	—	22,33	—	+0,56	2,934	
1114	3.4	Argus λ	10 49,71	—	2 49,32	9 1 49,64	—	49,47	—	+0,17	2,301	
1115	6	Pixid Naut ϵ	—	—	5 49,68	9 2 49,66	—	49,34	—	+0,32	2,536	
1116	6.7	81 Canceri π^1	1 5,45	7 5,56	3 5,46	9 3 5,52	—	5,38	—	+0,14	3,293*	
1117	6	Canceri	—	5 0,78	—	9 4 0,80	—	0,39	—	+0,41	3,443	
1118	5	18 Ursæ Maj. e	8 2,82	3 2,60	—	9 4 2,77	—	2,88	—	-0,11	4,380	
1119	6	21 Hydræ K^1	—	5 8,08	—	9 4 8,08	—	7,61	—	+0,47	2,964	
1120	5	Arg. in car G	—	5 39,65	4 39,50	9 4 39,47	—	38,51	—	+0,96	0,233	
1121	4.5	22 Hydræ θ	10 37,36	—	6 37,32	9 5 37,35	37,19	37,03	+0,16	+0,32	3,117	
1122	6	82 Canceri π^2	1 56,80	4 56,85	2 56,73	9 5 56,82	—	56,57	—	+0,25	3,326	
1123	5	Arg. in car a	5 33,05	—	2 33,26	9 6 33,10	—	32,55	—	+0,55	1,584	
1124	5	Arg. in car i	5 27,03	—	5 26,85	9 7 26,89	—	27,43	—	-0,54	1,376	
1125	4	38 Lyncis p	8 22,00	1 22,00	2 21,92	9 8 22,00	22,07	21,45	-0,07	+0,55	3,767	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich. Catalogue.	A. S. Catalogue.	Difference from		Annual Precession	
	No. 1831		No. 1832		No. 1833					Green.	A. S. C.		
	i	u	i	u	i	u				i	u		i
1081	—	—	4 32	50,09	—	—	70 32	50,09	32 47,36	+	2,73	+12,955	
1082	—	—	3 1	52,62	—	—	74 1	52,62	1 45,26	+	7,36	12,986	
1083	—	—	5 1	59,65	—	—	61 1	59,65	2 0,07	—	0,42	13,046	
1084	2 50	25,01	3 50	26,88	—	—	118 50	26,13	50 29,71	—	3,58	13,077	
1085	5 5	27,76	1 5	27,42	—	—	117 5	27,70	5 26,80	+	0,90	13,104	
1086	—	—	5 0	8,64	—	—	72 0	8,64	0 4,79	+	3,85	13,123	
1087	—	—	5 26	11,34	—	—	61 26	11,34	26 12,18	—	0,84	13,245	
1088	—	—	5 8	6,02	—	—	72 8	6,02	8 3,90	+	2,12	13,267	
1089	5 25	10,11	11 25	11,17	13 25	9,78	83 25	10,37	25 12,61	25 9,81	-2,24	+ 0,56	13,307
1090	—	—	5 44	11,43	—	—	77 44	11,43	44 13,26	44 12,60	-1,83	- 1,17	13,322
1091	—	—	—	—	—	—	168	Invisible	21 9,78	—	—	13,339	
1092	6 18	18,23	4 18	15,44	11 18	17,63	41 18	17,33	18 16,41	18 17,50	+0,97	- 0,12	13,380
1093	—	—	5 2	16,49	—	—	74 2	16,49	2 19,01	—	2,52	13,395	
1094	3 46	42,08	—	—	—	—	73 46	42,08	46 46,92	—	4,89	13,416	
1095	5 2	20,65	2 2	22,54	—	—	117 2	21,19	2 17,40	+	3,79	13,426	
1096	—	—	—	—	—	—	168	Invisible	27 15,74	—	—	13,462	
1097	5 29	48,17	3 29	47,45	11 29	47,35	77 29	47,58	29 49,34	29 49,69	-1,76	- 2,11	13,487
1098	4 12	57,60	1 12	56,92	—	—	71 12	57,54	12 54,77	—	2,77	13,512	
1099	5 11	9,85	5 11	9,31	8 11	8,70	42 11	9,19	11 8,32	11 11,48	+0,87	- 2,29	13,668
1100	5 53	33,85	—	—	—	—	64 53	33,85	53 26,80	—	7,05	13,719	
1101	5 34	57,41	6 34	56,67	—	—	148 34	57,01	34 59,70	—	2,69	13,721	
1102	4 26	31,64	5 26	32,29	—	—	148 26	32,00	26 37,91	—	5,91	13,875	
1103	5 52	56,76	5 52	57,05	—	—	50 52	56,91	52 52,91	+	4,00	13,903	
1104	5 14	28,74	—	—	—	—	84 14	28,74	14 28,57	+	0,17	13,987	
1105	5 25	55,86	5 25	55,70	—	—	136 25	55,78	26 0,43	—	4,65	14,067	
1106	—	—	6 39	34,39	—	—	78 39	34,39	39 38,06	39 39,50	-3,67	- 5,11	14,082
1107	—	—	5 40	50,69	—	—	62 40	50,69	40 51,76	—	1,07	14,096	
1108	—	—	5 51	16,31	—	—	71 51	16,31	51 11,34	+	4,97	14,141	
1109	5 16	46,61	—	—	—	—	67 16	46,61	16 48,47	16 44,68	-1,86	+ 1,93	14,146
1110	5 43	38,07	5 43	33,15	—	—	155 43	35,61	43 38,21	—	2,60	14,156	
1111	—	—	4 19	33,38	—	—	67 19	33,38	19 32,65	+	0,73	14,207	
1112	—	—	—	—	—	—	115 11	—	11 3,78	—	—	14,208	
1113	—	—	4 6	32,13	—	—	98 6	32,13	6 32,85	—	0,72	14,251	
1114	5 45	27,36	5 45	28,03	—	—	132 45	27,70	45 23,98	+	3,72	14,280	
1115	5 41	3,51	2 41	3,55	—	—	119 41	3,52	40 59,77	+	3,75	14,341	
1116	1 19	56,79	4 19	55,04	—	—	74 19	55,39	19 47,95	+	7,44	14,355	
1117	—	—	5 1	43,21	—	—	68 1	43,21	1 42,16	+	1,05	14,411	
1118	5 17	31,26	6 17	31,35	—	—	35 17	31,30	17 27,17	+	4,13	14,412	
1119	1 25	33,32	4 25	32,53	—	—	96 25	32,69	25 28,63	+	4,06	14,419	
1120	5 55	37,78	5 55	37,28	—	—	161 55	37,53	55 41,61	—	4,08	14,456	
1121	5 58	54,01	6 58	52,81	8 58	53,29	86 58	53,58	58 52,50	58 52,25	+1,08	+ 1,33	14,509
1122	—	—	5 21	59,09	—	—	74 21	59,09	21 53,71	+	5,38	14,528	
1123	5 16	50,61	—	—	5 16	49,07	148 16	49,34	17 6,50	—	16,66	14,568	
1124	4 37	42,60	4 37	42,89	—	—	151 37	42,74	37 52,91	—	10,17	14,623	
1125	5 29	30,43	2 29	30,01	6 29	30,35	52 29	30,33	29 30,27	29 24,00	+0,06	+ 6,33	14,672

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^h Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831.	No. 1832	No. 1833				Green.	A. S.	
1126	6	23 Hydræ K ²	—	6 21,19	—	9 8 21,19	—	20,90	—	+0,29	2,978
1127	6	24 Hydræ	—	5 27,18	—	9 8 27,18	—	27,53	—	-0,35	2,940
1128	7	Leonis	—	6 44,06	—	9 8 44,07	—	45,71	—	+0,36	3,265
1129	5	Arg. in vel l	6 59,80	6 60,07	—	9 8 59,92	—	59,64	—	+0,28	2,363
1130	6	83 Cancri g	—	6 35,74	2 35,66	9 9 35,73	—	35,48	—	+0,25	3,369
1131	4.5	40 Lyncis r	7 47,99	—	6 47,84	9 10 47,94	48,13	47,89	-0,19	+0,05	3,701
1132	7	Leonis	—	5 23,39	—	9 11 23,41	—	25,09	—	-1,68	3,523
1133	2	Argus β	4 19,45	1 19,80	7 19,73	9 11 19,59	—	20,46	—	-0,87	0,729
1134	5.6	26 Hydræ M ²	—	5 41,07	—	9 11 41,07	—	40,54	—	+0,53	2,800
1135	5	Draconis	—	2 26,26	5 25,55	9 12 26,46	—	29,24	—	-2,78	9,478
1136	5.6	27 Hydræ	—	—	5 16,98	9 12 16,98	—	16,63	—	+0,35	2,929
1137	2	Argus r	5 35,64	4 36,16	6 36,01	9 12 35,81	—	33,38	—	+1,93	1,609
1138	5	Pixid Naut θ	10 3,75	4 3,69	—	9 14 3,73	—	3,34	—	+0,39	2,659
1139	7	Hydræ	—	6 49,98	—	9 14 49,98	—	49,25	—	+0,73	3,160
1140	5	1 Leonis x	9 51,40	1 51,49	—	9 14 51,41	—	51,25	—	+0,16	3,516
1141	7	Leonis	—	6 17,46	—	9 15 17,47	—	17,94	—	-0,47	3,397
1142	5.6	Pixid Naut λ	—	5 56,52	—	9 15 56,50	—	56,32	—	+0,18	2,599
1143	7	Leonis	—	6 13,16	—	9 16 13,17	—	13,12	—	+0,05	3,341
1144	3	Argus x	6 55,04	—	4 55,23	9 16 55,09	—	55,01	—	+0,08	1,854
1145	6	28 Hydræ A	—	5 0,21	—	9 17 0,21	—	59,37	—	+0,84	3,001
1146	4	23 Ursæ Maj h	8 11,57	—	3 11,43	9 18 11,56	11,74	10,80	-0,18	+0,76	4,831
1147	2	30 Hydræ a	15 19,91	26 19,97	50 20,08	9 19 20,01	19,97	19,59	+0,04	+0,42	2,948
1148	5	24 Ursæ Maj d	5 23,50	—	5 27,30	9 19 28,00	—	29,37	—	-1,37	5,512
1149	6.7	2 Leonis ω	—	4 27,32	—	9 19 27,33	27,38	26,60	-0,05	+0,73	3,217
1150	6.7	3 Leonis	—	7 32,18	—	9 19 32,18	—	31,76	—	+0,42	3,203
1151	5.6	31 Hydræ τ ¹	—	6 37,41	—	9 20 37,41	—	36,59	—	+0,82	3,038
1152	3	25 Ursæ Maj θ	5 34,44	2 34,45	—	9 21 34,46	34,58	33,90	-0,12	+0,56	4,057*
1153	4.5	4 Leonis λ	5 7,62	—	6 7,43	9 22 7,52	7,45	7,29	+0,07	+0,23	3,441
1154	5	5 Leonis ζ	2 53,06	4 53,04	6 53,04	9 22 53,05	53,09	52,98	-0,04	+0,07	3,249
1155	6	6 Leonis η	—	6 56,90	—	9 22 56,90	—	55,95	—	+0,95	3,224
1156	5	Arg. in car n	4 5,21	1 5,28	4 5,70	9 23 5,33	—	8,36	—	-3,03	1,320
1157	6	32 Hydræ τ ²	1 24,68	6 25,11	—	9 23 25,05	—	24,62	—	+0,23	3,062
1158	5	10 Leonis Min b	—	3 54,43	6 54,38	9 23 54,45	54,54	54,48	-0,09	-0,03	3,706
1159	4.5	Argus α	4 5,99	1 5,72	5 5,61	9 24 5,75	—	5,71	—	+0,04	2,369
1160	5	Arg. in vel N	8 7,42	3 7,29	—	9 26 7,36	—	5,86	—	+1,50	1,822
1161	6	33 Hydræ	—	5 9,81	—	9 26 9,81	—	9,14	—	+0,67	2,993
1162	6.7	7 Leonis	2 41,49	6 41,55	1 41,41	9 26 41,55	—	41,19	—	+0,36	3,292
1163	6.7	8 Leonis	—	7 45,97	1 45,99	9 27 45,99	—	45,61	—	+0,38	3,323
1164	7	9 Leonis	—	6 12,34	—	9 28 12,36	—	11,65	—	+0,71	3,459
1165	5.6	10 Leonis	—	6 20,34	—	9 28 20,34	20,27	19,67	+0,07	+0,67	3,178
1166	7	11 Leonis	—	5 50,70	—	9 28 50,71	—	50,50	—	+0,21	3,289
1167	5	Arg. in car h	10 34,61	—	2 34,74	9 29 34,62	—	34,62	—	-0,00	1,738
1168	5.6	2 Sextantis b	—	6 41,34	—	9 29 41,34	—	41,02	—	+0,32	3,145
1169	5	35 Hydræ	10 16,71	—	3 16,46	9 31 16,70	—	16,03	—	+0,67	3,063
1170	6	13 Leonis	—	6 57,64	—	9 31 57,65	—	57,28	—	+0,37	3,473

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	1831		1832		1833					Green. A. S. C		
	No.	"	No.	"	No.	"				"	"	
1126	3	20 24,41	2	39 23,96	—	—	95 39 24,23	39 21,45	+	2,78	+14,673	
1127	—	—	5	2 49,17	—	—	98 2 49,17	2 48,17	+	1,00	14,680	
1128	—	—	5	17 56,20	—	—	77 47 56,20	47 57,32	—	1,12	14,695	
1129	4	52 23,13	—	—	—	—	127 52 23,13	52 23,79	—	5,66	14,713	
1130	—	—	5	35 13,37	—	—	71 35 13,37	35 9,65	+	3,72	14,746	
1131	5	54 13,75	3	54 13,00	7	54 11,93	54 54 12,75	54 5,01	+4,44	+ 7,74	14,817	
1132	—	—	5	7 22,29	—	—	64 7 22,29	—	—	—	14,854	
1133	5	1 29,66	—	—	—	—	159 1 29,66	1 45,81	—	16,15	14,855	
1134	—	—	5	16 11,12	—	—	101 16 11,12	16 7,61	+	3,51	14,870	
1135	—	—	—	—	7	56 27,48	7 56 27,48	56 34,30	—	6,82	14,904	
1136	—	—	5	50 48,06	—	—	98 50 48,06	50 42,96	+	5,10	14,905	
1137	2	34 20,50	5	34 19,78	—	—	148 34 19,98	34 24,37	—	4,39	14,924	
1138	5	15 18,89	2	15 16,58	—	—	115 15 18,20	15 19,17	—	0,97	15,009	
1139	2	3 52,39	3	3 51,69	—	—	84 3 51,97	3 53,14	—	1,17	15,052	
1140	—	—	8	5 55,70	2	5 54,97	63 5 55,55	5 56,45	—	0,90	15,054	
1141	—	—	3	29 33,58	—	—	69 29 33,58	29 17,26	+	16,32	15,089	
1142	—	—	5	7 5,24	—	—	118 7 5,24	7 8,68	—	3,44	15,118	
1143	—	—	5	41 38,47	—	—	72 41 38,47	41 38,04	+	0,43	15,133	
1144	5	17 45,69	2	17 47,57	—	—	144 17 46,23	17 51,37	—	5,14	15,175	
1145	—	—	6	23 50,75	—	—	94 23 50,75	23 45,13	+	5,62	15,177	
1146	7	12 37,72	2	12 35,01	—	—	26 12 37,12	12 32,98	-0,40	+ 4,19	15,242	
1147	10	56 1,76	42	56 0,43	66	56 0,78	97 56 0,74	56 1,40	-2,52	- 0,66	15,310	
1148	5	26 21,25	—	—	—	—	19 26 21,24	26 19,20	+	2,04	15,314	
1149	—	—	5	13 1,08	—	—	80 13 1,08	12 57,35	12 52,12	+8,73	+ 8,96	15,316
1150	—	—	5	5 0,60	—	—	81 5 0,60	4 53,85	+	6,75	15,321	
1151	—	—	5	2 20,05	—	—	92 2 20,05	2 15,48	+	4,57	15,382	
1152	3	33 41,97	8	33 43,11	—	—	37 33 42,66	33 46,31	-1,71	- 3,65	16,934*	
1153	2	17 44,73	5	17 41,89	—	—	66 17 42,69	17 44,76	-2,07	- 2,14	15,466	
1154	1	57 35,18	6	37 35,40	7	57 33,68	77 57 34,52	57 33,61	-1,54	+ 0,91	15,508	
1155	2	32 53,36	3	32 54,10	—	—	79 32 53,80	32 48,58	+	5,22	15,511	
1156	2	12 11,20	3	12 12,16	—	—	154 12 11,78	12 19,17	—	7,39	15,526	
1157	—	—	6	26 54,90	—	—	90 26 54,90	26 51,91	+	2,99	15,538	
1158	5	51 38,91	—	—	5	51 39,12	52 51 39,01	51 35,48	-2,57	+ 3,53	15,564	
1159	5	44 8,98	3	44 7,30	—	—	129 44 8,35	44 7,51	+	0,34	15,577	
1160	9	17 45,43	4	17 44,49	—	—	146 17 45,15	17 48,45	—	3,30	15,687	
1161	1	10 14,20	4	10 12,54	—	—	95 10 12,87	10 4,91	+	7,96	15,688	
1162	1	52 32,55	4	52 32,59	—	—	74 52 32,58	52 25,78	+	6,80	15,717	
1163	—	—	5	48 46,88	—	—	72 48 46,88	48 44,52	+	2,36	15,775	
1164	—	—	5	34 48,31	—	—	64 34 48,31	34 43,99	+	4,32	15,798	
1165	2	24 55,71	3	24 52,97	—	—	82 24 54,14	24 50,79	+0,35	+ 3,35	15,806	
1166	—	—	5	53 54,57	—	—	74 53 54,57	53 50,49	+	4,08	15,833	
1167	5	28 56,13	3	28 56,38	—	—	148 28 56,22	29 2,97	—	6,75	15,875	
1168	5	35 48,42	1	35 47,76	—	—	84 35 48,31	35 44,06	+	4,25	15,878	
1169	5	23 7,11	—	—	7	23 4,97	90 23 5,99	23 1,36	+	4,63	15,963	
1170	3	19 31,99	2	19 33,54	—	—	63 19 32,61	19 37,97	—	5,36	15,998	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion	
			No. 1831 No. 1832 No. 1833						Green.	A. S.		
			s.	s.	s.							
1171	4	14 Leonis	o	8 10,66	4 10,71	6 10,78	9 32 10,72	10,73	10,18	-0,01	+0,54	3,219
1172	5	38 Hydræ	x	5 15,34	—	6 15,42	9 32 15,38	—	15,21	—	+0,17	2,874
1173	7	Leonis	—	—	5 57,64	2 57,52	9 33 57,62	—	56,87	—	+0,75	3,373
1174	6	16 Leonis	ψ	3 34,50	4 34,56	—	9 34 34,53	—	34,28	—	+0,25	3,277
1175	3	17 Leonis	ε	14 18,06	2 18,12	14 18,12	9 36 18,11	18,21	17,73	-0,10	+0,38	3,426
1176	6	Antl. Pneum	θ	—	5 43,14	—	9 36 43,12	—	42,95	—	+0,17	2,669
1177	6	18 Leonis	—	—	5 19,90	—	9 37 19,91	—	19,57	—	+0,34	3,242
1178	7	19 Leonis	—	—	6 23,71	—	9 38 23,72	—	23,34	—	+0,38	3,238
1179	4.5	29 Ursæ Maj	υ	10 58,46	1 58,35	5 58,16	9 38 58,40	58,46	59,06	-0,06	-0,66	4,356*
1180	7	20 Leonis	—	2 25,18	4 25,25	1 24,90	9 40 25,19	—	25,00	—	+0,19	3,377
1181	5	30 Ursæ Maj.	Φ	7 37,13	—	6 37,07	9 40 37,15	—	37,79	—	-0,64	4,153
1182	5	Arg. in car	l	5 37,67	—	6 38,39	9 40 38,00	—	36,40	—	+1,60	1,648
1183	6	4 Sextantis	s	—	5 45,50	—	9 41 45,50	—	43,59	—	+1,91	3,136
1184	6	22 Leonis	g	—	5 19,96	—	9 42 19,97	—	19,14	—	+0,83	3,424
1185	6	6 Sextantis	t	—	6 46,11	—	9 42 46,11	—	45,86	—	+0,25	3,023
1186	3.4	Argus	υ	5 54,01	1 54,31	2 54,34	9 42 54,08	—	54,07	—	+0,01	1,505
1187	3	24 Leonis	μ	6 11,59	—	6 11,59	9 43 11,60	11,77	11,02	-0,17	+0,58	3,448
1188	5	39 Hydræ	υ ¹	6 24,04	—	4 24,10	9 43 24,06	—	24,09	—	-0,03	2,880
1189	7	7 Sextantis	A	—	6 32,14	—	9 43 32,14	—	31,80	—	+0,34	3,111
1190	6	8 Sextantis	d	—	6 11,54	—	9 44 11,54	—	11,43	—	+0,11	2,972
1191	7	9 Sextantis	—	—	6 19,73	—	9 45 19,73	—	19,30	—	+0,43	3,143
1192	6	10 Sextantis	—	—	6 31,35	—	9 47 31,35	—	31,32	—	+0,03	3,193
1193	5.6	27 Leonis	υ	—	8 10,76	1 10,87	9 49 10,78	10,77	10,44	+0,01	+0,34	3,238
1194	6	11 Sextantis	—	—	6 13,33	—	9 49 13,33	—	12,90	—	+0,43	3,184
1195	6.7	Sextantis	—	—	6 0,06	—	9 51 0,06	—	59,92	—	+0,14	3,120
1196	4	Argus	Φ	5 58,78	2 58,90	6 58,58	9 50 58,63	—	58,59	—	+0,04	2,095
1197	4.5	29 Leonis	π	7 19,77	—	6 19,83	9 51 19,81	19,90	19,68	-0,09	+0,13	3,179
1198	6.7	Leonis	—	—	6 26,12	—	9 53 26,13	—	25,97	—	+0,16	3,362
1199	6	Hydræ	—	—	6 24,80	—	9 51 24,80	—	24,64	—	+0,16	2,914
1200	7	Leonis	—	—	8 8,36	1 8,20	9 55 8,35	—	7,02	—	+0,43	3,221
1201	7	13 Sextantis	e	—	5 26,15	—	9 55 26,15	—	25,63	—	+0,52	3,117
1202	7	Leonis	—	1 33,46	6 33,50	—	9 56 33,50	—	33,27	—	+0,23	3,272
1203	5.6	40 Hydræ	υ ²	—	3 56,85	—	9 56 56,84	—	56,70	—	+0,14	2,920
1204	5	21 Leonis Min	d	10 30,12	—	2 29,54	9 57 30,03	—	29,56	—	+0,47	3,564
1305	6	14 Sextantis	C	—	2 0,14	—	9 58 0,14	—	59,47	—	+0,67	3,145
1206	3.4	30 Leonis	η	8 10,08	3 10,15	2 9,92	9 58 10,07	9,93	9,52	+0,14	+0,55	3,283
1207	5	31 Leonis	A	5 59,20	3 59,04	4 59,02	9 58 59,11	58,99	58,74	+0,12	+0,37	3,197
1208	5	15 Sextantis	f	2 20,19	4 20,34	6 20,38	9 59 20,33	—	21,32	—	-1,49	3,073
1209	1	32 Leonis	z	7 25,19	36 25,08	40 25,09	9 59 25,11	25,14	24,73	-0,03	+0,38	3,221
1210	6	16 Sextantis	—	—	6 26,52	—	10 0 26,53	—	25,83	—	+0,70	3,150
1211	6	17 Sextantis	g ¹	—	—	4 46,85	10 1 46,85	—	46,99	—	-0,14	2,980
1212	4.5	41 Hydræ	λ ²	11 24,06	3 24,26	—	10 2 24,09	24,13	23,60	+0,04	+0,49	2,934
1213	6	18 Sextantis	g ²	—	—	6 34,86	10 2 34,86	—	34,66	—	+0,20	2,981
1214	6	34 Leonis	—	—	6 35,80	—	10 2 35,82	—	35,23	—	+0,59	3,233
1215	6	Sextantis	—	—	6 54,51	—	10 2 54,51	—	—	—	—	2,994

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
	' "	' "	' "				' "	' "	
1171	7 20 51,93	1 20 50,18	13 29 51,60	79 20 51,64	20 50,14	20 48,11	+1,50	+ 3,53	+16,010
1172	5 34 22,51	3 34 19,27	—	103 34 21,30	—	34 25,77	—	— 4,47	16,015
1173	1 2 32,74	4 2 31,78	—	69 2 31,98	—	2 34,01	—	— 2,03	16,103
1174	5 12 52,03	—	—	75 12 52,03	—	12 49,76	—	— 2,27	16,135
1175	9 27 24,28	10 27 23,33	10 27 22,59	65 27 23,40	27 22,44	27 19,47	+0,96	+ 3,93	16,224
1176	5 0 11,81	—	—	117 0 11,81	—	0 10,91	—	— 0,90	16,247
1177	—	5 25 10,80	—	77 25 10,80	—	25 5,51	—	— 5,29	16,277
1178	2 39 27,39	3 39 29,62	—	77 39 28,73	—	39 29,21	—	— 0,48	16,331
1179	5 10 40,08	3 10 38,27	5 10 38,62	30 10 39,10	10 34,06	10 37,42	+5,04	+ 1,68	16,359
1180	1 2 25,16	4 2 26,84	—	68 2 26,51	—	2 29,90	—	— 3,39	16,433
1181	7 9 24,15	—	5 9 24 11	35 9 24,14	—	9 22,00	—	— 2,14	16,443
1182	5 44 4,62	—	6 44 3,35	151 44 4,03	—	44 8,53	—	— 4,50	16,446
1183	—	5 52 29,47	—	84 52 29,47	—	52 15,19	—	— 14,28	16,499
1184	3 48 50,35	2 48 52,98	—	64 48 51,40	—	48 47,85	—	— 3,55	16,523
1185	2 27 31,81	3 27 32,17	—	93 27 32,03	—	27 30,18	—	— 1,85	16,550
1186	6 17 40,52	—	—	154 17 40,52	—	17 38,42	—	— 2,10	16,560
1187	5 12 19,22	5 12 18,06	11 12 19,03	63 12 18,84	12 21,52	12 19,23	-2,68	- 0,39	16,570
1188	5 3 40,14	1 3 39,70	—	104 3 40,07	—	3 37,85	—	— 2,22	16,582
1189	—	5 45 57,33	—	86 45 57,33	—	45 54,61	—	— 1,72	16,588
1190	—	5 19 1,68	—	97 19 1,68	—	19 4,70	—	— 3,02	16,621
1191	—	5 16 1,08	—	84 16 1,98	—	15 55,08	—	— 6,00	16,675
1192	1 16 32,56	4 16 30,36	—	80 16 30,80	—	16 21,89	—	— 8,91	16,781
1193	3 45 26,72	2 45 26,10	—	76 45 26,47	45 26,75	45 21,55	-0,28	+ 4,92	16,860
1194	5 53 16,96	5 53 16,37	—	80 53 16,67	—	53 14,08	—	— 2,59	16,862
1195	—	—	—	85 43 —	—	43 54,70	—	—	16,946
1196	5 46 12,74	1 46 9,72	5 46 10,01	143 46 11,23	—	46 16,53	—	— 5,30	16,946
1197	5 9 12,76	—	5 9 12,15	81 9 12,45	9 11,49	9 9,61	+0,96	+ 2,84	16,961
1198	—	5 14 37,26	—	67 14 37,26	—	14 36,64	—	— 0,62	17,058
1199	1 29 23,21	3 29 22,71	—	102 29 22,83	—	29 24,85	—	— 2,02	17,103
1200	—	5 33 42,34	—	77 33 42,34	—	33 41,68	—	— 0,66	17,136
1201	—	5 59 4,70	—	85 59 4,70	—	59 3,99	—	— 0,50	17,149
1202	—	5 25 42,44	—	73 25 42,44	—	25 42,23	—	— 0,21	17,200
1203	—	5 15 10,82	—	102 15 10,82	—	15 6,14	—	— 4,68	17,218
1204	3 56 24,50	2 56 23,06	—	53 56 23,93	—	56 22,11	—	— 1,82	17,241
1205	—	5 37 18,20	—	83 37 18,20	—	37 17,25	—	— 0,95	17,264
1206	1 25 17,04	4 25 14,36	2 25 14,83	72 25 14,67	25 17,44	25 16,64	-2,57	- 1,77	17,271
1207	4 10 55,15	—	—	79 10 55,15	10 55,67	10 53,15	-0,52	+ 2,00	17,308
1208	—	—	3 33 15,61	89 33 15,61	—	33 11,33	—	— 4,28	17,325
1209	10 12 55,92	5 12 53,08	5 6 12 54,05	77 12 53,78	12 52,46	12 49,82	+1,32	+ 3,96	17,327
1210	5 0 28,81	—	—	83 0 28,81	—	0 27,01	—	— 1,80	17,372
1211	1 34 58,25	8 35 7,91	—	97 34 58,25	—	35 0,90	—	— 2,65	17,431
1212	5 31 35,70	2 31 36,25	—	101 31 35,86	31 37,00	31 33,81	-1,14	+ 2,05	17,457
1213	1 35 23,03	—	—	97 35 23,93	—	35 24,94	—	— 1,91	17,465
1214	—	5 49 8,36	—	75 49 8,36	—	—	—	—	17,465
1215	—	5 29 28,57	—	96 29 28,57	—	29 24,44	—	— 4,13	17,478

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832		Green ^h Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833					Green.	A. S.	
			s.	s.	s.	h. m.	s.			s.	s.	
1216	7	19 Sextantis	—	7 3,74	—	10 4 3,74	—	3,24	—	+0,50	+3,180	
1217	7	Leonis	—	6 14,33	—	10 5 14,35	—	14,23	—	+0,12	3,328	
1218	6	21 Sextantis	5 46,18	—	—	10 5 46,18	—	46,12	—	+0,06	2,986	
1219	3.4	33 Ursæ Maj. λ	9 55,97	5 56,00	—	10 6 56,02	56,12	55,91	-0,10	+0,11	3,675	
1220	6	Leonis	—	5 6,58	1 6,57	10 7 6,60	—	5,84	—	+0,76	3,281	
1221	4.5	36 Leonis ζ	6 20,00	—	13 20,02	10 7 20,03	20,02	19,67	+0,01	+0,36	3,358	
1222	6	37 Leonis	—	5 39,24	1 39,16	10 7 39,24	—	39,29	—	-0,05	3,232	
1223	4	Arg. in vel g	9 42,10	—	3 41,77	10 7 42,01	—	42,11	—	-0,10	2,516	
1224	6	22 Sextantis x	—	6 17,06	—	10 9 17,06	—	16,99	—	+0,07	2,989	
1225	4.5	Argus ω	6 44,34	—	2 44,41	10 9 44,31	—	43,72	—	+0,59	1,440	
1226	6	Antl. Pneum	—	6 26,20	—	10 10 26,18	—	25,88	—	+0,35	2,739	
1227	6	40 Leonis	—	6 35,02	—	10 10 35,04	—	34,57	—	+0,47	3,296	
1228	2	41 Leonis γ	6 41,99	4 41,93	16 42,05	10 10 42,03	41,96	41,61	+0,07	+0,42	3,300	
1229	5	Arg. in car q	4 29,23	2 28,93	6 29,43	10 11 29,22	—	27,62	—	+1,60	1,991	
1230	3	34 Ursæ Maj μ	7 17,48	—	6 17,31	10 12 17,44	17,53	17,43	-0,09	+0,01	3,620	
1231	6	23 Sextantis h	—	7 21,49	—	10 12 21,49	—	21,44	—	+0,05	3,101	
1232	6	42 Leonis	3 47,78	3 47,87	—	10 12 47,83	—	47,52	—	+0,31	3,239	
1233	6	43 Leonis z	—	—	—	10 14 —	—	12,73	—	—	3,145	
1234	5	Arg. in vel T	8 40,67	5 40,77	—	10 14 40,69	—	38,89	—	+1,80	3,215	
1235	4.5	Arg. in vel r	6 8,14	—	5 8,18	10 15 8,13	—	8,17	—	-0,04	2,558	
1236	4.5	30 Leonis Min f	6 15,89	—	4 15,57	10 16 15,79	15,91	14,90	-0,12	+0,89	3,473	
1237	6	44 Leonis b1	1 23,92	6 23,88	—	10 16 23,89	—	23,49	—	+0,40	3,167	
1238	4	42 Hydræ μ	7 58,36	—	4 58,28	10 17 58,32	58,31	57,66	+0,01	+0,66	2,903	
1239	6	26 Sextantis i	—	5 2,66	—	10 18 2,66	—	2,12	—	+0,54	3,067	
1240	4.5	31 Leonis Min g	5 8,76	—	6 8,66	10 18 8,74	8,62	7,33	-0,08	+1,41	3,511	
1241	6	27 Sextantis	2 17,90	5 17,75	—	10 18 17,79	—	17,62	—	+0,17	3,083	
1242	6	45 Leonis	—	6 46,41	—	10 18 46,43	—	46,14	—	+0,29	3,175	
1243	4.5	Antl. Pneum α	4 28,54	—	6 28,61	10 19 28,56	28,48	28,61	+0,08	-0,05	2,737	
1244	5	36 Ursæ Maj	6 49,10	4 49,18	2 49,01	10 19 49,15	49,07	50,16	+0,08	-1,01	3,935	
1245	6	Sextantis g	2 13,10	4 13,42	—	10 20 13,31	—	13,33	—	-0,52	3,040	
1246	6	28 Sextantis k	—	5 56,05	—	10 20 56,05	—	56,32	—	-0,27	3,050	
1247	5	Arg. in car J	—	6 3,08	—	10 21 2,91	—	0,76	—	+2,15	1,218	
1248	6	30 Sextantis l	—	5 42,34	—	10 21 42,34	—	42,02	—	+0,32	3,070	
1249	5.6	Antl. Pneum	—	—	6 44,09	10 21 44,05	—	43,85	—	+0,20	2,762	
1250	7	31 Sextantis u	—	—	6 50,26	10 21 50,26	—	49,45	—	+0,81	3,097	
1251	6	Antl. Pneum δ	—	—	6 52,04	10 21 52,00	—	51,76	—	+0,24	2,751	
1252	6	46 Leonis i	1 13,23	5 13,32	—	10 23 13,31	—	13,37	—	-0,06	3,215	
1253	7	32 Sextantis x	—	6 34,84	—	10 23 34,84	—	34,29	—	+0,55	3,121	
1254	4	47 Leonis p	14 57,63	4 57,74	5 57,71	10 23 57,67	57,61	57,47	+0,06	+0,20	3,166	
1255	5	37 Ursæ Maj. m	5 16,72	—	5 16,61	10 24 16,73	16,96	16,71	-0,23	+0,02	3,935	
1256	5.6	48 Leonis	5 2,02	2 2,22	—	10 26 2,08	1,94	1,56	+0,14	+0,52	3,141	
1257	6	44 Hydræ n	—	5 2,10	—	10 26 2,08	—	1,61	—	+0,47	2,843	
1258	4	Arg. in car p	7 4,24	—	4 4,08	10 26 4,15	—	4,79	—	-0,64	2,114	
1259	6	49 Leonis	—	6 13,31	—	10 26 13,31	—	12,76	—	+0,55	3,157	
1260	6	1 Hyd & Crat	—	6 4,92	—	10 28 4,90	—	4,70	—	+0,20	2,924	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
1216	—	4 33 30,96	—	84 33 30,96	—	33 24,97	+	5,99	+17,528
1217	—	5 59 53,40	—	67 59 53,40	—	0 0,85	—	7,45	17,577
1218	3 9 47,07	2 9 45,91	—	97 9 46,61	—	9 38,96	+	7,65	17,600
1219	6 15 1,46	13 14 59,91	8 14 59,02	46 14 59,99	15 0,96	15 1,36	-0,91	-1,37	17,647
1220	—	5 25 34,66	—	71 25 34,66	—	25 33,47	+	1,19	17,655
1221	6 44 55,96	—	7 44 56,44	65 44 56,22	44 58,76	44 49,87	-2,54	+ 6,85	17,664
1222	2 26 14,31	3 26 15,12	—	75 26 14,80	—	26 11,71	+	3,09	17,678
1223	5 17 32,76	—	5 17 29,02	131 17 30,89	—	17 25,05	+	5,84	17,681
1224	—	5 13 57,59	—	97 13 57,59	—	13 55,34	+	2,25	17,745
1225	5 12 16,09	—	—	159 12 16,39	—	12 28,09	—	12,00	17,765
1226	—	5 9 14,40	—	118 9 14,40	—	9 14,54	—	0,14	17,792
1227	—	5 40 49,26	—	69 40 49,26	—	40 41,10	+	8,16	17,797
1228	6 18 42,29	11 18 40,07	8 18 41,35	69 18 41,00	18 41,80	18 40,22	-0,80	+ 0,78	17,802
1229	4 29 41,48	—	4 29 43,23	159 29 42,35	—	29 47,65	—	5,30	17,834
1230	5 39 31,21	—	6 39 30,46	47 39 30,80	39 33,02	39 28,95	-2,22	+ 1,85	17,865
1231	—	5 52 6,27	—	86 52 6,27	—	51 58,48	+	7,79	17,868
1232	2 10 45,57	5 10 47,89	—	74 10 47,16	—	10 41,83	+	5,33	17,885
1233	—	5 36 25,47	—	82 36 25,47	—	36 23,56	+	1,94	17,941
1234	5 12 1,78	—	5 12 0,22	145 12 1,00	—	11 58,62	+	2,38	17,959
1235	5 48 26,43	—	6 48 26,19	130 48 26,30	—	48 22,69	+	3,61	17,978
1236	4 21 4,12	13 21 7,28	8 21 7,04	55 21 6,70	21 4,01	21 3,29	+2,69	+ 3,41	18,019
1237	1 21 51,24	4 21 54,48	—	80 21 53,83	—	21 47,43	+	6,40	18,025
1238	5 58 53,03	—	5 58 52,49	105 58 52,76	58 52,34	58 50,37	+0,42	+ 2,39	18,085
1239	—	5 8 15,18	—	90 8 15,18	—	8 10,55	+	4,63	18,083
1240	5 26 5,86	—	5 26 5,55	52 26 5,71	26 4,77	26 3,63	+0,94	+ 2,08	18,090
1241	—	5 32 6,04	—	93 32 6,04	—	32 7,13	—	1,69	18,097
1242	—	5 23 1,81	—	79 23 1,81	—	22 59,59	+	2,22	18,115
1243	5 12 56,66	2 12 55,37	5 12 55,38	120 12 55,91	12 51,67	12 52,02	+4,24	+ 3,89	18,112
1244	5 9 37,73	—	8 9 38,45	33 9 38,17	9 41,63	9 37,66	-3,46	+ 0,51	18,154
1245	—	4 53 5,40	—	92 53 5,40	—	—	—	—	18,170
1246	—	5 52 53,75	—	91 52 53,75	—	52 52,37	+	1,33	18,196
1247	4 10 37,87	—	—	163 10 37,87	—	11 5,40	—	27,53	18,201
1248	3 46 41,76	2 46 43,82	—	89 46 42,54	—	46 35,24	+	7,34	18,223
1249	—	5 48 21,67	—	118 48 21,67	—	48 22,43	—	0,41	18,225
1250	—	5 59 23,80	—	86 59 23,80	—	59 17,10	+	6,40	18,228
1251	—	5 44 59,59	—	119 44 59,59	—	44 56,94	+	2,65	18,230
1252	—	4 0 14,48	—	75 0 14,48	—	0 8,25	+	6,23	18,278
1253	—	4 29 44,96	—	84 29 44,96	—	29 38,16	+	6,80	18,291
1254	5 49 53,31	—	5 49 57,31	79 49 57,81	49 52,80	49 50,25	+5,61	+ 7,56	18,305
1255	5 3 20,46	—	6 3 19,81	32 3 20,10	3 20,02	3 18,08	+0,08	+ 2,02	18,315
1256	—	5 11 3,32	—	82 11 3,32	11 2,30	10 59,69	+1,02	+ 3,63	18,378
1257	—	5 52 51,15	—	112 52 51,15	—	52 51,16	—	0,01	18,378
1258	6 49 21,06	6 49 21,53	—	150 49 21,74	—	49 24,98	—	3,24	18,381
1259	—	4 29 4,64	—	80 29 4,64	—	28 58,00	+	0,64	18,384
1260	3 28 37,07	2 28 37,61	—	105 28 37,29	—	28 30,97	+	6,32	18,440

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Greenw. A. S. January 1, 1832		Greenw. Catal.	A. S. Catal.	Difference from		Annual Precession
			No. 1831	No. 1832	No. 1833	h. m. s.	s.			Green.	A. S.	
1261	4	37 Leonis Min <i>l</i>	9 14,74	1 14,89	3 14,83	10 29 14,79	14,86	14,58	-0,07	+0,21	+3,401	
1262	6	Antl. Pucum	—	5 21,20	—	10 29 21,17	—	21,14	—	+0,03	2,811	
1263	6.7	50 Leonis	—	4 53,45	1 53,47	10 29 53,48	—	53,26	—	+0,22	3,225	
1264	5	Arg in vel <i>p</i>	6 15,96	—	6 15,62	10 30 15,75	—	15,79	—	-0,04	2,514	
1265	5	2 Hyd & Crat ϕ^j	7 24,15	—	4 24,27	10 30 24,19	—	24,27	—	-0,08	2,922	
1266	5	Ursæ Maj.	1 54,76	3 54,98	5 54,44	10 30 54,61	—	54,89	—	-0,28	4,459	
1267	6	33 Sextantis <i>m</i>	—	6 51,63	—	10 32 51,63	—	50,95	—	+0,68	3,061	
1268	5	Chamæl <i>y</i>	—	—	—	10 33 Invisibile	—	23,66	—	—	0,801	
1269	5.6	40 Leonis Min <i>n</i>	5 47,60	2 47,48	—	10 33 47,57	—	47,01	—	+0,56	3,321	
1270	6	34 Sextantis	1 57,34	6 56,98	—	10 33 57,03	—	56,33	—	+1,70	3,106	
1271	5.6	41 Leonis Min <i>z</i>	—	6 16,22	—	10 34 16,24	—	15,53	—	+0,71	3,287	
1272	7	35 Sextantis	—	5 37,36	—	10 34 37,36	—	36,39	—	+0,37	3,116	
1273	5	Argus θ^1	5 18,13	—	3 17,90	10 36 18,00	—	15,97	—	+2,03	2,106	
1274	4.5	42 Leonis Min <i>n</i>	8 30,30	1 30,54	3 30,35	10 36 30,34	30,37	29,64	-0,03	+0,70	3,361	
1275	6	36 Sextantis <i>n</i>	—	4 30,39	—	10 36 30,39	—	29,61	—	+0,78	3,096	
1276	2.3	Argus θ^2	15 58,96	—	3 59,41	10 36 59,02	—	57,15	—	+1,87	2,117	
1277	6	37 Sextantis θ^1	1 21,07	6 20,85	2 20,58	10 37 20,82	20,60	20,04	+0,22	+0,78	3,128	
1278	6	51 Leonis <i>mz</i>	—	6 21,04	—	10 37 21,06	—	20,93	—	+0,13	3,238	
1279	6	52 Leonis <i>k</i>	—	6 31,01	1 30,92	10 37 31,01	—	30,66	—	+0,35	3,195	
1280	7	38 Sextantis θ^2	—	1 34,81	—	10 38 34,81	34,75	33,92	+0,06	+0,89	3,127	
1281	2	Argus <i>n</i>	6 34,11	—	6 34,17	10 38 34,09	—	31,60	—	-0,51	2,300	
1282	6	3 Hyd & Crat b^1	—	6 38,68	—	10 38 38,67	—	38,28	—	+0,39	2,931	
1283	3	Argus μ	7 33,72	—	4 33,96	10 39 33,82	—	33,50	—	+0,32	2,548	
1284	6	53 Leonis <i>sl</i>	1 24,69	5 25,23	2 25,37	10 40 25,22	—	25,07	—	+0,15	3,160	
1285	6	44 Leonis Min	—	1 38,86	5 38,63	10 40 38,70	—	37,66	—	+1,04	3,218	
1286	6	40 Sextantis <i>p</i>	—	1 46,30	5 46,37	10 40 46,36	—	45,48	—	+0,88	3,043	
1287	4	4 Hyd & Crat <i>v</i>	2 20,50	3 20,60	3 20,45	10 41 20,50	20,49	20,01	+0,01	+0,49	2,945	
1288	6	41 Sextantis <i>r</i>	—	5 52,65	—	10 41 52,65	—	52,40	—	+0,25	3,005	
1289	4.5	46 Leonis Min <i>p</i>	8 53,76	—	4 53,83	10 43 53,80	53,60	53,47	+0,20	+0,33	3,375	
1290	5	Chamæl δ^2	—	—	—	10 44 Invisibile	—	5,82	—	—	0,689	
1291	5	45 Ursæ Maj <i>w</i>	6 16,93	—	4 16,59	10 44 16,84	—	16,38	—	+0,46	3,488	
1292	5.6	6 Hyd & Crat b^3	1 16,61	2 16,82	—	10 45 16,74	—	16,63	—	+0,11	2,919	
1293	4.5	54 Leonis	8 30,29	—	5 30,22	10 46 30,28	30,36	30,16	-0,08	+0,12	3,271	
1294	5	Arg. in car <i>u</i>	6 42,05	—	5 42,11	10 46 42,03	—	43,18	—	-1,10	2,396	
1295	6	55 Leonis <i>u</i>	—	5 3,93	—	10 47 3,93	3,75	3,39	+0,18	+0,54	3,080	
1296	7	56 Leonis	—	5 17,95	—	10 47 17,95	17,84	17,60	+0,11	+0,35	3,120	
1297	6	50 Leonis Min	—	5 26,77	—	10 47 26,79	—	26,07	—	+0,72	3,275	
1298	7	57 Leonis	—	—	5 33,46	10 47 33,46	—	33,34	—	+0,12	3,078	
1299	5	Antl. Pneu	7 54,23	—	5 54,42	10 48 54,29	—	54,41	—	-0,12	2,769	
1300	4	7 Hyd & Crat <i>z</i>	6 35,65	—	6 35,88	10 51 35,76	35,88	35,23	-0,07	+0,53	2,905*	
1301	2	48 Ursæ Maj. <i>\beta</i>	5 39,04	1 39,23	5 38,84	10 51 39,03	39,03	39,16	-0,00	-0,13	3,680	
1302	5	58 Leonis <i>d</i>	6 52,96	—	5 53,02	10 51 52,98	52,89	52,68	+0,09	+0,30	3,099	
1303	5.6	59 Leonis <i>c</i>	—	7 2,27	—	10 52 2,27	2,23	1,95	+0,04	+0,32	3,116	
1304	5.6	61 Leonis <i>s</i>	—	6 15,71	—	10 53 15,71	—	14,96	—	+0,75	3,058	
1305	1.2	50 Ursæ Maj. <i>\alpha</i>	6 17,49	19 17,38	21 17,20	10 53 17,37	17,46	17,38	-0,09	-0,01	3,811	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N P D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831		No. 1832		No. 1833					Green.	A. S. C.	
	°	'	°	'	°	'				"	"	
1261	5	9 14,62	2	9 13,18	6	9 14,21	57 9 14,21	9 12,05	9 10,55	+2,16	+ 3,66	+18,488
1262				5 32 37,15			116 32 37,15		32 36,67		+ 0,48	18,492
1263				5 59 58,79			72 59 58,79		0 0,49		- 1,70	18,510
1264	4	21 16,74	1	21 18,98			137 21 17,19		21 22,17		- 4,98	18,523
1265	5	0 24,89			6	0 24,44	106 0 24,65		0 19,23		+ 5,42	18,527
1266	5	2 59,12					20 2 59,12		2 53,20		+ 5,83	18,543
1267	3	51 40,12	2	51 38,30			90 51 39,40		51 35,88		+ 3,52	18,608
1268							167 Invisible		44 15,04			18,628
1269				5 47 36,80			62 47 36,80		47 40,09		- 3,29	18,638
1270				5 32 31,48			85 32 31,48		32 25,34		+ 6,14	18,642
1271				5 56 2,76			65 56 2,76		56 1,50		+ 1,26	18,653
1272				5 22 23,20			84 22 23,20		22 17,72		+ 5,48	18,664
1273	5	35 16,14					153 35 16,14		35 25,70		- 9,56	18,718
1274	0	26 2,68	7	26 3,58	8	26 4,30	58 26 3,60	26 6,46	26 3,52	-2,86	+ 0,08	18,723
1275				5 37 49,04			86 37 49,04		37 45,50		+ 3,54	18,724
1276	5	30 53,58			5	30 52,17	153 30 52,87		30 52,86		+ 0,01	18,739
1277	1	44 33,75	9	44 35 77			82 44 35,57	44 37,10	44 35,54	-1,53	+ 0,03	18,750
1278							70		13 25,95			18,750
1279							74		55 9,48			18,755
1280							82	46 8,90	46 5,96			18,787
1281	5	48 6,43					148 48 6,43		48 13,26		- 6,83	18,789
1282				5 24 45,81			106 24 45,81		24 41,11		+ 4,70	18,790
1283	6	31 59,47					138 31 59,47		31 45,50		+ 13,97	18,818
1284				5 34 0,31			78 34 0,34		33 58,62		+ 1,72	18,843
1285	1	8 23 70	4	8 27,25			61 8 26,94		8 28,80		- 1,66	18,849
1286				4 8 12,26			98 8 12,26		8 11,08		+ 1,18	18,853
1287	5	19 1,95					105 19 1,95	18 59,30	18 56,20	+2,65	+ 5,75	18,871
1288				5 0 31,92			98 0 31,92		0 32,17		- 0,25	18,886
1289	5	52 59,77			7	52 57,92	54 52 58,69	52 53,03	52 51,03	+5,66	+ 7,66	18,944
1290							Invisible		39 20,61			18,953
1291	3	53 3,86	2	55 3,62			45 55 3,77		55 2,60		+ 1,17	18,955
1292				5 14 9,10			109 14 9,10		13 59,07		+ 10,03	18,984
1293	5	21 23,52	5	21 22,43	5	21 22,15	64 21 22,70	21 22,56	21 17,92	+0,14	+ 4,78	19,018
1294	5	57 42,71					147 57 42,71		57 48,17		- 5,46	19,025
1295				5 22 7,05			88 22 7,05	22 9,22	22 4,70	-2,17	+ 2,35	19,033
1296				5 55 8,92			82 55 8,92	55 12,21	55 7,37	-3,29	+ 1,55	19,040
1297						5	36 13,30		36 16,40		- 3,10	19,043
1298						3	40 17,70		40 18,40		- 0,70	19,047
1299	5	14 11,41					126 14 11,41		14 3,66		+ 7,75	19,084
1300	5	24 19,85					107 24 19,85	24 21,37	24 19,00	-1,52	+ 0,85	19,154
1301	1	43 7,35	7	43 7,20	2	43 7,14	32 43 7,20	43 8,92	43 6,90	-1,72	+ 0,30	19,155
1302	5	23 59,19					85 23 59,19	23 56,18	23 50,71	+3,01	+ 8,48	19,161
1303				5 59 50,20			82 59 50,20	59 52,01	59 49,45	-1,81	+ 0,75	19,165
1304	4	34 50,89	4	34 53,96			91 34 52,42		34 51,49		+ 0,93	19,196
1305	5	20 39,41	22	20 37,41	39	20 38,31	27 20 38,06	20 37,23	20 38,45	+0,83	- 0,39	19,196

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^b January 1, 1832		A. S. Catal.	Difference from		Annual Preces- sion
						1832					
			No. 1831	No. 1832	No. 1833	h. m. s.	s.		Green.	A. S.	
1306	5	60 Leonis <i>b</i>	7 21,07	—	4 21,06	10 53 21,07	—	20,62	—	+0,45	3,216
1307	6	8 Hyd & Crat <i>v</i>	—	6 17,28	—	10 54 17,26	—	16,58	—	+0,68	2,884
1308	7	Leonis	—	6 39,18	—	10 54 39,18	—	38,90	—	+0,28	3,069
1309	6	62 Leonis <i>p</i> ¹	—	—	2 0,68	10 55 0,68	—	0,35	—	+0,33	3,074
1310	4.5	63 Leonis <i>x</i>	7 21,11	1 21,07	3 20,94	10 56 21,03	20,95	20,42	+0,08	+0,61	3,086*
1311	5	9 Hyd & Crat <i>x</i> ¹	8 15,09	3 15,20	6 15,11	10 57 15,11	—	14,90	—	+0,21	2,889
1312	5.6	Hyd & Crat <i>x</i> ²	—	3 49,61	—	10 57 49,59	—	49,66	—	-0,07	2,890
1313	5.6	65 Leonis <i>p</i> ²	6 19,94	—	1 20,07	10 58 19,96	—	19,68	—	+0,28	3,086
1314	6	67 Leonis	—	6 47,58	—	10 59 47,61	—	47,36	—	+0,25	3,234
1315	3.4	52 Ursæ Maj <i>ç</i>	8 11,17	4 11,39	—	11 0 11,26	11,40	11,59	-0,14	-0,33	3,419
1316	5	10 Hyd & Crat	6 37,21	—	6 37,01	11 0 37,10	—	37,06	—	+0,04	2,892
1317	7	66 Leonis <i>p</i> ³	1 39,45	6 39,14	—	11 0 39,18	—	38,86	—	+0,32	3,066
1318	7	Leonis	—	6 54,81	—	11 2 54,82	—	54,74	—	+0,08	3,158
1319	4	11 Hyd & Crat <i>β</i>	6 24,50	—	6 24,45	11 3 24,46	24,40	24,48	+0,06	+0,02	2,937
1320	3	68 Leonis <i>δ</i>	4 10,11	7 10,18	1 9,96	11 5 10,15	9,98	9,50	+0,17	+0,65	3,193
1321	5.6	69 Leonis <i>p</i> ⁴	9 10,25	—	—	11 5 10,25	—	9,44	—	+0,81	3,073
1322	6.7	Leonis	2 18,05	6 17,95	1 17,91	11 5 17,97	—	17,51	—	+0,46	3,118
1323	3	70 Leonis <i>θ</i>	7 25,01	—	6 25 12	11 5 25,08	25,14	24,61	-0,06	+0,47	3,161
1324	5.6	72 Leonis <i>ι</i>	—	6 15,50	—	11 6 15,52	—	14,36	—	+1,16	3,207
1325	5.6	73 Leonis <i>κ</i>	—	5 4,21	—	11 7 4,22	—	3,78	—	+0,44	3,146
1326	6	Leonis	—	6 11,01	1 11,01	11 7 11,03	—	10,80	—	+0,23	3,143
1327	5	74 Leonis <i>φ</i>	7 7,47	—	6 7,01	11 8 7,26	—	6,73	—	+0,53	3,054
1328	5.6	75 Leonis <i>q</i>	—	6 38,70	1 38,73	11 8 38,70	38,73	38,24	-0,03	+0,46	3,053
1329	4	53 Ursæ Maj. <i>ξ</i>	6 12,38	—	6 12,22	11 9 12,33	12,43	11,67	-0,10	+0,66	3,221*
1330	4	54 Ursæ Maj. <i>ν</i>	3 23,19	3 23,06	6 23,11	11 9 23,15	23,22	23,08	-0,07	+0,07	3,266
1331	5	55 Ursæ Maj. <i>p</i>	3 57 19	4 57,24	6 57,14	11 9 57,20	—	56,84	—	+0,36	3,304
1332	6	76 Leonis	1 17,66	6 17,75	—	11 10 17,73	17,67	17,06	+0,06	+0,67	3,081
1333	3.4	12 Hyd & Crat <i>δ</i>	2 57,12	5 56,95	6 56,91	11 10 56,94	56,87	56,73	+0,07	+0,21	2,998
1334	4	77 Leonis <i>σ</i>	5 28,61	3 28,41	2 28,38	11 12 28,50	28,36	27,91	+0,14	+0,59	3,071
1335	4	Centauri <i>π</i>	4 22,49	1 22,87	6 22,80	11 13 22,63	—	21,80	—	+0,33	2,702
1336	7	Leonis	7 41,91	—	—	11 14 41,91	—	41,71	—	+0,20	3,073
1337	6	13 Hyd & Crat <i>λ</i>	1 2,70	5 2,73	—	11 15 2,71	—	2,70	—	+0,01	2,984
1338	4	78 Leonis <i>ι</i>	9 9,97	—	4 9,74	11 15 9,90	9,72	9,38	+0,18	+0,52	3,121
1339	5.6	79 Leonis <i>ρ</i>	—	6 25,16	1 25,13	11 15 25,16	25,13	24,65	+0,03	+0,51	3,079
1340	5	14 Hyd & Crat <i>ε</i>	7 7,87	—	5 7,95	11 16 7,93	—	7,80	—	+0,13	3,023
1341	6.7	Leonis	—	6 15,89	—	11 16 15,90	—	15,73	—	+0,17	3,124
1342	4	15 Hyd & Crat <i>γ</i>	3 29,88	2 30,17	5 29,91	11 16 29,94	29,77	29,26	+0,17	+0,68	2,992
1343	6	81 Leonis	—	6 50,66	—	11 16 50,66	—	50,55	—	+0,11	3,147
1344	7	82 Leonis	—	5 1,13	—	11 17 1,13	—	0,92	—	+0,21	3,087
1345	7	80 Leonis	—	5 12,00	—	11 17 12,00	—	11,60	—	+0,40	3,089
1346	6	16 Hyd & Crat <i>κ</i>	—	6 42,29	—	11 18 42,28	—	41,67	—	+0,61	3,020
1347	4	84 Leonis	9 18,04	1 18,00	4 17,83	11 19 17,98	17,92	17,48	+0,06	+0,50	3,084
1348	7	Leonis	—	5 19,04	—	11 19 19,04	—	18,68	—	+0,36	3,065
1349	6	85 Leonis	2 56,49	6 56,51	—	11 20 56,52	—	55,60	—	+0,92	3,135
1350	3.4	1 Draconis <i>λ</i>	—	5 19,87	6 19,87	11 21 20,04	20,28	20,57	-0,24	-0,53	3,695

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	1831		1832		1833					Green.	A. S. C.	
	No.	"	No.	"	No.	"						
1306	5 55	11,22					68 55 11,22	55 10,38		+ 0,84	+19,198	
1307			5 55	25,93			115 55 25,93	55 20,96		+ 4,97	19,222	
1308			5 50	45,10			89 50 45,10	50 34,55		+ 10,55	19,231	
1309			5 5	52,89			89 5 52,89	5 47,99		+ 4,90	19,240	
1310	5 45	26,74	10 45	24,99			81 45 25,57	45 27,43	45 23,43	-1,86 +	2,14 19,272	
1311	4 23	18,01					116 23 18,01	23 16,16		+ 1,85	19,294	
1312	1 22	52,66	3 22	51,53			116 22 51,81	22 49,24		+ 2,57	19,307	
1313	5 8	3,13					87 8 3,13	7 57,68		+ 5,45	19,319	
1314			5 26	2,14			64 26 2,14	25 57,44		+ 4,70	19,352	
1315	5 35	32,29					44 35 32,29	35 29,75	35 28,81	+2,54 +	3,48 19,361	
1316	5 10	16,92					117 10 16,92	10 14,29		+ 2,63	19,371	
1317			5 25	29,63			90 25 29,63	25 23,05		+ 6,58	19,372	
1318	5 41	19,84					74 41 19,84	41 14,52		+ 5,32	19,422	
1319	5 54	34,77			5 54	35,09	111 54 34,91	54 35,34	54 33,02	-0,43 +	1,89 19,433	
1320	5 33	22,64	16 33	20,12	5 33	21,72	68 33 20,91	33 24,88	33 21,88	-3,97 -	0,97 19,469	
1321			5 9	23,76			89 9 23,76	9 19,99		+ 3,77	19,469	
1322			5 1	16,76			81 1 16,76	1 12,66		+ 4,10	19,472	
1323	5 39	7,77			5 39	8,88	73 39 8,32	39 12,04	39 8,15	-3,72 +	0,17 19,475	
1324			5 59	23,56			65 59 23,56	59 23,42		+ 0,14	19,498	
1325			5 47	37,67			75 47 37,67	46 31,41		+ 6,26	19,508	
1326			5 14	12,05			76 14 12,05	14 12,20		- 0,15	19,511	
1327	5 44	4,93			1 44	4,54	92 44 4,86	43 57,23		+ 7,58	19,529	
1328			5 4	2,33			87 4 2,33	4 0,96	3 58,98	+1,37 +	3,35 19,539	
1329	4 31	34,00	2 31	32,95	5 31	34,48	57 31 34,03	31 36,54	31 37,61	-2,51 -	3,58 20,190*	
1330	1 59	26,29	4 59	24,64	5 59	25,58	55 59 25,27	59 24,83	59 20,52	+2,44 +	4,75 19,554	
1331	2 53	41,12	3 53	39,08			50 53 39,90	53 33,85		+ 6,05	19,564	
1332	2 25	45,85	3 25	45,87			87 25 45,86	25 42,63		+ 3,23	19,571	
1333	2 52	13,31	4 52	8,58			103 52 10,16	52 14,40		-4,24 +	2,65 19,583	
1334	5 3	1,01			5 3	0,63	83 3 0,82	3 4,71	3 3,66	-3,89 -	2,84 19,611	
1335	5 34	15,92			5 34	14,51	143 34 15,21	34 23,97		- 8,76	19,627	
1336			4 56	47,89			88 56 47,89	56 46,06		+ 1,83	19,650	
1337			4 51	25,53			107 51 25,53	51 23,27		+ 2,26	19,657	
1338	4 32	44,24	7 32	41,61	5 32	42,09	78 32 42,35	32 46,97	32 42,23	-4,62 +	0,12 19,658	
1339			5 40	15,36			87 40 15,36	40 16,04	40 9,32	-0,68 +	6,04 19,663	
1340	5 56	24,15	1 56	21,95			99 56 23,78	56 13,41		+ 10,37	19,675	
1341	1 38	49,61	5 38	48,36			77 38 48,57	38 47,81		+ 1,26	19,677	
1342	7 45	42,08			2 45	40,90	106 45 41,82	45 43,82	45 39,54	-2,00 +	2,28 19,681	
1343			6 37	12,59			82 37 12,59	37 10,84		+ 1,75	19,687	
1344			5 46	30,44			85 46 30,44	46 22,02		+ 8,42	19,689	
1345			4 13	0,33	1 13	0,89	85 13 0,44	12 52,64		+ 7,80	19,692	
1346			5 26	3,40			101 26 3,40	26 1,28		+ 2,12	19,716	
1347	6 13	9,80	1 13	9,24	6 13	10,68	86 13 10,17	13 10,19	13 6,00	-0,02 +	4,17 19,726	
1348	1 46	33,35	4 16	35,13			90 46 34,77	46 32,41		+ 2,36	19,726	
1349			5 39	30,42			73 39 30,42	39 32,19		- 1,77	19,750	
1350			5 44	36,85	5 44	36,73	19 44 36,79	44 35,72	44 35,38	+1,07 +	1,41 19,756	

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832		Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion	
			No.	1831	No.	1832	No.			1833	Green.		A. S.
			s.	s.	s.	h.	m.			s.	s.		s.
1351	6	86 Leonis	—	6 42,72	—	11 21 42,74	—	41,61	—	+1,13	+3,146		
1352	4.5	87 Leonis e	7 44,03	—	5 44,04	11 21 44,03	44,03	43,97	-0,00	+0,06	3,060		
1353	7	Hyd & Crat	5 23,94	2 24,20	—	11 23 24,01	—	23,86	—	+0,15	3,047		
1354	5.6	17 Hyd & Crat	—	6 57,65	—	11 23 57,63	—	53,56	—	-0,93	2,955		
1355	7	Hyd & Crat	—	5 15,80	—	11 24 15,80	—	15,16	—	+0,64	3,043		
1356	4	19 Hyd & Crat ζ	7 45,51	—	5 45,38	11 24 45,44	45,41	44,66	+0,03	+0,78	2,945		
1357	6	89 Leonis H	2 46,06	7 46,10	1 46,27	11 25 46,11	—	45,77	—	+0,34	3,082		
1358	6	90 Leonis C	—	4 57,51	—	11 25 57,53	—	57,19	—	+0,34	3,131		
1359	6	Ursæ Min.	5 26,51	1 26,77	—	11 27 26,56	—	26,48	—	+0,08	3,172		
1360	4	Centauri λ	8 4,88	—	6 4,85	11 28 4,77	—	2,89	—	+1,88	2,717		
1361	4	21 Hyd & Crat θ	8 9,89	—	4 10,05	11 28 9,94	10,00	9,49	-0,06	+0,45	3,039		
1362	4.5	91 Leonis υ	6 21,07	—	6 20,98	11 28 21,03	21,12	20,65	-0,09	+0,38	3,068		
1363	6.7	1 Virginis ω	1 47,57	7 47,80	2 47,71	11 29 47,76	—	47,51	—	+0,25	3,096		
1364	7	Virginis	—	6 49,37	—	11 29 49,37	—	48,93	—	+0,44	3,063		
1365	5.6	24 Hyd & Crat	—	4 8,59	—	11 30 8,58	—	8,31	—	+0,27	3,030		
1366	5.6	92 Leonis	3 2,36	3 2,19	—	11 32 2,28	—	2,17	—	+0,11	3,135		
1367	6.7	Virginis	7 20,51	—	—	11 35 20,51	—	20,30	—	+0,21	3,054		
1368	4	27 Hyd & Crat ζ	8 15,53	4 15,52	1 15,70	11 36 15,54	15,49	14,98	+0,05	+0,56	3,025		
1369	5	2 Virginis ξ ¹	9 37,36	1 37,30	4 37,43	11 36 37,37	37,19	36,47	+0,18	+0,90	3,090		
1370	4	63 Ursæ Maj. x	5 8,99	1 9,07	6 8,72	11 37 8,91	9,00	8,21	-0,09	+0,70	3,220		
1371	4.5	3 Virginis υ	5 13,29	4 13,29	4 13,45	11 37 13,34	13,43	13,20	-0,09	+0,14	3,086		
1372	5.6	4 Virginis ξ ²	1 17,12	5 17,10	—	11 39 17,10	—	16,18	—	+0,92	3,088		
1373	4	93 Leonis E	2 19,05	4 18,88	6 18,65	11 39 18,81	18,80	18,39	+0,01	+0,42	3,115		
1374	6	Hyd & Crat	—	6 17,05	—	11 40 17,03	—	16,48	—	+0,55	3,012		
1375	2.3	94 Leonis β	3 29,07	23 29,23	6 29,14	11 40 29,21	29,18	28,90	+0,03	+0,31	3,064*		
1376	3.4	5 Virginis β	8 56,79	17 56,98	1 56,84	11 41 56,92	56,90	56,43	+0,02	+0,49	3,124*		
1377	6	Virginis B	6 27,18	—	—	11 42 27,18	—	26,71	—	+0,47	3,060		
1378	4	28 Hyd & Crat β	6 26,65	6 26,52	—	11 44 26,57	26,42	26,51	+0,15	+0,06	3,009		
1379	2	64 Ursæ Maj γ	11 57,42	11 57,45	23 57,45	11 44 57,52	57,42	57,21	+0,10	+0,31	3,192		
1380	6	6 Virginis A	6 25,85	2 26,08	—	11 46 25,91	—	25,60	—	+0,31	3,081		
1381	6	29 Hyd & Crat	—	6 8,66	—	11 47 8,64	—	8,39	—	+0,25	3,029		
1382	6	30 Hyd & Crat η	—	5 27,85	—	11 47 27,84	—	27,75	—	+0,09	3,047		
1383	7	Virginis	—	6 37,42	1 37,29	11 49 37,40	—	37,09	—	+0,31	3,073		
1384	5	Chancel ε	—	—	—	11 51 Invisible	—	19,72	—	—	2,842		
1385	5.6	7 Virginis δ	1 20,69	7 20,67	—	11 51 20,67	—	20,48	—	+0,19	3,072		
1386	5	8 Virginis π	13 15,84	9 15,79	—	11 52 15,81	15,15	15,76	+0,06	+0,05	3,074		
1387	5.6	31 Hyd & Crat	—	—	6 16,42	11 52 16,40	—	15,89	—	+0,51	3,053		
1388	7	Virginis	—	6 26,04	—	11 52 26,04	—	25,78	—	+0,26	3,067		
1389	6	1 Comæ Ber	5 7,58	1 7,57	—	11 53 7,58	—	6,71	—	+0,87	3,085		
1390	7	Virginis	6 9,92	1 10,07	3 10,01	11 55 9,97	—	10,11	—	-0,14	3,071		
1391	6	2 Comæ Ber	—	6 40,03	—	11 55 40,05	—	39,32	—	+0,73	3,079		
1392	4.5	9 Virginis δ	15 39,12	5 39,16	—	11 56 39,13	39,02	38,85	+0,11	+0,28	3,071		
1393	7	Virginis	1 24,01	3 24,08	—	11 57 24,06	—	23,81	—	+0,25	3,067		
1394	4.5	Crucis η	6 10,93	—	—	11 58 10,93	—	11,33	—	-0,40	3,046		
1395	3	Centauri δ	6 41,32	—	6 41,46	11 59 41,35	—	41,69	—	-0,34	3,065		

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession		
	No. 1831		No. 1832				No. 1833	Green.		A. S. C.	
	' "	' "	' "				' "				
1351	4 39	57,47	—	—	70 39 57,47	39 52,76	+	4,71	+ 19,762		
1352	5 4	40,05	1 4	39,02	5 4 38,95	92 4 39,51	4 39,05	4 36,76	+ 0,46	+ 2,75	19,762
1353	5 32	23,44	—	—	—	95 32 23,44	—	32 20,19	+	3,25	19,786
1354	—	—	5 20	25,61	—	118 20 25,61	—	20 24,55	+	1,06	19,794
1355	5 54	1,79	—	—	—	96 54 1,79	—	53 59,27	+	2,52	19,798
1356	5 55	43,98	7 55	42,52	5 55 42,22	120 55 42,89	55 41,64	55 43,61	+ 1,25	— 0,72	19,805
1357	2 0	24,91	4 0	26,02	—	86 0 26,25	—	0 22,93	+	3,32	19,818
1358	—	—	5 16	30,92	—	72 16 30,92	—	16 27,90	+	3,02	19,820
1359	—	—	5 17	25,84	—	61 17 25,84	—	17 24,01	+	1,83	19,839
1360	5 5	28,74	—	—	2 5 25,25	152 5 27,74	—	5 29,73	—	1,99	19,847
1361	6 52	21,51	—	—	5 52 23,54	98 52 22,43	52 25,69	52 19,79	— 3,26	+ 2,64	19,848
1362	6 53	50,85	2 53	51,85	6 53 52,52	89 53 51,70	53 49,33	53 46,61	+ 2,37	+ 5,09	19,850
1363	4 56	11,52	—	—	—	80 56 11,52	—	56 10,60	+	0,92	19,868
1364	3 30	24,12	2 30	23,95	—	91 30 24,05	—	30 22,55	+	1,50	19,868
1365	—	—	5 17	38,95	—	102 17 38,95	—	16 36,23	+	2,67	19,872
1366	5 42	46,97	—	—	—	67 42 46,97	—	42 47,32	—	0,35	19,892
1367	1 44	37,60	4 44	37,60	—	95 44 37,61	—	44 32,05	+	5,56	19,926
1368	5 25	0,99	15 24	58,30	—	107 24 59,04	25 1,87	24 59,51	— 2,53	— 0,47	19,934
1369	5 48	31,78	—	—	3 48 31,03	80 48 31,50	48 29,96	48 23,29	+ 1,54	+ 8,21	19,937
1370	5 17	19,27	—	—	—	41 17 19,27	17 21,71	17 20,92	— 2,44	— 1,65	19,942
1371	5 31	42,61	—	—	—	82 31 42,61	31 44,01	31 41,88	— 2,30	+ 2,27	19,943
1372	1 40	19,20	4 40	17,75	—	80 40 18,04	—	40 13,26	+	4,78	19,960
1373	5 50	48,45	—	—	—	68 50 48,45	50 52,09	50 49,32	— 3,04	— 0,87	19,960
1374	—	—	5 48	58,18	—	115 48 58,18	—	48 54,84	+	3,34	19,968
1375	7 29	19,61	13 29	20,01	22 29 20,65	74 29 20,27	29 20,38	29 15,64	— 0,11	+ 4,63	19,969
1376	8 17	19,24	7 17	19,87	5 17 18,77	87 17 19,35	17 19,68	17 17,60	— 0,33	+ 1,75	19,980
1377	4 23	57,85	1 23	57,45	—	94 23 57,77	—	23 53,75	+	4,02	19,983
1378	6 58	20,90	—	—	—	122 58 20,90	58 24,04	58 21,95	— 3,14	— 4,05	19,996
1379	7 22	18,72	21 22	19,16	21 22 20,07	35 22 18,72	22 15,55	22 18,27	+ 3,17	+ 0,45	19,999
1380	3 37	20,56	2 37	20,20	—	80 37 20,42	—	37 15,19	+	5,23	20,007
1381	5 32	22,35	—	—	—	117 32 22,35	—	32 23,99	—	1,64	20,011
1382	1 12	54,25	—	—	—	106 12 54,25	—	12 50,07	+	4,18	20,012
1383	—	—	5 34	59,78	—	85 34 59,78	—	34 55,96	+	3,82	20,022
1384	—	—	—	—	—	167 Invisible	—	17 15,24	—	—	20,028
1385	5 24	34,43	—	—	—	85 24 34,43	—	24 30,57	+	3,86	20,028
1386	9 26	55,08	—	—	7 26 55,32	82 26 55,18	26 50,30	26 53,79	+ 4,88	+ 1,39	20,031
1387	2 43	23,52	—	—	—	108 43 23,52	—	43 18,65	+	4,87	20,031
1388	—	—	4 49	40,12	—	90 49 40,12	—	49 40,47	—	0,35	20,031
1389	—	—	4 58	7,49	—	66 58 7,49	—	58 10,00	—	2,51	20,033
1390	—	—	5 30	6,81	—	83 30 6,81	—	30 5,72	+	1,09	20,038
1391	1 36	14,37	4 36	15,89	—	67 36 15,57	—	36 12,97	+	2,60	20,039
1392	12 20	4,51	—	—	10 20 3,83	80 20 4,20	20 0,44	20 0,55	+ 3,76	+ 3,65	20,040
1393	3 11	43,60	—	—	—	92 11 43,60	—	11 39,33	+	4,27	20,041
1394	5 40	29,26	—	—	—	153 40 29,26	—	40 37,32	—	8,06	20,042
1395	9 47	15,07	7 47	13,33	—	139 47 14,35	—	47 16,33	—	1,98	20,043

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Greenh January 1, 1832			A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	h. m.	s.	s.		Green.	A. S.	
1396	4.5	1 Corvi α	3 45,83	4 45,82	12 45,91	11 59 45,85	45,79	45,65	+0,06	+0,20	+3,067	
1397	6	10 Virginis r	3 4,73	4 5,18	1 5,10	12 1 5,00		4,53		+0,47	3,068	
1398	4	2 Corvi ϵ	6 30,07		6 29,88	12 1 29,96	29,89	29,64	+0,07	+0,32	3,071	
1399	7	11 Virginis s		6 29,79	2 29,75	12 1 29,78		29,91		-0,13	3,067	
1400	6	3 Corvi		5 26,07		12 2 26,05		25,54		+0,51	3,074	
1401	4	Centauri ρ	6 54,57		6 54,77	12 2 54,57		53,42		+1,15	3,088	
1402	6	4 Comæ Ber		6 19,39		12 3 19,40		19,15		+0,25	3,059	
1403	6	5 Comæ Ber		6 36,26		12 3 36,27		35,75		+0,52	3,060	
1404	5	Draconis	1 13,20	5 13,52	3 12,10	12 4 13,47		14,00		-0,53	2,949	
1405	6	12 Virginis t	2 52,63	4 52,60		12 4 52,61		52,07		+0,54	3,063	
1406	3	Crucis δ	7 16,82		6 17,31	12 6 17,00		14,77		+2,23	3,125	
1407	3	69 Ursæ Maj. δ	8 4,40		4 4,29	12 7 4,40	4,38	3,59	+0,02	+0,81	3,003	
1408	3	4 Corvi γ	6 10,69		9 10,68	12 7 10,69	10,69	10,37	0,00	+0,32	3,080	
1409	5	6 Comæ Ber		4 28,32	6 28,05	12 7 28,17		27,68		+0,49	3,056	
1410	5	7 Comæ Ber h	5 50,28	1 49,99		12 7 50,23		49,88		+0,35	3,047	
1411	5	Chamæl β				12 8		46,03			3,313	
1412	6	13 Virginis n	6 3,77	6 3,74	2 3,91	12 10 3,78	3,78	3,50	0,00	+0,28	3,068	
1413	6.7	14 Virginis H		6 41,89		12 10 41,88		41,62		+0,26	3,077	
1414	6	8 Comæ Ber		6 49,84		12 10 49,86		49,16		+0,70	3,040	
1415	3.4	15 Virginis η	5 18,73		5 18,41	12 11 18,57	18,69	18,77	-0,12	-0,20	3,068	
1416	6	10 Comæ Ber		6 23,28		12 11 23,30		22,57		+0,73	3,031	
1417	5.6	16 Virginis c	1 49,34	4 49,60	5 49,27	12 11 49,42	48,47	48,47	+0,95	+0,95	3,026*	
1418	5.6	5 Corvi ζ		4 52,21		12 11 52,20		51,95		+0,25	3,095	
1419	5	11 Comæ Ber s	5 13,48		6 13,51	12 12 13,52		12,86		+0,66	3,044	
1420	6	Corvi		1 16,01	4 15,99	12 12 15,99		15,65		+0,34	3,054	
1421	4	Crucis ϵ		4 21,35	4 21,20	12 12 21,19		19,52		+1,67	3,189	
1422	6	17 Virginis		6 59,74	1 59,68	12 13 59,74		58,80		+0,94	3,059	
1423	5	12 Comæ Ber e	9 3,18		5 2,92	12 14 3,11		2,94		+0,17	3,027	
1424	5.6	6 Corvi		6 37,39		12 14 37,37		37,35		+0,02	3,106	
1425	5	13 Comæ Ber f	6 52,42		6 52,26	12 15 52,36		51,62		+0,74	3,021	
1426	4	Crucis α_1	1 14,71		6 14,97	12 17 14,20		12,41		+1,79	3,257	
1427	1	Crucis α_2	5 18,98	2 18,75	4 19,29	12 17 18,99		17,62		+1,37	3,258	
1428	5	14 Comæ Ber b	2 59 72	1 59 56	2 59 26	12 17 59,52		58,90		+0,62	3,012	
1429	5	15 Comæ Ber c		6 33,57		12 18 33,59		32,90		+0,69	3,008	
1430	4.5	16 Comæ Ber a	5 34,86		3 34,67	12 18 34,80	34,82	34,13	+0,02	+0,67	3,011	
1431	5	Centauri σ	5 59,68	3 59,89		12 18 59,75		0,23		-0,48	3,196	
1432	6.7	Virginis		4 14,83		12 19 14,83		14,55		+0,28	3,075	
1433	4	Centauri u	1 28,25	4 28,20		12 19 28,18		28,44		-0,26	3,156	
1434	7	Virginis		6 45,10		12 19 45,11		44,63		+0,48	3,057	
1435	5.6	17 Comæ Ber d		2 30,83	4 30,79	12 20 30,82		30,30		+0,52	3,008	
1436	6	18 Comæ Ber			6 2,32	12 21 2,55		2,02		+0,33	3,011	
1437	3	7 Corvi δ	2 10,80	4 11,07		12 21 10,98	11,06	11,25	-0,08	-0,27	3,102	
1438	6.7	Virginis			6 25,75	12 21 25,74		26,11		-0,37	3,095	
1439	2.3	Crucis γ	5 54,05	1 54,06		12 21 54,04		54,51		-0,47	3,257	
1440	6	19 Virginis			4 1,63	12 22 1,64		1,28		+0,36	3,044	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession	
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.		
	" "	" "	" "				" "	" "		
1396	5 47 25,70	6 47 23,91	6 47 24,59	113 47 24,68	47 28,41	47 27,60	-3,76	- 2,92	+ 20,043	
1397	—	5 9 31,44	—	87 9 31,44	—	9 30,44	+	1,00	20,042	
1398	—	—	6 41 3,00	111 41 3,90	41 4,50	41 7,18	-0,60	- 3,28	20,042	
1399	—	5 15 28,92	—	83 15 28,92	—	15 23,52	+	5,40	20,042	
1400	—	6 39 57,44	—	112 39 57,44	—	39 55,98	+	1,46	20,042	
1401	5 25 56,83	—	—	141 25 56,83	—	26 7,32	—	10,49	20,041	
1402	—	5 11 31,35	—	63 11 31,35	—	11 34,81	—	3,46	20,041	
1403	—	5 31 13,86	—	68 31 13,86	—	31 17,91	—	4,05	20,040	
1404	—	5 26 59,50	—	11 26 59,50	—	26 57,43	+	2,07	20,039	
1405	—	5 48 5,77	—	78 48 5,77	—	48 4,23	+	1,54	20,038	
1406	6 48 46,84	—	6 43 46,13	147 48 46,48	—	48 55,31	—	8,83	20,035	
1407	6 2 2,70	—	6 2 0,83	32 2 1,76	2 0,22	2 4,41	+1,54	- 2,65	20,033	
1408	1 36 29,53	4 36 29,14	—	106 36 29,22	36 30,42	36 25,74	-1,20	+	3,48	20,033
1409	—	5 9 51,84	—	74 9 51,84	—	9 49,73	+	2,11	20,032	
1410	—	4 7 15,25	2 7 13,17	65 7 14,65	—	7 9,43	+	5,12	20,031	
1411	—	—	—	168 Invisible	—	22 54,24	—	—	20,028	
1412	—	5 51 13,14	—	89 51 13,14	51 9,83	51 7,83	+3,31	+	5,31	20,024
1413	—	5 58 43,54	—	97 58 43,54	—	58 43,42	+	0,12	20,021	
1414	—	3 1 51,23	—	66 1 51,23	—	1 42,88	+	8,35	20,021	
1415	7 43 59,26	8 43 57,10	7 43 59,79	89 43 58,64	43 56,93	43 53,70	+1,71	+	4,94	20,019
1416	—	5 36 6,33	—	60 36 6,33	—	36 6,33	+	0,05	20,018	
1417	—	6 45 2,04	—	85 45 2,04	45 4,50	45 0,87	-2,46	+	1,17	20,016
1418	—	5 16 50,42	—	111 16 50,42	—	16 50,69	—	0,27	20,017	
1419	4 16 41,10	—	—	71 16 41,10	—	16 39,40	+	1,70	20,015	
1420	—	—	5 37 57,23	102 37 57,23	—	37 57,91	—	0,68	20,014	
1421	5 23 21,07	—	—	149 23 21,07	—	23 24,15	—	3,08	20,014	
1422	—	5 45 31,28	—	83 45 31,28	—	45 28,92	+	2,36	20,006	
1423	2 13 13,77	3 13 11,99	—	63 13 12,70	—	13 15,45	—	2,75	20,005	
1424	—	6 54 23,67	—	113 54 23,67	—	54 23,79	—	0,12	20,002	
1425	5 58 4,40	1 58 3,65	—	62 58 4,27	—	58 7,27	—	3,00	19,995	
1426	—	—	—	152 11 —	—	11 32,97	—	—	19,987	
1427	6 10 0,96	4 10 3,03	5 10 0,39	152 10 1,32	—	10 7,97	—	6,65	19,986	
1428	—	5 47 59,96	—	61 47 59,96	—	47 58,53	+	1,43	19,982	
1429	—	5 17 47,38	—	60 17 47,38	—	47 47,15	+	0,23	19,978	
1430	5 14 33,38	—	11 14 34,43	62 14 34,10	14 34,62	14 31,73	-0,52	+	2,37	19,978
1431	4 17 54,66	—	4 17 56,13	139 17 55,40	—	17 41,83	+	13,57	19,975	
1432	—	5 41 1,33	—	93 41 1,33	—	40 59,64	+	1,69	19,973	
1433	2 6 32,60	2 6 34,76	—	128 6 33,68	—	6 25,29	+	8,39	19,971	
1434	—	6 40 23,05	—	84 40 23,05	—	40 16,82	+	6,23	19,969	
1435	—	5 9 20,49	—	63 9 20,49	—	9 21,99	—	1,50	19,963	
1436	—	4 56 41,99	—	64 56 41,99	—	57 33,76	—	51,77	19,959	
1437	3 34 47,62	11 34 46,52	—	105 34 46,76	34 45,23	34 41,19	+1,53	+	5,57	19,958
1438	—	5 27 36,40	—	102 27 36,40	—	27 37,81	—	1,41	19,956	
1439	5 10 16,13	—	—	146 10 16,13	—	10 4,66	+	11,47	19,952	
1440	—	—	5 21 9,09	79 21 9,09	—	21 1,66	+	7,43	19,951	

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832		Green ^b Catal.	A S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	h. m. s.	s.			Green.	A. S.	
			s.	s.	s.							
1441	4	Muscae γ	—	1 33,22	—	12 22 33,11	—	34,34	—	-1,23	+3,452	
1442	5.6	21 Comæ Ber g	—	6 36,82	—	12 22 36,84	—	36,73	—	+0,11	3,066	
1443	6.7	Virginis	—	5 1,24	—	12 23 1,23	—	0,87	—	+0,36	3,078	
1444	4.5	8 Corvi η	6 25,62	—	—	12 23 25,62	25,45	25,47	+0,17	+0,15	3,105	
1445	6	20 Virginis	—	2 32,78	3 32,63	12 24 32,70	—	32,36	—	+0,34	3,040	
1446	5.6	21 Virginis q	1 6,85	5 7,03	—	12 25 7,00	—	6,79	—	+0,21	3,090	
1447	6	22 Comæ Ber	—	—	6 11,49	12 25 11,52	—	11,22	—	+0,30	2,999	
1448	2.3	9 Corvi β	7 34,73	—	—	12 25 34,73	34,66	34,05	+0,07	+0,08	3,129	
1449	4.5	8 Canum Ven d	5 44,90	—	—	12 25 44,90	44,90	44,84	0,00	+0,06	2,864*	
1450	3.4	5 Draconis α	3 15,95	2 15,80	—	12 26 15,93	15,84	14,92	+0,09	+1,01	2,600*	
1451	4.5	23 Comæ Ber k	1 28,40	4 28,49	2 28,34	12 26 28,45	28,73	28,04	-0,28	+0,41	3,001	
1452	5.6	24 Comæ Ber l	—	2 41,61	3 41,87	12 26 41,79	—	41,52	—	+0,27	3,014	
1453	4	Muscae a	2 16,05	4 15,77	—	12 27 15,81	—	16,19	—	-0,38	3,463	
1454	6.7	25 Virginis f	2 8,78	2 8,67	—	12 28 8,72	—	8,20	—	+0,52	3,082	
1455	6	25 Comæ Ber	—	1 32,76	5 32,67	12 28 32,70	—	31,59	—	+0,81	3,014	
1456	5	Centauri τ	5 33,27	—	5 33,12	12 28 33,16	—	33,30	—	-0,14	3,249	
1457	5.6	Hyd & Crat d	—	6 48,79	—	12 28 48,77	—	48,56	—	+0,21	3,150	
1458	7	Virginis	—	6 48,66	1 48,57	12 29 48,66	—	48,50	—	+0,16	3,060	
1459	6.7	Virginis	—	3 5,60	2 5,50	12 30 5,55	—	5,41	—	+0,14	3,079	
1460	4	26 Virginis α	—	6 35,18	—	12 30 35,17	35,01	34,51	+0,16	+0,66	3,090	
1461	6	26 Comæ Ber m	—	—	6 45,52	12 30 45,54	—	44,96	—	+0,58	2,996	
1462	5	Centauri l	10 48,74	—	1 48,67	12 30 48,74	—	48,68	—	+0,06	3,213	
1463	3	Centauri γ	7 17,58	—	—	12 32 17,58	—	18,27	—	-0,69	3,276	
1464	6	27 Virginis	—	—	6 6,51	12 33 6,51	—	5,39	—	+1,12	3,030	
1465	4	29 Virginis γ_1	3 9,13	6 8,97	5 9,20	12 33 9,08	9,13	8,83	-0,05	+0,25	3,022*	
1466	4	Virginis γ^2	—	4 9,14	—	12 33 9,14	—	9,07	—	+0,07	3,022*	
1467	6	28 Virginis	1 16,73	3 17,08	2 17,07	12 33 17,01	—	16,77	—	+0,24	3,090	
1468	5	30 Virginis ρ	—	6 22,88	—	12 33 22,89	—	22,31	—	+0,58	3,030	
1469	6	31 Virginis δ_1	—	—	5 26,42	12 33 26,43	—	26,15	—	+0,28	3,042	
1470	6	Hyd & Crat e	—	5 4,67	—	12 35 4,65	—	4,26	—	+0,39	3,173	
1471	4	Muscae β	4 4,72	3 4,68	—	12 36 4,67	—	5,49	—	-0,82	3,564	
1472	6	33 Virginis	—	5 50,67	—	12 37 50,68	—	50,43	—	+0,25	3,027	
1473	2	Crucis β	8 58,02	—	3 58,41	12 37 58,10	—	58,84	—	-0,74	3,430	
1474	6	27 Comæ Ber n	7 15,17	—	—	12 38 15,17	—	14,89	—	+0,28	2,998	
1475	6	34 Virginis	2 45,95	4 46,17	—	12 38 46,10	—	46,05	—	+0,05	3,017	
1476	6.7	Virginis	3 53,28	5 52,96	—	12 38 53,03	—	52,73	—	+0,30	3,089	
1477	6	35 Virginis	3 18,23	4 18,41	—	12 39 18,34	—	17,98	—	+0,36	3,050	
1478	6	29 Comæ Ber o	2 29,14	4 29,03	—	12 40 29,07	—	28,18	—	+0,89	3,005	
1479	6	30 Comæ Ber	—	5 5,81	—	12 41 5,83	—	5,57	—	+0,26	2,939	
1480	6.7	Virginis	1 38,97	5 39,08	—	12 42 39,06	—	38,71	—	+0,35	3,109	
1481	6	37 Virginis	—	6 4,08	—	12 43 4,09	—	3,74	—	+0,35	3,051	
1482	5.6	31 Comæ Ber p	—	6 30,57	—	12 43 30,59	—	30,26	—	+0,33	2,932	
1483	5	Centauri n	10 9,83	—	4 9,78	12 44 9,81	—	9,54	—	+0,27	3,277	
1484	6	38 Virginis	1 35,82	5 35,44	—	12 44 35,50	—	35,21	—	+0,29	3,080	
1485	5	Centauri o	7 46,87	—	4 47,88	12 44 47,21	—	47,01	—	+0,20	3,455	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from			Annual Precession.	
	1831		1832		1833					Green.	A	S. C		
	No.	"	No.	"	No.	"								
1441	—	—	1 12	13,63	3 12	14,24	161 12	14,09	12 10	50	+	3,50	+19,946	
1442	—	—	5 30	13,12	—	—	64 30	13,12	30 10	38	+	2,74	19,946	
1443	—	—	—	—	5 7	28,79	94 7	28,79	7 27	31	+	1,48	19,943	
1444	4 15	56,11	2 15	56,17	—	—	105 15	56,13	15 52,33	15 47,73	+3,80	+	8,40	19,939
1445	—	—	4 46	29,66	—	—	78 46	29,66	46 34	12	—	4,46	19,929	
1446	—	—	8 31	23,97	—	—	98 31	23,97	31 22	75	+	1,22	19,923	
1447	—	—	—	—	4 47	21,85	64 47	21,85	47 17	49	+	4,36	19,923	
1448	3 27	59,83	3 27	59,25	—	—	112 27	59,54	27 58,06	27 59,40	+1,48	+	0,14	19,919
1449	—	—	5 43	39,71	—	—	47 43	39,71	43 42,66	43 37,95	-3,89	+	1,76	19,917
1450	6 17	5,13	—	—	—	—	19 17	5,13	17 4,41	17 6,63	+0,72	—	1,50	19,912
1451	1 26	37,34	4 26	39,35	—	—	66 26	38,95	26 40,75	26 38,89	-1,80	+	0,06	19,910
1452	—	—	5 41	49,36	—	—	70 41	49,36	41 46	08	+	3,28	19,908	
1453	5 12	24,80	—	—	1 12	24,90	158 12	24,82	12 26	26	—	1,44	19,902	
1454	—	—	2 54	19,92	3 54	21,32	91 54	20,76	54 15	94	—	4,82	19,893	
1455	—	—	—	—	4 59	0,61	71 59	0,61	59 0	60	+	0,01	19,889	
1456	5 36	52,32	—	—	—	—	137 36	52,32	36 49	12	+	3,20	19,888	
1457	—	—	5 12	32,04	—	—	116 12	32,04	12 25	64	+	6,40	19,886	
1458	—	—	—	—	6 13	10,35	87 13	10,35	13 7	08	+	3,27	19,874	
1459	—	—	—	—	5 26	52,01	93 26	52,01	26 53	18	—	1,17	19,871	
1460	—	—	—	—	—	—	97	—	4 12,03	4 6,44	—	—	19,866	
1461	—	—	5 0	41,11	—	—	68 0	41,11	0 43	33	—	2,22	19,864	
1462	3 3	39,12	1 3	40,40	—	—	129 3	39,44	3 36	94	+	2,50	19,863	
1463	5 2	2,16	—	—	2 2	4,33	138 2	2,78	2 0	79	+	1,99	19,845	
1464	—	—	5 38	58,91	—	—	78 38	58,91	39 1	26	—	2,35	19,835	
1465	—	—	9 31	38,42	—	—	90 31	38,42	31 36,32	31 35,55	+2,10	+	2,87	19,835
1466	—	—	9 31	38,42	—	—	90 31	38,42	31 36	55	+	1,87	19,835	
1467	—	—	5 34	28,47	—	—	96 34	28,47	34 25	49	+	2,98	19,833	
1468	—	—	3 50	9,54	3 50	11,35	78 50	10,45	50 8	45	+	2,00	19,832	
1469	—	—	—	—	5 16	12,37	82 16	12,37	16 6	41	—	—	19,831	
1470	—	—	5 24	0,32	—	—	117 24	0,32	23 56	27	+	4,05	19,810	
1471	6 11	13,44	—	—	—	—	157 11	13,44	11 10	39	+	3,05	19,796	
1472	4 31	12,53	1 31	11,59	—	—	79 31	12,34	31 7	15	+	5,19	19,771	
1473	6 16	2,07	2 46	1,53	—	—	148 46	1,94	46 2	16	—	0,22	19,770	
1474	—	—	5 30	9,17	—	—	72 30	9,17	30 9	16	—	0,01	19,766	
1475	—	—	5 7	20,13	—	—	77 7	20,13	7 12	28	+	7,85	19,758	
1476	—	—	5 21	52,29	—	—	95 22	52,29	22 48	08	+	4,21	19,756	
1477	—	—	5 30	31,92	—	—	85 30	31,92	30 28	10	+	3,82	19,750	
1478	—	—	6 57	33,25	—	—	74 57	33,25	57 26	80	+	6,45	19,732	
1479	—	—	5 31	50,26	—	—	61 31	50,26	31 45	59	+	4,67	19,723	
1480	—	—	4 25	20,21	—	—	99 25	20,21	25 17	27	+	2,94	19,698	
1481	—	—	5 1	41,36	—	—	86 1	41,36	1 37	04	+	4,32	19,691	
1482	—	—	5 32	33,40	—	—	61 32	33,40	32 34	12	—	0,72	19,684	
1483	4 15	48,69	—	—	—	—	129 15	48,69	15 43	01	+	5,68	19,674	
1484	—	—	5 38	15,61	—	—	92 38	15,61	38 11	47	+	1,14	19,666	
1485	5 15	46,82	—	—	—	—	146 15	46,82	15 44	35	+	2,47	19,663	

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832		Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	h. m.	s.			Green.	A. S.	
			s.	s.	s.							
1486	5	35 Comæ Ber <i>q</i>	5 1,14	1 1.09	5 1,10	12 45	1,13		1,17		-0,04	+2,962
1487	6	41 Virginis		5 23,72		12 45	23,73		23,41		+0,32	3,006
1488	5.6	40 Virginis ↓		5 37,52		12 45	37,51	37,44	37,55	+0,07	-0,04	3,108
1489	3	77 Ursæ Maj. <i>ε</i>	4 36,94	1 36,97		12 46	36,95	36,92	36,24	+0,03	+0,71	2,655
1490	7	42 Virginis			1 41,30	12 46	41,30		40,74		+0,56	3,027
1491	3.4	43 Virginis <i>δ</i>	5 8,64	5 8,78	2 8,59	12 47	8,69	8,59	8,38	+0,10	+0,31	3,004*
1492	2.3	12 Canum Ven <i>a</i>	4 9,60	1 9,83		12 48	9,65	9,58	9,11	+0,07	+0,54	2,841
1493	4.5	36 Comæ Ber <i>r</i>	6 36,67		10 36,75	12 50	36,73	36,77	36,31	-0,04	+0,42	2,971
1494	4	Muscoe <i>δ</i>	1 51,98	5 51,21		12 50	51,35		49,74		+1,61	3,902
1495	6	44 Virginis <i>κ</i> ¹	3 0,68	3 0,62	1 0,66	12 51	0,65		0,41		+0,24	3,083
1496	6.7	46 Virginis <i>κ</i> ³		6 57,30		12 51	57,30		56,91		+0,39	3,081
1497	5	37 Comæ Ber	9 13,65		4 13,52	12 52	13,62		14,14		-0,52	2,882
1498	6	38 Comæ Ber		5 50,82		12 52	50,83		50,39		+0,44	2,969
1499	3.4	47 Virginis <i>ε</i>	7 49,06		7 48,83	12 53	48,95	49,01	48,78	-0,06	+0,17	3,003
1500	6	48 Virginis <i>κ</i> ⁴	6 15,35	2 15,43		12 55	15,37		15,05		+0,32	3,083
1501	5	Centauri <i>ξ</i> ³			5 8,85	12 57	8,79		9,93		-1,14	3,447
1502	6.7	Virginis		6 35,83		12 57	35,82		35,67		+0,15	3,151
1503	5	14 Canum Ven <i>f</i>				12 57			52,40			2,820
1504	5	39 Comæ Ber <i>l</i>			6 9,62	12 58	9,65		9,55		+0,10	2,932
1505	6	40 Comæ Ber			3 11,69	12 58	11,72		10,71		+1,01	2,922
1506	5.6	49 Virginis <i>g</i>			2 6,32	12 59	6,31	6,37	6,39	-0,06	-0,08	3,127
1507	4	41 Comæ Ber <i>u</i>			4 6,59	12 59	6,62	6,83	6,17	-0,21	+0,45	2,883
1508	6	Comæ Ber			6 50,73	12 59	50,76		51,29		-0,53	2,881
1509	4.5	1 Hydræ Con ↓	2 1,51		1 1,17	13 0	1,42	1,37	0,90	+0,05	+0,52	3,209
1510	6	50 Virginis			4 58,28	13 0	58,27		58,21		+0,06	3,126
1511	4.5	51 Virginis <i>θ</i>	1 15,52		3 15,64	13 1	15,60	15,67	15,54	-0,07	+0,06	3,097
1512	5	Centauri <i>ω</i>			4 49,10	13 1	49,04		49,04		0,00	3,393
1513	4.5	42 Comæ Ber <i>v</i>		3 48,78		13 1	48,79	48,95	48,48	-0,16	+0,31	2,950
1514	5	53 Virginis	7 8,00			13 3	8,00		7,40		+0,60	3,167
1515	6	43 Comæ Ber <i>ω</i>		5 1,69		13 4	1,71		0,48		+1,23	2,787*
1516	6	Virginis	4 11,44	2 11,25		13 4	11,38		11,27		+0,11	2,987
1517	6	55 Virginis		6 12,59		13 5	12,57		12,40		+0,17	3,197
1518	7	Virginis		5 24,21	2 24,36	13 5	24,25		24,17		+0,08	3,053
1519	6	57 Virginis	2 54,68	4 54,64		13 6	54,65		54,15		+0,50	3,201
1520	6	59 Virginis <i>e</i>			6 26,24	13 8	26,25		26,11		+0,14	2,997
1521	6	58 Virginis		6 39,63		13 8	39,62		39,43		+0,19	3,135
1522	6	60 Virginis <i>r</i>		5 7,47		13 9	7,46		7,09		+0,37	3,024
1523	4.5	61 Virginis	12 37,98		2 37,98	13 9	37,98	38,15	37,37	-0,17	+0,61	3,106*
1524	4.5	2 Hydræ Con <i>γ</i>	6 48,40		6 48,36	13 9	48,36	48,45	48,14	-0,09	+0,22	3,232
1525	5	20 Canum Ven <i>h</i>	5 60,00	1 59,97	3 59,80	13 9	59,94		59,40		+0,54	2,713
1526	5	21 Canum Ven	4 4,89	1 5,33		13 11	4,98		4,69		+0,29	2,573
1527	3	Centauri <i>ι</i>		6 10,87	6 10,86	13 11	10,83	10,96	11,62	-0,13	-0,79	3,362
1528	7	62 Virginis		6 31,23		13 11	31,22		31,16		+0,06	3,143
1529	6	64 Virginis <i>z</i>		6 41,77		13 13	41,78		41,46		+0,32	3,023
1530	6	63 Virginis		5 2,09		13 14	2,08		1,57		+0,51	3,196

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
	" "	" "	" "				" "	" "	
1486	2 50 20,10	3 50 21,95	—	07 50 21,21	—	50 21,46	—	0,25	+ 19,659
1487	—	5 40 0,41	—	76 40 0,41	—	39 55,58	+	4,83	19,658
1488	—	6 37 25,25	—	98 37 25,25	37 29,17	37 26,99	-3,92	- 1,74	19,649
1489	5 7 36,10	—	—	33 7 36,10	7 36,53	7 39,47	-0,43	- 3,37	19,631
1490	—	—	—	81 —	—	16 5,60	—	—	19,630
1491	1 41 15,89	5 41 17,59	—	85 41 17,31	41 16,10	41 13,93	+1,21	+ 3,38	19,622
1492	9 46 22,80	—	—	50 46 22,80	46 22,44	46 21,21	+0,36	+ 1,59	19,604
1493	6 40 57,75	—	6 40 56,65	71 40 57,20	40 59,14	40 50,56	-1,94	+ 6,64	19,558
1494	5 38 24,10	—	—	160 38 24,10	—	38 21,47	+	2,63	19,554
1495	—	5 54 10,55	—	92 54 10,55	—	54 9,07	+	1,48	19,550
1496	—	5 28 45,18	—	92 28 45,18	—	27 42,97	+	2,21	19,532
1497	5 19 20,92	—	—	58 19 20,92	—	18 20,59	+	0,33	19,526
1498	—	5 58 6,79	—	71 58 6,79	—	58 4,28	+	2,51	19,514
1499	4 8 7,16	2 8 6,54	7 8 6,62	78 8 6,78	8 8,96	8 5,77	-2,18	+ 1,01	19,495
1500	—	4 45 23,82	—	92 45 23,82	—	45 21,90	+	1,92	19,465
1501	—	—	5 0 11,97	139 0 11,97	—	0 4,43	+	7,54	19,425
1502	—	5 0 52,74	—	104 0 52,74	—	0 52,34	+	0,40	19,415
1503	1 17 57,90	4 17 58,37	—	53 17 58,28	—	18 0,88	—	2,60	19,409
1504	—	5 56 31,73	—	67 56 31,73	—	56 34,72	—	2,99	19,403
1505	—	5 28 45,28	—	66 28 45,28	—	28 47,89	—	2,61	19,402
1506	—	1 50 25,41	4 50 27,43	99 50 27,03	50 23,14	50 19,47	+3,89	+ 7,56	19,382
1507	1 28 18,02	6 28 16,26	9 28 17,42	61 28 17,04	28 17,10	28 16,51	-0,06	+ 0,53	19,382
1508	—	—	4 32 30,38	61 32 30,38	—	32 21,77	+	8,61	19,365
1509	—	5 13 2,56	—	112 13 2,56	13 1,37	13 2,52	+1,19	+ 0,04	19,362
1510	—	—	5 25 52,46	99 25 52,46	—	25 58,62	—	6,16	19,340
1511	—	4 38 24,85	—	94 38 24,85	38 23,99	38 22,75	+0,86	+ 2,10	19,333
1512	5 28 12,85	—	—	132 28 12,85	—	28 12,89	—	0,04	19,320
1513	—	3 33 46,53	2 33 46,31	71 33 46,44	34 48,01	34 46,95	-1,57	- 0,51	19,320
1514	3 17 27,80	2 17 27,16	—	105 17 27,54	—	17 19,61	+	7,93	19,290
1515	—	3 16 2,86	—	61 16 2,86	—	16 0,82	+	2,04	18,428*
1516	—	5 32 52,91	—	77 32 52,91	—	32 51,33	+	1,58	19,264
1517	—	4 2 43,40	—	109 2 43,40	—	2 41,48	+	1,92	19,239
1518	—	5 38 53,03	—	87 38 53,03	—	38 51,11	+	1,92	19,235
1519	2 2 47,52	3 2 46,97	—	109 2 47,19	—	2 44,08	+	3,11	19,197
1520	—	5 41 46,56	—	79 41 46,56	—	41 42,70	+	3,86	19,158
1521	—	1 39 32,58	4 39 34,41	99 39 34,04	—	39 27,10	+	6,94	19,153
1522	—	5 38 29,43	—	83 38 29,43	—	38 28,98	+	0,45	19,141
1523	5 22 25,67	—	—	107 22 25,67	22 27,38	22 25,41	-1,71	+ 0,26	20,208*
1524	6 16 56,26	3 16 56,18	—	112 16 56,23	16 56,96	16 52,28	-0,73	+ 3,95	19,123
1525	2 32 27,11	3 32 23,86	—	48 32 25,16	—	32 23,65	+	1,51	19,117
1526	4 25 58,91	1 25 54,87	—	39 25 58,11	—	25 59,43	—	1,32	19,089
1527	—	5 19 25,99	—	125 49 25,99	—	49 19,47	+	6,52	19,086
1528	—	1 25 9,64	4 25 10,27	100 25 10,15	—	25 2,56	+	7,59	19,077
1529	—	1 57 38,18	4 57 41,24	83 57 40,63	—	57 39,15	+	1,48	19,018
1530	—	5 51 9,97	—	106 51 9,97	—	51 5,85	+	4,12	19,009

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832		Green ^d Catal.	A. S. Catal.	Difference from		Annual Preces- sion	
			No. 1831	No. 1832	No. 1833	h.	m.			s.	Green.		A. S.
			s.	s.	s.						s.		s.
1531	6	65 Virginis	4 37,07	1 37,04	—	13	14	37,06	—	37,25	—0,19	+3,098	
1532	6	66 Virginis	—	6 49,08	—	13	15	49,07	—	48,71	+0,36	3,100	
1533	1	67 Virginis	15 21,29	9 21,32	40 21,19	13	16	21,25	21,14	21,00	+0,11	+0,25	3,147
1534	3	79 Ursæ Maj.	9 8,79	—	4 8,32	13	17	8,67	8,62	7,33	+0,05	+1,34	2,419
1535	4	68 Virginis	6 51,42	—	—	13	17	51,42	51,36	50,77	+0,06	+0,65	3,161
1536	5.6	69 Virginis	3 30,37	3 30,32	—	13	18	30,34	—	29,95	+0,39	3,189	
1537	5	80 Ursæ Maj.	—	4 28,91	—	13	18	28,95	—	28,71	+0,24	2,407	
1538	5.6	70 Virginis	—	6 12,78	—	13	20	12,79	—	12,80	—0,01	2,948	
1539	var.	Hydræ Con u	—	5 33,17	—	13	20	33,16	—	33,01	+0,15	3,257	
1540	7	Virginis	2 38,04	5 38,04	1 38,12	13	20	38,06	—	37,79	+0,27	3,072	
1541	6	71 Virginis	—	1 53,47	9 53,74	13	20	53,73	—	53,46	+0,27	2,972	
1542	4	Centauri	8 20,12	—	2 20,01	13	21	20,09	—	20,04	+0,05	3,437	
1543	6	73 Virginis	—	—	5 0,18	13	23	0,16	—	59,66	+0,50	3,220	
1544	6	Centauri	—	—	6 12,59	13	23	12,56	—	12,64	—0,08	3,327	
1545	6	74 Virginis	4 14,39	5 14,42	—	13	23	14,41	14,68	14,06	—0,27	+0,35	3,113
1546	6	75 Virginis	4 53,85	2 53,85	—	13	23	53,85	—	53,38	+0,47	3,191	
1547	6	76 Virginis	—	6 7,87	—	13	24	7,86	—	7,33	+0,53	3,146	
1548	7	77 Virginis	—	1 38,46	7 38,65	13	24	38,62	—	38,34	+0,28	3,125	
1549	6	78 Virginis	1 37,58	5 37,57	—	13	25	37,57	—	36,40	+1,17	3,029	
1550	4	79 Virginis	9 8,44	1 8,43	6 8,47	13	26	8,45	8,46	8,39	—0,01	+0,06	3,066
1551	6	80 Virginis	6 47,40	1 47,47	4 47,38	13	26	47,40	—	47,16	+0,24	3,107	
1552	6	Hydræ Con f	—	2 30,35	4 30,31	13	27	30,30	—	30,78	—0,48	3,307	
1553	6	Centauri	4 17,10	2 17,03	—	13	29	17,07	—	17,28	—0,21	3,345	
1554	3	Centauri	5 17,97	—	3 18,22	13	29	18,03	—	18,64	+0,61	3,731	
1555	6	1 Bootis	7 38,98	1 38,98	—	13	32	38,98	—	38,35	+0,63	2,868	
1556	5.6	82 Virginis	5 48,29	4 48,26	4 48,24	13	32	48,26	48,37	48,09	—0,11	+0,17	3,140
1557	6	2 Bootis	—	6 5,37	—	13	33	5,36	—	4,87	+0,49	2,840	
1558	6	84 Virginis	—	6 37,47	—	13	34	37,47	—	36,91	+0,56	3,027	
1559	7	Virginis	—	5 10,31	—	13	35	10,31	—	10,20	+0,11	3,112	
1560	6	83 Virginis	—	6 26,80	—	13	35	26,85	—	25,79	+1,06	3,216	
1561	7	Virginis	1 8,69	1 8,94	5 9,17	13	36	9,05	—	8,90	+0,15	3,133	
1562	5	1 Centauri	9 10,11	—	—	13	36	10,11	—	10,85	—0,74	3,410	
1563	6	Hydræ Con g	—	—	6 15,51	13	36	15,48	—	15,60	—0,12	3,325	
1564	6	85 Virginis	—	—	6 33,13	13	36	33,11	—	32,62	+0,49	3,213	
1565	6	86 Virginis	—	6 59,96	—	13	36	59,96	—	59,57	+0,39	3,180	
1566	7	Solitarii	—	—	3 4,43	13	38	4,41	—	4,09	+0,32	3,252	
1567	6	87 Virginis	—	—	3 18,18	13	38	18,16	—	17,38	+0,78	3,238	
1568	6	3 Bootis	—	—	—	13	38	—	—	51,82	—	2,789	
1569	5	4 Bootis	6 16,70	2 16,85	—	13	39	16,82	—	16,30	+0,52	2,883	
1570	4	Centauri	—	—	—	13	39	—	—	27,83	—	3,553	
1571	7	88 Virginis	1 31,22	6 31,39	—	13	39	31,36	—	30,95	+0,41	3,127	
1572	4	Centauri	5 32,09	—	—	13	39	32,09	—	32,11	—0,02	3,567	
1573	5	2 Centauri	3 44,42	3 44,40	—	13	39	44,40	—	44,32	+0,08	3,442	
1574	5.6	89 Virginis	—	6 45,45	—	13	40	45,44	—	45,19	+0,25	3,245	
1575	2.3	85 Ursæ Maj.	2 54,68	4 54,55	16 54,49	13	40	54,59	54,71	54,08	—0,12	+0,51	2,353*

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from..		Annual Precession.					
	No.	1831	No.				1832	No.		1833	Green.	A. S. C.		
	"	"	"				"	"		"	"	"		
1531	—	—	5 2	32,98	—	94. 2	32,98	—	2	32,56	+	0,42	+ 18,992	
1532	—	—	5 16	58,82	—	94 16	58,82	—	16	56,66	+	2,16	18,959	
1533	21 16	58,21	14 16	56,55	41 16	100 16	57,57	16	53,96	16 51,02	+ 3,61	+	6,55	18,944
1534	7 11	43,28	—	—	—	84 11	43,28	11	42,34	11 41,00	+ 0,94	+	2,28	18,921
1535	4 49	51,55	2 49	49,08	—	101 49	50,73	49	50,32	49 50,75	+ 0,41	—	0,02	18,901
1536	—	—	5 6	2,69	—	105. 6	2,69	—	—	5 54,77	+	7,92	18,882	
1537	—	—	5 8	5,29	—	34. 8	5,29	—	—	8 6,02	—	0,73	18,881	
1538	—	—	—	—	5 19	19,56	19,56	—	—	19 12,90	+	6,66	19,800*	
1539	—	—	—	—	5 24	34,68	34,68	—	—	24 34,83	—	0,15	18,821	
1540	3 27	28,12	—	—	—	90 27	28,12	—	—	29 18,61	+	9,51	18,818	
1541	—	—	5 18	27,03	—	78. 18	27,03	—	—	18 27,53	—	0,50	18,810	
1542	5 32	10,45	—	—	—	128 32	10,45	—	—	32 7,82	+	2,63	18,797	
1543	4 51	33,44	—	—	—	107 51	33,44	—	—	51 33,95	—	0,51	18,746	
1544	—	—	5 41	50,25	—	118 41	50,25	—	—	41 49,47	+	0,78	18,739	
1545	—	—	5 23	10,36	—	95 23	10,36	23	9,47	23 2,87	+ 0,89	+	7,49	18,739
1546	—	—	4 29	46,47	—	104 29	46,47	—	—	29 42,76	+	3,73	18,718	
1547	—	—	—	—	6 17	47,96	47,96	—	—	17 44,26	+	3,70	18,711	
1548	1 45	24,43	4 15	23,26	—	96 45	23,26	—	—	45 18,55	+	4,94	18,695	
1549	—	—	5 28	32,75	—	85 28	32,75	—	—	28 27,54	+	5,21	18,664	
1550	6 44	4,98	—	—	7 44	5,67	5,67	44	2,73	43 58,04	+ 2,62	+	7,31	18,647
1551	—	—	5 32	15,89	—	94 32	15,89	—	—	32 11,26	+	4,63	18,626	
1552	—	—	4 38	7,40	—	115 38	7,40	—	—	38 4,13	+	3,27	18,603	
1553	4 41	53,03	—	—	—	118 41	53,03	—	—	41 54,29	—	1,26	18,544	
1554	7 36	27,72	—	—	—	142 36	27,72	—	—	36 23,39	+	4,33	18,514	
1555	5 11	30,99	—	—	—	69 11	30,99	—	—	11 29,76	+	1,23	18,431	
1556	—	—	5 51	4,20	—	97 51	4,20	51	8,06	51 4,90	- 3,86	—	0,20	18,425
1557	—	—	5 38	56,79	—	66 38	56,79	—	—	38 56,61	+	0,18	18,415	
1558	—	—	4 36	33,61	—	85 36	33,61	—	—	36 28,41	+	5,20	18,362	
1559	—	—	5 38	59,17	—	94 38	59,17	—	—	38 54,84	+	4,33	18,343	
1560	—	—	—	—	5 19	53,70	53,70	—	—	19 47,62	+	6,08	18,334	
1561	—	—	—	—	6 17	15,67	15,67	—	—	17 12,24	+	3,43	18,308	
1562	5 11	26,06	—	—	—	122 11	26,06	—	—	11 26,67	—	0,61	18,307	
1563	—	—	3 16	10,57	2 16	11,78	11,05	—	—	16 7,67	+	3,38	18,305	
1564	—	—	5 55	17,44	—	104 55	17,44	—	—	55 9,17	+	8,27	18,294	
1565	—	—	6 34	52,05	—	101 34	52,05	—	—	34 49,91	+	2,14	18,278	
1566	—	—	5 24	39,51	—	108 24	39,51	—	—	24 40,88	—	1,37	18,239	
1567	—	—	—	—	5 0	56,38	56,38	—	—	0 49,55	+	6,83	18,231	
1568	—	—	5 27	3,16	—	63 27	3,16	—	—	27 6,71	—	3,55	18,208	
1569	5 42	8,30	5 42	8,77	—	71 42	8,53	—	—	42 6,53	+	2,00	18,195	
1570	—	—	—	—	—	130	—	—	—	50 44,66	—	—	18,189	
1571	—	—	—	—	5 59	43,40	43,40	—	—	59 43,33	+	0,16	18,186	
1572	5 37	57,83	—	—	—	131 37	57,83	—	—	37 54,48	+	3,35	18,186	
1573	3 36	28,26	3 36	28,41	—	123 36	28,83	—	—	36 31,62	—	2,79	18,179	
1574	—	—	5 17	37,37	—	107 17	37,37	—	—	17 31,04	+	6,33	18,141	
1575	14 50	47,50	2 50	49,98	6 50	50,16	48,45	50	44,00	50 41,88	+ 4,45	+	6,57	18,134

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832			Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion			
			No.	1831	No.	1832	No.	1833			h.	m.		s.	s.	s.
1576	7	Solitarii														
1577	4	5 Bootis	3	22,48	3	22,52	1	22,41	13	41	1,36	22,46	22,01	+0,04	+0,49	2,897
1578	6	6 Bootis					6	45,98	13	41	46,01		46,18		-0,17	2,835
1579	4.5	3 Centauri			7	9,51	3	9,52	13	42	9,49	9,38	10,20	+0,11	-0,71	3,430
1580	5	4 Centauri	5	34,00	1	34,12			13	43	34,02		33,88		+0,14	3,419
1581	6	Hydræ Con			5	47,55			13	44	47,64		47,75		-0,11	3,378
1582	3	Centauri	6	6,34					13	45	6,34		6,53		-0,19	3,690
1583	6	7 Bootis					6	11,21	13	45	11,23		10,54		+0,69	2,867
1584	6	90 Virginis	2	5,17	5	4,93			13	46	5,00		5,00		0,00	3,075
1585	7	Virginis	2	10,46	5	10,08			13	46	10,20		10,68		-0,48	3,144
1586	4.5	10 Draconis	5	31,41	1	31,42			13	46	31,42	31,36	30,23	+0,06	+1,19	1,751
1587	3	8 Bootis	6	41,25	1	41,18	9	41,21	13	46	41,24	41,35	40,90	-0,11	+0,34	2,859
1588	5	Centauri	5	5,75	1	5,70			13	48	5,74		5,73		+0,01	3,600
1589	5	Centauri			5	20,71			13	48	20,68		20,92		-0,24	3,654
1590	5	9 Bootis	2	53,90	4	53,19			13	48	53,84		53,62		+0,22	2,739
1591	6	3 Hydræ Con S ¹					5	6,71	13	49	6,68		5,76		+0,92	3,342
1592	6	4 Hydræ Con S ²					6	36,73	13	50	36,70		36,88		-0,18	3,346
1593	7	Virginis			6	7,49			13	51	7,49		7,45		+0,04	3,098
1594	7	Virginis			5	14,14			13	51	14,13		13,70		+0,43	3,148
1595	5	Centauri	6	17,32					13	51	17,32		17,32		0,00	3,686
1596	1	Centauri	5	2,70			4	2,92	13	52	2,75		3,71		-0,96	4,134
1597	5.6	Hydræ Con			6	50,93			13	52	50,91		50,86		+0,05	3,384
1598	4.5	93 Virginis	6	6,29	1	6,14	1	6,28	13	53	6,28	6,18	6,07	+0,10	+0,21	3,042
1599	6	11 Bootis			2	33,15	4	33,18	13	53	33,20		32,23		+0,97	2,728
1600	6.7	Virginis	3	22,31	2	22,14			13	55	22,24		21,95		+0,29	3,230
1601	7	Virginis	2	28,21	6	28,38			13	55	28,33		28,19		+0,14	3,164
1602	5	Centauri	6	49,55			4	49,65	13	55	49,57		49,13		+0,44	3,617
1603	4.5	5 Hydræ Con	5	49,55	1	49,52	4	49,64	13	56	49,57	49,55	49,41	+0,02	+0,16	3,384
1604	2	5 Centauri	4	49,51	3	49,51	3	49,89	13	56	49,60	49,47	50,07	+0,13	-0,47	3,491*
1605	6	94 Virginis			3	24,74	2	24,51	13	57	24,64		24,38		+0,26	3,161
1606	6	95 Virginis			6	50,43			13	57	50,42		50,02		+0,40	3,166
1607	3.4	11 Draconis	4	50,64	2	50,91	7	50,06	13	59	50,47	50,72	49,82	-0,25	+0,65	1,625
1608	6.7	96 Virginis	1	4,60	6	4,23			14	0	4,27		3,85		+0,42	3,180
1609	5	Octantis							14	0	invariable		59,86			8,440
1610	6	Virginis	3	40,82	3	40,90			14	1	40,85		40,22		+0,63	3,255
1611	5.6	12 Bootis			4	44,16	2	44,16	14	2	44,18		43,96		+0,22	2,737
1612	5	6 Hydræ Con	8	9,92					14	3	9,92		9,85		+0,07	3,408
1613	7	97 Virginis	1	36,92	6	37,13			14	3	37,09		36,49		+0,60	3,178
1614	6	Virginis			5	46,14			14	3	46,15		45,98		+0,17	3,029
1615	4	98 Virginis	8	56,76	1	56,68	3	56,74	14	3	56,75	56,79	56,73	-0,04	+0,02	3,183
1616	6	14 Bootis					5	0,51	14	6	0,53		0,24		+0,29	2,897
1617	6	Virginis			2	9,66	2	9,57	14	6	9,59		9,14		+0,45	3,287
1618	6	15 Bootis			2	37,29	3	37,79	14	6	37,60		37,32		+0,28	2,933
1619	4	99 Virginis	3	13,04	2	13,06			14	7	13,04	12,98	12,83	+0,06	+0,21	3,132
1620	1	16 Bootis	9	0,03	9	0,03	13	0,03	14	8	0,04	0,10	59,75	-0,06	+0,29	2,731*

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
							Green. A. S. C.		
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
1576	—	—	5 1 56,56	110 1 56,56	—	1 50,94	+	5,02	+18,131
1577	5 21 54,57	—	—	73 21 54,57	21 55,77	21 51,22	-1,20	+ 3,35	18,118
1578	—	5 53 44,01	—	67 53 44,01	—	53 53,75	—	9,74	18,102
1579	4 9 27,74	1 9 26,94	—	122 9 27,58	9 22,80	9 23,87	+4,78	+ 3,71	18,088
1580	—	5 5 38,66	—	121 5 38,66	—	5 33,97	+	4,69	18,035
1581	—	—	3 44 6,38	117 44 6,38	—	44 6,13	+	0,25	17,988
1582	—	4 27 26,69	—	136 27 26,69	—	27 20,34	+	6,35	17,976
1583	—	—	5 14 8,40	71 14 8,40	—	14 8,39	+	0,01	17,973
1584	—	5 40 22,92	—	90 40 22,92	—	40 20,32	+	2,60	17,938
1585	—	5 13 42,73	—	97 13 42,73	—	13 41,25	+	1,48	17,934
1586	3 26 45,42	3 26 42,62	—	24 26 44,02	26 41,29	26 44,69	+2,73	- 0,67	17,919
1587	5 45 26,36	—	5 45 25,47	70 45 25,91	45 24,10	45 21,94	+1,81	+ 3,97	17,914
1588	4 16 33,38	2 16 34,45	—	131 16 33,73	—	16 27,90	+	5,83	17,859
1589	4 58 45,28	1 58 46,06	—	133 58 45,43	—	58 36,30	+	9,13	17,849
1590	—	5 40 53,06	—	61 40 53,06	—	40 51,15	+	1,91	17,826
1591	—	5 8 52,11	—	114 8 52,11	—	8 49,57	+	2,54	17,819
1592	—	—	6 11 9,98	114 11 9,98	—	11 7,18	+	2,80	17,758
1593	—	—	5 43 37,88	92 43 37,88	—	43 36,05	+	1,83	17,737
1594	—	2 20 23,98	3 20 26,94	97 20 25,76	—	20 22,73	+	3,03	17,732
1595	5 47 10,07	—	—	134 47 10,07	—	47 4,04	+	6,03	17,731
1596	7 33 26,79	—	—	149 33 26,79	—	33 20,66	+	6,13	17,700
1597	—	6 36 51,12	—	116 36 51,12	—	36 48,40	+	2,72	17,666
1598	2 38 18,09	3 38 19,80	5 38 19,94	87 38 19,51	38 20,11	38 18,37	-0,60	+ 1,14	17,655
1599	—	5 47 52,98	—	61 47 52,98	—	47 54,59	—	1,61	17,637
1600	—	4 9 35,37	1 9 35,68	104 9 35,43	—	9 34,21	+	1,22	17,561
1601	—	—	5 26 47,14	98 26 47,14	—	26 45,84	+	1,30	17,557
1602	5 22 14,68	—	—	130 22 14,68	—	22 6,54	+	8,14	17,542
1603	5 52 8,76	—	—	115 52 8,76	52 5,23	52 6,33	+3,53	+ 2,43	17,499
1604	5 32 24,13	—	—	125 32 24,13	—	32 21,94	+	2,19	17,499
1605	—	—	5 5 6,74	98 5 6,74	—	5 5,53	+	1,21	17,474
1606	—	5 30 26,53	—	98 30 26,53	—	30 23,63	+	2,90	17,456
1607	5 49 15,29	—	1 49 12,22	24 49 14,79	49 9,40	49 8,91	+5,39	+ 5,88	17,367
1608	—	5 32 5,98	—	99 32 5,98	—	32 2,55	+	3,43	17,359
1609	—	—	—	172 Invisible	—	52 58,81	—	—	17,326
1610	—	2 30 15,65	4 30 17,52	105 30 16,90	—	30 12,86	+	4,04	17,289
1611	—	4 6 31,73	—	64 6 31,73	—	6 29,91	+	1,92	17,241
1612	4 27 59,17	2 27 57,77	—	116 27 58,74	—	27 53,73	+	5,01	17,223
1613	—	—	5 6 20,59	99 6 20,59	—	6 20,01	+	0,58	17,202
1614	—	4 47 43,97	—	86 47 43,97	—	47 42,39	+	1,58	17,195
1615	4 29 18,22	—	—	99 29 18,22	—	29 20,70	-	2,48	17,187
1616	—	4 14 55,86	—	76 14 55,86	—	14 54,82	+	1,04	17,093
1617	—	5 24 44,94	—	107 24 44,94	—	24 44,14	+	0,80	17,087
1618	—	3 6 15,64	2 6 16,99	79 6 16,18	—	6 11,69	+	4,49	17,065
1619	5 11 46,43	—	—	95 11 46,43	11 40,62	11 37,21	+5,81	+ 9,22	17,038
1620	14 55 21,59	12 56 26,36	44 56 26,92	69 56 26,36	56 21,62	56 18,73	+4,74	+ 7,63	18,962*

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^b January 1, 1832			A. S. Catal.	Difference from		Annual Preces- sion	
			No. 1831	No. 1832	No. 1833	h.	m.	s.		Green.	A. S.		
			s.	s.	s.	s.	s.	s.		s.	s.		
1621	6	Bootis			6 10,33	14	8	10,35		9,94	+0,41	+2,814	
1622	4.5	Lupi	6 41,45	1 41,26		14	8	41,42		40,94	+0,48	3,786	
1623	4	19 Bootis	1 59,57	4 59,71		14	9	59,70	59,51	59,03	-0,19	+0,67	2,265*
1624	4	100 Virginis	5 2,07	3 2,05		14	10	2,06	2 23	1,72	-0,17	+0,54	3,228
1625	4.5	21 Bootis		6 12,73	2 12,58	14	10	12,76	12,72	12,28	+0,04	+0,48	2,143
1626	6	102 Virginis		5 53,55		14	10	53,55		53,59		-0,04	3,087
1627	6	18 Bootis				14	11			8,01			2,891
1628	6	20 Bootis			4 48,25	14	11	48,27		47,78		+0,49	2,845
1629	6	103 Virginis			6 19,51	14	13	19,51		19,33		+0,18	3,083
1630	6	7 Hydræ Con		6 26,37		14	13	26,36		25,90		+0,46	3,442
1631	6	2 Libræ	1 24,12	3 23,91	2 23,90	14	14	23,94		23,06		+0,88	3,211
1632	6	Bootis		1 7,19	4 7,28	14	15	7,27		6,99		+0,28	2,947
1633	6	Solitarii				14	15			14,53			3,399
1634	5	1 Lupi	6 23,69			14	15	23,69		23,42		+0,27	3,797
1635	5	2 Lupi	5 25,26			14	15	25,26		24,77		+0,49	3,802
1636	5.6	Bootis		6 50,04		14	15	50,05		50,17		-0,12	2,982
1637	5.6	8 Hydræ Con		6 21,67		14	18	21,65		21,48		+0,17	3,483
1638	6.7	104 Virginis N ¹		5 35,92		14	18	35,91		35,37		+0,54	3,139
1639	4	23 Bootis	6 28,64		8 28,77	14	19	28,77	28,52	27,74	+0,25	+1,03	2,015*
1640	5	105 Virginis	8 33,35		4 33,50	14	19	33,40		33,06		+0,34	3,088
1641	6	106 Virginis N ²		5 50,61		14	19	50,60		50,26		+0,34	3,150
1642	5	Lupi	5 21,17		1 21,44	14	21	21,20		21,68		-0,48	3,979
1643	4	25 Bootis	6 35,35			14	24	35,35	35,34	35,29	+0,01	+0,06	2,592
1644	3	Centauri	4 52,49	2 52,72		14	24	52,57		52,53		+0,04	3,764
1645	6	26 Bootis			5 54,41	14	24	54,44		53,73		+0,71	2,733
1646	3.4	27 Bootis		6 18,79	2 18,74	14	25	18,80	18,72	18,59	+0,08	+0,21	2,426
1647	7	Solitarii		2 24,46	3 24,11	14	25	24,42		24,15		+0,27	3,351
1648	5	Lupi	6 38,14			14	26	38,14		39,17		-1,03	3,975
1649	4.5	Apodis				14	27	Invisible		26,03			6,937
1650	5	28 Bootis	4 21,81	2 21,61		14	27	21,75		21,36		+0,39	2,597
1651	6.7	Libræ			3 4,52	14	28	4,51		6,06		-1,55	3,233
1652	4	5 Ursæ Min	1 58,45	4 58,65		14	27	58,73	58,78	59,43	-0,05	-0,70	-0,269
1653	4	Centauri		4 14,54		14	28	14,48		20,70		-6,22	+4,470
1654	1	Centauri	3 16,25			14	28	16,25		22,60		-6,35	4,470
1655	4	Circini			3 2,74	14	29	2,74		4,71		-1,97	4,742
1656	7	3 Libræ			3 41,33	14	29	41,30		40,90		+0,40	3,434
1657	3	Lupi	6 48,10			14	30	48,10		48,65		-0,55	3,933
1658	6	Bootis			5 40,57	14	32	40,59		40,39		+0,20	2,858
1659	3.4	29 Bootis	6 49,89			14	32	49,89	49,94	49,24	-0,05	+0,65	2,813
1660	3.4	30 Bootis	5 7,75	1 7,95	1 7,78	14	33	7,78	7,74	7,18	+0,04	+0,60	2,855
1661	5	Centauri	5 24,57	1 24,91		14	33	24,62		24,88		-0,26	3,636
1662	5	31 Bootis		6 24,08		14	33	24,09		23,90		+0,19	2,938
1663	7	4 Libræ		4 31,96		14	33	31,94		31,95		-0,01	3,442
1664	6	32 Bootis			2 39,79	14	33	39,80		39,68		+0,12	2,886
1665	4.5	107 Virginis	2 13,02	3 13,14	1 13,05	14	34	13,08	13,02	12,89	+0,06	+0,19	3,140

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	1831		1832		1833					Green.	A. S. C	
	No.	"	No.	"	No.	"				"	"	
1621					2 18	10,14	70 18	10,14	18 5,48	+	4,66	+16,954
1622	4 16	43,08					135 16	43,08	16 34,43	+	8,65	16,971
1623	4 8	14,28	1 8	12,47			43 8	14,00	8 13,34	-0,73	+ 0,66	16,908
1624	1 35	34,43	5 35	34,08			102 35	34,14	35 36,66	-2,52	+ 0,98	16,907
1625			3 51	17,69	1 51	17,49	37 51	17,64	51 18,57	-0,93	- 0,94	16,897
1626					3 29	5,22	91 29	5,22	29 2,52	+	2,70	16,867
1627					1 12	54,77	76 12	54,77	12 53,27	+	1,50	16,855
1628			5 55	6,26			72 55	6,26	55 7,12	-	0,86	16,823
1629			5 12	56,68			91 12	56,68	12 55,77	+	0,91	16,751
1630					5 58	28,10	116 58	28,10	58 44,36	-	16,26	16,746
1631			5 56	33,63			100 56	33,63	56 29,91	+	3,72	16,700
1632					4 47	6,53	80 47	6,53	47 4,16	+	2,37	16,664
1633			4 2	20,09	1 2	19,94	114 2	20,06	2 18,35	+	1,71	16,659
1634	5 27	22,45					134 27	22,45	27 15,98	+	6,47	16,652
1635	4 36	55,17	3 36	57,37			134 36	56,11	36 49,25	+	6,86	16,651
1636			5 24	46,11			83 24	46,11	24 46,95	-	0,84	16,629
1637			5 43	52,40			118 43	52,40	43 52,96	-	0,56	16,506
1638			2 21	31,56	3 21	33,26	95 21	33,78	21 24,56	+	9,22	16,494
1639	5 22	13,10			4 22	11,95	37 22	12,59	22 12,17	+0,42	- 0,81	16,988*
1640			5 28	12,64			91 28	12,64	28 13,02	-	0,38	16,446
1641			5 8	29,17			96 8	29,17	8 23,42	+	5,75	16,431
1642	4 42	24,57					139 42	24,57	42 16,98	+	7,64	16,356
1643	5 33	14,75			6 53	15,35	58 53	15,08	53 15,49	+0,51	- 0,41	16,189
1644			5 24	51,02			131 24	51,02	24 48,12	+	2,90	16,176
1645			1 58	45,34	4 58	45,43	66 58	45,41	59 46,41	-	1,00	16,173
1646	4 57	12,85	3 57	13,01	1 57	11,97	50 57	12,80	57 9,94	-0,06	+ 2,86	16,151
1647			5 41	53,35			109 41	53,35	41 49,08	+	4,27	16,148
1648					5 41	12,91	138 41	12,91	41 13,83	-	0,92	16,084
1649							168	Invisible	19 8,54			16,049
1650	5 31	19,44					59 31	19,44	31 15,73	+	3,71	16,045
1651					5 35	8,67	101 35	8,67	35 21,03	-	12,36	16,007
1652			3 33	24,31	2 33	22,03	13 33	23,10	33 26,10	-2,70	- 3,96	16,007
1653			4 7	26,79			150 7	26,79	7 28,11	-	1,32	15,996
1654	3 8	8,88	5 8	8,27	1 8	8,16	150 8	8,50	8 3,11	+	5,39	15,995
1655					5 14	10,29	151 14	10,29	13 48,08	+	22,21	15,958
1656			5 17	45,12			114 17	45,12	17 44,47	+	0,65	15,924
1657			5 39	38,91			136 39	38,91	39 37,67	+	1,24	15,865
1658			2 44	22,25	3 44	23,61	75 44	23,06	44 17,10	+	5,96	15,763
1659			4 51	24,36	1 51	23,91	72 51	24,27	51 25,45	- 1,18	+ 0,36	15,755
1660	5 32	49,63			5 32	49,52	75 32	49,57	32 48,21	+ 1,36	+ 6,70	15,738
1661	5 26	43,09					124 26	43,09	26 31,17	+	11,52	15,724
1662	1 6	55,18			4 6	54,57	81 6	54,69	6 52,45	+	2,24	15,724
1663					5 16	33,29	114 16	35,29	16 32,93	+	2,33	15,717
1664							77 36		36 38,38			15,709
1665			5 55	28,34			94 55	28,31	55 22,71	+ 5,63	- 21,63	15,680

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832		Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion		
			No.	1831	No.	1832	No.			1833	Green.		A. S.	
				s.		s.				s.			s.	
1666	4.5	34 Bootis	1	2,32	4	2,37	—	14 36 2,37	2,41	1,77	-0,04	+0,60	+2,635	
1667	5.6	10 Hydræ Con	—	—	—	—	5 17,99	14 36 17,96	—	17,61	—	+0,35	3,455	
1668	7	Libræ	—	—	—	—	4 40,42	14 36 40,40	—	40,03	—	+0,37	3,383	
1669	6	5 Libræ	p	1 42,91	—	—	4 42,77	14 36 42,78	—	42,70	—	+0,08	3,290	
1670	4.5	35 Bootis	o	2 24,35	5	24,30	—	14 37 24,32	24,33	23,83	-0,01	+0,49	2,798	
1671	5.6	11 Hydræ Con	—	—	—	—	—	14 37 —	—	36,85	—	—	3,462	
1672	3	36 Bootis	s	6 39,00	—	—	1 38,98	14 37 39,00	38,97	38,52	+0,03	+0,48	2,621	
1673	7	Libræ	—	—	—	—	—	14 37 —	—	41,36	—	—	3,387	
1674	4	109 Virginis	z	—	5	45,83	—	14 37 45,84	45,77	45,27	+0,07	+0,57	3,029	
1675	5.6	12 Hydræ Con	—	—	—	—	—	14 37 57,48	—	56,30	—	+1,18	3,471	
1676	6	13 Hydræ Con	—	—	—	—	—	14 38 —	—	9,00	—	—	3,481	
1677	5.6	7 Libræ	μ	—	5	7,44	—	14 40 7,48	7,40	7,17	+0,03	+0,26	3,273	
1678	5	6 Libræ	—	5 26,64	1	27,00	—	14 40 26,70	—	26,41	—	+0,29	3,511	
1679	5	Lupi	o	6 42,88	—	—	—	14 40 42,88	—	42,39	—	+0,49	3,268	
1680	6	8 Libræ	α ¹	—	5	24,52	—	14 41 24,50	24,45	24,39	+0,05	+0,11	3,304	
1681	3	9 Libræ	α ²	4 35,86	2	35,99	17	36,01	14 41 35,97	35,92	35,91	+0,05	+0,06	3,305
1682	6	Libræ	—	—	—	—	3 11,96	14 42 11,94	—	11,64	—	+0,30	3,335	
1683	6	11 Libræ	d	—	—	—	4 18,74	14 42 18,74	—	18,62	—	+0,12	3,092	
1684	7	10 Libræ	σ	—	—	—	5 26,81	14 42 26,79	—	26,00	—	+0,79	3,345	
1685	6	Bootis	—	—	—	—	3 44,43	14 42 44,47	—	43,97	—	+0,50	2,579	
1686	3.4	37 Bootis	ξ	6 38,51	—	—	—	14 43 38,51	38,59	38,22	-0,08	+0,29	2,753	
1687	6	12 Libræ	—	—	6	35,76	—	14 44 35,74	—	35,77	—	-0,03	3,458	
1688	6	13 Libræ	ξ ¹	1 16,25	—	—	—	14 45 16,25	—	15,96	—	+0,29	3,243	
1689	3.4	Lupi	β	6 33,97	—	—	—	14 47 33,97	—	34,00	—	-0,03	3,383	
1690	6	Libræ	—	—	—	—	5 40,69	14 47 —	—	37,96	—	—	3,404	
1691	5	15 Libræ	ξ ²	4 40,01	3	39,90	2	39,99	14 47 39,96	39,96	39,08	0,00	+0,88	3,237
1692	7	14 Libræ	—	—	—	—	5 43,39	14 47 43,36	—	43,25	—	+0,11	3,479	
1693	3	Centauri	x	5 16,17	—	—	—	14 48 16,17	—	16,12	—	+0,05	3,857	
1694	5.6	16 Libræ	c	—	5	25,26	—	14 48 25,26	—	25,24	—	+0,02	3,125	
1695	6	15 Hydræ Con	η	—	—	—	1 44 20	14 48 41,17	—	43,8.	—	+0,34	3,524	
1696	6	1 Serpentis	—	—	—	—	—	14 48 —	—	56,44	—	—	3,060	
1697	7	17 Libræ	1	7,93	5	7,81	—	14 49 7,83	—	7,80	—	+0,03	3,234	
1698	6	Bootis	—	—	—	—	—	14 49 —	—	21,50	—	—	2,792	
1699	7	18 Libræ	—	—	5	49,23	—	14 49 49,22	49,18	48,98	-0,03	+0,24	3,234	
1700	3	7 Ursæ Min.	β	2 17,28	4	17,12	10	15,70	14 51 16,46	16,73	16,73	-0,27	-0,27	-0,286
1701	4.5	19 Libræ	δ	6 0,54	—	—	—	14 52 0,54	0,47	0,21	+0,07	+0,33	+3,193	
1702	7	Libræ	—	—	2	12,64	4	12,66	14 53 12,65	—	12,47	—	+0,18	3,179
1703	7	Libræ	—	—	—	—	3 33,87	14 53 33,86	—	33,85	—	+0,01	3,183	
1704	5	Lupi	π	4 43,43	1	43,15	—	14 53 43,37	—	43,32	—	+0,05	4,031	
1705	3.4	20 Libræ	γ	5 15,49	1	15,53	1	15,50	14 54 15,49	15,48	15,13	+0,01	+0,36	3,490
1706	5	110 Virginia	—	2 25,08	1	25,23	—	14 54 25,13	—	24,98	—	+0,15	3,024	
1707	5.6	41 Bootis	ω	—	—	—	—	14 54 —	—	44,31	—	—	2,624	
1708	3	42 Bootis	β	1 37,05	4	37,36	2	36,86	14 55 37,21	37,11	36,74	+0,10	+0,47	2,261
1709	7	Libræ	—	—	—	—	—	14 56 —	—	27,60	—	—	3,456	
1710	5	43 Bootis	ι	5 14,90	—	—	—	14 57 14,90	—	14,25	—	+0,65	2,580	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
	" "	" "	" "				" "	" "	
1666	4 45 14,91	4 45 12,70	—	62. 45 13,80	45 14,41	45 12,31	-0,61	+ 1,49	+ 15,579
1667	—	—	—	114 —	—	43 24,99	—	—	15,566
1668	—	—	—	110 —	—	27 25,29	—	—	15,545
1669	—	—	—	104 —	—	44 44,28	—	—	15,543
1670	—	4 19-10,15	1 19 12,95	72. 19 10,71	19 11,95	19 10,73	-1,24	- 0,02	15,504
1671	—	—	5 54 52,16	114 54 53,16	—	54 47,52	+	4,64	15,493
1672	—	1 12 48,60	4 12 48,07	62 12 48,18	12 48,31	12 48,39	-0,13	- 0,21	15,490
1673	—	5 36 54,26	—	110 36 54,26	—	36 51,59	+	2,62	15,489
1674	—	—	4 23 38,47	87 23 38,47	23 42,11	23 35,73	-3,64	+ 2,74	15,485
1675	—	—	1 22 43,97	115 22 43,97	—	22 40,83	+	3,14	15,475
1676	—	—	—	115 55 —	—	56 9,78	—	—	15,463
1677	—	1 26 34,20	2 26 34,50	103 26 34,40	26 37,36	26 35,76	-2,96	- 1,36	15,353
1678	3 15 17,39	3 15 17,92	—	117 15 17,65	—	15 15,45	+	2,20	15,335
1679	4 52 25,53	—	—	132 52 25,53	—	52 20,43	+	5,15	15,321
1680	—	—	—	105 20 —	17 36,13	17 33,46	—	—	15,280
1681	8 29 20,38	6 20 23,44	13 20 23,65	105 20 22,64	20 18,23	20 14,27	+4,41	+ 8,37	15,270
1682	—	—	5 5 10,08	107 5 10,08	—	5 3,99	+	6,09	15,236
1683	—	—	5 35 38,90	91 35 38,90	—	35 36,07	+	2,83	15,229
1684	—	—	1 39 26,55	107 39 26,55	—	39 21,52	+	5,03	15,222
1685	—	—	—	60 4 —	—	41 3,36	—	—	15,204
1686	—	5 11 55,39	—	70 11 55,39	11 54,04	11 51,96	+1,35	+ 3,43	15,152
1687	—	5 56 56,11	—	113 56 56,11	—	56 57,35	-	1,22	15,099
1688	—	4 12 29,17	—	101 12 29,17	—	12 20,99	+	8,18	15,060
1689	7 27 1,72	—	—	132 27 1,72	—	26 57,40	+	4,32	14,927
1690	—	—	5 39 9,98	110 39 9,98	—	39 11,05	+	58,93	14,923
1691	5 43 36,88	—	—	100 43 36,88	43 35,98	43 31,85	+0,90	+ 5,03	14,921
1692	—	—	5 45 35,77	114 45 35,77	—	45 30,42	+	5,35	14,918
1693	5 25 25,73	—	—	131 25 25,73	—	25 25,08	+	0,65	14,886
1694	—	—	3 39 22,21	93 39 22,21	—	39 18,53	+	3,38	14,876
1695	—	—	—	116 58 —	—	58 34,94	—	—	14,859
1696	—	—	—	89 29 —	—	29 3,63	—	—	14,845
1697	—	5 28 31,64	—	100 28 31,64	—	28 25,91	+	5,73	14,835
1698	—	—	1 55 46,91	72 55 46,91	—	55 45,71	-	1,80	14,820
1699	—	5 27 48,77	—	100 27 48,77	27 48,27	27 47,55	+0,50	+ 1,22	14,794
1700	—	3 9 31,71	11 9 36,87	15 9 35,76	9 27,90	9 33,53	+7,86	+ 2,23	14,701
1701	—	1 50 46,30	4 50 46,25	97 50 46,26	50 50,56	50 46,27	-4,30	- 0,01	14,664
1702	—	—	2 54 26,78	96 54 26,78	—	54 20,11	+	6,67	14,592
1703	—	—	—	97 10 —	—	10 19,73	—	—	14,571
1704	5 23 13,76	—	3 23 12,59	136 23 13,32	—	23 7,87	+	5,45	14,563
1705	5 36 58,96	—	—	114 36 58,96	36 55,98	36 54,84	+2,98	+ 7,12	14,530
1706	1 14 38,25	4 14 37,71	—	87 14 37,81	—	14 36,45	+	1,36	14,519
1707	—	—	2 —	64 19 26,09	—	19 21,23	+	4,86	14,499
1708	5 56 33,45	1 56 32,79	—	48 56 33,34	56 35,03	56 33,89	-1,69	- 0,55	14,446
1709	—	—	3 —	112 39 48,06	—	39 46,49	+	1,57	14,396
1710	—	6 23 25,89	—	62 23 25,89	—	23 32,55	-	6,66	14,347

lxxviii *Comparison of the Observed Places of the Principal Fixed Stars*

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^b January 1, 1832		A S. Catal.	Difference from		Annual Preces- sion	
			No. 1831			No. 1832			Green.	A. S.		
			s.	s.	s.	h. m.	s.					
1711	6	21 Libræ	v ¹	1 16,30	4 16,18	2 16,11	14 57 16,17	16,24	15,79	-0,07	+0,38	+3,328
1712	6.7	22 Libræ	v ²	—	—	5 27,30	14 57 27,28	—	26,51	—	+0,77	3,338
1713	5	Lupi	λ	1 33,92	—	—	14 57 33,92	—	34,05	—	-0,13	3,990
1714	5	44 Bootis	i	—	2 15,28	1 15,20	14 58 15,25	—	33,87	—	+1,38	1,955*
1715	5	45 Bootis	e	7 55,48	1 55,46	—	14 59 55,48	—	54,71	—	+0,77	2,617
1716	.6	Solitarii	—	—	—	2 4,59	15 0 4,56	—	4,22	—	+0,34	3,475
1717	4	Lupi	ζ	4 15,60	1 15,93	—	15 0 15,66	—	16,85	—	-1,19	4,254
1718	5	Lupi	κ	—	—	—	15 0 —	—	18,06	—	—	4,121
1719	6	46 Bootis	b.	—	—	—	15 1 —	—	8,36	—	—	2,585
1720	.6	Bootis	—	—	—	—	15 1 —	—	16,22	—	—	2,610
1721	5.6	24 Libræ	γ ¹	1 39,45	5 39,69	—	15 2 39,64	39,64	39,74	0,00	-0,10	3,400
1722	3	Triang Aus	γ	—	1 22,50	4 22,31	15 3 22,20	—	21,93	—	+0,27	5,444
1723	6.7	25 Libræ	δ ²	—	—	4 46,01	15 3 45,99	—	45,81	—	+0,18	3,399
1724	5	Circini	β	6 26,02	—	—	15 4 26,02	—	27,20	—	-1,18	4,618
1725	7	26 Libræ	τ	—	2 5,75	—	15 5 5,74	—	5,66	—	+0,08	3,365
1726	6.7	Scorpii	—	—	—	1 39,88	15 6 39,86	—	39,60	—	+0,26	3,456
1727	6	3 Serpentis	—	—	—	—	15 6 —	—	50,67	—	—	2,973
1728	5	Lupi	μ	4 53,61	1 53,58	—	15 6 53,60	—	54,06	—	-0,46	4,119
1729	6	4 Serpentis	—	—	—	4 16 25	15 7 16,25	—	15,29	—	+0,96	3,051
1730	5	48 Bootis	κ	6 27,90	1 27,84	—	15 7 27,89	—	27,50	—	+0,39	2,510
1731	4.5	2 Lupi	φ	—	5 38,09	—	15 7 38,07	37,91	37,79	+0,13	+0,34	3,620
1732	2.3	27 Libræ	β	2 58,72	1 58,52	—	15 7 58,65	58,71	58,43	-0,06	+0,22	3,218
1733	3.4	49 Bootis	δ	1 44,07	2 43,83	1 43,50	15 8 43,83	43,90	43,39	-0,07	+0,44	2,408
1734	5	Lupi	δ	—	—	3 22,67	15 10 22,62	—	22,64	—	-0,02	3,896
1735	5	Lupi	ν	—	—	2 29,12	15 10 29,05	—	28,85	—	+0,20	4,137
1736	5.6	5 Serpentis	—	—	—	—	15 10 —	—	43,65	—	—	3,026
1737	6	Bootis	—	—	—	—	15 10 —	—	52,56	—	—	2,685
1738	5	Lupi	φ ¹	2 10,33	—	1 10,61	15 11 10,40	—	10,34	—	+0,06	3,777
1739	4.5	Lupi	ε	—	3 18,06	1 18,08	15 11 18,03	—	18,52	—	-0,49	4,026
1740	6	28 Libræ	υ	—	2 22,99	—	15 11 22,98	—	22,99	—	-0,01	3,381
1741	7	29 Libræ	φ ²	—	—	2 40,52	15 11 40,50	—	38,70	—	+1,30	3,332
1742	5	Lupi	φ ²	6 27,12	—	—	15 12 27,12	—	27,16	—	-0,04	3,797
1743	6	6 Serpentis	—	—	—	—	15 12 —	—	27,92	—	—	3,045
1744	6	30 Libræ	ο ²	—	—	—	15 13 —	—	40,31	—	—	3,327
1745	6	7 Serpentis	—	—	—	5 26,13	15 14 26,15	—	25,74	—	+0,41	2,833
1746	6	Libræ	—	1 39,65	3 39,95	1 39,96	15 14 39,89	—	39,69	—	+0,20	3,277
1747	5.6	31 Libræ	ε	—	6 6,21	—	15 15 6,21	—	6,00	—	+0,21	3,240
1748	5.6	9 Serpentis	—	—	—	5 0,16	15 18 0,18	—	59,67	—	+0,51	2,776
1749	4	51 Bootis	μ	8 8,65	—	—	15 18 8,65	8,78	8,21	-0,13	+0,44	2,275
1750	6	32 Libræ	ζ ¹	2 47,68	3 47,61	2 47,58	15 18 47,61	—	47,87	—	-0,26	3,362
1751	5.6	10 Serpentis	—	—	—	5 9,66	15 20 9,66	—	9,16	—	+0,50	3,024
1752	7	Libræ	—	—	—	—	15 20 —	—	41,35	—	—	3,375
1753	4	3 Cor Bor	β	—	7 54,37	—	15 20 54,39	54,34	54,23	+0,05	+0,16	2,483
1754	3.4	13 Ursæ Min	γ ²	1 3,03	—	—	15 21 3,03	3,81	2,76	-0,78	+0,27	-0,179
1755	6	34 Libræ	ζ ³	—	1 12,70	—	15 21 12,69	—	12,07	—	+0,62	+3,363

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in					Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.		
	1831		1832		1833				Green.	A. S. C			
	No.	"	No.	"	No.								
1711	—	—	1 36	1 60	2 36	1 85	105 36	1 77	35 58,50	35 55,83	+ 3,27	+ 5,94	+ 14,347
1712	—	—	5 49	42,81	—	—	105 49	42,81	—	49 35,42	—	+ 7,39	14,336
1713	—	—	—	—	5 37	37,33	134 37	37,33	—	37 29,42	—	+ 7,91	14,330
1714	—	—	—	—	5 41	18,81	41 41	18,81	—	41 19,33	—	— 0,52	14,285
1715	5 28	20,19	—	—	—	—	64 28	20,19	—	28 19,63	—	+ 0,56	14,183
1716	—	—	4 20	12,46	—	—	113 20	12,46	—	20 10,94	—	+ 1,52	14,175
1717	5 27	10,61	—	—	—	—	141 27	10,61	—	27 8,47	—	+ 2,14	14,164
1718	3 5	27,13	2 5	26,61	—	—	138 5	26,92	—	5 29,46	—	— 2,54	14,162
1719	—	—	—	—	3	—	63 2	58,66	—	2 58,86	—	— 0,20	14,107
1720	—	—	—	—	2	—	64 14	38,27	—	14 38,81	—	— 0,54	14,099
1721	—	—	5 8	56,41	—	—	109 8	56,41	8 58,93	8 57,99	— 2,52	— 1,58	14,014
1722	—	—	—	—	5 2	54,17	158 2	54,17	—	2 54,01	—	+ 0,16	13,974
1723	—	—	—	—	3	—	109 0	21,37	—	0 27,56	—	— 6,19	13,945
1724	—	—	5 9	53,23	—	—	148 9	53,23	—	9 52,39	—	+ 0,84	13,904
1725	—	—	—	—	2	—	107 8	7,99	—	8 2,08	—	+ 5,91	13,861
1726	—	—	—	—	3	—	111 46	18,71	—	46 16,55	—	+ 2,16	13,762
1727	—	—	4 25	54,59	—	—	84 25	54,59	—	25 49,02	—	+ 5,57	13,749
1728	3 14	58,99	—	—	1 14	59,86	137 14	59,21	—	14 48,18	—	+ 11,03	13,748
1729	—	—	—	—	3 0	1,09	89 0	1,09	—	0 1,64	—	— 0,55	13,723
1730	4 12	32,02	—	—	1 12	31,67	60 12	31,95	—	12 19,33	—	+ 12,62	13,709
1731	—	—	—	—	5 31	29,45	119 31	29,45	31 25,78	31 28,71	+ 3,67	+ 0,74	13,700
1732	5 45	23,49	—	—	—	—	98 45	23,49	45 26,80	45 23,04	— 3,31	+ 0,45	13,678
1733	—	—	6 3	15,79	—	—	56 3	15,79	3 14,04	3 10,71	+ 1,75	+ 5,08	13,628
1734	—	—	1 2	1,59	4 2	0,72	130 2	0,89	—	1 50,70	—	+ 10,19	13,525
1735	—	—	—	—	5 18	30,89	137 18	30,89	—	18 29,28	—	+ 1,61	13,518
1736	—	—	—	—	3	—	87 35	34,98	—	35 35,00	—	— 0,02	13,500
1737	—	—	—	—	2	—	68 48	24,79	—	48 28,00	—	— 3,21	13,490
1738	2 38	43,80	—	—	2 38	45,10	125 38	44,45	—	38 40,00	—	+ 4,45	13,473
1739	—	—	5 4	37,56	—	—	134 4	37,56	—	4 33,35	—	+ 4,21	13,465
1740	—	—	—	—	—	—	107 32	—	—	32 26,38	—	—	13,458
1741	—	—	—	—	—	—	104 56	—	—	56 2,86	—	—	13,441
1742	5 14	56,97	—	—	—	—	126 14	56,97	—	14 54,86	—	+ 2,11	13,390
1743	—	—	5 40	3,25	—	—	88 40	3,25	—	40 2,88	—	+ 0,37	13,387
1744	—	—	—	—	1 31	45,39	104 31	45,39	—	31 26,28	—	+ 19,11	13,309
1745	—	—	4 49	33,25	2 49	31,58	76 49	32,69	—	49 27,35	—	+ 5,34	13,259
1746	—	—	—	—	2 45	50,54	101 45	50,54	—	45 47,86	—	+ 2,68	13,214
1747	—	—	—	—	2 42	49,15	99 42	49,15	—	42 39,97	—	+ 9,18	13,216
1748	—	—	5 58	29,75	—	—	73 58	29,75	—	58 27,97	—	+ 1,78	13,023
1749	5 1	49,15	—	—	1 1	47,31	52 1	48,84	1 46,87	1 41,52	+ 1,97	+ 7,32	13,018
1750	—	—	2 7	28,86	3 7	27,46	106 7	28,02	—	7 20,87	—	+ 7,15	12,971
1751	—	—	—	—	4 34	6,44	87 34	6,44	—	34 1,76	—	+ 4,68	12,860
1752	—	—	—	—	—	—	106 41	18,12	—	41 19,76	—	—	12,844
1753	5 18	40,07	—	—	—	—	60 18	40,07	18 38,28	18 35,42	+ 1,79	+ 4,65	12,828
1754	5 34	4,46	—	—	—	—	17 31	4,46	35 5,20	34 3,43	— 0,74	+ 1,03	12,813
1755	—	—	4 1	38,09	—	—	106 1	38,09	—	1 33,67	—	+ 4,42	12,810

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832			Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No.	1831	No.	1832	No.	1833			Green.	A. S.	
				s.		s.		s.				s.	
1756	3	12 Draconis	1	12,08	4	12,24	—	15 21 12,24	12,08	11,80	+0,16	+0,44	+1,319
1757	5	Triang Aus ε	—	—	—	—	15 21 —	—	27,44	—	—	—	5,349
1758	6.7	Libræ	—	—	—	—	15 22 58,58	—	58,35	—	—	+0,23	3,426
1759	6	35 Libræ ζ ⁴	—	158,46	2	58,66	15 23 26,63	26,64	25,97	-0,01	+0,66	—	3,370
1760	4	Lupi γ	3	58,73	—	—	15 23 58,73	—	58,67	—	+0,06	—	3,957
1761	6	11 Serpentis	—	—	—	4	19,25	15 24 19,25	—	18,57	—	+0,68	3,079
1762	6	36 Libræ	—	—	—	—	15 24 27,28	—	26,94	—	—	+0,34	3,608
1763	4	37 Libræ f ¹	7	0,43	—	—	15 25 0,43	0,37	0,02	+0,06	+0,41	—	3,242
1764	4.5	38 Libræ γ	3	8,24	3	8,35	1 8,48	15 26 8,26	8,46	8,58	-0,20	-0,32	3,333
1765	4.5	4 Cor Bor θ	—	—	2	9,65	3 9,43	15 26 9,56	9,48	8,91	+0,08	+0,65	2,416
1766	5	Lupi ι	—	—	—	2	46,11	15 26 46,05	—	46,01	—	+0,04	4,012
1767	3	13 Serpentis δ	—	—	—	1	47,10	15 26 47,11	47,04	46,60	+0,07	+0,51	2,862
1768	5	39 Libræ	—	—	—	—	15 26 —	—	50,46	—	—	—	3,615
1769	7	Scorpii	—	—	—	—	15 27 —	—	24,52	—	—	—	3,574
1770	2	5 Cor Bor α	1	34,71	15	34,62	22 34,63	15 27 34,65	34,65	34,15	0,00	+0,50	2,526
1771	6	15 Serpentis	—	—	—	—	15 27 —	—	55,69	—	—	—	2,721
1772	6	14 Serpentis A ¹	—	—	—	—	15 27 —	—	56,74	—	—	—	3,068
1773	7	Libræ	—	—	—	—	15 28 —	—	3,90	—	—	—	3,619
1774	4.5	40 Libræ	6	21,56	—	—	15 28 21,56	21,68	21,20	-0,12	+0,36	—	3,657
1775	6	16 Serpentis	—	—	—	3	25,75	15 28 25,76	—	25,17	—	+0,59	2,871
1776	6	18 Serpentis τ ²	—	—	—	—	15 28 —	—	45,00	—	—	—	2,752
1777	5	6 Cor Bor μ	1	4,91	1	5,03	—	15 29 4,97	—	4,85	—	+0,12	2,195
1778	6	41 Libræ φ	—	—	—	—	15 29 —	—	14,91	—	—	—	3,427
1779	5	Lupi σ	—	—	—	—	15 29 —	—	40,77	—	—	—	4,093
1780	5.6	42 Libræ ζ	—	—	3	21,76	—	15 30 21,75	22,02	21,63	-0,27	+0,12	3,524
1761	5	43 Libræ α	7	16,88	—	—	2 17,02	15 32 16,91	16,98	16,75	-0,07	+0,16	3,438
1782	5	7 Cor Bor ζ	4	3,32	—	—	—	15 33 3,32	—	2,77	—	+0,55	2,256
1783	6	19 Serpentis τ ³	—	—	—	3	15,96	15 33 15,98	—	15,47	—	+0,51	2,749
1784	5.6	20 Serpentis ζ	—	—	—	3	53,48	15 33 53,50	—	52,94	—	+0,56	2,812
1785	5	21 Serpentis ι	6	3,93	—	—	—	15 34 3,93	—	3,13	—	+0,80	2,672
1786	6	22 Serpentis	—	—	—	1	21,52	15 34 21,54	—	20,89	—	+0,65	2,698
1787	4.5	44 Libræ η	2	38,15	4	38,25	2 38,14	15 34 38,19	38,19	37,83	0,00	+0,36	3,359
1788	7	Libræ	—	—	2	0,55	—	15 34 0,54	—	—	—	—	3,346
1789	6	23 Serpentis ↓	—	—	—	—	2 35,43	15 35 35,45	—	35,22	—	+0,23	3,010
1790	6	8 Cor Bor γ	—	—	—	—	15 35 —	—	40,48	—	—	—	2,522
1791	2.3	24 Serpentis α	—	—	11	59,95	20 59,94	15 35 59,95	59,93	59,77	+0,02	+0,18	+2,936
1792	5	15 Ursæ Min. θ	—	—	—	—	15 36 —	—	35,25	—	—	—	-1,990
1793	6	26 Serpentis	—	—	—	4	4,37	15 37 4,39	—	3,29	—	+1,10	+2,720
1794	6	25 Serpentis A ²	—	—	—	3	24,96	15 37 24,96	—	24,49	—	+0,47	3,092
1795	4.5	27 Serpentis λ	6	17,86	—	—	3 17,89	15 38 17,87	17,86	17,61	+0,01	+0,26	2,917
1796	3.4	28 Serpentis β	6	26,22	—	—	—	15 38 26,22	26,05	25,87	+0,17	+0,35	2,757
1797	4.5	5 Lupi ζ	4	18,28	1	18,43	—	15 40 18,30	18,22	18,21	+0,08	+0,09	3,782
1798	3	Triang Aus β	1	25,63	1	25,75	—	15 40 25,66	—	26,22	—	+0,56	5,208
1799	3.4	32 Serpentis μ	—	—	—	—	5 51,75	15 40 51,75	51,66	51,61	+0,09	+0,14	3,124
1800	5	1 Scorpii δ	—	—	—	—	2 53,61	15 40 53,58	—	53,09	—	+0,49	3,585

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	1831		1832		1833					Green.	A. S. C.	
	No.	"	No.	"	No.	"						
1756	5	26 37,72	—	—	—	—	26 35,58	26 35,88	+2,14	+ 1,84	+12,806	
1757	—	—	—	—	5	44 31,34	155 44 31,34	44 31,57	—	— 0,23	12,797	
1758	—	—	—	—	2	—	109 5 27,75	5 28,56	—	— 0,81	12,691	
1759	—	—	—	—	3	—	106 16 35,40	16 31,48	—0,47	+ 3,92	12,659	
1760	2	35 40,85	5	35 41,59	—	—	130 35 41,38	35 35,00	—	+ 6,38	12,624	
1761	—	—	—	—	4	36 39,57	90 36 39,57	36 37,79	—	+ 1,78	12,599	
1762	—	—	—	—	3	—	117 28 13,82	28 22,48	—	— 8,66	12,591	
1763	—	—	—	—	4	28 59,00	99 28 59,60	28 55,62	+3,40	+ 3,98	12,552	
1764	3	13 21,85	—	—	1	13 22,53	104 13 22,02	13 21,96	+0,06	+ 6,74	12,475	
1765	—	—	—	—	3	4 8,72	58 4 8,72	4 9,16	—0,44	+ 1,36	12,472	
1766	—	—	—	—	2	0 32,11	132 0 32,11	0 27,26	—	+ 4,85	12,433	
1767	—	—	—	—	3	—	78 53 37,36	53 33,96	—2,51	+ 3,40	12,430	
1768	—	—	—	—	2	—	117 34 15,08	34 15,86	—	— 0,78	12,427	
1769	—	—	—	—	3	—	115 43 3,08	42 58,70	—	+ 4,38	12,388	
1770	8	42 52,53	13	42 52,48	24	42 54,15	62 42 53,38	42 49,48	—0,84	+ 3,90	12,375	
1771	—	—	—	—	1	46 46,25	71 46 46,25	46 43,31	—	+ 2,94	12,351	
1772	—	—	—	—	3	—	89 59 57,15	59 50,84	—	+ 6,31	12,350	
1773	—	—	—	—	—	—	117 38	38 42,88	—	—	12,343	
1774	4	13 5,05	3	13 5,09	—	—	119 13 5,06	13 1,72	+1,29	+ 3,34	12,324	
1775	—	—	—	—	—	—	79 25	25 16,65	—	—	12,317	
1776	—	—	—	—	—	—	73 19	19 7,87	—	—	12,294	
1777	—	—	1	25 40,82	4	25 43,14	50 25 42,68	25 35,02	—	+ 7,66	12,270	
1778	—	—	—	—	3	—	108 44 27,25	44 28,58	—	+ 1,33	12,261	
1779	1	5 48,21	—	—	2	5 49,66	134 5 49,21	5 36,60	—	+ 12,61	12,233	
1780	—	—	—	—	3	—	113 15 52,12	15 53,28	—1,16	+ 3,51	12,184	
1781	—	—	—	—	5	7 39,48	109 7 39,48	7 32,86	+0,84	+ 6,62	12,050	
1782	1	48 51,22	1	48 49,97	3	48 51,04	52 48 50,77	48 48,02	—	+ 2,75	11,994	
1788	—	—	1	25 37,19	5	25 38,85	73 25 38,57	25 36,47	—	+ 2,10	11,980	
1784	—	—	—	—	1	36 28,18	76 36 28,18	36 26,04	—	+ 2,14	11,937	
1785	5	47 3,11	—	—	—	—	69 47 3,11	46 58,00	—	+ 5,11	11,924	
1786	—	—	—	—	3	—	70 59 43,88	59 39,47	—	+ 4,41	11,904	
1787	4	7 53,06	2	7 55,89	—	—	105 7 55,34	7 46,92	+4,01	+ 8,42	11,885	
1788	—	—	—	—	3	—	104 29 52,13	29 40,71	—	+ 11,42	11,858	
1789	—	—	—	—	3	56 26,46	86 56 26,46	56 19,69	—	+ 6,77	11,817	
1790	—	—	—	—	—	—	63 10 0,63	9 56,58	—	+ 4,05	11,810	
1791	7	2 23,27	5	2 22,38	19	2 23,28	83 2 23,14	2 22,55	—0,87	+ 0,59	11,788	
1792	1	5 41,41	—	—	5	5 42,04	12 5 41,94	5 41,81	—	+ 0,13	11,734	
1793	—	—	—	—	3	—	72 12 8,99	12 2,05	—	+ 6,94	11,712	
1794	—	—	—	—	3	—	91 16 17,26	16 16,27	—	+ 0,99	11,688	
1795	5	6 54,83	—	—	—	—	82 6 54,83	6 53,26	+0,85	+ 1,57	11,625	
1796	—	—	—	—	5	2 47,97	74 2 47,97	2 41,14	—1,67	+ 6,83	11,615	
1797	—	—	3	6 30,10	2	6 27,18	123 6 28,93	6 26,67	+4,49	+ 2,26	11,488	
1798	1	54 2,06	—	—	4	54 2,88	152 54 2,71	53 51,07	—	+ 11,64	11,477	
1799	3	54 32,71	—	—	1	54 32,86	92 54 33,50	54 30,15	—2,11	+ 3,35	11,442	
1800	—	—	—	—	3	—	115 14 2,30	13 59,79	—	+ 2,51	11,441	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832			Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion	
			No. 1831		No. 1832	No. 1833		h. m. s.			s.	s.		s.
			s.	s.	s.	s.	s.							
1801	4	35 Serpentis κ	—	—	—	3 11,04	15 41 11,06	—	10,84	10,42	+0,22	+0,64	+2,697	
1802	6	34 Serpentis ω	—	—	—	3 49,28	15 41 49,28	—	—	48,90	—	+0,38	3,016	
1803	3	37 Serpentis ϵ	—	—	—	2 26,79	15 42 26,79	—	26,82	26,65	-0,03	+0,14	2,972	
1804	6	36 Serpentis b	—	—	—	—	15 42 —	—	—	30,64	—	—	3,118	
1805	4.5	10 Cor Bor δ	—	—	—	—	15 42 —	—	33,11	32,54	—	—	2,516	
1806	5	2 Scorpii A^1	2 32,60	—	—	—	15 43 32,60	—	32,64	31,97	-0,04	+0,63	3,579	
1807	5	45 Libræ λ	1 35,75	—	—	—	15 43 35,75	—	35,75	35,33	0,00	+0,42	3,463	
1808	6	Scorpii f^1	—	—	—	—	15 43 —	—	—	52,77	—	—	3,561	
1809	5	38 Serpentis ρ	—	—	—	3 53,44	15 43 53,46	—	53,35	52,52	+0,11	+0,94	2,632	
1810	6	Scorpii f^2	—	—	—	1 57,19	15 43 57,17	—	—	56,84	—	+0,33	3,549	
1811	4.5	46 Libræ θ	5 16,39	—	—	—	15 44 16,39	—	16,41	16,59	-0,02	-0,20	3,390	
1812	6	3 Scorpii A^2	—	—	—	3 35,53	15 44 35,51	—	—	35,16	—	+0,35	3,579	
1813	5	11 Cor Bor κ	—	4 53,96	—	—	15 44 53,98	—	—	53,79	—	+0,19	2,256	
1814	7	47 Libræ	—	1 18,74	—	2 18,62	15 45 18,64	—	—	18,22	—	+0,42	3,448	
1815	6.7	4 Scorpii	—	—	—	3 21,94	15 45 21,91	—	—	21,72	—	+0,19	3,604	
1816	4	5 Scorpii ρ	2 31,89	—	—	2 32,16	15 46 32,01	—	31,88	31,66	+0,13	+0,55	3,679	
1817	6	Serpentis	—	—	—	—	15 47 —	—	—	10,67	—	—	2,643	
1818	3.4	6 Scorpii π	5 42,31	—	—	—	15 48 42,31	—	42,46	42,24	-0,15	+0,07	3,606	
1819	3	41 Serpentis γ	—	—	—	3 41,34	15 48 41,36	—	41,85	41,79	-0,49	-0,43	2,741	
1820	5	48 Libræ \downarrow	—	—	—	3 47,55	15 48 47,53	—	47,45	47,42	+0,03	+0,11	3,343	
1821	5	Lupi η	—	—	—	—	15 49 —	—	—	6,85	—	—	3,943	
1822	6	Serpentis	—	—	—	3 29,81	15 49 29,82	—	—	29,52	—	+0,20	2,769	
1823	3	7 Scorpii δ	5 24,79	1 24,81	—	—	15 50 24,79	—	24,80	24,85	-0,01	-0,06	+3,527	
1824	4	16 Ursæ Min ζ	—	—	—	—	15 50 —	—	13,92	13,23	—	—	-2,384	
1825	4.5	13 Cor Bor ϵ	4 38,11	—	—	—	15 50 38,11	—	38,18	38,08	-0,07	+0,03	+2,484	
1826	5.6	49 Libræ	—	2 54,72	—	3 54,80	15 50 54,75	—	—	54,26	—	+0,49	3,342 ^c	
1827	6	50 Libræ	—	—	—	2 44,30	15 51 44,30	—	—	43,89	—	+0,41	3,226	
1828	6	3 Herculis	—	—	—	3 31,19	15 52 31,19	—	—	30,84	—	+0,35	2,971	
1829	6	Scorpii	—	—	—	5 12,63	15 53 12,62	—	—	12,25	—	+0,37	3,607	
1830	6	5 Herculis r	—	—	—	4 41,59	15 53 41,61	—	—	41,18	—	+0,43	2,692	
1831	5	Normæ δ	2 38,50	—	—	1 39,11	15 54 38,77	—	—	39,02	—	-0,25	4,197	
1832	4.5	44 Serpentis π	—	—	—	2 3,42	15 55 3,45	—	3,92	3,19	-0,47	+0,26	2,577	
1833	4.5	51 Libræ ξ	5 8,54	—	—	—	15 55 8,54	—	8,33	8,23	+0,21	+0,31	3,288	
1834	6	43 Serpentis	—	—	—	2 27,64	15 55 27,64	—	—	26,92	—	+0,72	2,659	
1835	4	Lupi θ	—	—	—	—	15 55 —	—	—	34,74	—	—	3,909	
1836	2	8 Scorpii β	8 40,95	6 40,78	—	—	15 55 40,87	—	40,90	40,93	-0,03	-0,06	3,469	
1837	4.5	9 Scorpii ω_1	5 59,68	—	—	—	15 56 59,68	—	59,68	59,74	0,00	-0,06	3,490	
1838	4.5	10 Scorpii ω_2	—	—	—	—	15 57 —	—	34,04	33,67	—	—	3,496	
1839	5	6 Herculis υ	—	—	—	—	15 57 —	—	33,98	33,19	—	—	1,856	
1840	6	Scorpii m	—	—	—	3 54,21	15 57 54,19	—	—	53,69	—	+0,50	3,626	
1841	6	11 Scorpii	—	—	—	4 17,44	15 58 17,43	—	—	17,17	—	+0,26	3,319	
1842	3.4	13 Draconis θ	—	—	—	3 44,46	15 58 44,59	—	—	45,30	—	-1,31	1,147	
1843	6	45 Serpentis g^1	—	—	—	3 36,93	15 59 36,95	—	—	36,29	—	+0,66	2,857	
1844	6	46 Serpentis g^2	—	—	—	4 5,75	16 0 5,77	—	—	5,20	—	+0,57	2,853	
1845	5	Triang Aus δ	4 13,30	1 13,22	—	—	16 0 13,27	—	—	14,59	—	-1,32	5,363	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
1801	—	—	5 20 3,31	71 20 3,31	20 3,33	19 58,82	-0,02	+ 4,49	+ 11,418
1802	—	—	1 17 6,04	87 17 6,04	—	17 2,71	—	+ 3,33	11,373
1803	1 0 44,29	—	3 0 44,24	85 0 44,25	0 38,95	0 36,66	+5,30	+ 7,59	11,327
1804	—	—	3 —	92 34 31,05	—	34 29,19	—	+ 1,86	11,323
1805	5 24 42,18	—	—	63 24 42,48	24 44,02	24 39,27	-1,54	+ 3,21	11,319
1806	2 49 9,52	—	—	114 49 9,52	49 3,89	49 2,91	+5,63	+ 6,61	11,250
1807	—	—	2 39 31,83	109 39 31,83	39 29,55	39 23,72	+2,28	+ 8,11	11,246
1808	—	—	3 —	114 1 29,54	—	1 29,46	—	+ 0,38	11,225
1809	—	—	3 30 39,01	68 30 39,01	30 42,59	30 38,07	-3,58	+ 0,94	11,223
1810	—	—	3 —	113 28 10,99	—	28 10,89	—	+ 0,10	11,220
1811	—	3 13 47,89	2 13 46,58	106 13 47,37	13 45,88	13 42,46	+1,49	+ 4,91	11,196
1812	—	—	—	114 44 —	—	44 14,84	—	—	11,174
1813	2 48 59,65	—	—	53 48 59,65	—	48 57,83	—	+ 2,32	11,148
1814	—	—	3 —	108 52 45,36	—	52 43,88	—	+ 1,48	11,121
1815	—	—	—	115 45 —	—	45 45,41	—	—	11,117
1816	—	—	5 42 56,20	118 42 56,20	42 48,47	42 55,26	+7,73	+ 0,94	11,033
1817	—	—	3 11 28,85	69 11 28,85	—	11 28,92	—	- 0,07	10,983
1818	5 37 24,84	—	—	115 37 24,84	37 21,35	37 21,32	+3,49	+ 3,52	10,873
1819	6 47 1,29	—	—	73 47 1,29	47 4,47	47 1,42	-3,18	+ 0,17	12,182*
1820	—	—	5 47 13,99	103 47 13,99	—	47 13,44	—	+ 0,55	10,866
1821	—	—	4 54 24,49	137 54 24,49	—	54 24,43	—	+ 0,06	10,851
1822	—	—	3 —	75 5 56,94	—	5 56,27	—	+ 1,07	10,813
1823	6 8 12,79	—	—	112 8 12,79	8 9,35	8 3,85	+3,44	+ 8,94	10,747
1824	—	2 41 32,68	1 41 33,89	11 41 33,08	41 34,20	41 35,38	-1,12	- 2,30	10,747
1825	—	—	4 37 49,57	62 37 49,57	37 51,22	37 48,16	-1,65	+ 1,41	10,728
1826	—	—	3 —	106 1 55,47	—	1 53,83	—	+ 1,64	10,710
1827	—	—	3 —	97 55 44,80	—	55 41,40	—	+ 3,40	10,649
1828	—	—	3 —	85 5 50,89	—	5 45,70	—	+ 5,19	10,590
1829	—	—	3 —	115 23 23,91	—	23 19,53	—	+ 4,38	10,540
1830	—	—	3 12 42,98	71 42 42,98	—	42 41,96	—	+ 1,02	10,502
1831	4 42 33,43	1 43 32,49	—	134 42 33,24	—	42 27,26	—	+ 5,98	10,434
1832	5 43 21,85	—	—	66 43 21,85	43 26,97	43 21,74	-2,12	+ 3,11	10,400
1833	—	—	5 51 11,97	100 51 11,97	51 11,88	51 6,15	+0,09	+ 5,82	10,395
1834	—	—	3 —	81 32 41,90	—	32 36,62	—	+ 5,28	10,371
1835	—	1 20 14,29	5 20 13,64	126 20 13,75	—	20 10,86	—	+ 2,89	10,364
1836	2 20 19,29	1 20 17,73	2 20 18,88	109 20 18,81	20 17,76	20 14,23	+1,05	+ 4,58	10,355
1837	—	—	4 12 30,75	110 12 30,75	12 23,47	12 20,12	+5,28	+ 10,63	10,267
1838	—	—	4 24 31,62	110 24 31,62	24 23,67	24 19,54	+4,95	+ 12,08	10,214
1839	4 29 33,22	—	—	43 29 33,22	29 37,33	29 35,03	-4,11	- 1,81	10,211
1840	—	—	3 52 11,39	115 52 11,39	—	52 7,13	—	+ 4,26	10,180
1841	—	—	3 —	102 17 10,42	—	17 8,88	—	+ 1,54	10,159
1842	6 59 4,00	1 59 3,40	—	30 59 3,92	59 4,21	59 2,04	-0,29	+ 1,91	10,118
1843	—	—	4 39 13,87	79 39 13,87	—	39 3,07	—	+ 10,80	10,058
1844	—	—	—	79 27 55,48	—	27 48,27	—	+ 7,21	10,022
1845	4 14 30,27	1 14 42,01	—	153 14 30,82	—	14 36,33	—	+ 3,49	10,016

lxxxiv *Comparison of the Observed Places of the Principal Fixed Stars*

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^b January 1, 1832		A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	s.	s.		Green.	A. S.	
			s.	s.	s.						
1846	6	47 Serpentis	—	—	2 22,61	h. m. s.	6.	s.	s.	s.	
1847	5.6	7 Herculis	—	—	—	16 0 —	—	22,06	+0,56	+2,885	
1848	6	Scorpii	—	—	—	16 0 —	—	29,23	—	2,703	
1849	6	12 Scorpii	—	—	3 54,25	16 0 —	—	37,07	—	3,709	
1850	5	13 Scorpii	—	—	—	16 1 54,23	—	53,73	+0,50	3,685	
1851	4	14 Scorpii	6 14,69	—	—	16 1 —	—	58,80	—	3,673	
1852	5	15 Scorpii	1 49,46	—	3 49,68	16 2 14,69	14,66	14,74	+0,03	-0,05	3,469
1853	6	16 Scorpii	—	—	—	16 2 49,62	—	49,37	+0,25	3,265	
1854	6	Scorpii	—	2 48,70	—	16 3 —	—	1,17	—	3,234	
1855	6	48 Serpentis	—	—	—	16 3 48,69	—	48,69	0,00	3,515	
1856	6	10 Herculis	—	—	—	16 3 —	—	62,14	—	2,708	
1857	6	17 Scorpii	—	—	3 34,07	16 4 —	—	28,57	—	2,549	
1858	6	9 Herculis	—	—	4 57,29	16 4 34,06	—	33,80	+0,26	3,304	
1859	3	1 Ophiuchi	—	5 33,11	13 33,00	16 4 57,29	—	56,88	+0,41	2,956	
1860	5	18 Scorpii	6 29,92	—	—	16 5 33,03	32,95	32,60	+0,08	+0,43	3,135
1861	7	Scorpii	1 11,04	—	3 10,75	16 6 29,92	—	29,43	+0,49	3,231	
1862	5	Normæ	5 18,44	—	—	16 6 —	—	10,95	-0,14	3,469	
1863	5	Apodis	—	—	—	16 7 18,44	—	18,08	+0,36	4,453	
1864	5.6	Scorpii	—	—	3 53,83	16 7 Invis.	—	59,46	—	3,860	
1865	6	Ophiuchi	—	—	4 5,70	16 7 53,81	—	53,61	+0,20	3,700	
1866	6	Scorpii	—	—	—	16 8 5,70	—	—	—	3,141	
1867	6	17 Herculis	—	—	3 7,08	16 8 —	—	55,83	—	3,764	
1868	7	Scorpii	—	3 18,26	—	16 9 7,10	—	6,94	+0,16	2,553	
1869	3	2 Ophiuchi	5 26,38	—	—	16 9 18,24	—	17,74	+0,50	3,494	
1870	6	18 Cor Bor	—	—	2 1,06	16 9 26,38	26,32	26,16	+0,06	+0,22	3,156
1871	5.6	19 Scorpii	—	—	3 32,54	16 10 1,09	—	0,72	+0,37	2,395	
1872	4	20 Scorpii	1 59,56	1 59,37	1 59,30	16 10 32,52	32,59	32,20	+0,13	+0,32	3,590
1873	5	50 Serpentis	4 34,13	—	2 34,33	16 10 59,40	59,42	59,31	-0,02	+0,09	3,626
1874	5	4 Ophiuchi	6 17,12	3 17,04	—	16 13 34,20	—	34,55	-0,35	3,038	
1875	3.4	20 Herculis	5 30,73	—	1 30,71	16 14 17,09	17,05	17,00	+0,04	+0,09	3,495
1876	4	22 Herculis	—	—	5 41,47	16 14 30,73	30,71	30,49	+0,02	+0,24	2,643
1877	5	5 Ophiuchi	—	—	3 31,36	16 14 41,52	41,73	41,60	-0,21	-0,08	1,797
1878	7	Scorpii	—	—	2 31,56	16 15 31,34	31,63	31,37	-0,29	-0,03	3,573
1879	5	19 Cor Bor	—	—	3 33,02	16 15 31,54	—	31,61	-0,07	3,577	
1880	5	20 Cor Bor	—	—	5 2,09	16 15 33,05	—	33,13	-0,08	2,339	
1881	5	21 Cor Bor	—	—	—	16 16 2,12	—	1,84	+0,28	2,252	
1882	5	7 Ophiuchi	3 17,82	3 17,91	2 18,05	16 16 —	—	9,14	—	2,255	
1883	5	51 Serpentis	1 40,02	—	—	16 17 17,91	—	17,08	+0,83	3,461	
1884	5	3 Ophiuchi	—	—	3 43,63	16 17 40,02	—	40,24	-0,22	2,758	
1885	1	21 Scorpii	—	14 7,33	14 7,24	16 18 43,63	7,09	43,02	+0,61	3,237	
1886	5	Apodis	—	—	—	16 19 7,26	—	7,25	+0,17	-0,09	3,659
1887	5	25 Herculis	—	—	4 24,97	16 19 Invisible	—	29,54	—	3,362	
1888	6	22 Scorpii	—	—	—	16 19 25,01	—	24,76	+0,25	2,130	
1889	5	Normæ	5 25,38	—	—	16 20 —	—	0,47	—	3,626	
1890	7	Scorpii	—	—	3 4,95	16 20 25,38	—	25,46	-0,08	3,895	
			—	—	—	16 21 4,92	—	4,88	+0,04	3,664	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in-					Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	1831		1832		1833				Green.	A. S. C.	
	No.	"	No.	"	No.						
1846	—	—	3	—	81 0 47,38	—	0 47,81	—	0,43	+10,001	
1847	—	—	3	—	72 29 56,64	—	29 57,37	—	0,73	9,991	
1848	—	—	3	—	118 52 52,90	—	57 48,91	+	1,99	9,984	
1849	—	—	3	—	117 58 16,30	—	58 16,88	—	0,58	9,887	
1850	2 28 55,72	—	—	—	117 28 55,72	—	28 55,61	+	0,11	9,880	
1851	—	—	5	0 55,32	109 0 55,32	0 59,03	0 59,78	-3,71	—	4,46	9,860
1852	—	—	4	37 23,39	99 37 23,39	—	37 16,75	+	6,64	9,815	
1853	—	—	1	6 21,91	98 6 21,91	—	6 18,35	+	3,56	9,800	
1854	—	—	4	57 51,16	110 57 51,16	—	57 48,92	+	2,24	9,740	
1855	—	—	3	—	72 53 34,90	—	53 31,77	+	2,13	9,734	
1856	—	—	3	—	66 3 54,83	—	3 55,52	—	0,69	9,687	
1857	—	—	3	—	101 24 6,58	—	24 7,55	—	0,97	9,682	
1858	—	—	3	32 38,34	84 32 38,34	—	32 33,59	+	4,75	9,652	
1859	6 15 16,59	1 15 16,34	6	15 17,81	93 15 17,13	15 18,17	15 13,51	-1,04	+	3,62	9,607
1860	5 55 3,12	—	—	—	97 55 3,12	—	55 0,71	+	2,41	10,064*	
1861	—	—	4	40 48,35	109 40 48,35	—	40 42,64	+	5,71	9,482	
1862	—	—	5	44 6,08	139 44 6,08	—	44 1,01	+	5,07	9,475	
1863	—	—	—	—	168 29 Invisible	—	29 58,87	—	—	9,433	
1864	—	—	4	11 16 50	118 11 16,50	—	11 11,82	+	4,68	9,427	
1865	—	—	4	31 48,91	93 31 48,91	—	31 51,98	—	3,07	9,410	
1866	—	—	3	—	120 29 28,32	—	29 27,31	+	1,01	9,347	
1867	—	—	3	—	66 27 14,40	—	27 10,93	+	3,47	9,330	
1868	—	—	3	—	109 48 5,90	—	48 1,43	+	4,47	9,318	
1869	5 16 33 86	—	—	—	94 16 33,86	16 35,50	16 29,77	-1,64	+	4,09	9,306
1870	—	—	5	25 47,71	60 25 47,71	—	25 40,37	+	7,34	9,260	
1871	—	—	4	45 21,06	113 45 21,06	45 22,45	45 22,50	-1,39	—	1,44	9,222
1872	5 10 58,01	1 10 58,11	—	—	115 10 58,03	10 54,35	10 49,07	+3,68	+	8,96	9,187
1873	4 34 9,63	1 34 7,61	—	—	88 34 9,23	—	34 8,87	+	0,36	8,984	
1874	—	1 38 12,68	3	38 15,22	109 38 14,59	38 11,65	38 9,22	+2,94	+	5,37	8,930
1875	2 26 50,03	—	4	26 51,06	70 26 50,72	26 48,81	26 43,08	+1,91	+	7,64	8,910
1876	—	—	5	16 56,82	43 16 56,82	16 58,83	17 11,25	-2,01	—	14,43	8,893
1877	—	—	5	3 7,42	113 3 7,42	3 8,68	3 2,95	-1,26	+	4,47	8,833
1878	—	—	1	0 35,91	113 0 35,91	—	0 35,37	+	0,54	8,833	
1879	5 42 45,41	—	—	—	58 42 45,41	—	42 45,15	+	0,26	8,827	
1880	—	—	5	18 5,07	53 18 5,07	—	18 0,21	+	4,86	8,790	
1881	—	—	3	—	55 51 3,65	—	51 2,51	+	1,14	8,780	
1882	—	1 3 59,35	4	4 2,01	108 4 1,48	—	3 59,39	+	2,03	8,694	
1883	—	—	4	31 28,69	75 34 28,69	—	34 17,76	+	10 93	8,662	
1884	5 59 15,45	—	—	—	97 59 15,45	—	59 17,51	—	2,06	8,580	
1885	5 3 4,58	10 3 5,20	13	3 4,74	116 3 4,88	3 3,03	3 1,25	+1,85	+	3,63	8,550
1886	—	—	—	—	167 8 Invisible	—	8 23,92	—	—	—	8,533
1887	4 13 8,88	—	2	13 9,49	52 13 9,08	—	13 8,56	+	5,22	8,522	
1888	—	—	3	—	114 44 17,24	—	44 9,98	+	7,26	8,479	
1889	—	1 19 48,62	4	19 49,21	124 19 49,09	—	19 45,33	+	3,76	8,447	
1890	—	—	3	—	116 9 46,50	—	9 46,81	—	0,81	8,394	

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833				Green.	A. S.	
			s.	s.	s.				s.	s.	
1891	4.5	8 Ophiuchi ϕ	6 32,08	2 32,09	—	16 24 32,07	31,94	32,04	+0,13	+0,03	+3,422
1892	3	14 Draconis η	3 43,80	—	—	16 21 43,80	43,69	43,06	+0,11	+0,74	0,792
1893	5	9 Ophiuchi ω	—	—	3 11,16	16 22 11,14	11,37	11,31	-0,23	-0,17	3,537
1894	4	10 Ophiuchi λ	—	—	3 26,87	16 22 26,87	26,05	26,44	-0,08	+0,43	3,018
1895	5	21 Ursæ Min. η	—	—	1 29,69	16 22 30,01	—	30,65	—	-0,64	-1,867
1896	2.3	27 Herculis β	—	8 0,08	—	16 23 0,10	0,11	59,42	-0,01	+0,65	+2,579
1897	5	30 Herculis g	2 7,38	—	1 7,45	16 23 7,43	—	7,53	—	-0,10	1,961
1898	5.6	28 Herculis n	—	—	5 20,10	16 24 20,10	—	19,50	—	+0,60	2,942
1899	4.5	29 Herculis h	—	—	4 44,97	16 24 44,98	44,85	44,84	+0,13	+0,14	2,811
1900	3.4	23 Scorpii τ	6 26,36	—	—	16 25 26,36	26,31	26,31	+0,05	+0,05	3,715
1901	5	12 Ophiuchi	1 32,68	—	4 32,55	16 27 32,57	—	32,01	—	+0,56	3,110
1902	3.4	13 Ophiuchi ζ	6 55,93	—	2 55 08	16 27 55,04	55,02	54,87	+0,02	+0,17	3,290
1903	4.5	15 Draconis d	—	—	3 20,13	16 28 20,31	20,80	20,66	-0,49	-0,35	-0,161
1904	6	33 Herculis	—	—	4 42,41	16 28 42,41	—	41,72	—	+0,69	+2,907
1905	4	35 Herculis σ	2 41,29	1 41,42	—	16 28 41,34	41,50	41,08	-0,16	+0,26	1,928
1906	2	Triang Aus a	—	—	3 58,34	16 30 58,17	—	58,96	—	-0,79	6,239
1007	5	24 Scorpii m	5 52,01	1 52,01	—	16 31 52,01	52,05	51,87	-0,04	+0,14	3,456
1908	6.7	Scorpii	—	—	5 30,68	16 34 30,66	—	30,44	—	+0,22	3,735
1909	3	40 Herculis ζ	6 57,33	15 57,31	—	16 34 57,32	57,33	56,86	-0,01	+0,16	2,246*
1910	4	Aræ η	5 19,43	—	—	16 35 19,43	—	20,06	—	-0,63	5,119
1911	6	25 Scorpii	—	—	4 34,93	16 36 34,90	—	35,32	—	-0,42	3,656
1912	6	16 Ophiuchi l	—	—	4 57,95	16 37 57,95	—	57,83	—	+0,12	3,039
1913	3	44 Herculis η	5 8,26	—	—	16 37 8,26	8,40	7,60	-0,14	+0,66	2,047
1914	5	43 Herculis i	2 46,34	—	3 46,34	16 37 46,34	—	45,73	—	+0,61	2,872
1915	3	26 Scorpii ϵ	—	3 18,10	1 18,14	16 39 18,08	18,51	17,83	-0,46	+0,25	3,870*
1916	5.6	45 Herculis e	—	—	4 30,44	16 39 30,45	—	30,11	—	+0,34	2,946
1917	6	18 Ophiuchi u	—	—	1 31,36	16 39 31,34	—	31,57	—	-0,23	3,035
1918	5	18 Draconis g	—	—	2 45,69	16 39 45,85	—	45,74	—	+0,11	0,387
1919	3.4	Scorpii μ^1	—	—	2 30,42	16 40 30,39	—	30,28	—	+0,11	4,041
1920	5	20 Ophiuchi r	7 32,89	—	—	16 40 32,89	—	32,57	—	+0,32	3,300
1921	4	Scorpii μ^2	—	—	—	16 40 —	—	58,39	—	—	4,040
1922	5	47 Herculis k	5 10,33	—	—	16 42 10,11	—	10,11	—	0,00	2,901
1923	6	21 Ophiuchi	—	1 54,08	3 54,11	16 42 54,10	—	53,84	—	+0,26	3,035
1924	6.7	Scorpii	2 30,83	—	3 30,70	16 43 30,74	—	30,59	—	+0,15	3,531
1925	5	50 Herculis s	5 5,79	—	—	16 44 5,79	—	5,15	—	+0,64	2,336
1926	5	52 Herculis	—	—	3 18,94	16 44 18,98	19,35	18,70	-0,37	+0,28	1,746
1927	6	49 Herculis	—	—	4 26,34	16 44 26,36	—	26,10	—	+0,26	2,723
1928	6.7	22 Ophiuchi	—	—	2 42,24	16 44 42,22	—	42,01	—	+0,21	3,611
1929	3.4	Aræ ζ	—	—	—	16 44 —	—	45,66	—	—	4,922
1930	6	51 Herculis X^2	—	—	—	16 44 —	—	47,19	—	—	2,480
1931	5	23 Ophiuchi q	3 37,33	—	—	16 45 37,33	—	37,15	—	+0,18	3,198
1932	4	25 Ophiuchi i	—	—	3 3,84	16 46 3,85	3,69	3,59	+0,16	+0,26	2,834
1933	4.5	Aræ ϵ	—	—	—	16 46 —	—	14,13	—	—	4,743
		Ophiuchi	1 20,62	1 20,51	1 20,48	16 46 20,63	—	20,11	—	+0,52	3,444
		Herculis	7 35,95	—	—	16 46 35,95	—	35,38	—	+0,57	2,276

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
1891	—	—	5 14 21,66	106 14 21,66	14 20,17	14 16,56	+1,49	+ 5,10	+ 8,358
1892	—	—	5 6 11,59	28 6 11,59	6 14,64	6 13,93	-3,05	- 2,34	8,336
1893	—	1 5 57,65	4 5 58,37	111 5 58,23	—	5 51,01	—	+ 7,22	8,306
1894	—	—	3 38 32,27	87 38 32,27	38 29,22	38 25,11	+3,05	+ 7,16	8,284
1895	—	—	3 —	13 51 44,18	—	51 39,20	—	+ 4,98	8,266
1896	6 8 17,19	6 8 16,45	1 8 16,25	68 8 16,78	8 20,18	8 18,27	-3,40	- 1,49	8,239
1897	2 44 37,68	—	1 44 37,62	47 44 37,66	—	44 36,41	—	+ 1,25	8,227
1898	—	—	3 —	84 6 52,52	—	6 49,94	—	+ 2,58	8,134
1899	—	—	3 8 40,86	78 8 40,86	8 41,76	8 39,43	-0,90	+ 1,43	8,100
1900	1 51 31,49	—	4 51 31,04	117 51 31,13	51 32,26	51 28,47	-1,13	+ 2,66	8,047
1901	—	—	3 57 35,27	91 57 35,27	—	57 28,14	—	+ 7,13	7,877
1902	6 13 15,72	—	—	100 13 15,72	13 11,51	13 6,65	+4,18	+ 9,07	7,847
1903	—	—	4 52 10,44	20 52 10,44	52 7,75	52 5,00	+2,69	+ 5,44	7,803
1904	—	1 32 34,74	3 32 41,18	82 32 41,18	—	32 35,91	—	+ 5,27	7,783
1905	5 12 45,04	1 12 48,47	3 12 45,52	47 12 45,57	12 45,20	12 42,25	+0,37	+ 3,32	7,781
1906	—	—	5 42 19,65	158 42 19,65	—	42 12,17	—	+ 7,48	7,607
1907	6 24 33,15	—	—	107 24 33,15	—	24 30,18	—	+ 2,97	7,528
1908	—	1 11 18,35	4 11 18,05	118 11 18,11	—	11 16,86	—	+ 1,25	7,314
1909	4 5 16,01	10 5 16,55	8 5 16,38	58 5 16,38	5 19,16	5 11,97	-2,78	+ 4,41	7,275
1910	—	—	5 43 41,41	148 43 44,41	—	43 39,24	—	+ 5,17	7,251
1911	—	—	5 12 56,48	115 12 56,48	—	12 48,09	—	+ 8,39	7,145
1912	—	—	5 39 48,56	88 39 48,56	—	39 50,08	—	- 1,52	7,112
1913	7 45 15,68	—	1 45 16,33	50 45 15,74	45 15,34	45 11,05	+0,40	+ 4,69	7,096
1914	—	—	3 6 18,40	81 6 18,40	—	6 11,49	—	+ 6,91	7,046
1915	5 58 45,71	—	—	123 58 45,71	—	58 46,30	—	- 0,59	6,923
1916	—	—	2 26 45,90	84 26 45,90	—	26 36,62	—	+ 9,28	6,904
1917	—	1 20 7,82	—	114 20 7,82	—	20 11,68	—	- 3,86	6,904
1918	—	—	6 5 35,66	25 5 35,66	—	5 26,39	—	+ 9,27	6,876
1919	—	—	5 14 59,90	127 44 59,90	—	44 55,89	—	+ 4,01	6,825
1920	4 28 43,48	—	1 28 45,73	100 28 43,93	—	28 36,08	—	+ 7,85	6,819
1921	—	—	3 43 17,95	127 43 17,95	—	43 15,06	—	+ 2,89	6,786
1922	5 27 21,19	—	—	82 27 21,19	—	27 16,61	—	+ 4,58	6,684
1923	—	—	3 29 26,35	88 29 26,35	—	29 20,69	—	+ 5,66	6,625
1924	—	—	2 7 41,34	110 7 41,34	—	7 33,61	—	+ 7,73	6,575
1925	4 54 10,26	1 54 10,35	—	59 54 10,28	—	54 2,02	—	+ 8,26	6,525
1926	—	—	5 43 10,99	43 43 10,99	43 13,34	43 12,32	-2,35	- 1,33	6,504
1927	—	—	3 —	74 44 18,80	—	44 11,34	—	+ 7,46	6,497
1928	—	—	3 —	113 13 40,21	—	13 38,52	—	+ 1,69	6,477
1929	5 42 51,30	—	—	145 42 51,30	—	42 45,07	—	+ 6,23	6,476
1930	—	—	3 —	65 3 22,85	—	3 10,91	—	+ 11,94	6,467
1931	1 52 18,40	—	4 52 20,50	95 52 20,08	—	52 15,80	—	+ 4,28	6,400
1932	—	1 33 7,32	4 33 8,78	79 33 8,49	33 6,10	32 53,12	+2,39	+ 10,37	6,362
1933	—	—	4 53 28,71	142 53 28,71	—	53 25,05	—	+ 3,66	6,353
1934	—	—	3 —	106 31 49,34	—	31 53,58	—	- 3,74	6,341
1935	5 0 54,41	—	—	58 0 54,41	—	0 55,80	—	- 1,39	6,317

lxxxviii *Comparison of the Observed Places of the Principal Fixed Stars*

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Greenb January 1, 1832		Greenb Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	1832				Green.	A. S.	
			s.	s.	s.	h. m.	s.					
1936	6.7	24 Ophiuchi	—	—	3 40,82	16 46 40,80	—	40,55	—	+ 0,25	+ 3,603	
1937	6.7	Scorpii	—	—	2 12,21	16 47 12,19	—	12,02	—	+ 0,17	3,512	
1938	5.6	54 Hercules	—	—	—	16 47 —	—	59,07	—	—	2,638	
1939	6	Ophiuchi	—	—	—	16 49 —	—	41,58	—	—	3,657	
1940	4	27 Ophiuchi <i>x</i>	5 43,40	2 43,28	—	16 49 43,36	43,36	43,04	0,00	+ 0,32	2,852	
1941	6	26 Ophiuchi <i>x</i>	—	—	2 52,93	16 49 52,90	—	51,81	—	+ 1,09	3,655	
1942	7	Ophiuchi	1 58,15	—	4 58,18	16 49 58,16	—	57,66	—	+ 0,50	3,481	
1943	6	Scorpii <i>p</i>	—	—	3 1,61	16 51 1,58	—	1,51	—	+ 0,07	3,862	
1944	6	29 Ophiuchi <i>s</i>	—	1 1,88	4 2,07	16 52 2,00	—	2,31	—	- 0,31	3,499	
1945	6	30 Ophiuchi <i>p</i>	—	—	2 12,35	16 52 12,35	—	12,29	—	+ 0,06	3,156	
1946	7	28 Ophiuchi	1 41,36	—	—	16 53 41,36	—	40,02	—	+ 0,44	3,677	
1947	5	Scorpii <i>k</i>	5 47,13	—	—	16 53 47,13	—	47,13	—	0,00	3,928	
1948	3	58 Hercules <i>e</i>	5 51,90	3 51,94	3 51,90	16 53 51,92	51,91	51,41	+ 0,01	+ 0,51	2,293	
1949	7	Scorpii	—	—	3 49,87	16 54 49,84	—	48,57	—	+ 1,27	3,541	
1950	5	19 Draconis <i>h</i>	—	—	3 6,72	16 55 6,88	—	6,05	—	+ 0,83	0,266	
1951	6	Ophiuchi	—	—	4 15,47	16 55 15,46	—	15,31	—	+ 0,15	3,314	
1952	5	59 Hercules <i>d</i>	1 24,15	—	—	16 55 24,15	—	23,61	—	+ 0,54	2,208	
1953	5.6	32 Ophiuchi	—	—	—	16 55 —	—	25,94	—	—	2,740	
1954	6	28 Scorpii	—	—	3 10 55	16 56 10,53	—	10,84	—	- 0,31	3,569	
1955	6	34 Ophiuchi	—	—	1 14,86	16 56 14,88	—	14,40	—	+ 0,48	2,752	
1956	6	Ophiuchi	—	—	—	16 56 —	—	52,76	—	—	3,083	
1957	5	60 Hercules	4 35,30	—	—	16 57 35,30	—	35,35	—	- 0,05	2,771	
1958	6.7	Ophiuchi	—	—	4 30,13	16 58 30,11	—	29,90	—	+ 0,21	3,471	
1959	6	Ophiuchi	—	—	5 33,93	16 59 33,93	—	33,63	—	+ 0,30	3,087	
1960	4	Scorpii <i>v</i>	5 7,53	—	—	17 0 7,53	—	7,97	—	- 0,44	4,272	
1961	2.3	35 Ophiuchi <i>v</i>	6 45,21	1 45,06	10 45,23	17 0 45,20	45,15	45,00	+ 0,05	+ 0,20	3,426	
1962	4	21 Draconis <i>μ</i>	3 51,61	—	—	17 1 51,61	51,58	51,29	+ 0,03	+ 0,32	1,242	
1963	5	Herculis	—	—	3 4,62	17 2 4,58	—	4,81	—	- 0,23	+ 2,123	
1964	4	22 Ursæ Min <i>ε</i>	—	—	1 —	17 3 26,89	27,43	24,26	- 0,54	+ 2,63	- 6,577*	
1965	6.7	29 Scorpii	—	—	—	17 3 —	—	47,14	—	—	+ 3,722	
1966	5	37 Ophiuchi.	6 32,86	—	—	17 4 32,86	—	32,71	—	+ 0,15	2,821	
1967	4.5	36 Ophiuchi <i>Δ</i>	3 1,68	—	3 1,60	17 5 1,63	1,62	1,96	+ 0,01	- 0,33	3,671*	
1968	7	30 Scorpii	—	—	1 54,65	17 5 54,62	—	55,28	—	- 0,66	3,671*	
1969	5.6	Scorpii <i>u</i>	—	—	3 8,52	17 6 8,48	—	8,57	—	- 0,09	3,895	
1970	3.4	64 Hercules <i>α</i>	8 59,46	24 59,48	20 59,43	17 6 59,47	59,44	59,02	+ 0,03	+ 0,45	2,729	
1971	6.7	31 Scorpii	—	—	3 13,87	17 7 13,84	—	13,69	—	+ 0,15	3,715	
1972	6.7	Scorpii	—	—	3 24,80	17 7 24,76	—	24,30	—	+ 0,46	3,892	
1973	5.6	39 Ophiuchi <i>o</i>	—	—	—	17 7 —	—	46,48	—	—	3,650	
1974	6	Ophiuchi	—	—	—	17 7 —	—	51,61	—	—	3,644	
1975	4.5	41 Ophiuchi <i>o</i>	3 59,70	—	—	17 7 59,70	58,99	59,65	—	+ 0,05	3,074	
1976	4	65 Hercules <i>δ</i>	—	—	3 8,10	17 8 8,13	8,07	7,06	+ 0,06	+ 1,07	2,460	
1977	3	22 Draconis <i>ζ</i>	—	—	—	17 8 —	19,40	18,48	—	—	0,153	
1978	3.4	67 Hercules <i>π</i>	5 11,99	—	—	17 9 11,99	12,06	11,74	- 0,07	+ 0,25	2,286	
1979	6.7	Ophiuchi	—	—	—	17 10 —	—	6,83	—	—	3,481	
1980	6	66 Hercules <i>ω</i>	—	—	3 43,28	17 10 43,29	—	43,06	—	+ 0,23	2,813	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
	"	"	"						
1936	—	—	4 52 30,21	112 52 30,21	—	52 29,29	+	0,92	+ 6,314
1937	—	—	3 —	109 15 58,22	—	15 55,83	+	2,39	6,270
1938	—	—	3 —	71 17 30,95	—	17 27,57	+	3,88	6,202
1939	—	—	5 49 48,88	114 49 48,88	—	49 41,83	+	6,55	6,063
1940	7 21 31,47	—	5 21 31,41	80 21 31,45	21 31,32	21 21,13	+ 0,13	+ 10,32	6,058
1941	—	—	4 43 33,04	114 43 33,04	—	43 29,44	+	3,60	6,048
1942	—	—	4 —	107 58 42,17	—	58 39,36	+	2,81	6,040
1943	—	—	3 —	121 53 5,19	—	53 1,89	+	3,30	5,952
1944	—	—	3 —	108 37 46,14	—	37 45,14	+	1,00	5,866
1945	—	—	3 —	98 57 47,63	—	57 41,15	+	6,48	5,852
1946	—	—	3 27 1,97	115 27 1,97	—	26 58,16	+	3,81	5,729
1947	5 52 38,09	—	—	123 52 38,09	—	52 38,56	—	0,47	5,721
1948	5 49 14,38	2 49 13,76	4 49 14,73	58 49 14,40	49 16,17	49 12,93	-1,77	+ 1,47	5,711
1949	—	—	3 15 7,10	110 15 7,10	—	14 57,14	+	9,96	5,634
1950	5 36 31,95	—	—	24 36 31,95	—	36 25,49	+	6,46	5,601
1951	—	—	3 —	100 50 43,48	—	50 38,98	+	4,50	5,596
1952	1 11 1,32	2 11 2,15	4 11 2,24	56 11 2,08	—	10 57,74	+	4,34	5,582
1953	—	—	3 —	75 39 33,09	—	39 27,77	+	10,32	5,580
1954	—	—	3 —	111 19 22,82	—	19 22,85	—	0,03	5,519
1955	—	—	—	76 10 —	—	10 54,70	—	—	5,512
1956	—	—	5 39 17,82	99 39 17,82	—	39 12,65	+	5,17	5,459
1957	5 1 20,69	—	—	77 1 20,69	—	1 14,66	+	6,03	5,399
1958	—	—	4 22 48,59	107 22 48,59	—	22 41,07	+	7,52	5,324
1959	—	—	3 —	90 51 4,59	—	50 59,93	+	4,66	5,233
1960	4 0 24,08	—	1 0 24,96	133 0 24,25	—	0 15,74	+	8,51	5,188
1961	5 30 34,54	—	—	105 30 34,54	30 33,63	30 26,28	+ 0,91	+ 8,26	5,134
1962	5 18 24,52	—	3 18 24,87	35 18 24,65	18 22,55	18 20,86	+ 2,10	+ 3,79	5,034
1963	—	—	4 50 29,86	53 50 29,86	—	50 28,92	+	0,94	5,018
1964	—	—	5 41 53,72	7 41 53,72	41 58,96	41 55,63	-0,24	+ 3,09	4,881
1965	—	—	3 —	116 46 28,26	—	46 23,95	+	4,31	4,877
1966	—	—	5 12 17,53	79 12 17,53	—	12 12,48	+	5,05	4,810
1967	5 20 52,23	—	—	116 20 52,23	20 49,60	20 46,96	+ 2,63	+ 5,27	6,021*
1968	—	—	3 17 42,88	116 17 42,88	—	17 41,59	+	1,29	5,936*
1969	—	—	3 —	122 27 43,56	—	27 47,18	—	3,62	4,678
1970	9 24 45,13	—	—	75 24 45,13	24 43,00	24 37,70	+ 2,13	+ 7,43	4,603
1971	—	—	3 —	116 26 5,03	—	26 1,30	+	3,73	4,585
1972	—	—	—	122 21 —	—	21 38,74	—	—	4,570
1973	—	—	—	114 5 —	—	5 39,65	—	—	4,538
1974	—	—	—	113 52 —	—	52 33,93	—	—	4,531
1975	5 14 54,35	—	—	90 14 54,35	14 56,82	14 53,37	-2,47	+ 0,98	4,518
1976	—	—	5 57 30,83	64 57 30,83	57 26,35	57 29,42	+ 4,48	+ 10,41	4,506
1977	—	—	5 4 41,17	24 4 41,17	4 42,09	4 46,84	-0,92	+ 0,33	4,483
1978	1 59 49,10	—	4 59 46,42	52 59 46,95	59 49,53	59 46,62	-2,63	+ 0,33	4,413
1979	—	—	1 34 20,67	107 34 20,67	—	34 19,56	+	1,11	4,388
1980	—	—	—	78 56 —	—	56 42,84	—	—	4,285

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.		Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^h January 1, 1832			A. S. Catal.	Difference from		Annual Preces- sion	
				No. 1831		No. 1832	No. 1833		s.		s.	s.		s.
				s.	s.	s.	h. m. s.	s.						
1981	4.5	40 Ophiuchi	ρ	5 56,48	—	2 56,52	17 10 56,48	56,39	56,05	+0,09	+0,43	3,567		
1982	4	68 Hercules	μ	—	—	3 7,32	17 11 7,36	7,45	7,38	-0,09	-0,02	2,211		
1983	3	Aræ	γ	—	—	—	17 11 —	—	18,46	—	—	5,019		
1984	3	Aræ	β	—	—	—	17 11 —	—	20,97	—	—	4,958		
1985	4.5	53 Serpentis	ν	—	—	3 23,05	17 11 23,04	23,10	23,23	-0,06	-0,19	3,362		
1986	3.4	42 Ophiuchi	θ	5 42,13	—	—	17 11 42,13	42,08	41,88	+0,05	+0,25	3,672		
1987	4.5	69 Hercules	e	1 52,84	—	2 52,74	17 11 52,80	52,96	52,34	-0,16	+0,46	2,066		
1988	6	43 Ophiuchi	γ	—	—	4 47,85	17 12 47,82	47,68	47,78	+0,14	+0,04	3,762		
1989	5.6	70 Hercules	a	—	—	—	17 4 —	—	58,76	—	—	2,467		
1990	6	Scorpii	—	—	1 39,66	2 39,66	17 14 39,64	—	39,41	—	+0,23	3,578		
1991	7	33 Scorpii	—	—	—	5 50,63	17 14 50,60	—	50,67	—	-0,07	3,654		
1992	4	Aræ	δ	2 57,60	2 57,52	—	17 15 57,53	—	59,12	—	-1,59	5,389		
1993	5.6	44 Ophiuchi	b	—	—	2 7,00	17 16 6,98	7,11	7,11	-0,13	-0,13	3,652		
1994	5	45 Ophiuchi	d	4 38,08	2 38,32	—	17 16 38,15	—	38,09	—	+0,06	3,817		
1995	6.7	Ophiuchi	—	—	—	3 55,60	17 16 55,57	—	55,17	—	+0,40	3,813		
1996	6	73 Hercules	—	—	—	4 5,03	17 17 5,06	—	4,56	—	+0,50	2,507		
1997	6	47 Ophiuchi	—	—	—	3 36,89	17 17 36,88	—	37,04	—	-0,16	3,356		
1998	5.6	Ophiuchi	—	—	—	1 43 58	17 17 43,58	—	43,24	—	+0,34	3,181		
1999	4	75 Hercules	ρ	6 53,45	—	4 53,30	17 17 53,39	53,50	53,63	-0,11	+0,36	2,067		
2000	4.5	49 Ophiuchi	σ	5 10,93	—	—	17 18 10,93	11,10	10,89	-0,17	+0,04	2,969		
2001	3	Aræ	α	—	—	1 52,32	17 18 52,25	—	52,76	—	-0,51	4,620		
2002	3.4	34 Scorpii	ν	—	1 20,78	1 21,13	17 19 20,92	—	21,18	—	-0,26	4,064		
2003	6	Herculis	—	—	—	—	17 19 —	—	33,85	—	—	2,583		
2004	6	Ophiuchi	—	—	—	2 15,84	17 20 15,84	—	15,67	—	+0,17	3,057		
2005	5	51 Ophiuchi	e^2	2 10,37	3 10,45	—	17 21 10,41	10,32	10,61	+0,09	-0,20	3,649		
2006	6.7	Sagittarii	—	—	—	4 18,92	17 21 18,89	—	18,82	—	+0,07	3,714		
2007	3	35 Scorpii	λ	3 12,79	1 12,62	3 12,78	17 22 12,74	12,65	12,77	+0,09	-0,03	4,060		
2008	6	Ophiuchi	h	—	—	5 56,18	17 22 56,18	—	56,05	—	+0,13	3,002		
2009	4.5	76 Hercules	λ	6 57,41	—	1 57,11	17 23 57,11	57,12	56,93	-0,01	+0,18	2,417		
2010	5	Scorpii	—	4 59,57	2 59,32	—	17 24 59,48	—	59,17	—	+0,31	4,119		
2011	7	52 Ophiuchi	—	—	1 12,38	2 12,63	17 25 12,53	—	12,33	—	+0,20	3,599		
2012	5	Scorpii	θ	—	—	3 15,68	17 25 15,62	—	15,37	—	+0,25	4,294		
2013	6	78 Hercules	—	—	—	4 13,86	17 25 13,89	—	13,38	—	+0,51	2,350		
2014	6	54 Ophiuchi	—	—	—	—	17 25 —	—	37,02	—	—	2,756		
2015	6	53 Ophiuchi	f	—	—	1 38,58	17 26 38,59	—	38,32	—	+0,27	2,842		
2016	2	23 Draconis	β	1 38,54	—	2 38,64	17 26 38,66	38,44	38,01	+0,22	+0,65	1,349		
2017	2	55 Ophiuchi	α	4 8,27	26 8,45	16 8,43	17 27 8,42	8,39	7,97	+0,03	+0,45	2,770		
2018	6.7	Serpentis	—	—	—	3 58,00	17 27 57,98	—	58,04	—	-0,06	3,434		
2019	5	55 Serpentis	ξ	4 58,41	1 58,03	—	17 27 58,34	—	58,11	—	+0,23	3,430		
2020	6	2 Sagittarii	—	—	—	3 39,42	17 28 39,40	—	39,60	—	-0,20	3,597		
2021	5	57 Ophiuchi	μ	6 43,63	—	—	17 28 43,03	43,02	42,59	+0,01	+0,44	3,254		
2022	5	24 Draconis	ρ^1	—	—	—	17 28 —	—	52,14	—	—	1,156		
2023	5	25 Draconis	ν^2	—	—	—	17 28 —	—	57,12	—	—	1,157		
2024	7	Sagittarii	—	—	—	2 4,81	17 29 4,77	—	4,38	—	+0,39	3,898		
2025	5	Pavonis	η	—	—	—	17 29 —	—	17,18	—	—	5,860		

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.	
	1831		1832		1833					Green.	A. S. C.		
	No.	"	No.	"	No.	"				"	"		
1981			2 55	26,66	3 55	30,68	110 55	29,07	55 28,31	55 25,25	+ 0,76	+ 3,82	+ 4,268
1982	5 42	50,05					56 42	50,05	42 49,93	42 48,49	+ 0,12	+ 1,56	4,248
1983	4 12	27,84					146 12	27,84		12 27,22		+ 0,62	4,241
1984					2 21	37,65	145 21	37,65		21 27,06		+ 10,50	4,237
1985							102 40		40 8,74	40 1,86			4,229
1986					4 49	29,39	114 49	29,39	49 23,70	49 20,25	+ 5,69	+ 9,14	4,208
1987	5 31	38,08					52 31	38,08	31 40,78	31 31,20	- 2,70	+ 6,88	4,184
1988			2 58	12,36	2 58	12,15	117 58	12,25	58 14,24	58 10,83	- 1,99	+ 1,42	4,110
1989					3 19	41,33	65 19	41,33		19 36,56		+ 4,77	4,005
1990					4 16	34,62	111 16	34,62		16 33,70		+ 0,92	3,950
1991							114 4			4 50,07			3,934
1992	5 31	51,67					150 31	51,67		31 50,02		+ 1,65	3,841
1993			4 0	41,66	1 0	43,56	114 0	42,64	0 44,33	0 38,05	- 1,69	+ 4,59	3,825
1994	5 42	25,02					119 42	25,02		42 23,67		+ 1,45	3,781
1995					4 34	21,31	119 34	21,31		34 8,66		+ 12,65	3,756
1996							66 52			52 36,46			3,739
1997							102 21			21 16,50			3,695
1998							94 55			55 47,30			3,686
1999	6 41	40,83					52 41	40,83	41 41,58	41 34,03	- 0,75	+ 6,80	3,668
2000	5 42	27,30			2 42	27,99	85 42	27,50	42 25,37	42 19,15	+ 2,13	+ 8,35	3,645
2001	1 43	56,17	5 43	57,83			139 43	57,55		43 55,90*		+ 1,65	3,590
2002					5 9	9,06	127 9	9,06		9 4,07		+ 4,99	3,548
2003					4 46	14,69	69 46	14,69		46 10,18*		+ 4,51	3,525
2004					4 31	36,41	89 31	36,41		31 31,91*		+ 4,50	3,467
2005	5 49	24,60					113 49	24,60	49 23,34	49 23,41	- 3,74	+ 1,19	3,389
2006			3 7	55,92	2 7	57,93	116 7	56,72		7 53,60*		+ 3,22	3,378
2007	5 58	17,32					126 58	17,32		58 10,08		+ 7,24	3,301
2008					5 8	35,50	87 8	35,50		8 33,00*		+ 2,50	3,236
2009	5 45	23,21	1 45	19,94	5 45	22,74	63 45	22,69	45 26,03	45 25,00	- 3,34	- 2,31	3,147
2010			1 30	22,50	4 30	23,94	128 30	23,65		30 14,85*		+ 8,80	3,062
2011			3 55	19,99	1 55	21,88	111 55	20,46		55 15,89		+ 4,57	3,042
2012	4 52	48,33			1 52	49,45	132 52	48,55		52 45,24		+ 3,31	3,039
2013					4 27	53,38	61 27	53,38		27 54,30		+ 1,08	3,036
2014							76			42 57,78			2,917
2015							80			17 31,16			2,915
2016	4 34	14,46	1 34	15,49			37 34	14,67	34 16,6	34 14,71	- 2,01	- 0,04	2,912
2017	8 13	39,49	29 18	40,55	23 18	40,56	77 18	40,43	18 39,55	18 34,38	+ 0,88	+ 6,05	2,872
2018							105			27 34,50*			2,802
2019	5 17	10,19					105 17	10,19		17 0,83		+ 9,33	2,802
2020			3 48	16,67	1 48	15,07	111 48	16,27		48 8,58		+ 7,69	2,743
2021	3 0	29,82	1 0	27,03	1 0	23,78	96 0	29,06	0 33,16	0 27,41	- 4,10	+ 1,65	2,737
2022					2 41	56,57	34 41	56,57		41 51,39		+ 5,18	2,717
2023					5 42	39,72	34 42	39,72		42 35,15		+ 4,67	2,710
2024							122			5 46,89*			2,708
2025					1 37	49,77	154 37	49,77		37 38,89*		+ 10,88	2,695

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832		Green ^h Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	h. m.	s.			Green.	A. S.	
			s.	s.	s.							
2026	6	79 Herculis			2 36,06	17 30 36,09		35,37		+0,72	+2,466	
2027	3	Scorpii κ	3 52,69	4 52,52		17 30 52,58		52,32		+0,26	4,139	
2028	4.5	56 Serpentis σ	2 58,75	3 58,74		17 31 58,74	58,64	58,37	+0,10	+0,37	3,369	
2029	7	Sagittarii			3 43,72	17 32 43,69		43,02		+0,67	3,767	
2030	5	27 Draconis f			1 38,85	17 32 39,02		38,45		+0,57	-0,290	
2031	7	Serpentis			4 2,08	17 33 2,06		1,57		+0,49	+3,435	
2032	5	58 Ophiuchi D		1 21,83	4 22,20	17 33 22,11	21,68	22,32	+0,43	-0,21	3,593	
2033	7	Ophiuchi		3 16,45		17 34 16,44		16,28		+0,16	3,607	
2034	6	Ophiuchi			4 26,52	17 34 26,54		26,25		+0,29	2,686	
2035	4	85 Herculis i	5 43,58	1 43,70		17 34 43,61	43,59	42,80	+0,02	+0,81	1,688	
2036	3	60 Ophiuchi β	5 10,59	2 10,79		17 35 10,74	10,66	10,34	+0,08	+0,40	2,960	
2037	4.5	Scorpii i	3 50,74	1 50,38		17 35 50,65		50,08		+0,57	4,185	
2038	5.6	84 Herculis			4 28,02	17 36 28,05		27,51		+0,54	2,465	
2039	5	3 Sagittarii p	5 59,32	1 59,21		17 36 59,30	59,42	59,04	-0,12	+0,26	3,768	
2040	7	Sagittarii			3 57,79	17 37 57,76		57,75		+0,01	3,743	
2041	5	28 Draconis w			3 55,92	17 37 56,12		56,09		+0,03	-0,367	
2042	5.6	Sagittarii		3 15,93		17 38 15,91		15,95		-0,04	+3,887	
2043	4	Telescopii γ		2 26,12	4 25,88	17 38 25,92	25,63	25,30	+0,29	+0,62	4,070	
2044	7	Sagittarii			2 28,17	17 38 28,15		28,05		+0,10	3,852	
2045	4	62 Ophiuchi γ	3 28,33	2 28,43		17 39 28,37	28,36	28,01	+0,01	+0,36	3,003	
2046	4	86 Herculis μ	6 53,29			17 39 53,29	53,34	52,56	-0,05	+0,73	2,366	
2047	7	Sagittarii				17 40		23,91			3,852	
2048	6	87 Herculis		3 0,50	2 0,72	17 42 0,61		0,08		+0,53	2,427	
2049	6.7	63 Ophiuchi π		1 34,12	4 34,09	17 44 34,07		33,81		+0,26	3,685	
2050	7	Serpentis		3 39,29	3 39,61	17 46 39,43		39,22		+0,21	3,445	
2051	6	Serpentis			3 55,96	17 46 55,95		55,48		+0,47	3,162	
2052	5	Sagittarii	5 18,27	4 18,38	3 18,46	17 48 18,34		18,04		+0,30	3,845	
2053	5.6	89 Herculis			5 38,90	17 48 38,93		38,62		+0,31	2,415	
2054	5	4 Sagittarii b	5 32,26	1 32,38		17 49 32,23		32,27		+0,01	3,656	
2055	4	64 Ophiuchi ν	5 46,84	1 46,77		17 49 46,83	46,85	46,56	-0,02	+0,27	3,297	
2056	7	5 Sagittarii i			5 53,91	17 49 53,88		54,52		-0,64	3,670	
2057	6.7	Sagittarii			4 0,85	17 50 0,83		0,42		+0,41	3,562	
2058	4	91 Herculis δ	3 29,68			17 50 29,68	29,72	28,99	-0,04	+0,69	2,052	
2059	3.4	32 Draconis ξ			3 37,33	17 50 37,43	37,66	36,81	-0,23	+0,62	1,020	
2060	4	92 Herculis ζ			4 14,25	17 51 14,29	14,31	13,92	-0,02	+0,37	2,320	
2061	5	57 Serpentis ζ		2 36,27		17 51 36,27		36,53		-0,26	3,154	
2062	7	6 Sagittarii				17 51		38,15			3,480	
2063	6	Sagittarii			1 43,85	17 51 43,83		43,77		+0,06	3,028	
2064	5	66 Ophiuchi η				17 51		56,49			2,970	
2065	5	94 Herculis ν				17 52		3,85			2,291	
2066	4	67 Ophiuchi σ			3 14,15	17 52 14,15	14,11	13,76	+0,04	+0,39	2,999	
2067	6	7 Sagittarii a		4 33,58		17 52 33,87		32,37		+1,50	3,670	
2068	5	93 Herculis E			1 34,32	17 52 34,34		33,98		+0,36	2,666	
2069	6	Sagittarii				17 52		36,14			3,573	
2070	6	Tauri Pon				17 52		40,32			2,921	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N.P.D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
2026	—	—	—	65 —	—	34 57,80	—	—	+2,572
2027	5 55 59,78	—	—	128 55 59,78	—	55 58,37*	+	1,41	2,552
2028	4 46 35,83	1 46 33,72	—	102 46 35,41	46 40,44	46 34,57	-5,03	+ 0,84	2,455
2029	—	4 47 37,60	4 47 35,21	117 47 36,41	—	47 36,47*	—	0,06	2,391
2030	—	2 45 28,55	2 45 29,78	21 45 29,16	—	45 28,12	—	1,04	2,386
2031	—	—	5 28 9,01	105 28 9,01	—	28 4,86*	+	4,15	2,363
2032	3 35 34,24	1 35 31,88	1 35 34,52	111 35 33,82	35 34,72	35 27,48	-0,90	+ 6,34	2,334
2033	—	—	—	112 —	—	6 35,04*	—	—	2,256
2034	—	—	—	73 —	—	57 51,99*	—	—	2,239
2035	—	—	5 54 0,95	43 54 0,95	54 2,44	53 58,83	-1,49	+ 2,12	2,212
2036	7 21 24,16	1 21 23,30	—	85 21 24,05	21 21,66	21 16,08	+2,39	+ 7,97	2,176
2037	5 3 17,21	—	—	130 3 17,21	—	3 3,12*	+	14,09	2, 21
2038	—	1 35 36,09	4 35 35,26	65 35 35,43	—	35 32,36	+	3,07	2,062
2039	2 45 25,71	2 45 25,77	—	117 45 25,74	45 25,82	45 23,77	-0,08	+ 1,97	2,020
2040	—	2 54 19,53	2 54 21,82	116 54 20,67	—	54 19,37*	+	1,30	1,935
2041	—	—	4 9 55,04	21 9 55,04	—	9 52,14	+	2,90	1,926
2042	—	—	—	121 —	—	38 8,23*	—	—	1,909
2043	5 58 47,46	—	—	126 58 47,46	—	58 45,33*	+	2,13	1,896
2044	—	—	—	120 31 —	—	31 43,04*	—	—	1,891
2045	4 13 23,30	—	1 13 21,13	87 13 22,87	13 22,12	18 17,21	+0,75	+ 5,63	1,802
2046	6 10 31,91	—	7 10 33,77	62 10 32,91	10 33,31	10 32,38	-0,40	+ 0,53	2,604*
2047	—	1 29 55,78	4 29 57,89	120 29 57,47	—	29 53,67*	+	3,80	1,723
2048	—	5 18 37,03	—	64 18 37,03	—	18 54,58	—	17,55	1,579
2049	—	4 50 45,67	1 50 47,20	114 50 45,98	—	50 40,13*	—	—	1,360
2050	—	3 46 30,20	2 46 32,53	105 46 31,33	—	46 25,50	—	—	1,177
2051	—	1 2 58,93	4 2 59,75	94 2 59,58	—	2 57,14	+	2,44	1,065
2052	6 13 39,0	—	—	120 13 39,06	—	13 34,60	+	4,46	1,034
2053	—	—	5 55 2,63	63 55 2,63	—	54 58,42	+	4,21	1,000
2054	6 47 28,50	—	—	113 47 28,50	47 32,04	47 25,99	-3,54	+ 2,51	0,925
2055	5 44 46,19	—	—	99 44 46,19	44 43,21	44 33,91	+2,98	+ 12,28	0,903
2056	—	—	6 15 42,26	114 15 42,26	—	15 41,01	+	1,25	0,893
2057	—	—	6 19 8,52	110 19 8,52	—	19 4,23	+	4,29	0,884
2058	5 43 19,35	—	—	52 43 19,35	43 21,38	43 17,17	-2,03	+ 2,18	0,838
2059	—	6 5 54,36	—	33 5 54,36	5 51,63	5 56,49	-0,27	- 2,13	0,824
2260	4 43 43,90	3 43 44,96	—	60 43 44,35	43 42,90	43 38,05	+1,45	+ 6,30	0,773
2061	—	—	5 40 16,58	93 40 16,58	—	40 15,29	+	1,29	0,743
2062	—	—	—	107 —	—	8 27,08	—	—	0,741
2063	—	—	—	112 —	—	45 59,96	—	—	0,734
2064	—	—	—	85 —	—	48 46,88	—	—	0,713
2065	—	—	3 47 32,98	59 47 32,98	—	47 26,45	+	6,53	0,701
2066	—	—	5 3 13,54	87 3 13,54	3 13,31	3 6,25	+0,23	+ 7,29	0,688
2067	—	—	—	114 —	—	16 17,19	—	—	0,663
2068	—	—	—	73 —	—	13 55,23	—	—	0,658
2069	—	—	—	110 —	—	43 36,07	—	—	0,657
2070	—	—	—	83 —	—	43 2,17	—	—	0,648

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.		Green Catal.	A. S. Catal.	Difference from		Annual Preces- sion.
			No. 1831		No. 1832	No. 1833				Green	A. S.	
			s.	s.	s.	h. m. s.	s.					
2071	2	33 Draconis	γ	8 42,42	28 42,53	15 42,06	17 52 42,42	42,56	42,17	-0,14	+0,25	+1,388
2072	5.6	68 Ophiuchi	k	—	—	2 13,95	17 53 13,95	—	13,54	—	+0,41	3,037
2073	4	Aræ	θ	—	—	—	17 53 —	—	34,09	—	—	4,665
2074	6.7	9 Sagittarii	—	—	—	—	17 53 —	—	34,50	—	—	3,673
2075	5	69 Ophiuchi	τ	5 56,31	1 56,41	—	17 53 56,33	—	55,75	—	+0,58	3,260
2076	5	Sagittarii	γ ¹	—	—	3 17,67	17 54 17,63	—	17,09	—	+0,54	3,825
2077	5.6	95 Hercules	B	—	1 22,55	—	17 54 22,57	—	22,16	—	+0,41	2,539
2078	7	Sagittarii	—	—	1 52,44	2 52,58	17 54 52,52	—	52,02	—	+0,50	3,674
2079	4	10 Sagittarii	γ ²	5 1,17	—	—	17 55 1,17	1,27	1,39	-0,10	-0,22	3,852
2080	5	96 Hercules	Q	4 12,39	1 12,24	—	17 55 12,36	—	11,69	—	+0,67	2,560
2081	6	97 Hercules	—	—	—	3 28,91	17 55 28,93	—	23,11	—	+0,82	2,503
2082	4.5	70 Ophiuchi	p	5 58,07	1 58,09	3 58,25	17 56 58,13	58,02	57,94	+0,11	+0,19	3,009
2083	7	Sagittarii	—	—	—	—	17 57 —	—	6,87	—	—	3,593
2084	5	Draconis	—	—	—	—	17 57 —	—	57,12	—	—	-2,710
2085	5	Sagittarii	—	—	5 26,55	—	17 57 26,53	—	26,50	—	+0,03	+3,792
2086	5	Telescopii	e	—	3 45,92	—	17 58 45,89	—	45,67	—	+0,22	4,450
2087	5.6	98 Hercules	—	—	—	4 57,56	17 58 57,59	—	57,07	—	+0,52	2,523
2088	6	Sagittarii	—	—	—	5 16,21	17 59 16,17	—	15,95	—	+0,22	3,863
2089	6	71 Ophiuchi	S ¹	—	3 16,45	—	17 59 16,46	—	16,04	—	+0,42	2,863
2090	4	72 Ophiuchi	S ²	5 23,03	1 23,14	—	17 59 23,02	23,31	22,85	-0,29	+0,17	2,843
2091	4	103 Hercules	α	6 59,57	—	5 59,35	18 0 59,46	59,58	59,24	-0,12	-0,22	2,335
2092	6	73 Ophiuchi	q	—	—	3 13,04	18 1 13,05	—	12,66	—	+0,39	2,975
2093	6	Sagittarii	—	—	1 28,00	4 28,36	18 1 28,27	—	23,14	—	+0,13	3,655
2094	5.6	102 Hercules	C	—	—	4 34,65	18 1 34,65	—	33,88	—	+0,77	2,561
2095	6	101 Hercules	P	—	5 33,33	—	18 1 33,35	—	37,78	—	+0,57	2,581
2096	3.4	13 Sagittarii	μ ¹	5 43,26	1 43,32	1 43,01	18 3 43,23	43,22	42,79	+0,01	+0,44	3,583
2097	6	14 Sagittarii	—	—	5 10,53	—	18 4 10,52	—	10,34	—	+0,18	3,601
2098	6	15 Sagittarii	μ ²	—	—	5 11,78	18 5 11,76	11,79	11,42	-0,03	+0,34	3,575
2099	6	16 Sagittarii	—	—	—	4 13,49	18 5 13,47	—	12,76	—	+0,71	3,566
2100	5	104 Hercules	A	6 34,95	—	—	18 5 34,95	—	34,70	—	+0,25	2,254
2101	4	Telescopii	β	1 15,77	4 15,73	—	18 6 15,71	15,56	16,05	+0,15	-0,34	4,067
2102	7	17 Sagittarii	—	—	3 35,19	1 35,01	18 6 35,13	—	35,17	—	-0,04	3,570
2103	5.6	Sagittarii	g	—	—	5 32,45	18 7 32,42	—	32,14	—	+0,28	3,751
2104	7	Clypei Sob	—	—	1 37,76	4 37,48	18 7 37,50	—	37,15	—	+0,35	3,515
2105	3.4	19 Sagittarii	δ	5 14,43	1 14,35	7 14,34	18 10 14,34	14,43	14,10	-0,09	+0,24	3,835
2106	6	Clypei Sob	—	—	—	4 28,76	18 10 28,74	—	23,04	—	+0,70	3,447
2107	5	105 Hercules	G	5 15,95	1 16,13	—	18 12 15,98	—	15,42	—	+0,56	2,463
2108	6	74 Ophiuchi	r	—	1 28,82	3 29,11	18 12 29,03	—	29,21	—	-0,18	2,991
2109	4	58 Serpentis	γ	—	—	—	18 12 —	37,23	36,92	—	—	3,092*
2110	3	20 Sagittarii	e	5 1,23	2 1,22	—	18 12 1,22	1,35	1,29	-0,13	-0,07	3,983
2111	5	36 Draconis	—	—	2 56,12	3 55,94	18 12 56,01	—	55,40	—	+0,61	0,291
2112	5.6	106 Hercules	—	—	5 11,70	—	18 13 11,72	—	10,92	—	+0,80	2,552
2113	4.5	1 Lyre	—	—	2 58,45	2 58,52	18 13 58,53	58,63	58,53	-0,10	0,00	2,099
2114	6	Sagittarii	—	—	—	3 13,34	18 14 13,80	—	13,76	—	+0,04	3,864
2115	4.3	Telescopii	α	1 30,89	4 30,95	—	18 14 30,90	—	30,85	—	+0,05	4,451

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N.P.D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession	
	1831		1832		1833					Green.	A. S. C.		
	No.	"	No.	"	No.	"				"	"		"
2071	18	29 14,85	35	29 16,03	15	29 17,08	38 29 15,96	29 18,40	29 18,56	-2,44	2,60	+0,642	
2072					4	40 58,35	88 40 58,25		40 58,68		0,43	0,601	
2073	5	5 32,51					140 5 32,51		5 28,87	+	3,64	0,576	
2074							114		21 17,46			0,573	
2075	5	10 17,71					98 10 17,71		10 18,59	-	0,88	0,540	
2076					5	34 45,42	119 34 45,42		34 41,16	+	4,26	0,511	
2077					1	23 49,55	68 23 49,55		23 48,79	+	0,76	0,500	
2078							114		23 51,32			0,460	
2079			3	25 2,81	1	25 5,32	120 25 3,43	24 53,25	24 58,26	+10,18	+	5,17	0,446
2080	3	9 37,34	2	9 38,24			69 9 37,70		9 34,72	+	2,98	0,428	
2081					4	4 18,13	67 4 18,13		4 17,15	+	0,98	0,404	
2082	5	27 12,93					87 27 12,93	27 14,61	27 15,21	-1,68	-	2,28	1,444*
2083			1	27 8,75	3	27 8,52	111 27 8,58		27 8,47	+	0,11	0,263	
2084			1	1 13,65	4	1 12,93	13 1 13,07		1 14,34	+	1,27	0,259	
2085	5	27 56,39					118 27 56,39		27 58,80	+	2,41	0,235	
2086	1	58 21,79	4				135 58 21,79		58 21,61	+	0,18	0,121	
2087					5	47 23,19	67 47 23,19		47 23,73	+	0,54	0,099	
2088					3	44 49,22	120 44 49,22		44 42,95	+	6,27	0,076	
2089					4	16 50,60	81 16 50,60		16 44,44	+	6,16	0,072	
2090	5	27 14,05					80 27 14,05	27 13,22	27 6,71	+0,83	+	7,84	+0,062
2091	5	15 18,20			3	15 19,85	61 15 18,82	15 18,50	15 15,75	+0,32	+	3,07	-0,080
2092			5	1 41,33			86 1 41,33		1 38,49	+	2,84	0,097	
2093					6	43 33,09	113 43 33,09		43 32,62	+	0,47	0,118	
2094					5	12 20,93	69 12 20,93		12 19,16	+	1,77	0,129	
2095			2	58 33,51			69 58 33,51		58 26,96	+	6,55	0,135	
2096	5	5 40,12					111 5 40,12	5 40,11	5 36,53	+0,01	+	3,59	0,314
2097					3	41 58,83	111 44 58,83		44 56,22	+	2,61	0,354	
2098			6	46 11,29			110 46 11,29	16 10,60	46 6,66	+0,69	+	4,63	0,443
2099			4	25 48,17	3	25 50,75	110 25 49,27		26 45,40	+	3,87	0,445	
2100	4	37 49,17	2	37 49,20			58 37 49,18		36 46,08	+	3,10	0,481	
2101	5	48 8,20					126 48 8,20		48 0,21	+	7,99	0,536	
2102					5	35 29,59	110 35 29,59		35 27,56	+	2,03	0,566	
2103			3	5 42,08	1	5 40,65	117 5 41,72		3 39,72	+	2,00	0,648	
2104					7	30 49,97	108 30 49,97		30 49,81	+	0,16	0,656	
2105	6	53 28,46					119 53 28,46	53 23,35	53 25,45	+5,11	+	3,01	0,884
2106			1	53 36,38	4	53 34,53	105 53 34,90					0,905	
2107	5	37 7,44					65 37 7,44		37 0,44	+	7,00	1,064	
2108			3	41 27,66			86 41 27,66		41 23,39	+	4,27	1,083	
2109	4	56 4,14			1	56 6,46	92 56 4,60	56 8,36	56 2,93	-3,76	+	1,67	0,414*
2110	5	27 18,19					124 27 18,19		27 5,18	+	13,01	1,127	
2111			5	39 34,09			25 39 34,09		39 29,43	+	6,66	1,129	
2112					6	6 11,65	68 6 11,65		6 10,22	+	1,43	1,145	
2113			6	0 23,12			54 0 23,12	0 23,39	0 21,98	-0,27	+	1,14	1,215
2114					5	50 0,67	120 50 0,67		49 55,29	+	5,38	1,232	
2115	4	3 6,65					136 3 6,65		2 54,74	+	11,91	1,255	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	NAMES.	Mean A. R. January 1, 1832, from Observations in				Mean A. R. January 1, 1832.			Green. Catal.	A. S. Catal.	Difference from		Annual Preces- sion.		
			No. 1831		No. 1832		No. 1833					Green.	A. S.			
			s.		s.		h.	m.	s.							
2116	6	107 Herculis			4	27,92	18	14	27,96	s.	27,58	s.	+0,38	+2,335		
2117	5.6	Herculis			3	8,61	18	15	8,64					2,497		
2118	6	21 Sagittarii		1	20,51	3	20,96	18	15	20,83		20,42	+0,41	3,570		
2119	5	Pavonis					18	15			40,89			5,615		
2120	5	Telescopii	1	52,22				18	15	52,22		51,68	+0,54	4,609		
2121	5.6	109 Herculis	F			2	32,46	18	16	32,18		32,48	0,00	2,538		
2122	4	22 Sagittarii	λ	6	36,22		3	36,34	18	17	36,25	36,17	+0,08	+0,12	3,704	
2123	6	Sagittarii				1	8,35	18	18	8,33		8,04	+0,29	3,495		
2124	5.6	59 Serpentis	d		5	37,01		18	18	37,01		36,72	+0,29	3,066		
2125	5	Clypei Sob		6	37,28			18	19	37,28				3,416		
2126	6	Sagittarii	v'		2	3,78		18	20	3,76		3,74	+0,02	3,935		
2127	6.7	Clypei Sob						18	20					3,417		
2128	6	Sagittarii				3	19,40	18	20	19,38		19,10	+0,28	3,522		
2129	6	60 Serpentis	c		1	56,68	3	56,82	18	20	56,79		55,83	+0,96	3,117	
2130	7	Sagittarii				3	27,85	18	21	27,83		27,91	-0,08	3,526		
2131	5	39 Draconis	b	4	27,59			18	21	27,59		27,39	+0,20	0,880		
2132	6	Sagittarii				3	35,79	18	21	35,77		35,49	+0,28	3,513		
2133	7	Sagittarii				3	59,81	18	21	59,79		59,49	+0,30	3,532		
2134	6	Herculis	H			4	37,85	18	22	37,82		37,31	+0,57	2,483		
2135		Sagittarii	v ²			3	56,44	19	22	56,40		56,10	+0,10	3,336		
2136	7	Sagittarii						18	22			56,79		3,666		
2137	6.7	Clypei Sob	s ⁴					18	23			7,63		3,124		
2138	4	Pavonis	ζ					18	23			23,81		7,054		
2139	6	61 Serpentis	e		4	16,76		18	23	16,76		16,86	-0,10	3,094		
2140	7	Sagittarii						18	23			19,44		3,512		
2141	6.7	24 Sagittarii			1	37,90		18	23	37,89		37,68	+0,21	3,664		
2142	6.7	Clypei Sob	s ⁴					18	24			2,65		3,423		
2143	4.5	44 Draconis	χ		3	4,98	1	3,71	18	24	4,73	4,98	-0,25	+0,60	-1,072*	
2144	7	Sagittarii				3	29,09	18	25	29,07		28,91	+0,13	+3,536		
2145	6	Clypei Sob	q			3	42,28	18	25	42,27		42,06	+0,21	3,329		
2146	6	Herculis				3	46,74	18	25	46,77		46,60	+0,17	2,491		
2147	5.6	1 Aquilæ	m			5	4,01	18	26	4,00		3,50	+0,50	3,263		
2148	3	23 Ursæ Min	δ	10	29,14	73	29,42		18	26	29,54	30,21	-0,67	+7,33	-10,168	
2149	7	Sagittarii				4	13,44	18	27	13,42		12,83	+0,59	+3,534		
2150	6.7	Sagittarii				3	50,91	18	27	50,89		50,65	+0,24	3,591		
2151	7	Clypei Sob						18	28			6,13		+3,483		
2152	6	Sagittarii						18	28			17,57		3,649		
2153	6	Herculis						18	28			30,54		2,492		
2154	6.7	Sagittarii						18	28			52,48		3,582		
2155	5	Pavonis		1	56,33	1	56,59		18	28	56,43		57,39	-0,96	5,914	
2156	1	3 Lyrae	a	15	15,21	52	15,18	45	14,96	18	31	15,13	15,14	-0,01	+0,34	2,010
2157	6	26 Sagittarii				4	36,30	18	31	36,77		36,63	+0,14	3,657		
2158	5	Pavonis	θ	1	5,35			2	5,68	18	32	5,46		9,31	-3,85	5,938
2159	7	Clypei Sob				4	8,53	18	32	8,51		8,10	+0,41	3,416		
2160	5	2 Aquilæ	o	6	4,60				18	33	4,60		4,41	+0,19	3,282	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N.P.D. January 1, 1832.			Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833						Green.	A. S. C.	
2116	—	—	5 12 15,99	61 12 15,99	12 13,89	—	—	+	2,10	-1,257	
2117	—	—	2 47 38,56	66 47 38,56	47 45,24	—	—	—	6,68	1,821	
2118	—	—	—	110 37 —	37 20,05	—	—	—	—	1,330	
2119	—	2 22 16,66	1 22 15,06	152 22 16,13	22 9,43	—	—	+	6,70	1,354	
2120	—	—	4 9 2,41	139 9 2,41	8 27,88	—	—	+	34,53	1,373	
2121	—	—	4 18 0,94	68 18 0,94	17 55,88	—	—	+	5,06	1,438	
2122	5 30 24,35	—	—	115 30 24,35	30 20,26	30 21,16	+4,09	+	3,19	1,527	
2123	—	—	—	107 46 —	46 51,56	—	—	—	—	1,574	
2124	—	2 53 52,64	2 53 54,11	89 53 53,37	53 42,13	—	—	+	11,24	1,617	
2125	6 39 56,36	—	3 39 57,54	104 39 56,75	39 54,19	—	—	+	2,56	1,707	
2126	—	—	5 5 25,45	123 5 25,45	5 21,46	—	—	+	3,99	1,741	
2127	—	—	—	104 —	41 8,52	—	—	—	—	1,759	
2128	—	—	4 49 35,10	108 49 35,10	49 34,14	—	—	+	0,96	1,764	
2129	—	4 —	4 5 14,99	92 5 14,99	5 7,92	—	—	+	7,07	1,819	
2130	—	—	2 0 24,19	109 0 24,19	0 19,14	—	—	+	5,05	1,864	
2131	6 17 40,86	—	—	31 17 40,86	17 39,66	—	—	+	1,20	1,871	
2132	—	—	—	108 —	30 29,78	—	—	—	—	1,875	
2133	—	—	1 14 6,46	109 14 6,46	14 2,18	—	—	+	4,28	1,910	
2134	—	1 14 21,76	4 14 23,81	66 14 23,40	14 23,64	—	—	—	0,24	1,968	
2135	—	—	—	123 —	7 51,25	—	—	—	—	1,991	
2136	—	—	5 13 25,31	114 13 25,31	13 22,56	—	—	+	2,75	1,993	
2137	—	—	2 58 49,44	104 58 49,44	58 43,53	—	—	+	5,91	2,009	
2138	—	—	2 33 19,91	161 33 19,91	33 15,20	—	—	+	4,71	2,022	
2139	—	—	—	91 —	6 51,33	—	—	—	—	2,024	
2140	—	—	—	108 —	28 55,76	—	—	—	—	2,026	
2141	—	1 8 56,01	2 8 55,73	114 8 55,82	8 50,38	—	—	+	5,44	2,052	
2142	—	—	2 58 17,66	104 58 17,66	58 11,78	—	—	+	5,88	2,089	
2143	5 20 30,57	4 20 31,34	—	17 20 30,91	20 30,10	20 28,93	+0,81	+	1,98	2,104	
2144	—	1 23 30,06	4 23 31,65	109 23 31,33	23 28,46	—	—	+	2,87	2,214	
2145	—	1 6 1,65	3 6 2,03	101 6 1,94	5 59,94	—	—	+	2,00	2,233	
2146	—	—	3 30 9,21	66 30 9,21	30 10,27	—	—	—	1,06	2,242	
2147	—	—	5 21 12,90	98 21 12,90	21 9,17	—	—	+	3,73	2,265	
2148	5 24 46,07	1 24 44,20	—	3 24 45,76	24 48,50	24 48,92	-2,74	—	3,16	2,357	
2149	—	—	2 20 21,44	109 20 21,44	20 21,95	—	—	—	0,51	2,365	
2150	—	—	—	111 —	31 35,13	—	—	—	—	2,419	
2151	—	—	—	107 —	21 55,14	—	—	—	—	2,441	
2152	—	—	—	113 —	38 17,60	—	—	—	—	2,457	
2153	—	—	—	66 —	31 28,47	—	—	—	—	2,480	
2154	—	—	3 10 56,53	111 10 56,53	10 50,67	—	—	+	5,86	2,508	
2155	5 0 56,67	—	—	155 0 56,67	1 19,74	—	—	—	23,07	2,508	
2156	40 22 5,92	65 22 5,72	51 22 5,78	51 22 5,78	22 5,85	22 4,22	-0,07	+	1,56	2,718	
2157	—	—	—	113 —	58 48,32	—	—	—	—	2,745	
2158	—	—	1 14 7,22	155 14 7,22	14 33,00	—	—	—	25,78	2,786	
2159	—	—	1 42 55,65	104 42 55,65	42 47,99	—	—	+	7,66	2,791	
2160	5 12 20,22	1 12 17,96	—	99 12 19,85	12 15,06	—	—	+	4,79	2,873	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.			Green Catal.	A. S. Catal.	Difference from		Annual Preces- sion.
			No. 1831			No. 1833					Green.	A. S.	
			s.	s.	s.	h.	m.	s.					
2161	5.6	3 Aquilæ	—	4 22,23	—	18 34 22,23	—	22,32	—	—0,09	+3,264		
2162	6	Sagittarii	—	—	5 30,00	18 34 29,97	—	29,66	—	+0,31	3,689		
2163	4.5	27 Sagittarii	3 9,61	4 9,58	—	18 35 9,58	9,55	9,89	+0,03	-0,31	3,745		
2164	6	28 Sagittarii	—	—	7 12,77	18 36 12,75	12,72	12,85	+0,03	-0,10	3,616		
2165	5.6	4 Aquilæ	—	—	4 21,30	18 36 21,30	—	21,23	—	+0,07	3,024		
2166	5	Pavouis	—	5 37,93	—	18 36 37,86	—	39,42	—	-1,56	5,588		
2167	5.6	6 Aquilæ	—	5 15,58	—	18 38 15,57	—	15,56	—	+0,01	3,182		
2168	5	110 Hercules	6 25,88	1 26,13	—	18 38 25,91	—	25,46	—	+0,45	2,578		
2169	5	4 Lyræ	2 46,58	—	—	18 38 46,58	—	45,89	—	+0,69	1,982		
2170	5	5 Lyræ	2 48,64	5 48,91	—	18 38 48,85	—	47,72	—	+1,13	1,985		
2171	5	6 Lyræ	—	—	4 58,98	18 38 59,00	—	58,72	—	+0,28	2,060		
2172	5	46 Draconis	—	—	5 22,28	18 39 22,39	22,57	22,20	-0,18	+0,19	1,162		
2173	5.6	111 Hercules	—	—	5 26,30	18 39 26,32	—	36,10	—	+0,22	2,640		
2174	6	29 Sagittarii	—	—	5 42,56	18 39 42,54	—	41,85	—	+0,69	3,560		
2175	6	30 Sagittarii	—	—	3 44,54	18 40 44,52	—	44,35	—	+0,17	3,609		
2176	6	31 Sagittarii	—	6 2,68	—	18 42 2,67	—	2,35	—	+0,32	3,602		
2177	3	10 Lyræ	9 52,84	27 52,86	24 52,61	18 43 52,78	52,79	52,21	-0,01	+0,57	2,211		
2178	6	33 Sagittarii	—	6 57,61	—	18 43 57,60	—	57,66	—	-0,06	3,586		
2179	5	32 Sagittarii	7 1,57	1 1,56	1 1,57	18 44 1,56	1,51	1,27	+0,05	+0,29	3,623		
2180	3	34 Sagittarii	5 50,74	—	—	18 44 50,74	50,71	50,73	+0,03	+0,01	3,722		
2181	5	35 Sagittarii	3 57,57	—	1 57,39	18 44 57,52	57,70	57,69	-0,18	-0,17	3,621		
2182	5.6	112 Hercules	—	—	3 5,85	18 45 5,83	—	5,36	—	+0,47	2,559		
2183	6	Sagittarii	—	—	2 50,49	18 45 50,47	—	50,04	—	+0,43	3,634		
2184	6	62 Serpentis	—	—	3 16,33	18 47 16,34	—	15,96	—	+0,38	2,921		
2185	6	36 Sagittarii	1 21,64	2 21,56	3 21,29	18 47 21,42	21,35	21,01	+0,07	+0,41	3,567		
2186	5	113 Hercules	5 39,81	—	—	18 47 39,81	—	39,44	—	+0,37	2,528		
2187	5	37 Sagittarii	—	1 41,00	—	18 47 41,89	42,20	42,10	-0,31	-0,21	3,578		
2188	4.5	63 Serpentis	—	—	3 52,26	18 47 52,27	52,14	51,91	+0,13	+0,36	2,977		
2189	5	Serpentis	—	—	3 53,58	18 47 53,59	53,50	53,33	+0,09	+0,26	2,977		
2190	5.6	9 Aquilæ	—	6 3,98	—	18 48 3,97	—	4,23	—	-0,26	3,207		
2191	5	12 Lyræ	—	5 37,99	—	18 48 38,02	—	37,28	—	+0,74	2,095		
2192	5	47 Draconis	—	—	1 43,20	18 48 43,32	42,91	42,45	+0,41	+0,87	0,878		
2193	6	64 Serpentis	—	—	1 50,05	18 48 50,06	—	49,62	—	+0,44	3,015		
2194	6	10 Aquilæ	—	—	5 4,28	18 51 4,30	—	4,05	—	+0,25	2,751		
2195	6.7	Sagittarii	1 30,20	5 30,09	—	18 51 30,10	—	29,83	—	+0,27	3,619		
2196	3.4	38 Sagittarii	6 55,20	—	—	18 51 55,20	55,26	55,02	-0,06	+0,18	3,823		
2197	7	Sagittarii	—	—	4 57,37	18 51 57,35	—	57,03	—	+0,32	3,430		
2198	3.4	13 Aquilæ	—	6 0,11	—	18 52 0,13	59,99	59,22	+0,14	+0,91	2,723		
2199	6.7	Sagittarii	—	—	3 10,76	18 52 10,73	—	10,53	—	+0,20	3,677		
2200	3	14 Lyræ	5 39,55	8 39,65	17 39,60	18 52 39,63	39,60	39,02	+0,03	+0,61	2,240		
2201	5.6	12 Aquilæ	—	—	3 42,62	18 52 42,61	—	42,54	—	+0,07	3,204		
2202	6	Sagittarii	—	—	2 37,57	18 53 37,53	—	37,49	—	+0,04	3,858		
2203	6	48 Draconis	1 53,88	2 54,13	—	18 53 54,08	—	53,75	—	+0,33	1,021		
2204	7	14 Aquilæ	—	—	3 3,77	18 54 3,76	—	3,72	—	+0,04	3,157		
2205	4.5	39 Sagittarii	4 36,44	1 36,52	2 36,83	18 54 36,56	36,18	36,68	+0,38	-0,12	3,592		

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N.P.D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession	
	1831		1832		1833					Green.	A. S. C.		
	No.	"	No.	"	No.	"				"	"		
2161	—	—	—	—	5 25	58,41	98 25	58,41	26	0,73	—	2,32	2,985
2162	—	—	—	—	4 10	20,57	1 15	10 20,57	10	11,06	—	9,51	2,995
2163	5 9	17,41	—	—	—	—	117 9	17,41	9 16,13	9 14,84	+1,28	+ 2,57	3,052
2164	—	—	1 33	31,30	4 33	33,51	112 33	33,07	33 37,34	33 31,77	-4,27	+ 1,30	3,143
2165	—	—	—	—	5 6	11,71	88 6	11,71	—	6 9,53	—	+ 2,18	3,157
2166	—	—	5 21	58,75	—	—	152 21	58,75	21	55,87	—	+ 2,88	3,176
2167	—	—	—	—	5 55	18,41	94 55	18,41	55	9,07	—	+ 9,34	3,321
2168	4 36	32,80	1 36	35,93	—	—	69 36	33,42	36	27,91	—	+ 5,51	3,337
2169	5 30	5,43	3 30	6,22	—	—	50 30	5,73	29	59,00	—	+ 6,73	3,368
2170	6 33	34,28	—	—	—	—	50 33	34,28	33	28,68	—	+ 5,60	3,371
2171	—	—	5 33	53,87	—	—	52 33	53,87	33	53,17	—	+ 0,70	3,386
2172	—	—	—	—	6 37	44,77	34 37	44,77	37 41,34	37 37,16	+2,93	+ 7,61	3,423
2173	—	—	—	—	4 59	58,16	71 59	58,16	59	54,10	—	+ 4,06	3,438
2174	—	—	—	—	5 30	29,33	110 30	29,33	30	23,80	—	+ 5,53	3,444
2175	—	—	—	—	5 20	46,88	112 20	46,88	20	42,94	—	+ 3,94	3,534
2176	—	—	3 6	37,85	1 6	36,61	112 6	37,54	6	31,61	—	+ 5,93	3,646
2177	7 49	38,81	36 49	40,46	22 49	39,36	56 49	39,90	49 39,04	49 35,84	+0,86	+ 4,06	3,807
2178	—	—	—	—	6 33	21,18	111 33	24,18	33	21,38	—	+ 2,80	3,811
2179	5 56	33,67	—	—	—	—	112 56	33,67	56 34,58	56 32,96	-0,91	+ 0,71	3,816
2180	5 29	46,40	—	—	—	—	116 29	46,40	29 47,68	29 47,09	-1,28	- 6,69	3,886
2181	2 52	19,86	3 52	20,45	—	—	112 52	20,21	52 22,08	52 17,06	-1,87	+ 3,15	3,896
2182	—	—	—	—	5 46	12,66	68 46	12,66	46	13,43	—	- 6,82	3,911
2183	—	—	2 22	40,61	3 22	42,66	113 22	41,84	22	42,95	—	- 1,11	3,971
2184	—	—	—	—	5 35	15,04	83 35	15,04	35	12,99	—	+ 2,05	4,096
2185	—	—	—	—	4 52	5,69	110 52	5,69	52 5,61	51 59,30	+0,08	+ 6,39	4,101
2186	5 33	40,20	—	—	—	—	67 33	40,20	33	40,09	—	+ 0,11	4,131
2187	—	—	—	—	4 19	9,40	111 19	9,40	18 59,92	19 6,25	+9,48	+ 3,15	4,131
2188	—	—	3 0	31,53	2 0	32,64	86 0	31,97	0 31,68	0 22,63	+0,29	+ 9,34	4,147
2189	—	—	1 0	34,06	—	—	86 0	34,06	0 36,58	0 24,41	-2,52	+ 9,65	4,149
2190	—	—	—	—	1 3	26,01	96 3	26,01	3	19,83	—	+ 6,18	4,164
2191	5 18	34,70	—	—	—	—	53 18	34,70	18	34,56	—	+ 0,14	4,214
2192	—	—	6 48	54,21	—	—	30 48	54,21	48 54,16	48 54,83	+0,03	- 0,62	4,225
2193	—	—	—	—	5 40	41,49	87 40	41,49	40	37,51	—	+ 3,98	4,229
2194	—	—	—	—	5 18	47,27	76 18	47,27	18	35,36	—	+ 11,91	4,421
2195	—	—	—	—	4 55	22,50	112 55	22,50	55	26,93	—	- 4,43	4,455
2196	5 6	44,50	3 6	44,98	—	—	120 6	44,66	6 38,79	6 40,58	+5,87	+ 4,08	4,491
2197	—	—	—	—	5 30	43,61	105 30	43,61	30	41,65	—	+ 1,96	4,495
2198	3 9	16,71	2 9	19,60	—	—	75 9	17,87	9 12,30	9 9,63	+5,57	+ 8,24	4,500
2199	—	—	—	—	5 4	16,96	115 4	16,96	4	5,42	—	+ 11,54	4,513
2200	5 32	7,89	25 32	7,61	17 32	8,36	57 32	8,21	32 9,45	32 6,27	-1,24	+ 1,94	4,558
2201	—	—	—	—	3 58	6,28	95 58	6,28	57	59,52	—	+ 6,76	4,560
2202	—	—	—	—	—	—	121	17	17	0,80	—	—	4,636
2203	—	—	6 24	19,69	—	—	32 24	19,69	24	21,67	—	- 1,98	4,667
2204	—	—	—	—	5 56	5,34	93 56	5,34	56	0,92	—	+ 4,42	4,675
2205	—	—	5 58	45,41	—	—	111 58	45,41	58 45,24	58 39,49	+0,17	+ 5,92	4,721

✶

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	NAMES.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.			Green. Catal.	A. S. Catal.	Difference from		Annual Preces- sion.
			No. 1831	No. 1832	No. 1833	h. m. s.	s.	s.			Green.	A. S.	
2206	5	Cor. Aust. γ	1 4,40	2 3,42	—	18 55 3,75	—	—	2,92	—	+0,83	+4,057	
2207	6	15 Aquilæ h	—	—	3 5,56	18 56 5,55	—	—	5,15	—	+0,40	3,165	
2208	4	40 Sagittarii τ	1 26,82	—	3 27,00	18 56 26,92	26,92	26,53	0,00	—	+0,39	3,755	
2209	5	52 Draconis ν	—	—	1 21,82	18 56 21,55	—	24,42	—	—	+0,13	-0,710	
2210	5	Cor. Aust. δ	2 38,68	—	—	18 56 38,68	—	38,63	—	—	+0,05	+4,185	
2211	6.7	Sagittarii	—	5 51,90	—	18 56 51,89	—	51,78	—	—	+0,11	3,611	
2212	7	Sagittarii	—	—	5 56,01	18 56 55,98	—	55,75	—	—	+0,23	3,783	
2213	3	16 Aquilæ λ	1 19,96	5 20,00	—	18 57 19,99	19,99	19,49	0,00	—	+0,50	3,184	
2214	3	17 Aquilæ ζ	14 41,42	28 41,48	21 41,38	18 57 41,44	41,36	40,87	+0,08	—	+0,57	2,754	
2215	7	Sagittarii	—	—	3 58,52	18 57 58,50	—	58,34	—	—	+0,16	3,669	
2216	5	Cor. Aust. α	1 2,17	2 2,48	—	18 58 2,36	—	1,66	—	—	+0,70	4,085	
2217	6.7	Sagittarii	1 24,44	—	2 24,32	18 58 24,36	—	23,79	—	—	+0,57	3,527	
2218	5	Cor. Aust. β	1 27,87	—	—	18 58 27,87	—	27,67	—	—	+0,20	4,138	
2219	5.6	18 Aquilæ	—	—	3 4,30	18 59 4,31	—	3,49	—	—	+0,82	2,821	
2220	4.5	41 Sagittarii π	1 46,22	3 46,23	2 46,01	18 59 46,15	46,20	46,07	-0,05	—	+0,08	3,571	
2221	7	Sagittarii	—	—	2 53,61	18 59 53,59	—	53,15	—	—	+0,44	3,540	
2222	6	19 Aquilæ	—	—	3 46,46	19 0 46,47	—	45,82	—	—	+0,65	2,937	
2223	7	Sagittarii	—	6 21,84	—	19 2 21,82	—	21,59	—	—	+0,23	3,410	
2224	6	Sagittarii	—	—	5 25,60	19 2 25,58	—	25,28	—	—	+0,30	3,586	
2225	6.7	Sagittarii	—	—	5 52,82	19 2 52,79	—	52,45	—	—	+0,34	3,701	
2226	5	20 Aquilæ B	5 34,08	1 33,99	—	19 3 34,06	—	33,53	—	—	+0,53	3,254	
2227	6	42 Sagittarii ν	—	4 14,07	2 14,12	19 5 14,08	14,17	14,32	-0,09	—	-0,24	3,681	
2228	6	21 Aquilæ C	—	5 14,61	—	19 5 14,61	—	14,29	—	—	+0,32	3,023	
2229	6.7	Sagittarii	—	1 19,13	4 19,01	19 5 19,00	—	18,27	—	—	+0,73	3,651	
2230	5	43 Sagittarii d	8 48,24	—	—	19 7 48,22	48,20	47,88	+0,02	—	+0,34	3,514	
2231	6	1 Sagittæ	—	—	5 2,96	19 8 2,98	—	2,54	—	—	+0,44	2,579	
2232	5	20 Lyre η	3 2,27	—	—	19 8 2,27	—	2,00	—	—	+0,27	2,038	
2233	6	22 Aquilæ	—	5 12,33	—	19 8 12,34	—	11,61	—	—	+0,73	2,967	
2234	5	53 Draconis π	2 29,45	—	—	19 8 29,45	—	28,92	—	—	+0,53	1,133	
2235	5	1 Vulpeculæ	6 59,82	—	—	19 8 59,82	—	59,68	—	—	+0,14	2,576	
2236	6	Sagittarii	—	—	—	19 9 —	—	25,31	—	—	—	3,430	
2237	5	25 Aquilæ ω	1 56,20	4 56,04	—	19 9 56,08	—	55,78	—	—	+0,30	2,813	
2238	6	23 Aquilæ	—	—	3 59,68	19 9 59,68	—	58,69	—	—	+0,99	3,051	
2239	6	24 Aquilæ	—	—	4 15,14	19 10 15,17	—	15,12	—	—	+0,05	3,067	
2240	4	Sagittarii β	—	—	3 32,69	19 10 32,63	—	32,68	—	—	-0,05	4,331	
2241	6	Sagittarii	—	—	3 33,85	19 10 33,83	—	33,69	—	—	+0,14	3,601	
2242	5	21 Lyre θ	—	—	3 31,95	19 10 32,00	—	31,40	—	—	+0,60	2,079	
2243	5	54 Draconis p	—	2 55,03	—	19 10 55,08	—	54,84	—	—	+0,24	1,077	
2244	4	Sagittarii β	—	—	—	19 11 —	—	4,34	—	—	—	4,346	
2245	6	26 Aquilæ f	—	—	2 34,74	19 11 34,73	—	34,57	—	—	+0,16	3,196	
2246	7	Sagittarii	—	—	—	19 11 —	—	46,09	—	—	—	3,519	
2247	6	28 Aquilæ Δ	—	—	—	19 11 —	—	49,20	—	—	—	2,796	
2248	5	44 Sagittarii ρ	1 55,38	7 55,60	—	19 11 55,56	—	55,69	—	—	-0,13	3,485	
2249	6	27 Aquilæ d	—	—	—	19 11 —	—	55,28	—	—	—	3,095	
2250	5.6	45 Sagittarii ρ	—	—	—	19 12 —	—	3,41	—	—	—	3,496	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N.P.D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	1831		1832		1833					Green.	A. S. C.	
	No.	' "	No.	' "	No.	' "				' "	' "	
2206	5	17 43,67	—	—	—	—	127 17 43,67	17 30,07		+ 13,60	—4,756	
2207	—	—	—	—	6 16 26,50	94 16 26,50	16 21,72	16 21,72		+ 4,78	4,847	
2208	3	54 24,42	2	54 24,58	—	—	117 54 24,48	54 23,84	54 24,98	-0,50	+ 0,64	4,876
2209	—	—	5	55 43,06	—	—	18 55 43,06	55 41,38	—	+ 1,68	4,885	
2210	—	—	—	—	5	44 53,05	130 44 53,05	44 46,12	—	+ 6,93	4,892	
2211	—	—	—	—	1	44 47,45	112 44 47,45	44 46,71	—	+ 0,74	4,912	
2212	—	—	—	—	—	—	118	53 13,06	—	—	4,917	
2213	5	7 42,77	—	—	—	—	95 7 42,77	7 32,39	7 38,21	+4,56	+ 10,38	4,952
2214	13	22 46,99	27	22 48,03	21	22 47,79	76 22 47,68	22 38,58	22 47,92	-0,24	+ 9,10	4,983
2215	—	—	—	—	—	—	114	54 39,74	—	—	5,006	
2216	3	9 21,28	2	9 22,66	—	—	123 9 21,83	9 17,99	—	+ 3,84	5,009	
2217	—	—	—	—	—	—	109 33	—	—	—	5,042	
2218	—	—	6	35 55,52	—	—	129 35 55,52	35 48,36	—	+ 7,16	5,046	
2219	—	—	1	10 52,56	4	10 51,97	79 10 52,09	10 48,71	—	+ 3,38	5,100	
2220	5	16 55,87	—	—	—	—	111 16 55,87	16 53,97	16 58,02	-2,15	+ 1,90	5,158
2221	—	—	1	3 44,36	4	3 46,65	110 3 46,19	3 34,56	—	+ 11,63	5,168	
2222	—	—	—	—	6	11 3,11	84 11 3,11	10 58,26	—	+ 4,85	5,244	
2223	—	—	3	51 22,05	2	51 20,53	104 51 21,44	51 16,89	—	+ 4,55	5,377	
2224	—	—	—	—	5	55 42,19	111 55 42,19	55 41,45	—	+ 0,74	5,382	
2225	—	—	3	10 47,91	3	10 47,58	116 10 47,74	10 46,90	—	+ 0,84	5,419	
2226	5	12 44,93	1	12 44,72	—	—	98 12 44,90	12 42,97	—	+ 1,93	5,478	
2227	—	—	7	32 17,95	—	—	115 32 17,95	32 13,30	32 16,16	+1,79	+ 4,65	5,618
2228	—	—	5	59 5,09	—	—	87 59 5,09	58 59,51	—	+ 5,58	5,620	
2229	—	—	—	—	4	27 29,73	114 27 29,73	27 20,32	—	+ 9,41	5,624	
2230	5	14 37,89	—	—	—	—	109 14 37,89	14 36,10	14 38,43	-0,54	+ 1,79	5,833
2231	—	—	1	3 18,88	4	3 20,71	69 3 20,34	3 16,29	—	+ 4,05	5,856	
2232	5	8 18,44	1	8 19,50	—	—	51 8 18,61	8 16,91	—	+ 1,70	5,857	
2233	—	—	—	—	5	27 20,92	85 27 20,92	27 12,21	—	+ 8,71	5,868	
2234	3	25 26,68	2	25 24,00	—	—	33 25 25,62	25 28,66	—	- 3,04	5,897	
2235	—	—	6	53 59,38	—	—	68 53 59,38	54 0,50	—	- 1,12	5,936	
2236	—	—	—	—	5	49 18,06	105 49 18,06	49 9,70	—	+ 8,36	5,429*	
2237	5	42 2,26	—	—	—	—	78 42 2,26	41 57,53	—	+ 4,73	6,013	
2238	—	—	—	—	5	12 49,18	80 12 49,18	12 47,18	—	+ 2,00	6,017	
2239	—	—	—	—	5	57 19,66	89 57 19,66	57 34,98	—	- 15,32	6,039	
2240	5	45 56,29	—	—	—	—	134 45 56,29	45 48,76	—	+ 7,53	6,060	
2241	—	—	—	—	3	42 24,52	112 42 24,52	42 23,89	—	+ 0,63	6,064	
2242	—	—	7	9 41,62	—	—	52 9 41,62	9 37,48	—	+ 4,14	6,065	
2243	—	—	3	34 56,05	—	—	32 34 56,05	34 55,01	—	+ 1,04	6,100	
2244	2	6 24,28	—	—	—	—	135 6 24,28	6 17,36	—	+ 6,92	6,104	
2245	—	—	—	—	3	43 24,33	95 43 24,33	43 14,86	—	+ 9,47	6,149	
2246	—	—	—	—	5	32 29,72	109 32 29,72	32 30,12	—	- 0,40	6,164	
2247	—	—	—	—	5	55 43,72	77 55 43,72	55 43,02	—	+ 0,70	6,171	
2248	—	—	—	—	—	—	108	9 18,19	—	—	6,178	
2249	—	—	—	—	—	—	91	11 47,96	—	—	6,178	
2250	—	—	—	—	—	—	108	36 36,64	—	—	6,189	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.		Green Catal.	A. S. Catal.	Difference from		Annual Preces- sion.	
			No.	1831	No.	1832	No.			1833	Green		A. S.
				s.		s.				h. m. s.			s.
2251	5.6	46 Sagittarii			2	6,42	19 12 6,40		5,97		+0,43	+3,439	
2252	4.5	Sagittarii		3 14,39			19 12 14,36		13,70		+0,66	4,170	
2253	3	57 Draconis	10 29,65	20 29,71			19 12 29,75	29,71	29,15	+0,04	+0,60	0,023	
2254	4	1 Cygni	5 12,97	1 13,04			19 13 12,99	13,10	12,56	-0,11	+0,43	1,381	
2255	6	Sagittarii		2 1,87	2	1,59	19 14 1,70		1,21		+0,49	3,747	
2256	6	47 Sagittarii			3	2,78	19 15 2,75		2,78		-0,03	3,654	
2257	6.7	48 Sagittarii			2	9,69	19 15 9,66		9,12		+0,24	3,651	
2258	6	49 Sagittarii			3	19,14	19 15 19,11		19,42		-0,31	3,639	
2259	6	3 Vulpeculæ			3	58,25	19 15 58,28		57,56		+0,72	2,453	
2260		50 Sagittarii		5 17,80			19 16 17,78		17,81		-0,03	3,581	
2261	6	Sagittarii			3	18,94	19 16 18,90		18,69		+0,21	3,799	
2262	6	Sagittarii		3 37,54	3	37,25	19 16 37,36		36,91		+0,45	3,415	
2263	6	2 Sagittæ			3	49,60	19 16 49,62		48,86		+0,76	2,691	
2264	7	Sagittarii			1	52,48	19 16 52,46		52,13		+0,33	3,403	
2265	5	31 Aquilæ	5 57,65	1 57,72			19 16 57,66		57,19		+0,47	2,871*	
2266	3.4	30 Aquilæ	6 1,67	24 1,74	12	1,67	19 17 1,71	1,59	1,33	+0,12	+0,38	3,007	
2267	5.6	2 Cygni			1	30,13	19 17 30,17		29,72		+0,45	2,361	
2268	5.6	32 Aquilæ			3	55,65	19 17 55,65		55,38		+0,27	3,063	
2269	6	4 Vulpeculæ					19 18 —		6,13		—	2,623	
2270	6	Sagittarii			2	18,76	19 18 18,74		18,59		+0,15	3,494	
2271	6	3 Cygni					19 18 —		28,50		—	2,491	
2272	4.5	60 Draconis					19 18 —	44,56	43,82		—	-1,057	
2273	7	Sagittarii		2 23,27	2	23,20	19 19 23,20		23,34		-0,14	+3,717	
2274	4	58 Draconis	4 47,13				19 19 47,13	47,60	47,06	-0,47	+0,07	0,326	
2275	6	35 Aquilæ		4 31,49			19 20 31,49		30,84		+0,65	3,033	
2276	6	Sagittarii		5 55,51			19 20 55,50		55,30		+0,20	3,566	
2277	4	6 Vulpeculæ	5 43,03	1 42,95	7	42,83	19 21 42,93	43,05	42,30	-0,12	+0,63	2,502	
2278	6	36 Aquilæ		3 52,34			19 21 52,83		51,45		+1,38	3,137	
2279	5.6	8 Vulpeculæ				5 56,31	19 21 56,34		55,93		+0,41	2,500	
2280	7	Sagittarii			3	11,58	19 22 11,55		11,26		+0,29	3,743	
2281	3	6 Cygni	5 56,89	9 56,94			19 23 56,93	56,84	56,47	+0,09	+0,46	2,416	
2282	7	Sagittarii			3	25,00	19 24 21,97		24,71		+0,26	3,629	
2283	6	Vulpeculæ			4	44,37	19 24 44,39		44,21		+0,18	2,600	
2284	5	10 Cygni		4 23,21			19 25 23,25	23,20	27,94	+0,05	+0,31	1,511	
2285	6.7	Sagittarii			3	32,49	19 25 32,46		32,48		-0,02	3,614	
2286	7	Sagittarii	1 39,42			5 39,09	19 25 39,13		39,01		+0,12	3,549	
2287	6	51 Sagittarii			1	49,22	19 25 49,19		49,21		-0,02	3,650	
2288	5	37 Aquilæ	5 51,90	1 51,68			19 25 51,86		51,45		+0,41	3,308	
2289	4.5	38 Aquilæ	1 53,12	3 53,03	1	52,96	19 25 53,03	52,94	52,54	+0,09	+0,49	2,915	
2290	4.5	52 Sagittarii	1 23,65	3 23,78			19 26 23,74	23,64	23,36	+0,10	+0,38	3,654	
2291	7	Sagittarii			3	38,31	19 26 38,29		37,87		+0,42	3,501	
2292	5.6	9 Vulpeculæ			3	12,14	19 27 12,16		11,56		+0,60	2,631	
2293	7	Sagittarii		4 18,31			19 27 18,29		17,89		+0,40	3,486	
2294	4	39 Aquilæ		1 51,04	3	51,03	19 27 51,03	51,06	50,99	-0,03	+0,04	3,229	
2295	5	41 Aquilæ	4 1,62				19 28 1,62		1,73		-0,11	3,104	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N.P.D. January 1, 1832.			Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No.	1831	No.	1832	No.	1833			Green	A. S. C.	
	"	"	"	"	"	"			"	"	
2251						106		15 42,50			-6,192
2252			4 55 22,09			130 55 22,09		55 14,16		+ 7,63	6,201
2253						22	38 2,68	38 2,92			6,234
2254	5 56 19,54					36 56 19,54	56 19,46	56 19,35	+0,08	+ 0,19	6,290
2255					3 10 55,14	118 10 55,14		10 55,69		- 0,55	6,351
2256		2 49 39,51		3 49 39,43	114 49 39,46			49 32,29		+ 7,17	6,436
2257				5 44 0,93	114 44 0,93			43 55,50		+ 5,43	6,445
2258				5 17 0,54	114 17 0,54			16 57,49		+ 3,05	6,459
2259		1 3 14,57		4 3 16,17	64 3 15,85			3 16,99		- 1,14	6,515
2260				5 6 5,43	112 6 5,43			6 0,88		+ 4,55	6,540
2261						120		3 57,65			6,540
2262						105		22 40,37			6,567
2263						73		23 55,10			6,585
2264						104		52 35,28			6,588
2265	5 24 29,13	4 24 28,80				78 24 28,99		24 25,91		+ 3,08	7,316*
2266	12 12 49,77	23 12 50,13	12 12 50,18			87 12 50,05	12 49,41	12 49,92	+0,61	+ 6,13	6,601
2267						60		42 0,93			6,642
2268		5 59 27,22				89 59 27,22		59 19,22		+ 8,00	6,676
2269				5 31 30,06	70 31 30,06			31 24,59		+ 5,47	6,692
2270				5 41 26,12	108 41 26,12			41 27,55		- 1,43	6,706
2271				5 23 8,76	65 23 8,76			23 6,10		+ 2,66	6,003*
2272	5 57 31,97					16 57 31,97	57 32,46	57 28,95	-0,49	+ 3,92	6,753
2273		1 19 17,73		4 19 18,45	117 19 18,31			19 13,94		+ 4,37	6,801
2274	5 36 31,73					24 36 31,73	36 30,69	36 28,91	+1,04	+ 2,82	6,836
2275				5 23 8,01	88 23 8,01			23 2,31		+ 5,70	6,889
2276		2 39 12,99		3 39 12,92	111 39 12,77			39 11,01		+ 1,76	6,921
2277	5 10 12,53	5 40 12,97	1 40 12,83			65 40 12,76	40 11,76	40 8,68	+1,00	+ 4,08	6,988
2278		4 7 53,70				93 7 53,70		7 49,66		+ 4,04	6,999
2279				5 34 21,63	65 34 21,63			34 18,31		+ 3,32	7,007
2280				4 19 58,18	118 19 58,18			19 19,52			7,024
2281	6 23 16,59	3 23 16,41				62 23 16,10	23 17,22	23 14,02	-0,82	+ 2,38	7,171
2282		2 12 51,30		3 12 52,48	114 12 52,01			12 50,40		+ 1,61	7,206
2283		2 25 19,99		3 25 21,97	69 25 21,18			25 19,04		+ 2,14	7,236
2284	5 37 29,06					38 37 29,06	37 30,11	37 29,00	-1,05	+ 0,06	7,298
2285				5 40 7,01	113 40 7,01			40 6,84		+ 0,17	7,299
2286				3 8 5,59	111 8 5,59			7 57,70		+ 7,89	7,308
2287				5 4 48,62	115 4 48,62			4 41,14		+ 7,48	7,321
2288	2 55 11,22	1 55 12,74				100 55 11,72		55 8,58		+ 3,14	7,325
2289	5 58 14,00	3 58 14,42		5 58 14,61	82 58 14,35		58 15,20	58 12,07	-0,85	+ 2,28	7,328
2290		5 14 49,48				115 14 49,48	14 45,79	14 43,41	+3,69	+ 6,07	7,374
2291		3 12 56,16		2 12 58,14	109 12 56,95			12 56,72		+ 0,23	7,387
2292		1 35 17,70		4 35 17,19	70 35 17,29			35 11,79		+ 5,50	7,436
2293		1 35 44,43		3 35 44,92	108 35 44,80			35 44,48		+ 0,32	7,442
2294	5 23 37,87					97 23 37,87	23 39,84	23 34,97	-1,97	+ 2,90	7,487
2295	5 39 7,79					91 39 7,79		39 3,53		+ 4,26	7,502

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	NAMES.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.		Green. Catal.	A. S. Catal.	Difference from		Annual Precession.				
			No. 1831	No. 1832	No. 1833	h.	m.			s.	Green.		A. S.			
			s.	s.	s.						s.		s.			
2296	5.6	9 Cygni			4 10,62	19 28	10,66		10,22		+0,44	+2,379				
2297	6	42 Aquilæ	P	4	52,63	19 28	52,64		52,18		+0,46	3,177				
2298	6	4 Sagittæ	e			4 41,11	19 29	41,13	41,17		-0,04	2,712				
2299	7	53 Sagittarii				4 43,40	19 29	43,38	43,15		+0,23	3,613				
2300	6.7	Sagittarii				4 0,80	19 30	0,78	0,85		-0,07	3,613				
2301	5	44 Aquilæ	σ	6	54,25		19 30	54,25	54,53		-0,28	2,960				
2302	5.6	54 Sagittarii	e	2	5,85	2	5,56	19 31	5,69	5,56	+0,13	3,437				
2303	4	13 Cygni	θ	3	55,93		2 55,85	19 31	55,94	56,15	-0,21	+0,30	1,611			
2304	6	45 Aquilæ				5	4,14	19 32	4,14	3,81	+0,33	3,090				
2305	4	5 Sagittæ	a	3	35,32	6	35,41		19 32	35,39	35,39	34,91	0,00	+0,48	2,678	
2306	5	61 Draconis	σ			5	39,98		19 32	40,07		39,04	+1,03	-0,110*		
2307	4	12 Cygni	φ				6 44,41	19 32	44,45	44,58	-0,13	-0,12	+2,365			
2308	5	55 Sagittarii	e	2	54,47	3	54,42		19 32	54,43		54,06	+0,37	3,432		
2309	5	6 Sagittæ	β	2	50,29	3	30,28		19 33	30,29		30,02	+0,27	2,691		
2310	6	Sagittarii					5 58,28	19 33	58,26			57,85	+0,41	3,416		
2311	6	47 Aquilæ	χ			4	39,96		19 34	39,96		39,44	+0,52	2,820		
2312	6.7	Sagittarii					5 45,35	19 34	45,31			45,41	-0,10	3,812		
2313	6	56 Sagittarii	f	1	33,52	3	33,51	4	33,36	19 36	33,43	33,41	-0,09	+0,02	3,516	
2314	6	10 Vulpeculæ	d				3 43,87	19 36	43,90	33,52		43,17	+0,73	2,490		
2315	6	Vulpeculæ					5	3,38	19 37	3,38		3,09	+0,29	2,454		
2316	5	15 Cygni		6	13,14	1	13,44		19 38	13,15		12,85	+0,30	2,154		
2317	3	50 Aquilæ	γ	24	16,40	53	16,42	13	16,43	19 38	16,43	16,40	+0,03	+0,14	2,849	
2318	6.7	Sagittarii					3 36,44	19 38	36,42			36,18	+0,24	3,373		
2319	7	Sagittarii					5 29,10	19 39	29,09			28,75	+0,34	3,342		
2320	6	Aquilæ					4 46,18	19 39	46,17			45,70	+0,47	3,310		
2321	3.4	18 Cygni	δ	6	43,22			19 39	43,22	43,55	-0,33	+0,03	1,868			
2322	4	7 Sagittæ	δ	4	53,90			19 39	53,90	53,94	-0,04	+0,28	2,672			
2323	5	17 Cygni	χ	4	3,24	1	3,05		19 40	3,20		2,63	+0,57	2,271		
2324	6	52 Aquilæ	π				3 47,19	19 40	47,20			46,73	+0,47	2,824		
2325	4	Pavonis	α				3	1,21	19 41	1,01		59,60	+1,41	7,109		
2326	5.6	51 Aquilæ	D				3 31,90	19 41	31,89			31,41	+0,48	3,307		
2327	5	8 Sagittæ	ζ			4	31,37		19 41	31,38		30,83	+0,55	2,659		
2328	5.6	57 Sagittarii		1	26,05	4	25,93		19 42	25,95		25,83	+0,12	3,494		
2329	12	53 Aquilæ	α	29	35,22	60	35,22	45	35,24	19 42	35,23	35,22	+0,01	+0,19	2,924*	
2330	5.6	54 Aquilæ	o				3 58,72	19 42	58,73			58,26	+0,47	2,856		
2331	4.5	Sagittarii	E	6	39,87			19 43	39,87			39,12	+0,75	4,162		
2332	5.6	12 Vulpeculæ	e				3 50,06	19 43	50,08			49,43	+0,65	2,578		
2333	4	55 Aquilæ	γ			3	54,91		19 43	54,91	54,73	54,24	+0,18	+0,67	3,056	
2334	6	56 Aquilæ	E				4	1,19	19 45	1,18		1,02	+0,16	3,258		
2335	6	58 Sagittarii	ω				4 32,44	19 45	32,41	32,36		32,18	+0,05	+0,23	3,671	
2336	5	59 Aquilæ	f	8	6,45			19 46	6,45			5,96	+0,49	2,899		
2337	6	58 Aquilæ				4	8,37	19 46	8,37			8,44	-0,07	3,071		
2338	5	13 Vulpeculæ		4	19,38	1	19,50		19 46	19,40		18,99	+0,41	2,545		
2339	5	59 Sagittarii	δ			3	37,84		19 46	37,83	37,74	37,63	+0,09	+0,20	3,693	
2340	3.4	60 Aquilæ	β	23	3,69	38	3,74	9	3,67	19 47	3,72	3,74	3,55	-0,02	+0,17	2,943

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N.P.D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession		
	No.	1831	No.	1832	No.	1833				Green.	A. S. C.			
2296	—	—	—	—	3 54	6,22	60 54	6,22	54	4,64	+	1,58	-7,515	
2297	—	—	4 1	0,68	1 0	58,95	95 1	0,33	0	50,89	+	9,44	7,570	
2298	—	—	1 54	29,61	3 54	30,56	73 54	30,32	54	27,59	+	2,73	7,637	
2299	—	—	1 48	2,83	3 48	4,86	113 48	4,35	47	59,60	+	4,75	7,638	
2300	—	—	—	—	4 48	17,83	113 48	17,83	48	12,50	+	5,33	7,661	
2301	2 58	45,55	6 58	49,28	—	—	84 58	48,04	58	39,71	+	8,33	7,735	
2302	—	—	—	—	6 40	14,86	106 40	14,86	40	9,90	+	4,96	7,749	
2303	5 9	56,79	1 9	59,25	—	—	40 9	57,20	9 53,06	9 50,72	+4,14	+	6,48	7,821
2304	—	—	—	—	5 0	13,64	91 0	13,64	0	8,89	+	4,75	7,828	
2305	5 21	59,52	—	—	1 22	0,86	72 21	59,74	22 0,64	21 54,00	-0,90	+	5,74	7,871
2306	—	—	6 37	30,67	—	—	20 37	30,67	37	31,69	—	—	—	5,764*
2307	4 13	47,70	1 13	45,52	3 13	47,98	60 13	47,63	13 44,19	13 39,34	+3,44	+	8,29	7,885
2308	—	—	2 30	37,67	3 30	36,51	106 30	36,97	30	31,17	+	5,80	7,895	
2309	5 54	24,19	1 54	24,23	—	—	72 54	24,20	54	22,92	+	1,28	7,945	
2310	—	—	2 51	8,70	2 51	11,15	105 51	9,92	51	0,98	+	8,94	7,980	
2311	—	—	—	—	5 33	44,87	78 33	44,87	33	39,77	+	5,10	8,037	
2312	—	—	—	—	5 17	52,80	121 17	52,80	17	49,12	+	3,68	8,042	
2313	—	—	5 9	31,71	—	—	110 9	31,71	9 27,07	9 25,54	+4,64	+	6,17	8,187
2314	—	—	—	—	5 37	33,76	64 37	33,76	37	23,55	+	10,21	8,203	
2315	—	—	—	—	5 15	38,51	63 15	38,51	—	—	—	—	—	8,229
2316	5 2	47,25	—	—	—	—	53 2	47,25	2 43,51	—	+	3,74	8,323	
2317	37 47	28,17	51 47	28,70	18 47	28,84	79 47	28,53	47 25,10	47 22,32	+3,43	+	6,21	8,326
2318	—	—	1 6	35,28	4 6	33,58	104 6	33,92	6 32,78	—	+	1,14	8,351	
2319	—	—	—	—	5 43	43,76	102 43	43,76	43	41,92	+	1,84	8,420	
2320	—	—	3 16	52,25	2 16	55,69	101 16	53,64	16 48,62	—	+	5,02	8,443	
2321	—	—	5 16	34,52	—	—	45 16	34,52	16 32,25	16 28,96	+2,27	+	5,56	8,443
2322	5 52	28,20	—	—	—	—	71 52	28,20	52 30,05	52 27,07	-1,85	+	1,18	8,455
2323	5 39	31,13	—	—	—	—	56 39	31,13	39 28,67	—	+	2,46	8,468	
2324	—	—	—	—	5 35	43,65	78 35	43,65	35 42,43	—	+	1,22	8,525	
2325	—	—	—	—	4 20	24,34	163 20	24,34	20 9,12	—	+	15,22	8,530	
2326	—	—	—	—	—	—	101	—	10 50,86	—	—	—	—	8,582
2327	4 16	24,01	—	—	—	—	71 16	24,01	16 19,53	—	+	4,48	8,583	
2328	—	—	2 27	53,23	1 27	53,44	109 27	53,30	27 44,17	—	+	9,13	8,653	
2329	70 34	8,21	60 34	8,24	50 34	8,18	81 34	8,21	34 10,65	34 5,48	-2,44	+	2,73	8,667
2330	—	—	—	—	5 0	0,82	80 0	0,82	59 53,97	—	+	6,85	8,698	
2331	—	—	2 18	5,82	3 18	4,03	132 18	4,75	18 2,90	—	+	1,85	8,718	
2332	—	—	4 48	38,63	—	—	67 48	38,63	48 37,88	—	+	0,75	8,765	
2333	5 25	10,73	—	—	—	—	89 25	10,73	25 10,10	25 5,24	+0,63	+	5,49	8,771
2334	—	—	—	—	4 0	11,86	99 0	11,86	0 6,61	—	+	5,25	8,857	
2335	—	—	—	—	5 44	14,14	116 44	14,14	44 15,46	44 11,55	-1,32	+	2,59	8,897
2336	3 58	3,05	5 58	2,95	—	—	81 58	2,99	57 55,72	—	+	7,27	8,943	
2337	—	—	1 9	36,44	5 9	39,40	90 9	38,96	9 31,17	—	+	7,73	8,946	
2338	2 21	9,81	2 21	11,90	1 21	9,44	66 21	10,58	21 8,43	—	+	2,15	8,961	
2339	5 36	25,08	—	—	—	—	117 36	25,08	36 26,50	36 21,41	-1,42	+	3,57	8,982
2340	26 0	22,86	19 0	23,28	12 0	22,11	84 0	22,84	0 24,90	0 21,49	-2,06	+	1,35	8,478*

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.	Green Catal.	A. S. Catal.	Difference from		Annual Preces- sion.		
			No. 1831		No. 1832				No. 1833			Green	A. S.
			s.	s.	s.				h.	m.			
2341	6	61 Aquilæ ϕ	—	—	3 16,96	19 48 16,97	—	16,59	—	+0,38	+2,837		
2342	6	10 Sagittæ	—	—	3 23,55	19 48 23,57	—	23,19	—	+0,38	2,723		
2343	6	61 Sagittarii g	—	—	3 24,88	19 48 24,86	—	25,26	—	-0,40	3,408		
2344	5.6	60 Sagittarii a	—	—	3 42,58	19 48 42,55	42,58	42,47	-0,03	+0,08	3,665		
2345	7	Sagittarii	—	—	3 36,55	19 49 36,53	—	36,13	—	+0,40	3,564		
2346	5	22 Cygni	3 51,52	6 51,73	—	19 49 51,68	—	51,22	—	+0,46	2,140		
2347	6	11 Sagittæ	—	—	2 7,87	19 50 7,89	—	7,53	—	+0,36	2,721		
2348	4.5	12 Sagittæ γ	6 17,18	6 17,21	—	19 51 17,20	17,25	16,74	-0,05	+0,46	2,660		
2349	6	Sagittarii	—	6 24,42	1 24,69	19 51 24,45	—	24,17	—	+0,28	3,574		
2350	5	14 Vulpeculæ f	7 58,15	—	—	19 51 58,15	—	57,53	—	+0,62	2,576		
2351	4	Pavonis δ	—	1 10,09	—	19 52 10,00	—	8,69	—	+1,31	5,794		
2352	4.5	62 Sagittarii c	—	3 19,16	4 19,03	19 52 19,08	19,11	18,82	-0,03	+0,26	3,700		
2353	6	13 Sagittæ χ	—	—	3 28,02	19 52 28,04	—	28,06	—	-0,02	2,706		
2354	6	63 Sagittarii	—	—	5 33,65	19 52 33,64	—	33,11	—	+0,53	3,364		
2355	5	Sagittarii L	2 40,00	4 39,70	—	19 53 39,78	—	38,85	—	+0,93	3,818		
2356	6.7	Sagittarii	—	4 46,55	—	19 53 46,54	—	46,51	—	+0,03	3,568		
2357	5	15 Vulpeculæ g	4 11,17	—	—	19 54 11,17	—	10,55	—	+0,62	2,462		
2358	5	Vulpeculæ	—	2 37,65	2 37,83	19 54 37,75	—	37,55	—	+0,20	2,538		
2359	6	16 Vulpeculæ h	—	—	3 53,86	19 54 53,89	—	54,00	—	-0,11	2,535		
2360	6	62 Aquilæ	—	—	3 43,83	19 55 43,83	—	43,80	—	+0,03	3,092		
2361	6	64 Sagittarii Y	—	—	2 48,59	19 55 48,58	—	48,30	—	+0,28	3,318		
2362	6	14 Sagittæ y	—	—	3 48,98	19 55 49,00	—	48,28	—	+0,72	2,742		
2363	5.6	63 Aquilæ τ	—	—	—	19 55 —	—	55,75	—	—	2,929		
2364	6	65 Sagittarii	—	—	2 5,52	19 56 5,51	—	4,96	—	+0,55	3,341		
2365	6	15 Sagittæ z	—	5 33,59	—	19 56 33,60	—	32,77	—	+0,83	2,686*		
2366	6	16 Sagittæ η	—	5 42,55	—	19 57 42,56	—	42,04	—	+0,52	2,656		
2367	7	Capricorni	—	4 0,28	2 0,24	19 59 0,26	—	59,76	—	+0,50	3,390		
2368	7	Capricorni	—	—	5 20,05	19 59 20,04	—	19,90	—	+0,14	3,284		
2369	6	64 Aquilæ	—	—	5 21,31	19 59 21,31	—	21,13	—	+0,18	3,092		
2370	5.6	17 Vulpeculæ i	—	—	5 40,10	19 59 40,07	—	40,56	—	-0,49	2,573		
2371	5	67 Draconis ρ	6 1,87	5 1,72	—	20 2 1,84	2,29	0,94	-0,45	+0,90	0,304		
2372	3.4	65 Aquilæ θ	6 38,14	6 38,17	6 38,15	20 2 38,15	38,12	37,98	+0,03	+0,17	3,095		
2373	6.7	1 Capricorni ξ'	—	4 39,08	—	20 2 39,08	—	38,85	—	+0,23	3,331		
2374	6	66 Draconis	4 51,70	—	—	20 2 51,70	—	50,97	—	+0,73	0,952		
2375	5	2 Capricorni ξ^a	—	—	6 4,10	20 3 4,09	—	3,59	—	+0,50	3,335		
2376	5	28 Cygni δ^a	3 11,57	—	—	20 3 11,57	—	10,94	—	+0,63	2,223		
2377	6	18 Vulpeculæ	—	—	4 33,02	20 3 33,05	—	32,70	—	+0,35	2,499		
2378	6	Sagittarii R	—	3 47,95	—	20 4 47,94	—	44,95	—	+2,99	3,747*		
2379	6	19 Vulpeculæ	—	—	3 47,24	20 4 47,27	—	46,08	—	+1,19	2,503		
2380	6	20 Vulpeculæ k	—	—	5 58,18	20 4 58,21	—	57,27	—	+0,94	2,511		
2381	5	67 Aquilæ ρ	—	1 30,24	4 30,26	20 6 30,27	—	30,06	—	+0,21	2,770		
2382	6.7	3 Capricorni	—	—	6 4,53	20 7 4,52	—	3,98	—	+0,54	3,327		
2383	5.6	21 Vulpeculæ l	—	6 20,59	—	20 7 20,61	—	20,33	—	+0,28	2,460		
2384	6	4 Capricorni	—	—	4 8,81	20 8 8,79	—	8,80	—	-0,01	3,533		
2385	5.6	22 Vulpeculæ m	—	—	4 14,93	20 8 14,96	—	14,52	—	+0,44	2,587		

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N.P.D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession		
	1831		1832		1833					Green.	A. S. C.			
	No.	"	No.	"	No.	"								
2341	—	—	5 0	57,80	—	—	79 0	57,80	—	—	—	—		
2342	—	—	—	—	5 48	14,67	73 48	14,67	48	9,71	+	4,96	9,122	
2343	—	—	—	—	4 55	49,13	105 55	49,13	55	36,57	+	12,56	9,123	
2344	—	—	—	—	5 38	32,78	116 38	32,78	38 31,28	38 32,06	-1,50	+	0,72	9,145
2345	—	—	—	—	5 39	32,07	112 39	32,07	—	39 32,20	—	—	0,13	9,215
2346	—	—	4 57	21,03	1 57	20,89	51 57	21,00	—	57 14,78	+	6,22	9,238	
2347	—	—	4 39	23,95	6 39	23,63	73 39	23,76	—	39 19,02	+	4,74	9,257	
2348	5 57	31,54	3 57	31,03	—	—	70 57	31,35	57 31,54	57 20,74	-0,19	+	10,61	9,347
2349	—	—	—	—	5 11	27,94	113 11	27,94	—	11 26,22	+	1,72	9,354	
2350	4 21	7,95	1 21	9,42	—	—	67 21	8,24	—	21 0,65	+	7,59	9,400	
2351	—	—	5 35	51,31	—	—	156 35	51,31	—	35 25,09	+	26,22	9,406	
2352	5 10	7,84	1 10	5,02	—	—	118 10	7,37	10 8,80	10 3,86	-1,43	+	3,51	9,424
2353	—	—	3 56	10,70	2 56	13,61	72 56	11,86	—	56 8,48	+	3,38	9,439	
2354	—	—	—	—	5 5	42,84	104 5	42,84	—	5 39,72	+	3,12	9,443	
2355	5 31	12,86	—	—	—	—	122 31	12,86	—	31 10,23	+	2,63	9,527	
2356	—	—	3 3	32,24	2 3	33,09	113 3	32,58	—	3 36,06	—	—	3,48	9,537
2357	4 42	19,52	—	—	1 42	17,95	62 42	19,21	—	42 16,97	+	2,21	9,571	
2358	2 39	36,53	5 39	39,62	—	—	65 39	38,74	—	39 41,43	—	—	2,69	9,605
2359	—	—	5 31	38,72	—	—	65 31	38,72	—	31 35,52	+	3,20	9,626	
2360	—	—	4 10	18,03	—	—	91 10	18,03	—	10 7,95	+	10,08	9,688	
2361	—	—	—	—	6 4	5,43	102 4	5,43	—	3 57,49	+	7,94	9,693	
2362	—	—	—	—	5 26	5,02	74 26	5,02	—	25 57,19	+	7,83	9,695	
2363	—	—	—	—	5 11	23,15	83 11	23,15	—	11 21,03	+	2,12	9,704	
2364	—	—	—	—	5 7	58,05	103 7	58,05	—	7 51,36	+	6,69	9,715	
2365	—	—	—	—	5 22	45,07	73 22	45,07	—	22 37,32	+	7,75	9,752	
2366	—	—	5 29	8,04	—	—	70 29	8,04	—	29 3,60	+	4,44	9,840	
2367	—	—	2 30	25,30	3 30	28,40	105 30	26,85	—	30 18,74	+	8,11	9,937	
2368	—	—	—	—	9 32	34,64	100 32	34,64	—	32 31,06	+	3,58	9,962	
2369	—	—	4 9	20,62	—	—	91 9	20,62	—	9 17,11	+	3,51	9,964	
2370	—	—	5 51	53,18	—	—	66 51	53,18	—	51 51,14	+	2,04	9,990	
2371	5 36	15,88	2 36	16,04	—	—	22 36	15,94	36 21,38	36 18,53	-5,44	—	2,59	10,173
2372	6 18	49,27	7 18	48,94	5 18	48,84	91 18	49,02	18 48,13	18 45,84	+0,89	+	3,18	10,212
2373	—	—	5 53	2,39	—	—	102 53	2,39	—	53 1,86	+	0,53	10,213	
2374	4 29	23,32	—	—	—	—	28 29	23,32	—	29 29,16	—	—	5,84	10,234
2375	—	—	—	—	5 6	8,62	103 6	8,62	—	6 5,25	+	3,37	10,244	
2376	4 38	58,67	—	—	—	—	53 38	58,67	—	38 58,19	+	0,48	10,256	
2377	—	—	2 35	15,86	2 35	17,22	63 35	16,53	—	35 14,78	+	1,75	10,282	
2378	—	—	5 31	35,99	—	—	117 31	35,99	—	31 26,39	+	9,60	11,129*	
2379	—	—	—	—	6 41	10,18	63 41	10,18	—	41 7,67	+	2,51	10,374	
2380	—	—	5 1	3,07	—	—	64 1	3,07	—	1 2,84	+	0,23	10,388	
2381	2 18	33,23	5 18	35,42	—	—	75 18	34,52	—	18 22,70	+	11,82	10,503	
2382	—	—	3 50	38,70	2 50	39,83	102 50	39,15	—	50 33,37	+	5,78	10,543	
2383	—	—	1 48	30,40	4 48	34,07	61 48	33,34	—	48 36,83	—	—	3,49	10,566
2384	—	—	4 19	18,85	—	—	112 19	18,85	—	19 13,46	+	5,39	10,623	
2385	—	—	—	—	5 59	57,77	66 59	57,77	—	59 55,43	+	2,34	10,632	

xcviii *Comparison of the Observed Places of the Principal Fixed Stars*

No.	Mag	NAMES.	Mean A. R. January 1, 1832, from Observations in				Mean A. R. January 1, 1832.			Green. Catal.	A. S. Catal.	Difference from		Annual Preces- sion.
			1831		1832		1833					Green	A. S.	
			No	s.	No	s.	No	h.	m.					
2386	4	5 Capricorni α^1	6 19,94	5 19,94	—	20 8 19,94	19,83	19,68	+0,11	+0,26	+3,330			
2387	4	31 Cygni σ^2	3 20,61	6 20,66	—	20 8 20,66	20,59	20,22	+0,07	+0,44	1,886			
2388	3	6 Capricorni α^2	16 43,72	7 43,69	—	20 8 43,71	43,68	43,61	+0,03	+0,07	3,331			
2389	4.5	23 Vulpeculæ η	5 48,55	2 48,83	—	20 8 48,64	48,60	48,25	+0,04	+0,39	2,484			
2390	6	18 Sagittæ	—	2 56,93	3 57,05	20 8 57,03	—	56,41	—	+0,62	2,632			
2391	4.5	33 Cygni	3 29,19	—	—	20 9 29,19	29,34	28,36	-0,15	+0,83	1,392			
2392	5	24 Vulpeculæ σ	—	—	3 35,77	20 9 35,80	—	35,78	—	+0,02	2,562			
2393	5.6	7 Capricorni σ	—	2 41,75	—	20 9 41,74	41,63	41,56	+0,11	+0,18	3,471			
2394	4.5	32 Cygni	—	—	5 16,35	20 10 16,42	16,78	16,81	-0,36	-0,39	1,852			
2395	7	Capricorni β^1	—	1 20,01	4 19,91	20 11 19,91	—	19,57	—	+0,34	3,376			
2396	5	8 Capricorni ν	—	5 20,52	—	20 11 20,51	—	20,40	—	+0,11	3,333			
2397	3.4	9 Capricorni β^2	—	2 34,01	3 34,02	20 11 34,00	34,01	33,79	-0,01	+0,21	3,375			
2398	2	Pavonis α	—	—	4 18,45	20 12 18,33	—	18,54	—	-0,21	4,811			
2399	4.5	1 Cephei ν	4 23,71	2 23,15	—	20 14 23,68	23,85	20,39	-0,17	+3,29	-1,882*			
2400	6	25 Vulpeculæ	—	6 50,27	—	20 14 50,28	—	50,02	—	+0,26	+2,575			
2401	3	37 Cygni γ	6 11,84	25 12,14	—	20 16 12,10	12,19	11,60	-0,09	+0,50	2,148			
2402	5	39 Cygni η	6 9,12	1 9,44	—	20 17 9,18	—	9,01	—	+0,17	2,387			
2403	5	10 Capricorni π	2 41,95	3 41,90	—	20 17 41,90	—	41,80	—	+0,10	3,443			
2404	5	11 Capricorni ρ	3 16,23	2 16,33	3 16,31	20 19 16,28	16,27	15,74	+0,01	+0,54	3,432			
2405	6.7	Capricorni	—	6 24,83	—	20 19 24,81	—	24,43	—	+0,38	3,424			
2406	6	Capricorni f	—	5 39,46	—	20 19 39,45	—	39,28	—	+0,17	3,532			
2407	7	Capricorni σ^1	—	2 14,38	—	20 20 14,37	—	14,41	—	-0,04	3,448			
2408	6	12 Capricorni σ^2	—	6 15,66	3 15,64	20 20 15,64	—	15,61	—	+0,03	3,448			
2409	5	69 Aquilæ G	6 52,04	—	—	20 20 52,04	—	51,70	—	+0,34	3,134			
2410	6	1 Delphini	—	—	6 15,56	20 22 15,57	—	14,92	—	+0,65	2,870			
2411	4.5	41 Cygni i	6 31,93	2 31,91	5 31,69	20 22 31,85	32,07	31,83	-0,22	+0,02	2,416			
2412	6	Capricorni	—	—	6 51,48	20 22 51,45	—	51,22	—	+0,23	3,586			
2413	6	Capricorni v	—	5 12,08	—	20 23 12,05	—	10,94	—	+1,11	3,268			
2414	7	Capricorni	—	5 50,41	—	20 24 50,40	—	49,78	—	+0,62	3,343			
2415	5	45 Cygni ω^2	6 51,33	—	—	20 24 51,33	—	50,76	—	+0,57	1,854			
2416	4	2 Delphini e	7 11,28	—	2 11,25	20 25 11,28	11,21	10,88	+0,07	+0,40	2,864			
2417	3	Indi α	—	7 43,25	—	20 25 43,21	—	43,37	—	-0,16	4,257			
2418	7	Capricorni	—	4 1,46	—	20 26 1,45	—	1,01	—	+0,44	3,399			
2419	6	3 Delphini η	—	—	5 0,36	20 26 0,37	—	59,93	—	+0,44	2,831			
2420	5	46 Cygni ω^3	—	—	1 8,05	20 26 8,13	—	7,64	—	+0,49	1,848			
2421	5	Pavonis ν	—	—	6 26,63	20 26 26,54	—	24,98	—	+1,56	5,610			
2422	5	2 Cephei θ	3 45,68	—	—	20 26 45,08	—	44,05	—	+1,03	1,016			
2423	5	4 Delphine ζ	6 27,30	—	4 27,31	20 27 27,32	—	26,78	—	+0,54	2,800			
2424	6	13 Capricorni τ^1	—	—	3 55,60	29 27 55,58	—	55,43	—	+0,15	3,369			
2425	5.6	70 Aquilæ H	—	—	7 58,69	20 27 58,69	—	58,61	—	+0,08	3,126			
2426	3	Pavonis β	2 42,96	3 43,12	—	20 29 43,02	—	43,45	—	-0,43	5,546			
2427	5	71 Aquilæ I	4 39,59	2 39,59	—	20 29 39,59	—	39,30	—	+0,29	3,099			
2428	4	6 Delphini β	—	—	5 40,46	20 29 40,48	40,31	39,91	+0,17	+0,57	2,803			
2429	5.6	5 Delphini ι	—	5 46,88	—	20 29 46,88	—	46,51	—	+0,37	2,866			
2430	6	14 Capricorni τ^2	—	—	3 52,36	20 29 52,34	—	52,20	—	+0,14	3,363			

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N.P.D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	1831		1832		1833					Green	A. S. C.	
	No.	"	No.	"	No.	"				"	"	
2386	5	1 12,86	—	—	—	—	103 1 12,86	1 16,06	1 10,14	-3,20	+ 2,72	-10,637
2387	5	45 50,78	—	—	—	—	43 45 50,21	45 53,34	45 48,48	-3,13	+ 1,73	10,641
2388	23	3 27,74	7	3 29,47	1	3 30,17	103 3 23,21	3 32,96	3 28,10	-4,75	+ 0,11	10,667
2389	—	—	—	—	—	—	5 41 45,05	62 41 45,05	41 47,42	-2,37	+ 2,46	10,674
2390	—	—	—	—	—	—	5 54 40,57	68 54 40,57	54 39,56	—	+ 1,01	10,684
2391	—	—	5	56 36,76	—	—	33 56 36,76	56 37,84	56 39,70	-1,08	- 2,94	10,727
2392	5	50 27,17	—	—	—	—	65 50 27,17	50 25,80	—	—	+ 1,37	10,733
2393	—	—	5	38 12,81	—	—	109 38 12,81	38 10,35	38 6,89	+2,46	+ 5,92	10,738
2394	—	—	4	47 50,95	3	47 51,22	42 47 51,07	47 54,13	47 55,00	-3,06	- 3,93	10,785
2395	—	—	—	—	—	—	105 18 23,15	18 18,79	—	—	—	10,858
2396	—	—	4	16 40,92	2	16 51,46	103 16 50,43	16 49,86	—	—	+ 0,57	10,859
2397	5	18 23,85	5	18 22,45	—	—	105 18 23,15	18 19,52	18 13,34	+3,63	+ 4,36	10,876
2398	5	15 49,93	—	—	—	—	147 15 49,93	15 46,81	—	—	+ 3,12	10,927
2399	5	47 49,50	3	47 51,65	—	—	12 47 50,31	47 53,52	47 57,12	-3,21	+ 6,81	11,092
2400	—	—	5	5 4,94	—	—	66 5 4,94	5 3,71	—	—	+ 1,23	11,117
2401	5	16 40,68	20	16 41,22	26	16 41,32	50 16 41,22	16 37,75	16 35,43	+3,47	+ 5,79	11,216
2402	4	20 49,13	1	20 51,46	—	—	58 20 49,57	20 50,01	—	—	- 0,44	11,285
2403	5	45 19,24	—	—	—	—	118 45 19,24	45 16,81	—	—	+ 2,43	11,322
2404	4	21 42,41	4	21 43,54	—	—	108 21 42,97	21 46,32	21 41,43	-3,35	+ 1,54	11,435
2405	—	—	5	58 58,88	—	—	107 58 58,88	59 0,27	—	—	- 1,39	11,445
2406	—	—	5	56 29,72	—	—	112 56 29,72	56 28,46	—	—	+ 1,26	11,463
2407	—	—	5	7 56,70	—	—	109 7 56,70	8 2,98	—	—	- 6,28	11,505
2408	—	—	2	7 56,55	3	7 58,23	109 7 57,56	7 49,66	—	—	+ 7,90	11,506
2409	7	26 14,91	—	—	—	—	93 26 14,94	26 12,74	—	—	+ 2,20	11,550
2410	—	—	5	39 43,66	—	—	79 39 43,66	39 34,74	—	—	+ 8,92	11,650
2411	6	11 18,92	5	11 19,31	6	11 18,90	60 11 19,03	11 15,28	11 11,63	+3,75	+ 7,40	11,671
2412	—	—	2	30 17,33	3	30 19,04	115 30 18,36	30 12,38	—	—	+ 5,98	11,691
2413	—	—	5	25 15,02	—	—	100 25 15,02	25 14,73	—	—	+ 0,29	11,715
2414	—	—	4	17 31,92	—	—	104 17 31,92	17 31,44	—	—	+ 0,48	11,831
2415	5	36 35,51	—	—	—	—	41 36 35,51	36 34,87	—	—	+ 0,64	11,836
2416	5	15 44,88	1	15 46,96	3	15 44,45	79 15 44,97	15 45,39	15 38,88	-0,42	+ 6,09	11,857
2417	—	—	5	52 7,51	—	—	137 52 7,51	52 9,06	—	—	- 1,55	11,892
2418	—	—	5	5 48,17	—	—	107 5 48,17	5 44,69	—	—	+ 3,48	11,915
2419	—	—	1	32 33,06	4	32 34,90	77 32 34,53	32 32,13	—	—	+ 2,40	11,915
2420	2	20 31,24	—	—	—	—	41 20 31,24	20 37,61	—	—	- 3,37	11,926
2421	—	—	1	20 31,30	3	20 31,74	157 20 31,63	20 31,72	—	—	- 0,09	11,938
2422	5	34 4,97	—	—	—	—	27 34 4,97	34 10,85	—	—	- 5,88	11,971
2423	5	54 1,38	—	—	—	—	75 54 1,38	53 52,12	—	—	+ 9,26	12,017
2424	—	—	5	43 25,44	3	43 25,10	105 43 25,10	43 16,73	—	—	+ 8,37	12,049
2425	—	—	2	7 31,11	—	—	93 7 31,11	7 23,97	—	—	+ 2,44	12,053
2426	5	47 43,28	—	—	—	—	156 47 43,23	47 51,88	—	—	- 8,60	12,169
2427	—	—	5	41 11,59	—	—	91 41 11,59	41 4,89	—	—	+ 6,70	12,170
2428	—	—	5	59 2,87	5	59 2,87	75 59 2,87	59 2,92	58 52,03	+0,05	+ 10,84	12,171
2429	—	—	5	12 12,15	—	—	79 12 12,45	12 11,94	—	—	+ 0,51	12,179
2430	—	—	5	32 19,25	105	32 19,25	105 32 19,25	32 8,85	—	—	+ 10,40	12,184

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.			Green Catal.	A. S. Catal.	Difference from		Annual Preces- sion.
			No.	1831	No.	1832	No.	1833			Green	A. S.	
			s.	s.	s.	h.	m.	s.			s.	s.	
2431	5.6	27 Velpculæ p							54,53			+2,554	
2432	5	15 Capricorni v	1 28,90	5 28,78		20 30 28,80	28,80	28,37	0,00	+0,43	3,427		
2433	5.6	1 Aquarii				20 30		48,18			3,070		
2434	4.5	8 Delphini θ	6 48,33			20 30 48,33	48,26	47,76	+0,07	+0,57	2,829		
2435	5.6	7 Delphini κ			3 58,31	20 30 58,32		57,72		+0,60	2,891		
2436	5.6	29 Vulpeculæ s			3 1,41	20 31 1,43		0,62		+0,82	2,671		
2437	6.7	Capricorni			4 5,80	20 31 5,78		5,85		-0,07	3,385		
2438	5.6	28 Vulpeculæ			3 12,87	20 31 12,90		12,31		+0,59	2,608		
2439	3.4	9 Delphini α	5 50,16	1 50,18	1 50,28	20 31 50,18	50,43	50,05	-3,25	+0,13	2,779		
2440	6	Cygni		5 4,88		20 32 4,90		4,45		+0,45	2,466		
2441	7	Capricorni		1 6,06	2 6,07	20 33 6,05		5,70		+0,35	3,423		
2442	6	10 Delphini			5 24,32	20 33 24,34		24,04		+0,30	2,807		
2443	5	11 Delphini δ	4 37,16			20 35 37,16		36,91		+0,25	2,800		
2444	1	50 Cygni α	31 42,50	50 42,44		20 35 42,48	42,46	41,97	+0,02	+0,51	2,040		
2445	4.5	16 Capricorni ψ	5 8,33			20 36 8,33	8,29	8,58	+0,04	-0,25	3,572		
2446	6	17 Capricorni		6 25,15		20 36 25,13		24,77		+0,36	3,490		
2447	6	30 Vulpeculæ		5 36,60		20 37 36,62		36,17		+0,45	2,594		
2448	7	Capricorni		1 27,96	7 28,08	20 38 28,04		28,05		-0,01	3,515		
2449	6	Capricorni			2 33,00	20 38 32,97		32,82		+0,15	3,512		
2450	4.5	2 Aquarii ε	6 34,29			20 38 34,89	34,69	34,51	+0,20	+0,38	3,252		
2451	4	3 Aquarii κ	3 52,04			20 38 52,04	52,16	51,70	-0,12	+0,34	3,170		
2452	4	12 Delphini γ	6 52,03			20 38 52,03	51,94	52,72	+0,09	-0,69	2,783		
2453	6	Capricorni			3 18,60	20 39 18,57		18,49		+0,08	3,578		
2454	4.5	Microscopii α		5 27,56		20 39 27,54	27,25	26,63	+0,29	+0,91	3,771		
2455	3	53 Cygni ε		5 25,06		20 39 25,08	24,89	24,52	+0,19	+0,56	2,393		
2456	5.6	13 Delphini λ	4 52,93	1 53,00		20 39 52,95		20,01			2,971		
2457	6.7	Capricorni			3 48,32	20 39 48,30		48,21		+0,09	3,414		
2458	5	54 Cygni λ	4 52,29	1 52,03	5 52,02	20 40 52,15		51,41		+0,74	2,330		
2459	5	Cephei χ		5 10,70		20 41 10,75		10,89		-0,14	1,500		
2460	6	Capricorni			3 20,40	20 41 20,37		17,34			3,595		
2461	6.7	Capricorni p			3 25,91	20 41 25,90		25,24		+0,66	3,306		
2462	6.7	Capricorni			1 29,87	20 41 29,84		29,81		+0,03	3,607		
2463	4	Indi β		5 37,16		20 41 37,10		37,20		-0,10	4,768		
2464	5.6	18 Capricorni ω			5 46,95	20 41 46,92		47,08		-0,16	3,599		
2465	3.4	3 Cephei γ	1 51,69	3 51,28		20 41 51,44	51,74	50,47	-0,30	+0,97	1,920		
2466	6	4 Aquarii		3 31,15	3 30,94	20 42 31,04		30,97		+0,07	3,179		
2467	7	Aquarii		2 49,30		20 42 49,29		48,51		+0,78	3,285		
2468	6	Capricorni m			3 9,36	20 43 9,33		8,81		+0,52	3,527		
2469	6	5 Aquarii		2 15,55		20 43 15,54		15,35		+0,19	3,177		
2470	4.5	6 Aquarii μ	6 35,47		1 35,29	20 43 35,43	35,29	35,07	+0,14	+0,36	3,239		
2471	6	Aquarii		4 53,97		20 43 53,96		53,45		+0,51	3,286		
2472	5	Octantis α				20 44		3,96			7,674		
2473	6	31 Vulpeculæ r		5 56,53		20 44 56,55		55,71		+0,84	2,568		
2474	6	19 Capricorni	1 17,88	3 17,50	2 17,98	20 45 17,69	17,89	17,43	-0,20	+0,26	3,405		
2475	7	Capricorni		5 47,75		20 46 47,68		47,29		+0,39	3,575		

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N.P.D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No.	1831	No.	1832	No.	1833				Green.	A. S. C.	
2431	—	—	—	—	5 7 4,89	64 7 4,89	—	6 58,50	—	+ 6,39	—12,189	
2432	—	—	—	—	4 43 23,37	108 43 23,37	43 27,13	43 19,27	-3,76	+ 4,10	12,226	
2433	—	—	3 5 58,08	—	2 6 1,54	90 5 59,46	—	5 51,15	—	+ 8,30	12,250	
2434	4 16 8,67	—	3 16 9,43	—	—	77 8 8,99	16 10,24	46 6,31	-1,25	+ 2,68	12,250	
2435	—	—	—	—	5 30 3,77	80 3 3,77	—	29 56,69	—	+ 7,08	12,261	
2436	—	—	—	—	4 23 1,48	69 23 1,48	—	22 56,97	—	+ 4,51	12,265	
2437	—	—	—	—	3 12 55,74	106 42 55,74	—	42 57,31	—	- 1,57	12,269	
2438	—	—	—	—	—	66 28 —	—	28 6,85	—	—	12,279	
2439	5 40 35,68	—	2 40 35,58	—	—	74 40 35,64	40 32,49	40 25,94	+3,15	+ 9,70	12,322	
2440	—	—	5 15 2,19	—	—	60 15 2,19	—	14 57,00	—	+ 5,19	12,339	
2441	—	—	1 42 14,62	—	4 42 13,99	108 42 14,12	—	42 15,73	—	- 1,61	12,407	
2442	—	—	3 0 31,62	—	2 0 35,58	76 0 35,01	—	0 30,66	—	+ 4,35	12,429	
2443	6 31 25,12	—	1 31 23,27	—	—	75 31 24,87	—	31 17,99	—	+ 6,88	12,581	
2444	70 19 1,75	—	56 19 1,72	—	40 19 1,61	45 19 1,70	19 0,6	18 57,07	+1,54	+ 4,63	12,588	
2445	—	—	—	—	5 52 6,55	115 52 6,55	52 5,03	52 1,86	+1,52	+ 4,69	12,615	
2446	—	—	5 7 7,24	—	—	112 7 7,24	—	7 4,10	—	+ 3,14	12,633	
2447	—	—	1 19 35,06	—	—	65 19 35,06	—	19 29,98	—	+ 5,08	12,716	
2448	—	—	—	—	4 27 25,09	113 27 25,09	—	27 25,54	—	- 0,45	12,772	
2449	—	—	1 20 36,16	—	5 20 38,31	113 20 37,98	—	20 32,06	—	+ 5,92	12,778	
2450	4 6 22,64	—	2 6 22,08	—	—	100 6 22,46	6 19,34	6 10,84	+3,12	+ 11,62	12,780	
2451	5 38 15,99	—	—	—	—	95 38 15,99	38 15,37	38 9,12	+0,62	+ 6,87	12,800	
2452	4 28 33,62	—	1 28 35,07	—	2 23 35,26	74 28 34,30	28 34,28	28 28,46	+0,02	+ 5,84	12,802	
2453	—	—	—	—	5 23 42,09	116 23 42,09	—	23 38,45	—	+ 3,64	12,829	
2454	—	—	2 23 41,40	—	3 23 43,11	124 23 42,43	—	23 37,88	—	+ 4,55	12,837	
2455	5 39 16,54	—	5 39 17,50	—	—	56 32 17,02	39 16,05	39 11,38	0,97	+ 5,64	12,838	
2456	—	—	1 36 16,59	—	4 36 16,71	84 36 16,69	—	36 15,83	—	+ 0,86	12,842	
2457	—	—	5 38 56,19	—	—	108 38 56,19	—	38 56,70	—	- 0,51	12,862	
2458	5 7 23,02	—	—	—	—	54 7 23,02	—	7 14,92	—	+ 8,10	12,935	
2459	—	—	6 1 13,84	—	—	33 1 13,84	—	1 6,54	—	+ 7,30	12,959	
2460	—	—	—	—	—	117 19 —	—	19 7,83	—	—	12,961	
2461	—	—	—	—	3 9 39,42	103 9 39,42	—	9 36,56	—	+ 2,86	12,971	
2462	—	—	—	—	5 51 50,51	117 51 50,51	—	51 50,17	—	+ 0,34	12,975	
2463	4 4 43,19	—	—	—	—	149 4 43,19	—	4 35,34	—	+ 7,85	12,981	
2464	—	—	3 32 25,56	—	3 32 26,19	117 32 25,87	—	32 22,87	—	+ 3,00	12,994	
2465	—	—	4 48 39,93	—	1 48 38,92	28 48 39,73	48 42,95	48 42,08	-3,22	- 2,35	13,813*	
2466	—	—	5 14 58,16	—	—	96 14 58,16	—	14 51,36	—	+ 6,80	13,044	
2467	—	—	4 3 46,86	—	—	102 3 46,89	—	3 42,88	—	+ 3,08	13,063	
2468	—	—	—	—	5 24 25,10	114 24 25,10	—	24 20,51	—	+ 4,59	13,085	
2469	—	—	—	—	5 7 53,29	96 7 53,29	—	7 47,34	—	+ 5,92	13,093	
2470	5 36 31,88	—	2 36 31,76	—	—	99 37 31,90	36 29,82	36 21,83	+2,08	+ 10,07	13,114	
2471	—	—	3 12 11,13	—	—	102 12 11,13	—	12 11,99	—	- 0,86	13,135	
2472	—	—	—	—	—	167 38 Invis.	—	38 4,89	—	—	13,137	
2473	—	—	5 31 36,72	—	—	63 31 36,72	—	31 35,95	—	+ 0,27	13,205	
2474	—	—	1 33 10,12	—	4 33 12,15	108 33 11,75	33 15,43	33 12,21	—	- 0,46	13,227	
2475	—	—	5 55 50,26	—	—	116 55 50,26	—	55 47,34	—	+ 2,92	13,324	

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	NAMES.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.			Green. Catal.	A. S. Catal.	Difference from		Annual Preces- sion.
			No. 1831	No. 1832	No. 1833	h.	m.	s.			Green.	A. S.	
			s.	s.	s.						s.	s.	
2476	6	Equulei		5 15,99		20	47	15,98		15,34		+0,64	+3,000
2477	5	57 Cygni	7 18,30			20	47	18,30		18,02		+0,28	2,115
2478	4.5	32 Vulpeculæ <i>q</i>	5 24,24		1 24,16	20	47	24,23	24,16	23,74	+0,07	+0,49	2,552
2479	6	16 Delphini <i>x</i>		2 37,61	3 38,03	20	47	37,87		37,13		+0,74	2,858
2480	6	17 Delphini <i>μ</i>			7 39,60	20	47	39,62		39,06		+0,56	2,837
2481	6	7 Aquarii			4 48,87	20	47	48,86		48,94		-0,08	3,249
2482	7	Capricorni			3 15,96	20	48	15,94		15,61		+0,33	3,365
2483	6	Equulei		5 23,78	1 23,93	20	49	23,81		23,43		+0,38	3,007
2484	6	20 Capricorni		5 3,07	1 2,82	20	50	3,02		2,91		+0,11	3,421
2485	6	18 Delphini <i>ν</i>			4 20,59	20	50	20,60		20,06		+0,54	2,891
2486	5.6	1 Equulei <i>e</i>			2 40,61	20	50	40,62		40,41		+0,21	3,005
2487	6	8 Aquarii <i>z</i> ¹			3 40,43	20	50	40,42		41,13		-0,71	3,308
2488	5.6	33 Vulpeculæ <i>x</i>			2 45,83	20	50	45,89		45,60		+0,29	2,678
2489	4	58 Cygni <i>ν</i>	6 54,59		1 51,78	20	50	54,63	54,88	54,36	-0,25	+0,27	2,229
2490	6	21 Capricorni	1 21,01		5 23,90	20	51	23,96		23,82		+0,14	3,390
2491	6	11 Aquarii <i>r</i>		2 42,80		20	51	42,80		42,74		+0,06	3,160
2492	6	Capricorni <i>n</i>		1 46,04	3 46,38	20	51	46,27		45,66		+0,56	3,578
2493	5	Cephei <i>K</i>	2 47,23	2 47,27		20	51	47,28					1,605
2494	6	9 Aquarii <i>s</i> ²		5 52,47		20	51	52,46		52,32		+0,14	3,315
2495	6	2 Equulei <i>λ</i>			6 55,68	20	53	55,69		55,33		+0,16	+2,957
2496	.5	76 Draconis		19 15,83		20	54	15,93		17,90		-1,97	-3,725
2497	7	22 Capricorni <i>η</i>	3 50,10		1 50,14	20	54	50,11	50,15	50,07	+0,07	+0,04	+3,430
2498	6	12 Aquarii		5 11,44		20	55	11,43		10,89		+0,54	+3,178
2499	5	Cephei <i>h</i>		3 55,47	1 55,50	20	55	55,48		46,96			-2,335
2500	6	3 Equulei <i>ξ</i>		5 12,65		20	56	12,86		12,24		+0,62	+2,987
2501	5.6	23 Capricorni <i>θ</i>			4 29,86	20	56	29,84	30,03	29,36	+0,19	+0,48	3,378
2502	7	Capricorni		5 7,10		20	57	7,09		6,62		+0,47	3,433
2503	6	4 Equulei		4 7,22	2 7,29	20	57	7,23		6,70		+0,55	2,979
2504	5.6	24 Capricorni <i>A</i>			3 17,54	20	57	17,51		17,05		+0,46	3,528
2505	4	62 Cygni <i>ξ</i>	1 49,18	1 49,64		20	58	49,42	49,43	48,72	0,01	+0,70	2,174
2506	5.6	25 Capricornix ¹			3 55,57	20	58	55,55	55,53	55,60	+0,02	-0,05	3,449
2507	6	27 Capricornix ²			5 26,26	20	59	56,24		56,06		+0,18	3,435
2508	.5	13 Aquarii <i>ν</i>	2 26,27			21	0	26,27	26,17	25,61	0,10	+0,63	3,270
2509	5	63 Cygni <i>f</i> ²		6 48,90		21	0	48,94		48,64		+0,30	2,059
2510	5	5 Equulei <i>γ</i>	4 10,37			21	2	10,37		10,10		+0,27	2,912
2511	6	3 Piscis Aust		5 18,93		21	3	18,90		19,02		-0,12	3,497*
2512	3	64 Cygni <i>ξ</i>	3 47,40	5 47,29		21	5	47,34	47,42	46,81	-0,06	+0,53	2,546
2513	7	Aquarii		5 58,77		21	5	58,76		58,31		+0,46	3,194
2514	6	28 Capricorni <i>φ</i>		4 3,54	4 3,65	21	6	3,58	3,46	3,34	+0,12	+0,24	3,428
2515	5.4	7 Equulei <i>δ</i>	1 17,96	4 18,05		21	6	18,04	17,97	17,80	0,07	+0,24	2,917
2516	5	29 Capricorni <i>S</i>		5 26,58		21	6	26,57	26,60	26,68	-0,03	-0,11	3,329
2517	4.5	8 Equulei <i>a</i>	1 25,63	4 25,42	1 25,58	21	7	25,46	25,59	25,28	0,13	+0,18	2,995
2518	5	4 Piscis Aust	5 44,28			21	7	44,28		43,43		+0,85	3,658
2519	5	65 Cygni <i>τ</i>	1 5,58	4 5,30		21	8	5,38		4,80		+0,58	2,373
2520	6	30 Capricorni <i>r</i>		2 31,36	3 31,58	21	8	31,47		31,65		-0,18	3,376

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N.P.D. January 1, 1832.			Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	1831		1832		1833							Green.	A. S. C.	
	No.	"	No.	"	No.	"	"	"	"					
2476	—	—	1 6 13,21	—	—	4 6 15,24	86 6 14,83	—	—	6 12,24	—	+	2,59	+13,356
2477	5 14 43,60	—	—	—	—	—	46 14 43,60	—	—	14 43,15	—	+	0,45	13,361
2478	6 34 34,65	—	—	—	—	2 34 34,94	62 34 34,72	34 38,02	—	34 31,79	—	+	2,98	13,366
2479	—	—	5 4 6,76	—	—	—	78 4 6,76	—	—	4 6,33	—	+	0,43	13,380
2480	—	—	3 54 52,57	—	—	2 54 52,55	76 54 52,56	—	—	54 42,90	—	+	9,66	13,382
2481	—	—	—	—	5 20 13,16	—	100 20 13,16	—	—	20 2,47	—	+	10,69	13,392
2482	—	—	—	—	4 40 19,19	—	106 40 19,19	—	—	40 18,85	—	+	0,34	13,421
2483	—	—	—	—	5 26 50,51	—	86 26 50,51	—	—	26 49,10	—	+	1,41	13,495
2484	—	—	4 40 54,16	—	—	1 40 54,75	109 40 54,28	—	—	40 46,30	—	+	7,98	13,536
2485	—	—	5 48 18,44	—	—	—	79 48 18,44	—	—	48 12,70	—	+	5,74	13,556
2486	—	—	—	—	5 20 45,94	—	86 20 45,94	—	—	20 42,62	—	+	3,32	13,578
2487	—	—	—	—	5 41 54,08	—	103 41 54,08	—	—	41 51,18	—	+	2,90	13,578
2488	—	—	2 19 5,25	—	—	3 19 6,14	68 19 5,77	—	—	19 5,99	—	—	0,22	13,584
2489	5 28 34,87	—	1 28 35,08	—	—	—	49 28 34,90	28 34,23	—	28 28,62	—	+	6,28	13,594
2490	—	—	—	—	5 10 46,70	—	108 10 46,70	—	—	10 46,44	—	+	0,26	13,623
2491	—	—	5 22 30,31	—	—	—	95 22 30,31	—	—	22 18,87	—	+	11,44	13,644
2492	—	—	1 31 53,19	—	—	5 31 53,57	117 31 53,49	—	—	31 52,02	—	+	1,47	13,646
2493	6 45 22,03	—	—	—	—	—	33 45 22,03	—	—	45 21,79	—	+	0,24	13,652
2494	—	—	—	—	2 10 51,47	—	104 10 51,47	—	—	10 47,90	—	+	3,57	13,654
2495	—	—	5 28 30,06	—	—	—	83 28 30,06	—	—	28 27,95	—	+	2,11	13,785
2496	4 5 49,32	—	1 5 50,53	—	—	6 5 48,77	8 5 49,13	—	—	5 55,81	—	—	6,68	13,823
2497	4 30 49,34	—	1 30 50,39	—	—	—	110 30 49,55	30 48,39	—	30 45,13	—	+	4,42	13,842
2498	—	—	5 28 57,98	—	—	—	96 28 57,98	—	—	28 53,96	—	+	4,02	13,864
2499	—	—	4 6 0,50	—	—	1 6 2,06	10 6 0,81	—	—	—	—	—	—	13,914
2500	—	—	3 9 38,52	—	—	1 9 39,91	85 9 38,87	—	—	9 32,92	—	+	5,95	13,929
2501	—	—	—	—	5 53 39,26	—	107 53 39,26	53 43,69	—	53 42,37	—	—	3,11	13,916
2502	—	—	5 50 48,62	—	—	—	110 50 48,62	—	—	50 45,26	—	+	3,36	13,985
2503	—	—	4 41 7,42	—	—	—	84 41 7,42	—	—	41 59,62	—	—	—	13,986
2504	—	—	—	—	5 40 19,72	—	115 40 19,72	—	—	40 11,02	—	+	8,70	13,996
2505	5 44 19,74	—	7 44 19,64	—	—	5 44 19,85	46 44 19,72	44 20,82	—	44 19,79	—	—	0,07	14,094
2506	—	—	5 51 46,64	—	—	—	111 51 46,64	51 45,23	—	51 40,97	—	+	5,67	14,099
2507	—	—	5 13 31,30	—	—	—	111 13 31,30	—	—	13 26,22	—	+	5,08	14,161
2508	5 2 48,46	—	—	—	—	—	102 2 48,46	2 49,92	—	2 43,06	—	—	5,40	14,192
2509	—	—	5 1 22,65	—	—	—	43 1 22,65	—	—	1 28,83	—	—	6,18	14,218
2510	5 32 28,77	—	—	—	—	—	80 32 28,77	—	—	32 23,81	—	+	4,96	14,300
2511	—	—	6 17 53,16	—	—	—	118 17 53,16	—	—	17 51,92	—	+	1,24	14,369
2512	5 27 31,51	—	—	—	6 27 32,97	—	60 27 32,30	27 30,04	—	27 27,72	—	+	4,58	14,520
2513	—	—	5 46 36,54	—	—	—	97 46 36,54	—	—	46 37,44	—	—	0,90	14,530
2514	—	—	4 20 36,06	—	—	—	111 20 36,06	20 37,12	—	20 32,58	—	—	3,48	14,535
2515	—	—	7 40 12,69	—	—	5 40 12,45	80 40 12,59	40 8,96	—	40 3,41	—	+	9,18	14,550
2516	—	—	5 51 52,31	—	—	—	105 51 52,31	51 55,13	—	51 45,53	—	—	6,78	14,558
2517	6 26 35,56	—	—	—	—	—	85 26 35,56	26 31,75	—	26 28,03	—	+	7,52	14,618
2518	4 52 5,40	—	—	—	—	—	122 52 5,40	—	—	52 58,83	—	+	6,57	14,634
2519	3 40 2,33	—	1 40 3,25	—	—	—	52 40 2,56	—	—	39 58,00	—	+	4,56	15,158*
2520	—	—	2 40 57,48	—	—	4 40 58,19	108 40 57,92	—	—	40 56,12	—	+	1,80	14,683

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	NAMES.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.			Green. Catal.	A. S. Catal.	Difference from		Annual Preces- sion.
			No. 1831	No. 1832	No. 1833	h.	m.	s.			Green	A. S.	
			s.	s.	s.						s.	s.	
2521	6.7	31 Capricorni	—	—	5 51,04	21	8	51,02	—	51,50	—0,48	+3,366	
2522	7	Aquarii	—	—	6 54,71	21	9	54,69	—	54,39	+0,30	3,312	
2523	4.5	67 Cygni	7 49,35	—	—	21	10	49,35	49,95	48,26	-0,10	+1,09	
2524	4.5	66 Cygni	—	3 0,71	—	21	11	0,71	0,89	59,77	-0,18	+0,94	
2525	6	16 Aquarii	—	4 15,75	—	21	12	15,75	—	15,16	+0,59	3,151	
2526	3	Pavonis	—	—	5 27,84	21	12	27,75	—	26,77	+0,98	5,086	
2527	6	9 Equulei	—	—	5 46,09	21	12	46,10	—	45,73	+0,37	2,964	
2528	5	32 Capricorni	4 53,06	2 53,06	—	21	12	53,06	53,01	53,13	+0,05	-0,07	
2529	7	Aquarii	—	—	4 57,40	21	12	57,40	—	57,28	+0,12	3,226	
2530	6	Capricorni	—	—	4 22,14	21	13	22,12	—	21,59	+0,53	3,452	
2531	6	17 Aquarii	—	5 55,58	—	21	13	55,57	—	55,83	-0,26	3,225	
2532	7	Capricorni	—	5 9,66	—	21	14	9,64	—	9,43	+0,21	3,408	
2533	5	Indi	3 13,31	—	—	21	14	13,31	—	13,56	-0,25	4,350	
2534	4	1 Pegasi	—	5 19,17	—	21	14	19,18	19,17	18,91	+0,01	+0,27	
2535	5.6	10 Equulei	—	5 33,28	—	21	14	33,29	—	33,01	+0,28	2,974	
2536	3	5 Cephei	3 34,12	5 33,80	9 33,12	21	14	33,66	33,89	33,07	-0,23	+0,59	
2537	6	33 Capricorni	—	—	3 37,40	21	14	37,38	—	37,09	+0,29	3,417	
2538	6	18 Aquarii	1 0,49	—	4 0,49	21	15	0,48	—	0,10	+0,38	3,281	
2539	5	6 Cephei	7 52,49	—	—	21	15	52,49	—	51,83	+0,69	1,257	
2540	6	19 Aquarii	—	—	3 11,01	21	16	11,00	—	10,59	+0,41	3,230	
2541	6	Pegasi	—	—	1 24,76	21	16	24,79	—	24,25	+0,54	2,687	
2542	6	21 Aquarii	—	5 31,19	—	21	16	31,18	—	31,11	+0,07	3,133	
2543	4	34 Capricorni	6 3,88	—	—	21	17	3,88	3,80	3,40	+0,08	+0,48	
2544	6	Pegasi	—	4 6,67	—	21	17	6,69	—	6,75	-0,06	2,653	
2545	6	35 Capricorni	—	5 42,83	—	21	17	42,81	—	42,31	+0,50	3,418	
2546	5.6	36 Capricorni	—	5 8,02	3 8,15	21	19	8,05	8,04	7,52	+0,01	+0,53	
2547	7	Capricorni	—	3 33,35	2 33,39	21	20	33,35	—	33,08	+0,27	3,373	
2548	7	Capricorni	—	5 43,10	—	21	20	43,08	—	42,69	+0,39	3,484	
2549	7	Aquarii	—	—	6 27,82	21	21	27,80	—	27,60	+0,20	3,297	
2550	5.6	2 Pegasi	2 20,63	4 20,54	1 20,33	21	22	20,55	—	20,57	-0,02	2,710	
2551	3	22 Aquarii	6 42,82	6 42,72	12 42,62	21	22	42,68	42,54	42,47	+0,14	+0,21	
2552	6	Capricorni	—	7 51,91	—	21	22	51,89	—	51,29	+0,60	3,469	
2553	5	71 Cygni	5 15,32	—	—	21	23	15,32	—	14,39	+0,93	2,200	
2554	6.7	Capricorni	—	5 6,55	—	21	25	6,54	—	6,02	+0,52	3,280	
2555	7	37 Capricorni	—	6 24,46	—	21	25	24,45	—	24,39	+0,06	3,386	
2556	7	38 Capricorni	—	2 27,46	3 27,48	21	25	27,46	—	27,38	+0,08	3,388	
2557	7	Capricorni	—	5 38,82	—	21	25	38,80	—	38,39	+0,41	3,443	
2558	5.6	8 Piscis Aust	—	4 23,81	—	21	26	23,79	—	25,51	+0,23	3,490	
2559	3	8 Cephei	2 27,66	—	1 26,91	21	26	27,59	27,74	26,69	-0,15	+0,90	
2560	5	39 Capricorni	6 40,01	—	3 39,79	21	27	39,94	39,99	39,78	-0,05	+0,16	
2561	5	73 Cygni	5 40,03	—	1 40,07	21	27	40,05	40,09	39,75	-0,04	+0,30	
2562	5	23 Aquarii	5 48,16	6 48,25	—	21	28	48,20	48,12	48,21	+0,08	-0,01	
2563	6	3 Pegasi	—	5 21,57	—	21	29	21,58	—	20,88	+0,70	2,984	
2564	5.6	5 Pegasi	—	4 54,07	2 53,90	21	29	54,03	—	53,27	+0,76	2,795	
2565	5	4 Pegasi	1 6,91	4 7,17	—	21	30	7,12	—	6,73	+0,39	2,997	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N.P.D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No.	1831	No.				1832	No.	
2521	—	—	—	4 9 39,38	108 9 39,38	9 37,00		+ 2,38	+14,703
2522	—	—	5 52 49,39	—	106 52 49,30	52 47,98		+ 1,32	14,763
2523	5 19 20,82	—	1 19 22,16	—	51 19 21,06	19 24,65	-1,01	- 3,59	14,829
2524	5 48 17,46	—	—	—	55 48 17,46	48 14,69	-0,04	+ 2,77	14,831
2525	—	—	1 16 9,75	4 16 11,25	95 16 10,95	16 0,10		+ 10,85	14,903
2526	3 7 3,34	—	1 7 4,40	1 7 5,02	156 7 3,87	7 10,90		- 7,03	14,911
2527	—	—	1 21 7,53	4 21 10,48	83 21 9,89	21 5,76		+ 4,10	14,933
2528	—	—	5 32 38,61	—	107 32 38,61	32 43,17	-4,56	+ 4,01	14,940
2529	—	—	—	5 2 13,65	100 2 13,65	—		—	14,944
2530	—	—	—	3 22 49,20	113 22 49,20	22 49,53		- 0,33	14,967
2531	—	—	—	3 1 53,38	100 1 53,38	1 44,94		+ 8,44	15,001
2532	—	—	3 54 56,84	—	115 54 56,84	54 52,36		+ 4,48	15,013
2533	5 22 48,49	—	—	—	145 22 48,49	22 34,03		+ 14,46	15,016
2534	5 54 37,74	—	—	—	70 54 37,74	54 36,23	+1,51	+ 4,33	15,024
2535	—	—	—	5 54 7,96	83 54 7,96	54 3,83		+ 4,13	15,037
2536	5 7 22,39	—	6 7 24,22	12 7 25,81	28 7 24,65	7 28,14	-3,49	- 7,39	15,040
2537	—	—	5 33 37,52	—	111 33 37,52	33 35,98		+ 1,54	15,040
2538	—	—	5 35 35,81	—	103 35 35,81	35 33,38		+ 2,43	15,063
2539	5 50 19,56	—	—	—	25 50 19,56	50 21,47		+ 1,91	15,116
2540	—	—	4 27 34,98	2 27 35,03	100 27 35,00	27 27,63		+ 7,37	15,130
2541	—	—	5 26 35,69	—	66 26 35,69	26 37,70		- 2,01	15,144
2542	—	—	—	5 16 20,16	94 16 20,16	16 12,91		+ 7,25	15,150
2543	6 7 59,07	—	—	—	113 7 59,07	8 1,87	-2,80	+ 2,41	15,180
2544	—	—	5 32 42,39	—	64 32 42,39	32 38,69		+ 3,70	15,185
2545	—	—	5 55 3,33	—	111 55 3,33	54 59,60		+ 3,73	15,217
2546	—	—	5 31 58,66	—	112 31 58,66	32 2,74	-4,08	- 4,00	15,298
2547	—	—	5 52 38,17	—	109 52 38,17	52 32,29		+ 5,88	15,378
2548	—	—	5 55 25,22	—	115 55 25,22	55 22,86		+ 2,56	15,387
2549	—	—	5 1 24,13	—	105 1 24,13	1 16,71		+ 7,42	15,429
2550	—	—	5 5 37,04	—	67 5 37,04	5 34,48		+ 2,56	15,479
2551	5 18 20,73	—	2 18 20,00	6 18 20,46	96 18 20,47	18 21,27	-1,25	+ 5,35	15,499
2552	—	—	5 19 40,35	—	115 19 40,35	19 35,41		+ 4,94	15,506
2553	6 11 48,83	—	—	—	44 11 48,83	11 50,31		+ 1,48	15,530
2554	—	—	3 13 29,78	—	104 13 29,78	13 30,94		- 1,16	15,630
2555	—	—	5 49 42,00	—	110 49 42,00	49 36,97		+ 5,03	15,647
2556	—	—	5 59 31,90	—	110 59 31,90	59 27,38		+ 4,52	15,650
2557	—	—	5 11 48,49	—	114 11 48,49	11 48,87		- 0,38	15,660
2558	—	—	5 54 56,42	—	116 54 56,42	54 53,61		+ 2,31	15,702
2559	6 10 37,42	—	—	10 10 35,92	20 10 36,82	10 32,93	+3,89	- 1,29	15,708
2560	5 12 56,01	—	—	—	110 12 56,01	12 49,89		+ 8,69	15,770
2561	5 9 0,11	—	—	2 8 56,41	45 8 59,06	8 52,17	+6,89	+ 5,37	15,771
2562	5 36 7,86	—	—	—	98 36 7,86	36 12,99	-5,13	+ 1,44	15,831
2563	—	—	5 7 56,02	—	84 7 56,02	7 48,57		+ 7,45	15,861
2564	—	—	5 25 58,12	—	71 25 58,12	25 55,25		+ 2,37	15,890
2565	5 59 0,40	—	1 59 2,34	—	84 59 0,72	58 51,19		+ 9,53	15,902

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.			Green Catal.	A. S. Catal.	Difference from		Annual Preces- sion.
			No. 1831	No. 1832	No. 1833						Green.	A. S.	
			s.	s.	s.	h.	m.	s.			s.	s.	
2566	4	40 Capricorni γ	5 46,72	—	5 46,50	21 30 46,68	46,53	46,26	+0,15	+0,42	+3,322		
2567	5.6	25 Aquarii d	—	5 1,86	—	21 31 1,86	—	1,91	—	-0,05	3,047		
2568	6	42 Capricorni d^1	—	—	3 24,42	21 32 24,40	—	24,21	—	+0,19	3,230		
2569	5	41 Capricorni	5 26,10	—	—	21 32 26,10	—	26,20	—	-0,10	3,426		
2570	5	43 Capricorni κ	4 15,96	6 16,22	—	21 33 16,11	16,24	15,67	-0,13	+0,44	3,353		
2571	5	9 Cephei	—	5 24,45	7 24,14	21 33 24,42	—	23,75	—	+0,67	1,610		
2572	6	26 Aquarii	—	5 36,06	—	21 33 36,06	—	36,02	—	+0,04	3,061		
2573	6	Capricorni	—	4 49,23	—	21 33 49,22	—	48,52	—	+0,70	3,364		
2574	5.6	7 Pegasi T^2	—	—	5 51,19	21 33 51,20	—	50,77	—	+0,43	3,000		
2575	6	44 Capricorni d^2	—	5 54,08	—	21 33 54,06	—	53,73	—	+0,33	3,284		
2576	6	45 Capricorni d^3	—	3 50,29	1 50,03	21 34 50,21	—	49,85	—	+0,36	3,288		
2577	4.5	9 Piscis Aust i	6 55,39	—	—	21 34 55,39	55,16	54,91	+0,23	+0,48	3,598		
2578	2.3	8 Pegasi	6 56,16	—	—	21 35 56,16	56,15	55,98	+0,01	+0,18	2,942		
2579	6	46 Capricorni c^1	—	3 2,66	—	21 36 2,65	—	1,76	—	+0,89	3,205		
2580	4.5	80 Cygni π^1	3 8,37	—	6 8,08	21 36 8,21	8,31	7,73	-0,10	+0,48	2,118		
2581	4.5	9 Pegasi g	—	2 33,42	3 33,53	21 36 33,60	33,56	33,63	+0,04	-0,03	2,835		
2582	5	78 Cygni μ	—	6 38,05	—	21 36 38,07	—	37,56	—	+0,51	2,652		
2583	4	10 Pegasi κ	—	2 2,36	2 2,62	21 37 2,52	2,57	2,37	-0,05	+0,15	2,706		
2584	6.7	47 Capricorni c^2	—	—	4 18,34	21 37 18,33	—	18,01	—	+0,32	3,206		
2585	5.6	48 Capricorni λ	—	—	3 29,16	21 37 29,15	29,40	28,38	-0,25	+0,77	3,236		
2586	3.4	49 Capricorni δ	6 45,70	6 45,64	5 45,71	21 37 45,67	45,58	45,42	+0,09	+0,25	3,304		
2587	5	10 Piscis Aust θ	5 51,60	—	—	21 37 51,60	—	51,72	—	-0,12	3,548		
2588	6	12 Pegasi W	—	—	4 20,93	21 38 20,95	—	20,49	—	+0,46	2,752		
2589	5.6	11 Pegasi	—	5 42,77	—	21 38 42,77	—	42,01	—	+0,76	3,042		
2590	4.5	11 Cephei τ	—	5 25,68	—	21 39 25,78	25,97	24,08	-0,19	+1,70	0,892		
2591	7	Aquarii	—	—	4 13,62	21 40 13,61	—	13,44	—	+0,17	3,151		
2592	7	Aquarii	—	—	6 35,93	21 40 35,91	—	35,66	—	+0,25	3,252		
2593	5	81 Cygni π^2	7 35,56	—	—	21 40 35,57	35,92	35,29	-0,35	+0,28	2,204		
2594	4.5	10 Cephei σ	—	4 35,94	—	21 40 36,00	36,35	35,06	-0,35	+0,94	1,727		
2595	5	78 Draconis z	—	4 58,68	—	21 40 58,81	—	57,93	—	+0,88	0,786		
2596	6	13 Pegasi	—	5 9,28	—	21 42 9,29	—	8,84	—	+0,45	2,844		
2597	5	14 Pegasi	6 24,92	—	—	21 42 24,92	—	25,01	—	-0,09	2,643		
2598	4	Gruis γ	4 44,06	—	—	21 43 44,06	—	43,25	—	+0,81	3,657		
2599	7	Aquarii	1 59,08	5 59,23	—	21 43 59,19	—	58,81	—	+0,38	3,131		
2600	5	51 Capricorni μ	6 7,73	1 7,79	9 7,83	21 44 7,79	7,90	7,26	-0,11	+0,53	3,259		
2601	6	15 Pegasi	—	5 0,19	—	21 45 0,18	—	59,51	—	+0,67	2,673		
2602	6.7	Aquarii p	—	5 23,73	—	21 45 23,73	—	23,58	—	+0,15	3,134		
2663	5.6	16 Pegasi B	—	5 25,51	—	21 45 25,52	—	25,14	—	+0,38	2,721		
2604	5	Indi δ	4 26,15	2 25,91	—	21 46 26,02	—	25,06	—	+0,96	4,151		
2605	5	17 Pegasi	2 45,27	3 45,10	2 44,88	21 48 45,09	—	44,42	—	+0,67	2,924		
2606	6.7	Aquarii	—	6 21,01	—	21 49 21,00	—	20,48	—	+0,52	3,359		
2607	6.7	Aquarii c	—	5 24,99	—	21 49 24,99	—	24,19	—	+0,80	3,147		
2608	9	12 Piscis Aust η	—	5 10,26	—	21 51 10,22	—	10,63	—	-0,41	3,467		
2609	6	18 Pegasi A	—	5 44,40	—	21 51 44,40	—	44,12	—	+0,28	2,995		
2610	6	28 Aquarii	—	5 29,07	—	21 52 29,07	—	28,81	—	+0,26	3,071		

No.	Mean N. P. D: reduced to January 1, 1832, from Observations in						Mean N.P.D. January 1, 1832.	Greenwich. Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	1831		1832		1833					Green	A. S. C.	
	No.	"	No.	"	No.	"						
2566	5 24	58,05					107 24 58,05	25 0,17	24 55,86	-2,12	+ 2,19	-15,936
2567			6 30	25,34			88 30 25,34		30 23,17		+ 2,17	15,950
2568							104 47		47 27,82			16,022
2569	5 1	4,59					114 1 4,59		1 0,43		+ 4,16	16,024
2570	6 37	39,88					109 37 39,88	37 38,76	37 34,59	+1,12	+ 5,29	16,067
2571	6 40	23,36					28 40 23,36		40 26,91		- 3,55	16,077
2572			4 28	33,69			89 28 33,69		28 27,31		+ 6,38	16,085
2573			5 23	2,67			140 23 2,67		22 57,72		+ 4,95	16,096
2574			5 4	57,15			85 4 57,15		4 46,66		+ 10,49	16,098
2575			4 9	52,68	1	9 52,00	105 9 52,54		9 42,36		+ 10,18	16,100
2576			5 30	56,19			105 30 56,19		30 49,98		+ 6,21	16,149
2577	4 47	14,07	1 47	16,20			123 47 14,49		47 11,07		+ 3,42	16,153
2578	5 53	29,36			5 53	30,87	80 53 30,11	53 29,45	53 24,80	+0,66	+ 5,31	16,206
2579			5 51	2,13			99 51 2,13		50 51,14		+ 10,99	16,211
2580	5 34	30,48			1 34	31,07	39 34 30,58	34 23,27	34 27,62	+2,31	+ 2,96	16,218
2581	2 24	55,90	3 24	57,23	2 24	58,36	73 24 57,19	25 1,17	24 56,68	-3,98	+ 0,51	16,239
2582	4 0	46,54					62 0 46,54		0 45,36		+ 1,18	16,242
2583			1 7	27,82	5 7	27,45	65 7 27,51	7 25,73	7 19,88	+1,78	+ 7,63	16,263
2584			5 2	51,88			100 2 51,88		2 44,10		+ 7,78	16,276
2585			5 8	12,23	3 8	10,57	102 8 11,61	8 11,72	8 7,34	-0,11	+ 4,27	16,285
2586	5 53	7,94					106 53 7,94	53 6,02	53 1,48	+1,92	+ 6,46	16,299
2587			2 40	17,56	3 40	14,97	121 40 16,01		40 10,55		+ 5,46	16,304
2588			5 49	16,64			67 49 16,64		49 15,56		+ 1,08	16,330
2589					5 5	11,56	88 5 11,56		5 3,36		+ 8,20	16,347
2590	6 27	40,74					19 27 40,74	27 41,50	27 44,71	-0,76	- 3,97	16,386
2591			4 10	43,21			96 10 43,21		10 40,88		+ 2,33	16,424
2592			5 30	3,07			103 30 3,07		30 6,04		- 2,97	16,442
2593	5 27	54,98			5 27	55,29	41 27 55,18		27 54,91		+ 0,22	16,444
2594	5 39	11,00					29 39 11,00	39 9,46	39 11,84	+1,54	- 0,84	16,444
2595	5 26	53,63					18 26 53,63		26 58,80		- 5,17	16,465
2596			29 27	80			73 29 27,89		29 26,52		+ 1,37	16,520
2597	1 36	21,25	36 18	77	4 36	20,37	60 36 20,04		36 14,09		+ 5,95	16,534
2598	5 9	1,10					128 9 1,10		9 57,43		+ 3,67	16,596
2599			6 46	44,31			94 46 44,31		46 40,00		+ 4,31	16,610
2600	5 29	18,69					104 29 18,69	29 18,54	29 12,61	+0,15	+ 6,08	16,617
2601			5 59	21,24			61 59 21,24		59 20 31		+ 0,93	16,660
2602			5 3	42,30			95 3 42,30		3 32,38		+ 9,92	16,679
2603			5 51	46,34			64 51 46,34		51 39,07		+ 7,27	16,681
2604	5 47	9,16					145 47 9,16		46 57,29		+ 11,87	16,727
2605			5 43	1,77			78 43 1,77		42 58,20		+ 3,57	16,840
2606			5 58	48,56			111 58 48,56		58 48,90		- 0,34	16,867
2607			5 13	1,97			96 13 1,97		12 57,42		+ 4,55	16,871
2608			5 15	20,12	2 15	17,24	119 15 19,30		15 15,26		+ 4,04	16,953
2609			5 5	1,96			84 5 1,96		4 56,05		+ 5,91	16,980
2610			4 11	58,53			90 11 58,53		11 49,78		+ 8,75	17,015

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	NAMES.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.	Green. Catal.	A. S. Catal.	Difference from		Annual Preces- sion.
			No.	No.	No.				Green	A. S.	
			1831	1832	1833				s.	s.	
2611	6	19 Pegasi	s.	5 48,96	s.	21 52 48,96	s.	48,93	s.	+0,03	+2,976
2612	5.6	20 Pegasi	—	—	6 54,57	21 52 54,58	—	54,02	—	+0,56	2,914
2613	6	29 Aquarii	—	4 14,56	5 14,47	21 53 14,51	—	15,45	—	-0,94	3,293
2614	5.6	30 Aquarii	2 26,00	4 26,12	—	21 54 26,08	—	25,83	—	+0,25	3,158
2615	5	31 Aquarii	6 37,41	1 37,33	2 37,31	21 54 37,39	37,37	36,94	+0,02	+0,45	3,104
2616	6	Aquarii	—	6 2,81	—	21 55 2,80	—	2,76	—	+0,04	3,431
2617	5.6	21 Pegasi	—	4 4,64	—	21 55 4,64	—	4,17	—	+0,47	2,939
2618	5.6	32 Aquarii	—	3 8,98	4 8,89	21 56 8,93	—	8,40	—	+0,53	3,089
2619	3	34 Aquarii	9 9,20	8 9,26	29 9,24	21 57 9,24	9,27	9,04	-0,03	+0,20	3,082
2620	5	22 Pegasi	5 12,39	—	—	21 57 12,39	—	12,25	—	+0,14	3,018
2621	7	Aquarii	—	5 16,34	—	21 57 16,34	—	12,80	—	+3,54	3,142
2622	4.5	33 Aquarii	8 21,47	1 21,55	—	21 57 21,48	21,44	21,48	+0,04	0,00	3,247
2623	2	Grus	4 36,56	—	—	21 57 36,56	—	36,66	—	-0,10	3,818
2624	6	23 Pegasi	—	4 58,55	—	21 57 58,57	—	57,94	—	+0,63	2,705
2625	5	17 Cephei	3 56,20	3 55,80	—	21 58 56,03	—	54,54	—	+1,49	1,699
2626	4	24 Pegasi	3 11,75	1 11,46	—	21 59 11,68	11,79	11,69	-0,11	-0,01	2,761
2627	5.6	35 Aquarii	—	5 45,75	—	21 59 45,74	45,63	45,37	+0,11	+0,37	3,303
2628	6	25 Pegasi	—	5 57,16	—	21 59 57,17	—	56,83	—	+0,34	2,813
2629	7	36 Aquarii	—	3 33,64	—	22 0 33,64	—	30,40	—	+3,24	3,174
2630	6	37 Aquarii	—	—	5 33,73	22 1 33,72	—	33,44	—	+0,28	3,204
2631	7	Aquarii	—	6 36,81	—	22 1 36,81	—	36,51	—	+0,30	3,123
2632	6	38 Aquarii	—	—	6 38,36	22 1 38,35	—	38,17	—	+0,18	3,213
2633	7	Aquarii	—	—	6 42,46	22 1 42,44	—	41,91	—	+0,53	3,336
2634	4	26 Pegasi	6 43,51	—	—	22 1 43,51	43,51	43,63	0,00	-0,12	3,006
2635	6.7	Aquarii	—	—	6 47,98	22 1 47,97	—	47,56	—	+0,41	3,127
2636	5	27 Pegasi	2 47,33	2 47,35	—	22 1 47,35	—	47,71	—	-0,36	2,650
2637	4	29 Pegasi	7 31,99	3 31,90	—	22 2 31,97	31,98	32,27	-0,01	-0,30	2,653
2638	6	28 Pegasi	—	2 34,17	1 34,00	22 2 34,12	—	33,54	—	+0,58	2,828
2639	7	Aquarii	—	5 19,24	—	22 3 19,23	—	17,54*	—	+1,69	3,205
2640	7	39 Aquarii	—	5 21,94	—	22 3 21,93	—	21,68	—	+0,25	3,243
2641	6	Pegasi	—	—	—	22 3 —	—	44,34	—	—	2,891
2642	5.6	Piscis Aust. ϕ	—	—	1 17,39	22 4 17,36	—	16,67	—	+0,69	3,384
2643	7	40 Aquarii	—	—	3 27,07	22 4 27,06	—	26,85	—	+0,21	3,214
2644	6	16 Piscis Aust. λ	—	—	3 46,57	22 4 46,54	—	45,98	—	+0,56	3,419
2645	6	41 Aquarii	—	5 0,79	—	22 5 0,77	—	0,47	—	+0,30	3,327
2646	4	21 Cephei	—	4 2,02	—	22 5 2,08	2,25	1,21	-0,17	+0,87	2,064
2647	7	Aquarii	—	4 6,49	—	22 5 6,49	—	5,55*	—	+0,94	3,128
2648	5	Grus μ^1	4 28,06	—	—	22 5 28,06	—	27,07	—	+0,99	3,649
2649	5	Grus μ^2	6 18,64	—	—	22 6 18,64	—	18,41	—	+0,23	3,651
2650	5	Lacerta m	—	5 40,54	—	22 6 40,56	—	40,25	—	+0,31	2,606*
2651	3	Tucanæ	—	5 55,23	—	22 6 55,16	—	55,95	—	-0,79	4,216
2652	6	Piscis Aust.	—	3 10,58	1 10,34	22 7 10,50	—	9,95	—	+0,55	3,587
2653	6	42 Aquarii	—	—	2 47,96	22 7 47,95	—	47,00	—	+0,95	3,221
2654	7	Aquarii	—	—	6 54,43	22 7 54,43	—	54,09	—	+0,34	3,095
2655	4.5	43 Aquarii	6 57,95	—	—	22 7 57,95	57,77	57,61	+0,18	+0,34	3,163

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N.P.D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession	
	1831		1832		1833					Green. A. S. C.			
	No.	"	No.	"	No.	"				"	"		
2611	—	—	5 32	47,63	—	—	82 32	47,63	32 42,19	+	5,44	17,030	
2612	—	—	2 40	50,04	4 40	50,29	77 40	50,21	40 48,05	+	2,16	17,034	
2613	—	—	—	—	5 46	11,32	107 46	11,32	46 4,27	+	7,05	17,050	
2614	—	—	5 19	50,05	—	—	97 19	50,05	19 46,65	+	3,40	17,104	
2615	5 57	43,42	—	—	—	—	92 57	43,42	57 47,37	57 42,84	-3,95	+ 0,58	17,112
2616	—	—	5 37	50,08	—	—	117 37	50,08	37 51,43	—	1,35	17,132	
2617	—	—	3 25	19,56	2 25	18,83	79 25	19,27	25 11,44	+	7,83	17,133	
2618	—	—	5 42	55,68	—	—	91 42	55,68	42 49,92	+	5,76	17,182	
2619	33 7	55,10	14 7	56,96	16 7	57,33	91 7	56,08	7 57,54	7 54,24	-1,46	+ 1,84	17,227
2620	—	—	—	—	5 45	35,94	85 45	35,94	45 29,18	+	6,76	17,229	
2621	—	—	2 10	7,17	3 10	7,93	96 10	7,63	—	—	—	17,229	
2622	—	—	—	—	5 40	54,00	104 40	54,00	40 51,38	40 41,57	+2,62	+ 12,43	17,236
2623	5 46	9,62	—	—	—	—	137 46	9,62	46 5,13	+	4,49	17,246	
2624	—	—	3 50	55,18	2 50	56,94	61 50	55,58	50 52,93	+	2,95	17,264	
2625	5 11	20,04	—	—	—	—	26 11	20,04	11 22,10	—	2,06	17,307	
2626	5 28	21,17	3 28	21,70	—	—	65 28	21,37	28 22,35	28 15,17	-0,98	+ 6,20	17,318
2627	—	—	5 20	17,38	—	—	109 20	17,38	20 11,44	+	5,94	17,342	
2628	—	—	2 6	40,71	3 6	40,92	69 6	40,84	6 33,58	+	2,26	17,351	
2629	—	—	5 0	27,90	—	—	99 0	27,90	0 28,74	—	0,84	17,375	
2630	—	—	4 38	39,11	—	—	101 38	39,11	38 34,55	+	4,56	17,420	
2631	—	—	5 42	53,51	—	—	94 42	53,51	42 46,64	+	6,87	17,423	
2632	—	—	5 23	14,37	—	—	102 23	14,37	23 9,11	+	5,26	17,424	
2633	—	—	—	—	5 3	13,65	112 3	13,65	3 11,50	+	2,15	17,426	
2634	5 37	36,05	—	—	5 37	33,36	84 37	34,70	37 31,06	37 26,45	+3,64	+ 8,25	17,428
2635	—	—	—	—	5 5	27,21	95 5	27,21	5 20,19	+	7,02	17,431	
2636	5 38	33,01	—	—	—	—	57 38	33,01	38 44,44	—	11,43	17,432	
2637	5 38	33,34	—	—	—	—	57 38	33,34	38 35,86	38 31,95	-2,52	+ 1,39	17,463
2638	—	—	—	—	4 50	43,52	69 50	43,52	50 37,14	+	6,58	17,464	
2639	—	—	4 53	27,15	1 53	29,56	101 53	27,15	—	—	—	17,495	
2640	—	—	—	—	5 1	8,87	105 1	8,87	0 58,36	+	10,51	17,498	
2641	—	—	4 47	5,42	—	—	74 47	5,42	47 1,72	+	3,70	17,514	
2642	—	—	2 0	38,05	5 0	34,03	116 0	38,05	0 32,87	+	5,18	17,537	
2643	—	—	—	—	5 45	8,64	102 45	8,64	45 2,36	+	6,28	17,544	
2644	—	—	5 35	40,60	—	—	118 35	40,60	34 57,95	—	—	17,557	
2645	—	—	3 54	22,19	2 54	21,59	111 54	21,95	54 19,59	+	2,36	17,568	
2646	5 37	28,58	1 37	29,07	2 37	29,44	82 37	28,86	37 30,52	37 32,61	-1,66	- 3,75	17,570
2647	—	—	—	—	4 16	50,05	95 16	50,05	—	—	—	17,571	
2648	5 10	45,36	—	—	—	—	132 10	45,36	10 41,07	+	4,29	17,586	
2649	5 27	31,29	—	—	—	—	132 27	31,29	27 28,72	+	2,57	17,622	
2650	5 6	56,93	—	—	—	—	51 6	56,93	6 54,94	+	1,99	17,838*	
2651	5 5	30,34	—	—	—	—	151 5	30,34	5 22,93	+	7,41	17,647	
2652	—	—	5 43	51,46	—	—	116 43	51,46	43 47,37	+	4,09	17,658	
2653	—	—	5 39	52,49	—	—	103 39	52,49	39 52,92	—	0,43	17,683	
2654	—	—	5 25	47,38	—	—	92 25	47,38	25 46,39	+	0,99	17,688	
2655	5 36	55,13	—	—	—	—	98 36	55,13	36 59,48	36 56,93	-4,35	- 1,80	17,691

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832.			Green Catal.	A. S. Catal.	Difference from		Annual Preces- sion.	
			No. 1831		No. 1832	h. m. s.		h. m. s.			Green.	A. S.		
			s.	s.	s.	s.	s.	s.			s.	s.		
2656	6	Aquarii	—	—	3 0,12	22 8 0,11	—	—	59,91	—	+0,20	+3,177		
2657	6.7	44 Aquarii	—	—	6 20,02	22 8 20,01	—	—	20,22	—	-0,21	3,136		
2658	5	1 Lacertæ	5 39,47	—	—	22 8 39,47	—	—	38,55	—	+0,92	2,590		
2659	4.5	23 Cephei	2 51,67	3 51,61	1 51,37	22 8 51,63	51,54	51,04	51,04	+0,09	+0,59	2,137		
2660	6	45 Aquarii	—	6 59,58	—	22 9 59,56	—	—	59,07	—	+0,49	3,224		
2661	6	46 Aquarii	—	5 21,37	1 21,36	22 11 21,37	21,35	21,61	21,61	+0,02	-0,24	3,161		
2662	5	30 Pegasi	6 0,43	—	—	22 12 0,43	—	—	0,05	—	+0,38	3,016		
2663	5	47 Aquarii	—	6 20,13	—	22 12 20,12	—	—	19,65	—	+0,47	3,318		
2664	4	48 Aquarii	—	6 58,71	—	22 12 58,71	—	—	58,67	—	+0,04	3,092		
2665	4.5	31 Pegasi	7 15,17	—	—	22 13 15,17	15,15	15,03	15,03	+0,02	+0,14	2,947		
2666	5.6	32 Pegasi	—	5 34,41	—	22 13 34,42	—	—	33,94	—	+0,48	2,757		
2667	5	2 Lacertæ	4 5,92	—	6 5,67	22 14 5,31	—	—	5,32	—	+0,49	2,458		
2668	6	49 Aquarii	—	4 8,49	—	22 14 8,48	—	—	8,36	—	+0,12	3,354		
2669	7	Aquarii	4 43,13	1 43,57	—	22 14 43,22	—	—	42,79	—	+0,43	3,152		
2670	5	Tucanæ	—	—	4 17,56	22 15 17,38	—	—	16,51	—	+0,87	4,383		
2671	6	51 Aquarii	—	5 21,67	—	22 15 21,67	—	—	20,95	—	+0,72	3,127		
2672	6	50 Aquarii	—	1 26,70	2 26,61	22 15 26,62	—	—	26,69	—	-0,07	3,219		
2673	7	Aquarii	—	—	5 55,73	22 15 55,73	—	—	55,31	—	+0,42	3,089		
2674	5	52 Aquarii	7 41,90	—	5 41,84	22 16 41,88	41,80	41,55	41,55	+0,08	+0,33	3,063		
2675	6	Piscis Aust.	—	—	4 52,54	22 16 52,51	—	—	52,63	—	-0,12	3,334		
2676	4	3 Lacertæ	1 58,03	5 57,65	3 57,55	22 16 57,61	58,10	57,16	57,16	-0,49	+0,45	2,341		
2677	6.7	53 Aquarii	—	6 26,42	—	22 17 26,41	—	—	26,66	—	-0,25	3,252		
2678	6.7	53 Aquarii	—	3 27,10	—	22 17 27,09	—	—	27,45	—	-0,36	3,252		
2679	5	4 Lacertæ	4 43,08	—	—	22 17 43,08	—	—	42,00	—	+1,08	2,413		
2680	5.6	34 Pegasi	—	4 4,20	—	22 18 4,20	—	—	3,86	—	+0,34	3,033		
2681	4	Gruis	3 11,32	—	—	22 19 11,32	—	—	11,64	—	+0,18	3,625		
2682	5.6	35 Pegasi	—	5 21,47	—	22 19 21,47	—	—	20,69	—	+0,78	3,030		
2683	5	Gruis	—	4 41,53	1 41,55	22 19 41,50	—	—	40,95	—	+0,55	3,627		
2684	4	55 Aquarii	5 10,80	—	1 10,83	22 20 10,83	10,82	10,53	10,53	+0,01	+0,30	3,077		
2685	6.7	Aquarii	—	4 2,31	—	22 21 2,31	—	—	1,29	—	+1,02	3,205		
2686	6	56 Aquarii	—	—	8 16,64	22 21 16,62	—	—	16,37	—	+0,25	3,222		
2687	6	37 Pegasi	—	—	5 28,54	22 21 28,53	—	—	28,24	—	+0,29	3,033		
2688	5	57 Aquarii	—	5 45,14	3 54,05	22 21 45,10	45,09	45,10	45,10	+0,01	0,00	3,182		
2689	4	17 Piscis Aust.	4 56,21	—	2 56,30	22 21 56,23	56,30	56,24	56,24	-0,07	-0,01	3,431		
2690	6	58 Aquarii	—	5 46,71	—	22 22 46,71	—	—	46,49	—	+0,22	3,183		
2691	4.5	27 Cephei	6 56,74	—	—	22 22 56,74	56,88	55,86	55,86	-0,14	+0,88	2,204		
2692	4	7 Lacertæ	3 23,09	—	2 22,31	22 24 23,02	23,17	22,88	22,88	-0,15	+0,14	2,436		
2693	6	39 Pegasi	—	4 28,91	—	22 24 28,92	—	—	28,38	—	+0,54	2,878		
2694	7	Aquarii	—	5 15,46	—	22 25 15,46	—	—	15,19	—	+0,27	3,167		
2695	6.7	60 Aquarii	—	4 23,40	2 23,33	22 25 23,38	—	—	22,95	—	+0,43	3,091		
2696	5	59 Aquarii	5 29,60	—	—	22 25 29,60	29,64	29,51	29,51	-0,04	+0,09	3,280		
2697	7	Aquarii	—	5 0,37	5 0,25	22 26 0,31	—	—	0,07	—	+0,24	3,071		
2698	4	62 Aquarii	6 43,46	—	—	22 26 43,46	43,42	43,13	43,13	+0,04	+0,33	3,077		
2699	7	61 Aquarii	—	5 45,72	—	22 26 45,71	—	—	44,75	—	+0,96	3,243		
2700	5	Octantis	—	—	—	22 28 —	—	—	19,87	—	—	6,904		

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
	" "	" "	" "				" "	" "	
2656	—	5 52 29,70	—	99 52 29,70	—	52 24,66	+	5,04	—17,692
2657	—	—	5 13 23,12	96 13 23,12	—	13 14,67	+	8,45	17,706
2658	6 5 6,72	—	—	53 5 6,72	—	5 2,23	+	4,49	17,719
2659	5 47 28,35	1 47 29,91	5 47 28,13	33 47 28,39	47 32,73	47 33,61	—	4,34	—5,22
2660	—	5 8 31,06	—	104 8 31,06	—	8 28,78	+	2,28	17,773
2661	—	5 39 37,85	—	98 39 37,85	39 21,52	39 33,27	+16,33	+	4,58
2662	5 3 10,27	—	—	85 3 10,27	—	3 3,19	+	7,08	17,854
2663	5 26 9,88	—	—	112 26 9,88	—	24 7,24	+	122,64	17,867
2664	4 13 52,85	1 13 51,47	—	92 13 52,57	13 51,26	13 44,87	+	1,31	+ 7,70
2665	—	5 38 14,78	4 38 15,14	78 38 14,94	38 18,26	38 12,53	—	3,32	+ 2,36
2666	—	5 30 45,01	—	62 30 45,01	—	30 44,65	+	0,36	17,916
2667	5 18 25,00	—	—	44 18 25,00	—	18 26,46	—	1,46	17,937
2668	—	5 36 31,96	—	115 36 31,96	—	36 23,62	+	8,34	17,938
2669	—	5 2 24,94	—	98 2 24,94	—	2 26,15	—	1,21	17,960
2670	4 49 0,37	—	2 49 0,85	155 49 0,53	—	48 46,22	+	14,31	17,981
2671	—	—	5 41 4,08	95 41 4,08	—	40 58,38	+	5,70	17,985
2672	—	4 22 42,05	1 22 42,58	104 22 42,15	—	22 35,85	+	6,30	17,989
2673	—	—	5 2 12,72	92 2 12,72	—	2 11,33	+	1,39	18,007
2674	6 28 21,88	—	—	89 28 21,88	28 21,12	28 17,18	+ 0,76	+	4,70
2675	—	5 31 59,62	—	114 31 59,62	—	31 56,52	+	3,10	18,043
2676	5 36 35,46	—	5 36 36,44	38 36 35,95	36 37,58	36 38,64	—	1,63	— 2,69
2677	—	4 35 30,15	3 35 32,41	107 35 31,12	—	35 32,82	—	1,70	18,065
2678	—	—	—	107	—	35 36,13	—	—	18,066
2679	5 22 21,57	—	—	41 22 21,57	—	22 24,62	—	3,05	18,076
2680	—	4 27 36,75	—	86 27 36,75	—	27 33,68	+	3,07	18,089
2681	4 21 0,79	—	2 21 0,19	134 21 0,59	—	21 0,71	—	0,12	18,130
2682	—	5 8 38,41	—	86 8 38,41	—	8 27,65	+	10,76	18,137
2683	3 36 22,26	2 36 22,57	—	134 36 22,38	—	36 14,72	+	7,66	18,149
2684	5 52 37,90	—	5 52 35,45	90 52 36,62	52 38,64	52 34,88	—	2,02	+ 1,74
2685	—	5 46 17,66	—	103 46 17,66	—	—	—	—	18,198
2686	—	5 26 31,86	—	105 26 31,86	—	26 24,50	+	7,36	18,208
2687	—	3 25 8,96	—	86 25 8,96	—	25 2,60	+	0,36	18,215
2688	5 32 3,71	—	—	101 32 3,71	32 5,90	31 55,22	—	2,19	+ 8,49
2689	5 12 14,90	—	—	123 12 14,90	12 12,70	12 14,18	+	2,20	+ 0,72
2690	—	5 45 47,63	—	101 45 47,63	—	45 42,01	+	5,62	18,262
2691	5 26 32,07	—	5 26 33,89	32 26 32,98	26 31,32	26 34,42	+ 1,66	—	1,44
2692	5 34 46,46	1 34 46,38	5 34 46,19	40 34 46,33	34 45,50	34 45,32	+ 0,83	+	1,01
2693	1 37 59,83	4 38 0,34	—	70 38 0,24	—	37 56,36	+	3,88	18,323
2694	1 28 22,61	4 28 21,80	—	100 28 21,96	—	28 19,79	+	2,17	18,350
2695	—	4 26 9,63	—	92 26 9,63	—	26 2,75	+	6,88	18,355
2696	5 33 55,19	—	—	111 33 55,19	33 45,04	33 52,29	+10,15	+	2,90
2697	—	—	5 16 2,81	90 16 2,81	—	15 54,11	+	8,70	18,377
2698	5 58 52,44	—	—	90 58 52,44	58 52,04	58 44,02	+ 0,40	+	8,42
2699	—	—	5 19 27,53	108 19 27,53	—	19 22,72	+	4,81	18,402
2700	—	—	—	173 Invisible	—	14 50,51	—	—	18,463

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion	
			No. 1831		No. 1832				Green.			A. S.
			s.	s.	s.				s.	s.		s.
2701	6	63 Aquarii α	—	5 3,33	6 3,29	22 29 3,30	3,26	3,27	+0,04	+0,03	+3,114	
2702	6.7	64 Aquarii	—	5 25,52	—	22 30 25,52	—	25,53	—	-0,01	3,166	
2703	6	40 Pegasi Q^1	—	4 45,40	—	22 30 45,41	—	44,74	—	+0,67	2,897	
2704	7	Aquarii	—	5 16,06	—	22 31 16,06	—	15,27	—	+0,79	3,159	
2705	4	18 Piscis Aust	6 21,02	—	4 20,94	22 31 20,98	20,98	20,88	0,00	+0,10	3,336	
2706	6	41 Pegasi Q^2	—	5 39,17	—	22 31 39,18	—	38,67	—	+0,51	2,897	
2707	5	31 Cephei	4 36,61	—	—	22 31 36,31	—	36,74	—	-0,43	1,447	
2708	3	Gruis β	6 35,88	—	—	22 32 35,88	—	35,85	—	+0,03	3,617	
2709	5	30 Cephei	4 42,82	—	—	22 32 42,82	—	42,45	—	+0,37	2,105	
2710	3	42 Pegasi ζ	6 5,16	—	5 5,27	22 33 5,21	5,23	4,90	-0,02	+0,31	2,981	
2711	5	43 Pegasi	—	—	—	22 33 —	—	52,35	—	—	2,802	
2712	7	65 Aquarii ι	1 10,68	4 10,72	1 10,82	22 34 10,73	—	10,43	—	+0,30	3,163	
2713	7	Aquarii	—	3 15,22	3 15,02	22 34 15,21	—	14,44	—	+0,77	3,147	
2714	6	67 Aquarii N	—	—	6 27,71	22 34 27,79	—	27,65	—	+0,05	3,135	
2715	6.7	66 Aquarii g^1	—	—	6 32,30	22 34 32,28	—	32,09	—	+0,19	3,243	
2716	3	44 Pegasi η	—	5 8,32	2 8,30	22 35 8,34	8,21	7,87	+0,13	+0,47	2,796	
2717	5	Gruis η	—	5 16,30	—	22 35 16,25	—	15,79	—	+0,46	3,743	
2718	6	20 Piscis Aust χ^1	—	5 18,83	—	22 36 18,82	—	18,54	—	+0,28	3,302	
2719	6	45 Pegasi	—	5 18,19	—	22 37 18,20	—	17,65	—	+0,55	2,910	
2720	5	46 Pegasi ξ	6 18,33	—	—	22 38 18,33	—	17,96	—	+0,37	2,975	
2721	4	Gruis ϵ	3 22,04	—	—	22 38 22,04	—	21,02	—	+1,02	3,670	
2722	4.5	47 Pegasi λ	6 26,81	—	2 26,80	22 38 26,81	26,83	27,23	-0,02	-0,42	2,873	
2723	6	68 Aquarii g^2	—	5 31,23	—	22 38 31,23	—	30,92	—	+0,36	3,242	
2724	6	69 Aquarii τ^1	1 46,99	6 47,38	3 47,19	22 38 47,28	48,09	46,85	-0,81	+0,43	3,192	
2725	6	70 Aquarii	—	7 39,56	2 39,65	22 39 39,57	—	39,97	—	-0,40	3,161	
2726	5.6	71 Aquarii τ^2	—	5 41,54	6 41,52	22 40 41,52	41,50	41,48	+0,02	+0,04	3,185	
2727	4	48 Pegasi μ	5 54,48	—	4 54,42	22 41 54,16	54,16	53,40	0,00	+0,76	2,872	
2728	5	22 Piscis Aust	5 10,22	—	—	22 43 10,22	—	10,11	—	+0,11	3,362	
2729	4	32 Cephei ν	1 43,48	—	—	22 43 43,48	43,16	42,27	+0,32	+1,21	2,118	
2730	4	73 Aquarii λ	7 50,88	—	3 50,82	22 43 50,85	50,78	50,40	+0,07	+0,45	3,133	
2731	5.6	49 Pegasi σ	—	5 53,71	—	22 43 53,71	—	53,03	—	+0,68	2,999	
2732	6	74 Aquarii K	—	5 37,90	2 37,61	22 44 37,82	—	37,09	—	+0,73	3,164	
2733	6	Pegasi	—	5 46,46	—	22 44 46,48	—	46,41	—	+0,07	2,945	
2734	5	Cephei e	1 50,12	—	—	22 44 50,12	—	—	—	—	2,297	
2735	3	76 Aquarii δ	6 43,73	—	1 43,72	22 45 43,73	43,75	43,53	-0,02	+0,20	3,196	
2736	6	78 Aquarii	—	5 49,33	—	22 45 49,33	—	48,93	—	+0,40	3,199	
2737	6	77 Aquarii	—	5 51,83	—	22 45 51,82	—	51,21	—	+0,61	3,199	
2738	6	1 Piscium	—	1 23,83	5 23,77	22 46 23,78	—	23,76	—	+0,02	3,067	
2739	7	Aquarii M^1	—	—	6 28,09	22 46 28,09	—	27,79	—	+0,30	3,112	
2740	5.6	50 Pegasi ρ	3 46,54	—	6 46,48	22 46 46,50	—	45,59	—	+0,91	3,010	
2741	1	24 Piscis Aust α	23 21,00	6 21,14	20 21,10	22 48 21,02	20,99	20,96	+0,03	+0,06	3,311	
2742	.6	51 Pegasi	—	5 13,14	—	22 49 13,15	—	12,28	—	+0,87	2,921	
2743	6	52 Pegasi	—	6 47,78	—	22 50 47,78	—	47,59	—	+0,19	2,992	
2744	6.7	2 Piscium α^1	—	6 51,09	—	22 50 51,09	—	50,60	—	+0,43	3,068	
2745	.5	Gruis ζ	5 55,34	—	—	22 50 55,34	—	54,78	—	+0,56	3,608	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in.						Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.
	1831		1832		1833					Green.	A. S. C.	
	No.	" "	No.	" "	No.	" "				" "	" "	
2701	2	5 36,26	5	5 35,69			95 5 35,85	5 32,65	5 25,23	+ 3,20	+ 10,62	18,482
2702			5	53 57,76			100 53 57,76		53 49,85		+ 7,91	18,528
2703			5	20 41,27			71 20 41,27		20 36,77		+ 4,50	18,539
2704			6	14 6,39			100 14 6,39		14 4,08		+ 2,31	18,555
2705	5	54 56,83					117 54 56,83	55 0,35	54 56,96	- 3,52	- 0,13	18,558
2706			4	11 27,65			71 11 27,65		11 20,07		+ 7,58	18,568
2707	5	13 38,47					17 13 38,47		13 38,49		- 0,02	18,569
2708	5	45 34,54					187 45 34,54		45 28,32		+ 6,22	18,599
2709	5	17 13,23					27 17 13,28		17 17,53		- 4,25	18,604
2710	5	2 41,42			5	2 41,05	80 2 41,42	2 35,03	2 29,85	+ 6,39	+ 11,57	18,615
2711	5	34 1,50					61 34 1,50		33 58,29		+ 3,21	18,641
2712			5	58 48,09			100 58 48,09		58 42,48		+ 5,61	18,650
2713			6	11 17,80			99 11 17,80		11 16,95		+ 0,85	18,652
2714			4	50 21,31	1	50 21,68	97 50 21,38		50 16,49		+ 4,89	18,659
2715			5	42 26,04	1	42 26,04	109 42 26,04		42 23,32		+ 2,72	18,662
2716	3	39 19,48	2	39 20,34	5	39 19,50	60 39 19,66	39 18,94	39 15,40	+ 0,72	+ 4,26	18,681
2717			3	22 53,19	3	22 51,33	144 22 52,26		22 48,55		+ 3,71	18,684
2718			5	7 2,68			116 7 2,68		6 58,22		+ 4,46	18,718
2719			5	30 59,96			71 30 59,96		30 51,61		+ 8,35	18,748
2720	5	41 8,06					78 41 8,06		41 2,79		+ 5,27	18,779
2721	5	11 52,35					142 11 52,35		11 51,82		+ 0,53	18,780
2722	3	18 56,97	3	18 56,30	5	18 58,00	67 18 57,25	18 58,68	18 49,83	- 1,43	+ 7,42	18,784
2723			5	29 16,27			110 29 16,27		29 11,94		+ 4,33	18,786
2724			5	56 26,19			104 56 26,19	56 22,90	56 20,01	+ 3,29	+ 6,18	18,794
2725			5	26 23,94			101 26 23,94		26 18,77		+ 5,17	18,821
2726			4	28 38,37			104 28 38,37	28 37,16	28 32,37	+ 1,21	+ 6,00	18,851
2727	5	16 59,25	2	16 58,83	5	16 59,28	66 16 59,13	17 0,47	16 59,80	- 1,34	- 0,67	18,887
2728	3	45 47,08					123 45 47,08		45 43,23		+ 3,85	18,924
2729	5	40 56,19					24 40 56,19	40 54,09	40 55,83	+ 2,10	+ 0,36	18,940
2730	5	28 13,56					98 28 13,56	28 17,48	28 14,46	- 3,92	- 0,90	18,943
2731	3	3 21,90	3	3 22,17			81 3 22,03		3 17,51		+ 4,52	18,944
2732			5	30 24,04			102 30 24,04		30 25,16		- 1,12	18,965
2733			5	2 52,60			74 2 52,60		2 52,35		+ 0,25	18,970
2734	4	11 41,56			1	11 41,83	29 11 41,61		11 47,39		- 6,28	18,971
2735	7	42 43,17					106 42 43,17	42 43,12	42 37,40	+ 0,05	+ 5,77	18,996
2736			5	5 40,48			98 5 40,48		5 40,40		+ 0,08	18,999
2737			5	9 39,62			107 9 39,62		9 33,23		+ 6,39	19,000
2738	1	49 45,40	4	49 45,66			89 49 45,61		49 39,96		+ 5,65	19,015
2739			2	52 51,13	4	52 51,43	95 52 51,33		52 50,27		+ 1,06	19,017
2740	1	4 41,56	5	4 41,86			82 4 41,81		4 37,35		+ 4,46	19,025
2741	48	30 38,18	20	30 39,07	32	30 40,27	120 30 39,01	30 36,81	30 35,74	+ 2,20	+ 3,27	19,068
2742					6	7 51,35	70 7 51,35		7 43,82		+ 7,53	19,091
2743			5	10 1,59			79 10 1,59		9 58,80		+ 2,79	19,133
2744			6	56 0,11			89 56 0,11		55 54,05		+ 6,06	19,134
2745	5	39 5,53					143 39 5,53		39 6,95		- 1,42	19,136

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832		Green ^b Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	h. m.	s.			Green.	A. S.	
			s.	s.	s.					s.	s.	
2746	6	3 Piscium α^2	4 1,05	—	—	22 52 1,05	—	0,94	—	+0,11	+3,073	
2747	7	" Piscium	—	6 9,48	—	22 52 9,48	—	8,96	—	+0,52	3,054	
2748	6	81 Aquarii	1 39,63	6 39,74	—	22 52 39,72	—	39,35	—	+0,37	3,122	
2749	7	" Piscium	—	6 10,41	—	22 53 10,41	—	—	—	—	3,050	
2750	6	82 Aquarii	—	5 49,18	—	22 53 49,18	—	49,06	—	+0,12	3,118	
2751	4	1 Andromedæ α	6 12,48	—	4 12,41	22 54 12,47	12,47	11,78	0,00	+0,69	2,734	
2752	5	4 Piscium β	6 19,84	—	—	22 55 19,84	19,71	19,35	+0,13	+0,49	3,049	
2753	2	53 Pegasi β	2 38,26	—	—	22 55 38,26	38,36	38,13	-0,10	+0,13	2,878	
2754	6	83 Aquarii δ^1	—	5 24,16	3 23,95	22 56 24,08	23,89	23,74	+0,19	+0,34	3,124	
2755	2	54 Pegasi α	17 23,89	18 23,93	30 23,94	22 56 23,93	23,94	23,66	-0,01	+0,27	2,975	
2756	7	85 Aquarii δ^3	—	5 7,95	—	22 57 7,94	7,53	8,05	+0,41	-0,11	3,124	
2757	5	Cephei f	2 10,43	—	—	22 57 10,43	—	—	—	—	2,243	
2758	5	Gruis δ	4 23,47	—	—	22 57 23,47	—	23,37	—	+0,10	3,422	
2759	5.6	86 Aquarii c^1	—	5 38,70	—	22 57 38,69	—	38,56	—	+0,13	3,233	
2760	5	55 Pegasi l	2 32,71	—	1 32,80	22 58 32,74	32,54	32,61	+0,20	+0,13	3,015	
2761	4.5	56 Pegasi h	6 56,27	—	—	22 58 56,27	56,39	56,24	-0,12	+0,03	2,907	
2762	6	Aquarii	—	5 15,09	—	22 59 15,08	—	14,59	—	-0,49	3,268	
2763	6	5 Piscium A	—	4 4,83	2 4,81	23 0 4,83	—	4,24	—	+0,59	3,061	
2764	4.5	88 Aquarii c^2	6 28,92	—	2 28,73	23 0 28,87	28,78	28,78	+0,09	+0,09	3,208	
2765	5	Gruis i	3 49,24	—	—	23 0 49,24	—	45,63	—	+3,61	3,424	
2766	5	89 Aquarii c^3	—	1 55,92	—	23 0 55,91	—	55,83	—	+0,08	3,216	
2767	5.6	57 Pegasi m	—	5 2,89	—	23 1 2,89	—	2,86	—	+0,03	3,022	
2768	5	33 Cephei π	2 34,33	—	—	23 2 34,33	—	34,11	—	+0,22	1,875	
2769	5.6	59 Pegasi p	—	4 15,51	—	23 3 15,51	—	15,53	—	-0,02	3,023	
2770	6	60 Pegasi	1 40,78	5 40,44	—	23 3 40,51	—	40,41	—	+0,10	2,910	
2771	5	7 Androm α	6 52,39	—	—	23 4 52,39	—	51,87	—	+0,52	2,709	
2772	5	90 Aquarii ϕ	6 37,20	4 37,42	2 37,15	23 5 37,26	37,29	36,94	-0,03	+0,32	3,106	
2773	5.6	91 Aquarii ψ^1	—	4 5,53	6 5,28	23 7 5,17	5,25	5,25	-0,08	-0,08	3,122	
2774	4	Tucanæ γ	4 34,51	—	—	23 7 34,51	—	33,63	—	+0,88	3,577	
2775	6	61 Pegasi	—	5 34,76	—	23 7 34,77	—	34,23	—	+0,54	2,911	
2776	5.6	92 Aquarii α	—	5 8,37	5 8,27	23 8 8,31	8,04	8,43	+0,27	-0,12	3,114	
2777	4.5	6 Piscium γ	5 27,37	7 27,53	—	23 8 27,46	27,46	27,26	0,00	+0,20	3,108*	
2778	5	93 Aquarii ψ^2	1 10,22	6 10,27	—	23 9 10,26	10,27	10,35	-0,01	-0,09	3,121	
2779	5	Ap. Sculp γ	6 44,40	—	—	23 9 44,40	—	44,18	—	+0,22	3,261	
2780	5	8 Androm	3 58,59	—	—	23 9 58,59	—	57,98	—	+0,61	2,745	
2781	5	95 Aquarii ψ^3	5 13,19	2 13,10	4 13,10	23 10 13,14	13,22	12,93	-0,08	+0,21	3,122	
2782	6	94 Aquarii Z	—	5 16,34	—	23 10 16,33	—	16,09	—	+0,24	3,142	
2783	6	96 Aquarii	—	5 41,35	5 41,31	23 10 41,32	41,28	41,06	+0,04	+0,26	3,098	
2784	6	7 Piscium b	—	5 47,13	—	23 11 47,13	—	46,77	—	+0,36	3,046	
2785	7	Aquarii	—	5 1,16	—	23 12 1,16	—	1,12	—	+0,04	3,101	
2786	6	Aquarii T	—	5 17,90	—	23 12 17,89	—	17,75	—	+0,14	3,215	
2787	5	62 Pegasi τ	5 19,99	—	—	23 12 19,99	—	19,45	—	+1,54	2,952	
2788	6	97 Aquarii	1 50,45	4 50,64	—	23 13 50,60	—	49,90	—	+0,70	3,145	
2789	5	98 Aquarii b^1	6 8,45	—	—	23 14 8,45	—	7,86	—	+0,59	3,170	
2790	6	65 Pegasi	—	7 19,23	—	23 14 19,24	—	19,19	—	+0,05	2,972	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.	
	"	"	"						
2746	4 42 53,08	3 42 54,40	—	90 42 53,65	—	42 47,62	—	—	—
2747	—	4 53 0,39	—	87 53 0,89	—	52 58,17	+	6,03	19,165
2748	—	4 57 36,86	2 57 37,86	97 57 37,19	—	57 31,73	+	2,22	19,168
2749	—	—	6 22 3,18	87 22 3,18	—	22 2,96	+	5,46	19,181
2750	—	5 28 24,22	—	97 28 24,22	—	28 19,12	+	0,22	19,194
2751	—	—	—	—	—	—	+	5,10	19,210
2751	5 34 28,16	2 34 25,33	5 34 27,41	48 34 27,37	34 59,68	34 31,53	- 3,26	- 4,16	19,220
2752	5 4 59,08	—	4 4 58,49	87 4 58,81	4 58,42	4 52,42	+ 0,39	+ 6,99	19,247
2753	5 49 35,54	—	—	62 49 35,54	49 37,42	49 32,38	- 1,88	+ 3,16	19,255
2754	—	4 35 51,46	3 35 53,73	98 35 51,46	35 56,67	35 49,01	- 5,21	+ 2,45	19,273
2755	37 41 50,14	25 41 50,06	26 41 50,20	75 41 50,14	41 48,33	41 44,79	+ 1,81	+ 5,35	19,273
2756	—	4 50 27,84	—	98 50 27,84	50 30,96	50 21,41	- 3,12	+ 6,43	19,291
2757	—	—	5 41 40,41	23 41 40,41	—	41 45,37	—	4,96	19,291
2758	4 25 30,79	—	—	134 25 30,79	—	25 28,48	+	2,31	19,297
2759	—	5 38 57,52	—	114 38 57,52	—	38 51,93	+	5,59	19,303
2760	5 29 47,50	—	5 29 46,32	81 29 46,32	29 47,04	29 46,04	- 1,03	+ 6,87	19,324
2761	—	5 26 13 82	5 26 14,82	65 26 14,32	26 12,53	26 6,90	+ 1,79	+ 7,43	19,333
2762	—	4 43 47,70	—	119 43 47,70	—	43 44,35	+	3,35	19,340
2763	—	3 47 1,93	—	88 47 1,93	—	47 1,90	+	0,03	19,350
2764	5 4 55,87	—	—	112 4 55,87	4 56,85	4 50,27	- 0,99	+ 5,60	19,368
2765	5 9 15,14	—	—	136 9 15,14	—	9 17,03	-	1,89	19,374
2766	4 21 56,15	1 21 55,90	—	113 21 56,10	—	21 54,07	+	2,03	19,378
2767	1 13 55,00	4 13 55,78	—	82 13 55,63	—	13 48,71	+	6,92	19,381
2768	5 31 18,69	—	—	15 31 18,69	—	31 15,35	-	1,66	19,415
2769	4 11 27,01	—	—	82 11 27,01	—	11 21,56	+	5,46	19,429
2770	—	4 3 28,04	1 3 29,50	64 3 28,33	—	3 24,88	+	3,45	19,438
2771	4 30 36,25	2 30 35,22	5 30 37,87	41 30 36,80	—	30 28,11	-	1,31	19,464
2772	5 57 10,83	—	—	96 57 10,83	57 11,63	57 8,39	+ 0,80	+ 2,44	19,479
2773	—	4 0 8,97	—	100 0 8,97	0 7,36	0 0,65	+ 1,61	+ 8,32	19,509
2774	5 9 18,35	—	—	149 9 18,25	—	9 28,90	-	5,55	19,518
2775	—	5 39 57,89	—	62 39 57,89	—	39 56,17	+	1,72	19,518
2776	—	4 38 25,14	—	98 38 25,14	38 29,57	38 20,81	- 4,43	+ 4,33	19,530
2777	5 38 1,85	6 38 2,10	5 38 2,41	87 38 2,12	38 3,21	38 1,63	- 1,09	+ 0,49	19,536
2778	4 5 57,00	—	—	100 5 57,00	5 53,87	5 48,46	+ 3,33	+ 8,54	19,550
2779	4 26 42,69	—	1 26 42,90	123 26 42,73	—	26 43,30	-	0,57	19,560
2780	3 54 5,44	2 54 5,46	—	41 54 5,45	—	54 7,74	-	2,29	19,565
2781	—	5 31 41,96	—	100 31 41,96	31 40,86	31 34,18	+ 1,10	+ 7,78	19,570
2782	—	5 22 18,99	—	104 22 18,99	—	22 10,34	+	8,65	19,571
2783	—	5 2 26,18	—	96 2 26,18	2 27,03	2 25,45	- 0,85	+ 0,73	19,578
2784	3 2 4,56	2 32 6,07	—	85 32 5,16	—	32 2,76	+	2,40	19,599
2785	—	5 49 25,50	—	96 49 25,50	—	49 20,08	+	5,42	19,603
2786	—	5 54 15,09	—	117 54 15,09	—	54 13,94	+	1,15	19,608
2787	5 10 39,51	—	—	67 10 39,51	—	10 36,43	+	3,08	19,609
2788	—	4 57 38 01	2 57 36,38	105 57 37,47	—	57 34,37	+	3,10	19,635
2789	5 0 59,15	—	—	111 0 59,15	—	0 57,88	+	1,27	19,641
2790	—	5 5 29 31	—	70 5 29,31	—	5 21,72	+	7,50	19,644

Comparison of the Observed Places of the Principal Fixed Stars

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. Green ^b January 1, 1832		A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833	1832			Green.	A. S.	
			s.	s.	s.	h. m. s.	s.		s.	s.	
2791	6	66 Pegasi	1 36 51	5 36,61	—	23 14 36,59	—	36,08	+0,51	+3,015	
2792	6.7	Piscium	—	5 54,93	—	23 14 54,93	—	54,51	+0,42	3,071	
2793	6	Aquarii	—	5 12,72	—	23 15 12,71	—	12,28	+0,43	3,177	
2794	5	68 Pegasi <i>v</i>	6 0,36	—	—	23 17 0,36	—	59,86	+0,50	2,965	
2795	5	99 Aquarii <i>b</i> ²	5 12,80	—	—	23 17 12,80	—	12,20	+0,60	3,166	
2796	5	4 Cassiopeæ <i>d</i>	1 24,25	—	1 24,40	23 17 24,39	24,35	23,67	+0,04	+0,72	2,615
2797	6	Aquarii	—	6 43,74	—	23 17 43,73	—	43,32	+0,41	3,171	
2798	5.6	8 Piscium <i>x</i> ¹	—	6 19,45	7 19,31	23 18 19,37	19,41	18,40	-0,04	+0,97	3,067
2799	6	9 Piscium <i>x</i> ²	—	5 38,54	—	23 18 38,54	—	38,27	+0,27	3,067	
2800	6	69 Pegasi	—	5 20,23	—	23 19 20,24	—	19,41	+0,83	2,962	
2801	5	10 Piscium <i>θ</i>	6 26,99	—	—	23 19 26,99	—	26,35	+0,64	3,046	
2802	5	Cephei <i>ε</i>	5 12,99	—	—	23 20 12,99	—	—	—	2,452	
2803	5	70 Pegasi <i>q</i>	1 39,78	—	—	23 20 39,78	—	39,40	+0,38	3,020	
2804	6.7	11 Piscium <i>w</i> ¹	—	6 49,67	—	23 20 49,67	—	49,00	+0,67	3,079	
2805	7	Piscium	—	1 51,10	5 51,00	23 20 51,09	—	50,25	+0,84	3,090	
2806	7	12 Piscium <i>w</i> ²	1 53,40	—	6 53,51	23 20 53,49	—	52,89	+0,60	3,076	
2807	5	Cassiopeæ	4 18,30	—	—	23 22 18,30	—	17,71	+0,59	2,722	
2808	7	13 Piscium <i>w</i> ³	3 20,47	1 20,67	—	23 23 20,51	—	19,83	+0,68	3,076	
2809	7	Aquarii	—	—	7 29,68	23 23 29,67	—	29,51	-0,14	3,115	
2810	5	101 Aquarii <i>b</i> ⁴	6 28,72	—	—	23 24 28,72	—	28,61	+0,11	3,151	
2811	5	71 Pegasi <i>γ</i>	5 4,01	—	—	23 25 4,01	—	3,65	+0,36	2,988	
2812	6.7	14 Piscium <i>w</i> ⁴	—	4 30,83	—	23 25 30,83	—	30,70	+0,13	3,076	
2813	5	Phoenicis <i>ι</i>	5 0,70	—	3 0,74	23 26 0,69	—	0,88	-0,19	3,256	
2814	6.7	Aquarii	—	5 52,15	—	23 26 52,15	—	51,67	+0,48	3,097	
2815	7	15 Piscium	—	5 53,48	—	23 26 53,48	—	52,76	+0,72	3,067	
2816	6	Aquarii	—	3 19,82	3 19,51	23 27 19,64	—	19,47	+0,17	3,169	
2817	6	16 Piscium	—	—	6 49,08	23 27 49,08	—	49,11	-0,03	3,064	
2818	6	Aquarii	—	4 56,84	—	23 28 56,83	—	56,38	+0,45	3,113	
2819	4.5	16 Androm	4 22,00	—	5 21,79	23 29 21,92	21,97	21,28	-0,05	+0,64	2,886
2820	6	75 Pegasi <i>λ</i> <i>s</i>	—	5 28,45	—	23 29 28,46	—	28,05	+0,41	3,012	
2821	5	Phoenicis <i>θ</i>	7 25,19	—	—	23 30 25,19	—	24,57	+0,62	3,257	
2822	5	102 Aquarii <i>ω</i> ¹	6 4,07	—	—	23 31 4,07	—	3,73	+0,34	3,114	
2823	4.5	17 Piscium <i>ι</i>	6 18,65	4 18,75	4 18,77	23 31 18,72	18,73	18,67	-0,01	+0,05	3,054
2824	5	19 Androm	3 9,32	—	—	23 32 9,32	—	8,97	+0,35	2,914	
2825	6	Aquarii	—	5 27,19	—	23 32 27,19	—	26,73	+0,46	3,104	
2826	3	35 Cephei <i>γ</i>	—	7 30,61	—	23 32 30,86	31,30	29,58	-0,44	+1,23	2,390
2827	5	103 Aquarii <i>A</i> ¹	—	2 51,51	—	23 32 51,50	—	51,00	+0,50	3,123	
2828	5	104 Aquarii <i>A</i> ²	—	—	—	23 33 —	—	1,83	—	3,122	
2829	5	16 Piscium <i>λ</i>	1 28,62	—	6 28,71	23 33 28,69	28,67	28,48	+0,02	+0,21	3,066
2830	5.6	105 Aquarii <i>ω</i> ²	—	4 0,54	—	23 34 0,53	—	0,55	-0,02	3,110	
2831	6	76 Pegasi	—	2 12,77	4 12,72	23 34 12,75	—	11,59	+1,16	3,027	
2832	5.6	77 Pegasi <i>O</i>	—	5 49,68	—	23 34 49,68	—	49,30	+0,38	3,044	
2833	5	106 Aquarii <i>A</i> ³	6 29,07	—	—	23 35 29,07	—	28,77	+0,30	3,118	
2834	5	78 Pegasi <i>i</i>	6 33,30	—	—	23 35 33,30	—	33,48	-0,18	2,991	
2835	7	Piscium	4 14,83	3 14,84	—	23 36 14,84	—	14,24	+0,60	3,053	

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in			Mean N. P. D. January 1, 1833.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession.	
	No. 1831	No. 1832	No. 1833				Green.	A. S. C.		
2791	3 36 17,32	—	3 36 15,42	78 36 16,37	—	36 18,57	—	2,20	—19,649	
2792	—	4 37 48,92	1 37 49,56	90 37 49,01	—	37 46,82	+	2,19	19,654	
2793	—	—	6 41 28,05	112 41 28,05	—	41 27,07	+	0,98	19,659	
2794	5 31 7,75	—	—	67 31 7,75	—	31 8,93	—	1,18	19,689	
2795	5 33 40,52	—	—	111 33 40,52	—	33 42,49	—	1,97	19,692	
2796	5 38 15,26	—	6 38 16,22	28 38 15,78	38 20,15	38 22,18	-4,37	-	6,40	19,696
2797	2 39 46,07	3 39 45,69	—	112 39 45,82	—	39 49,12	—	—	3,30	19,701
2798	—	7 39 49,98	5 39 49,19	89 39 49,65	39 47,05	39 42,03	+2,60	+	7,62	19,710
2799	—	4 48 0,08	—	89 48 0,08	—	47 50,43	—	—	9,65	19,715
2800	—	—	5 45 17,46	65 45 17,46	—	45 14,96	—	—	2,50	19,726
2801	5 32 18,43	—	—	84 32 18,43	—	32 35,27	—	—	16,84	19,728
2802	5 33 51,98	—	—	20 33 51,98	—	33 50,34	—	—	1,64	19,740
2803	5 9 53,77	—	—	78 9 53,77	—	9 49,07	—	—	4,70	19,746
2804	3 42 51,98	3 42 53,43	—	92 42 52,70	—	42 50,26	—	—	2,44	19,749
2805	—	1 26 49,40	4 26 51,23	95 26 50,86	—	26 40,13	—	—	10,73	19,749
2806	—	—	5 57 34,19	91 57 34,19	—	57 31,44	—	—	2,75	19,750
2807	5 22 34,54	—	—	32 22 34,51	—	22 36,79	—	—	2,25	19,771
2808	—	1 0 46,83	4 0 46,48	92 0 46,55	—	0 42,42	—	—	4,13	19,785
2809	—	2 28 11,62	3 28 10,77	102 28 11,11	—	28 11,09	—	—	0,02	19,787
2810	5 50 31,07	—	—	111 50 31,07	—	50 28,18	—	—	2,89	19,801
2811	6 25 35,03	—	—	68 25 35,03	—	25 28,08	—	—	6,95	19,809
2812	1 10 28,51	—	3 10 28,26	92 10 28,32	—	10 22,70	—	—	5,62	19,815
2813	5 32 30,59	—	—	133 32 30,59	—	32 28,00	—	—	2,59	19,821
2814	—	—	3 23 33,84	98 23 33,84	—	23 36,63	—	—	2,79	19,832
2815	—	3 36 53,24	2 36 52,53	89 36 52,96	—	36 49,32	—	—	3,64	19,832
2816	—	—	5 48 14,84	117 48 14,84	—	48 10,44	—	—	4,40	19,838
2817	—	5 49 42,70	—	88 49 42,70	—	49 37,97	—	—	4,73	19,844
2818	—	—	5 59 24,44	103 59 24,44	—	59 25,39	—	—	0,95	19,858
2819	6 27 6,93	1 27 6,18	4 27 7,40	44 27 7,06	27 3,76	27 0,13	+3,30	+	6,93	19,863
2820	—	5 31 43,25	—	72 31 43,25	—	31 49,04	—	—	5,79	19,864
2821	5 34 7,18	—	—	137 34 7,18	—	34 8,03	—	—	0,85	19,875
2822	5 9 5,08	—	—	105 9 5,08	—	8 54,54	—	—	10,54	19,882
2823	5 17 4,03	1 17 1,80	5 17 3,80	85 17 3,73	17 1,80	16 59,69	+1,93	+	4,04	19,335*
2824	5 35 42,28	—	—	46 35 42,28	—	35 44,72	—	—	2,44	19,894
2825	—	5 36 38,94	—	102 36 38,94	—	36 40,61	—	—	1,67	19,897
2826	4 18 15,84	1 18 14,43	—	13 18 15,56	18 17,35	18 20,20	-1,79	-	4,64	19,898
2827	4 57 16,50	2 57 14,23	—	108 57 15,74	—	57 11,09	—	—	4,65	19,901
2828	1 44 45,21	4 44 48,10	—	108 44 47,77	—	44 46,54	—	—	1,23	19,903
2829	3 8 36,73	1 8 35,89	1 8 36,91	89 8 36,52	8 36,54	8 37,92	-2,02	-	1,40	19,907
2830	—	5 28 25,20	—	105 28 25,20	—	28 18,52	—	—	6,68	19,913
2831	—	5 35 48,10	—	74 35 48,10	—	35 42,87	—	—	5,23	19,915
2832	—	4 36 4,02	—	80 36 4,02	—	36 57,96	—	—	6,06	19,921
2833	5 12 31,61	—	—	109 12 31,61	—	12 26,85	—	—	4,76	19,927
2834	5 34 3,11	—	—	61 34 3,11	—	34 6,02	—	—	2,91	19,928
2835	5 44 19,45	—	—	83 44 19,45	—	44 21,98	—	—	2,53	19,934

No.	Mag	Names.	Mean A. R. January 1, 1832, from Observations in			Mean A. R. January 1, 1832	Greenh Catal.	A. S. Catal.	Difference from		Annual Preces- sion
			No. 1831	No. 1832	No. 1833				Green.	A. S.	
			s.	s.	s.				h. m. s.	s.	
2836	6	107 Aquarii A ⁴	—	5 17,19	—	23 37 17,18	—	16,67	+0,51	+3,115	
2837	5	90 Androm ψ	6 43,99	—	—	23 37 43,99	—	44,83	-0,84	2,936	
2838	6	19 Piscium μ	—	5 48,74	8 48,65	23 37 48,69	48,69	43,57	+0,12	3,062	
2839	6	Aquarii Y	—	6 36,83	1 36,85	23 38 36,83	—	36,58	+0,25	3,097	
2840	5	5 Cassiopeæ τ	4 52,97	—	—	23 38 52,92	—	52,24	+0,73	2,873	
2841	5.6	20 Piscium π	1 18,42	5 18,46	—	23 39 18,45	—	17,93	+0,52	3,076	
2842	6.7	Aquarii	—	5 54,50	—	23 39 54,50	—	54,34	+0,16	3,083	
2843	5	Draconis	3 55,86	—	—	23 39 55,86	—	55,59	+0,27	2,793	
2844	5	Ap. Sculp δ	6 10,01	—	—	23 40 10,01	—	9,56	+0,45	3,133	
2845	6	21 Piscium	7 51,57	—	—	23 40 51,57	—	51,42	+0,15	3,068	
2846	6	79 Pegasi	—	3 9,94	3 9,97	23 41 9,98	—	9,21	-0,77	3,010	
2847	6	Aquarii	—	3 34,65	—	23 41 34,65	—	33,83	+0,82	3,089	
2848	6	Aquarii	1 53,25	5 53,33	—	23 41 53,31	—	52,56	+0,75	3,097	
2849	5	Octantis γ^1	—	—	—	23 41 —	—	56,11	—	3,926	
2850	6	108 Aquarii A ⁵	—	5 40,51	—	23 42 40,50	—	39,85	+0,65	3,105	
2851	7	80 Pegasi	—	5 47,27	—	23 42 47,27	—	47,34	-0,07	3,053	
2852	6	22 Piscium	—	5 22,12	—	23 43 22,12	—	21,74	+0,38	3,065	
2853	6	Aquarii	—	—	6 51,92	23 43 51,90	—	51,90	0,00	3,094	
2854	6	23 Piscium	—	—	5 52,29	23 43 52,31	—	52,16	+0,15	3,032	
2855	6	81 Pegasi ϕ	—	—	4 56,95	23 43 56,97	—	56,41	+0,56	3,037	
2856	6	82 Pegasi	—	2 3,56	4 3,50	23 44 3,53	—	3,28	+0,25	3,052	
2857	6.7	24 Piscium	1 17,82	—	6 17,94	23 44 17,92	—	17,73	+0,19	3,075	
2858	6.7	25 Piscium	—	5 28,78	—	23 44 28,78	—	28,07	+0,71	3,066	
2859	6.7	Piscium	4 10,96	2 10,96	—	23 46 10,96	—	10,88	+0,08	3,069	
2860	6	26 Piscium	6 32,44	5 32,44	—	23 46 32,44	—	32,13	+0,31	3,060	
2861	5	Octantis γ^2	—	—	—	23 48 —	—	1,78	—	3,651	
2862	5	Tucanæ η	6 42,75	—	—	23 48 42,75	—	41,64	+1,11	3,212	
2863	5.6	84 Pegasi \downarrow	1 12,59	4 12,78	—	23 49 12,73	—	12,61	+0,12	3,040	
2864	5	27 Piscium p	5 4,40	5 4,52	4 4,37	23 50 4,43	4,30	3,71	+0,13	+0,72	3,073
2865	4.5	28 Piscium α	5 41,40	7 41,54	1 41,46	23 50 41,46	41,28	41,34	+0,17	+0,12	3,062
2866	7	Piscium	—	4 3,60	—	23 51 3,60	—	2,88	+0,72	3,121*	
2867	5	Tucanæ ϵ	3 6,88	—	—	23 51 6,88	—	7,08	-0,20	3,189	
2868	5	Cassiopeæ	—	—	—	23 53 —	—	—	—	2,936	
2869	5	29 Piscium q	4 12,98	—	3 12,94	23 53 12,97	13,02	12,11	-0,05	+0,86	3,071
2870	4.5	30 Piscium r	6 20,73	1 20,65	3 20,59	23 53 20,65	20,74	20,27	-0,09	+0,38	3,073
2871	6	85 Pegasi	—	5 24,61	—	23 53 24,60	—	23,86	+0,74	3,109*	
2872	6	31 Piscium c^1	—	5 48,31	—	23 53 48,31	—	47,83	+0,48	3,063	
2873	6	32 Piscium c^2	—	5 54,81	—	23 53 54,87	—	54,32	+0,55	3,063	
2874	4	2 Ceti g	6 7,28	6 7,85	—	23 55 7,81	7,77	7,72	+0,04	+0,09	3,078
2875	6	3 Ceti p	—	6 54,01	—	23 55 54,01	—	53,96	+0,05	3,073	
2876	6.7	Piscium	—	6 27,35	—	23 56 27,35	—	26,91	+0,44	3,069	
2877	5	33 Piscium	7 44,20	—	4 44,17	23 56 44,19	44,23	43,89	-0,04	+0,30	3,079
2878	6	86 Pegasi k	1 4,78	4 5,09	—	23 57 5,03	—	5,08	-0,05	3,064	
2879	7	4 Ceti	2 7,84	1 7,83	—	23 59 7,84	—	7,51	+0,33	3,068	
2880	7	5 Ceti	4 35,85	4 36,05	—	23 59 35,95	—	35,77	+0,18	3,068	
2881	1	21 Androm α	29 43,24	23 43,15	15 43,13	23 59 43,19	43,20	42,72	-0,01	+0,47	+3,067

No.	Mean N. P. D. reduced to January 1, 1832, from Observations in						Mean N.P.D. January 1, 1832.	Greenwich Catalogue.	A. S. Catalogue.	Difference from		Annual Precession
	1831		1832		1833					Green.	A. S. C.	
	No.	"	No.	"	No.	"						
2836			5 36	46,35			109 36	46,35	36 42,32	+ 4,03	19,948	
2837	6 30	45,81					44 30	45,81	30 43,02	+ 2,79	19,947	
2838			4 26	38,78			87 26	38,78	26 35,36	+ 3,42	19,948	
2839			5 50	19,53			102 50	19,53	50 20,10	- 0,57	19,955	
2840	5 17	0,52					32 17	0,52	17 1,63	- 1,11	19,957	
2841			5 41	39,03			93 41	39,03	41 36,52	+ 2,51	19,960	
2842	2 18	44,22	4 18	43,14			97 18	43,50	18 44,26	- 0,76	19,965	
2843	5 7	34,06					23 7	34,06	7 37,99	- 3,92	19,965	
2844	6 3	29,40	1 3	28,34			119 3	29,25	3 23,21	+ 1,04	19,967	
2845	5 51	26,34					89 51	26,34	51 21,85	+ 4,49	19,972	
2846			4 5	32,13	1 5	32,37	62 5	32,18	5 27,30	+ 4,88	19,974	
2847					5 54	42,62	100 54	42,62	54 43,76	- 1,14	19,977	
2848			5 20	5,68			105 20	5,68	19 58,98	+ 6,70	19,979	
2849							172	Invisible	57 3,16		19,979	
2850			5 50	38,98			109 50	38,98	50 33,54	+ 5,44	19,985	
2851			5 37	2,75			81 37	2,75	36 59,36	+ 3,39	19,985	
2852	3 0	8,99	2 0	9,95			88 0	9,88	0 10,55	- 1,17	19,989	
2853			3 11	10,73	3 11	11,03	105 11	10,88	11 6,15	+ 4,73	19,992	
2854	1 15	45,65	2 15	44,91	2 15	45,42	69 15	45,26	15 37,75	+ 7,51	19,992	
2855	1 48	47,37			4 18	43,63	71 48	43,38	48 43,11	+ 0,27	19,993	
2856					5 59	17,54	79 59	17,54	59 12,45	+ 5,09	19,993	
2857					5 5	15,27	94 5	15,27	5 6,32	+ 8,45	19,995	
2858					5 50	33,71	88 50	33,71	50 31,42	+ 2,29	19,996	
2859			5 49	30,42			90 49	30,42	49 29,00	+ 1,42	20,006	
2860	1 51	46,25	4 51	44,59			83 51	44,92	51 44,20	+ 0,72	20,007	
2861							173	Invisible	6 4,85		20,015	
2862	1 13	41,11					155 13	41,11	14 25,84	- 44,73	20,018	
2863	3 47	30,76	2 47	32,01	1 47	32,67	65 47	31,49	47 27,32	+ 4,17	20,020	
2864	5 29	16,79					94 29	16,79	29 16,55	+ 4,49	20,023	
2865	5 3	57,54	4 3	58,57	3 3	58,98	84 3	58,10	3 59,82	- 1,72	20,026	
2866			5 49	33,05			96 49	33,05	49 30,48	+ 2,57	20,027	
2867			5 30	38,94			156 30	38,94	30 31,77	+ 7,17	20,027	
2868	5 42	47,29					29 42	47,29	42 45,48	+ 1,81	20,033	
2869	5 57	43,40					93 57	43,40	57 45,89	+ 2,71	20,034	
2870	5 56	51,37					96 56	51,37	56 52,03	- 0,66	20,034	
2871	4 48	22,25	1 48	22,54			63 48	22,31	48 34,67	- 12,36	18,884	
2872			6 58	41,93			81 58	41,93	58 39,45	+ 2,48	20,035	
2873			5 26	51,12			82 26	51,12	26 53,29	+ 2,17	20,035	
2874	5 16	15,56					108 16	15,56	16 15,52	+ 0,04	20,038	
2875			5 26	39,05			101 26	39,05	26 35,61	+ 3,44	20,039	
2876	2 26	9,06	1 26	10,54			91 26	9,55	26 6,71	+ 2,84	20,040	
2877	5 38	48,34					96 38	48,34	38 51,70	- 3,36	20,040	
2878			3 32	18,04	1 32	18,70	77 32	18,20	32 19,03	- 0,83	20,041	
2879			6 28	59,43			93 28	59,43	28 57,29	- 2,14	20,042	
2880	5 22	55,16					93 22	55,16	22 52,14	- 3,02	20,043	
2881	50 50	13,12	42 60	13,46	9 50	12,85	61 50	13,24	50 14,37	- 1,12	20,043	

ERRATA IN THE CATALOGUE OF FIXED STARS.

No.	Column.				
60	Mean A. R.	for	1",21	read	1",26
101	{ A. R. 1832	—	4",62	—	7",72
	{ Mean A. R.	—	4",62	—	7",72
196	{ A. R. 1831	—	24",11	—	23",61
	{ Mean A. R. }				
281	Mean A. R.	—	2h. 3m.	—	2h. 32m.
465	Mean A. R.	—	3h.	—	4h.
772	{ N.P.D. 1831	—	43",73	—	43",11
	{ Mean N.P.D.	—	41",74	—	41",46
813	{ N.P.D. 1831	—	4",08	—	4",70
	{ N.P.D. 1832	—	0",58	—	1",97
	{ Mean N.P.D.	—	2",33	—	3",33
989	{ N.P.D. 1832	—	55'	—	53'
	{ Mean N.P.D. }				
1110	{ N.P.D. 1832	—	33",15	—	33",75
	{ Mean N.P.D.	—	33",61	—	33",91
1211	{ N.P.D. 1831	—	34' 58",25	—	35' 7",25
	{ Mean N.P.D.	—	34' 58",25	—	35' 7",85
1333	{ N.P.D. 1832	—	8",58	—	9",14
	{ Mean N.P.D.	—	10",16	—	10",53
1741	{ A. R. 1833	}	The seconds in these columns to be transferred to No. 1744.		
	{ Mean A. R. }				
1752	Mean N.P.D.	for	106° 41'	read	106° 40'
1819	{ A. R. 1833	—	41",34	—	41",92
	{ Mean A. R.	—	41",36	—	41",94
1964	Mean A. R.	—	26",89	—	26",63
2205	Green. A. R.	—	36",18	—	36",78
2424	A. R.	—	29h.	—	20h.
2439	Diff. of A. R.	—	— 3",25	—	— 0",25
2536	{ N.P.D. 1831	—	22",39	—	23",48
	{ N.P.D. 1833	—	25",81	—	25",05
	{ Mean N.P.D.	—	24",65	—	24",49
2594	{ A. R. 1832	—	35",94	—	36",22
	{ Mean A. R.	—	36",00	—	36",28
2642	{ N.P.D. 1832	—	38",05	—	36",74
	{ Mean N.P.D.	—	38",05	—	34",81

N. B.—In addition to the above the differences should be corrected.

REMARKS UPON THE CATALOGUE OF FIXED STARS.

The casualties to which I have already alluded at the early part of this work as affecting the observations made with the Transit Instrument in 1832 and 1833, renders it desirable that a comparison should be made between the observations of these years and the observations of 1831, in which no uncertainty of any kind exists; for this purpose putting α α^1 &c. and β β^1 &c. to represent the errors in seconds of space of Azimuth and Collimation respectively, and selecting from the catalogue those Stars situated near the Pole (as affording large co-efficients) which have been observed in each of the three years, we have as follows.

From observations made in 1831 1832 1833.

NAMES.	N.P.D.	A.R.	Mean } Place. }			Mean } Place. }			Mean } Place. }			
			h. m. s.	+ A	+ C	h. m. s.	+ A	+ C	h. m. s.	+ A	+ C	
Draconis	11,27	12 4	13,20	+ ,31	α + ,50	β 13,69	+ ,31	α^1 + ,50	β^1 12,45	+ ,31	α^{11} + ,50	β^{11}
7 Urs. Min.	β 15,10	14 51	17,18	+ ,22	+ ,38	17,28	+ ,22	+ ,38	15,99	+ ,22	+ ,38	
Urs. Maj.	20, 3	10 30	54,76	+ ,16	+ ,29	55,06	+ ,16	+ ,29	54,60	+ ,16	+ ,29	
Cassiopei	23,21	2 15	20,50	+ ,14	+ ,25	20,43	+ ,14	+ ,25	19,82	+ ,14	+ ,25	
11 Draconis	α 24,49	13 59	50,64	+ ,13	+ ,24	50,97	+ ,13	+ ,24	50,19	+ ,13	+ ,24	
50 Ursæ. Maj.	α 27,21	10 53	17,49	+ ,11	+ ,22	17,40	+ ,11	+ ,22	17,31	+ ,11	+ ,22	
5 Cephei	α 28, 8	21 14	34,12	+ ,10	+ ,21	33,88	+ ,10	+ ,21	33,27	+ ,10	+ ,21	
29 Ursæ. Maj.	ν 30 11	9 38	58,46	+ ,09	+ ,20	58,40	+ ,09	+ ,20	58,27	+ ,09	+ ,20	
18 Cassiopeæ	ω 34,23	0 31	1,42	+ ,08	+ ,18	1,28	+ ,08	+ ,18	1,18	+ ,08	+ ,18	
64 Ursæ. Maj.	γ 35 22	11 44	57,42	+ ,07	+ ,17	57,50	+ ,07	+ ,17	57,54	+ ,07	+ ,17	
33 Draconis	γ 38 29	17 52	42,42	+ ,06	+ ,16	42,58	+ ,06	+ ,16	42,15	+ ,06	+ ,16	
33 Persei	α 40,44	3 12	22,05	+ ,06	+ ,15	22,14	+ ,06	+ ,15	21,94	+ ,06	+ ,15	
51 Persei	μ 42, 1	4 2	35,48	+ ,05	+ ,15	35,52	+ ,05	+ ,15	35,22	+ ,05	+ ,15	
13 Aurigæ	α 44, 1	4 4	17,46	+ ,05	+ ,14	17,48	+ ,05	+ ,14	17,22	+ ,05	+ ,14	
26 Persei	β 49,41	2 57	16,03	+ ,04	+ ,13	15,89	+ ,04	+ ,13	16,15	+ ,04	+ ,13	

Similarly we have the following observations made near to the South Horizon in the years 1831 1832 and 1833.

NAMES.	N.P.D.	A.R.	Mean } Place. }			Mean } Place. }			Mean } Place. }			
			h. m. s.	+ A	+ C	h. m. s.	+ A	+ C	h. m. s.	+ A	+ C	
Columbæ	α 124,10	5 33	34,14	- ,06	α + ,12	β 34,10	- ,06	α^1 + ,12	β^1 34,23	- ,06	α^{11} + ,12	β^{11}
Columbæ	γ 125,18	5 51	34,86	- ,06	+ ,12	34,98	- ,06	+ ,12	35,02	- ,06	+ ,12	
Eridani	ϵ 133,43	3 13	13,19	- ,07	+ ,14	13,29	- ,07	+ ,14	13,67	- ,07	+ ,14	
Arg. in pup. L.	δ^1 134,53	7 8	12,17	- ,08	+ ,14	11,82	- ,08	+ ,14	11,86	- ,08	+ ,14	
Argus	ϕ 143,45	9 50	58,78	- ,10	+ ,17	58,87	- ,10	+ ,17	58,50	- ,10	+ ,17	
Argus	δ 144, 5	8 40	3,76	- ,10	+ ,17	3,93	- ,10	+ ,17	4,21	- ,10	+ ,17	
Eridani	α 148, 5	1 31	27,19	- ,12	+ ,18	27,27	- ,12	+ ,18	27,29	- ,12	+ ,18	
Argus	ι 148,33	9 12	35,64	- ,12	+ ,18	36,11	- ,12	+ ,18	35,90	- ,12	+ ,18	
Arg. in Car.	η 150,29	10 11	29,23	- ,13	+ ,20	28,88	- ,13	+ ,20	29,32	- ,13	+ ,20	
Crucis	α^2 152, 9	12 17	18,98	- ,14	+ ,21	18,69	- ,14	+ ,21	19,16	- ,14	+ ,21	
Arg. in Car.	ν 154,11	9 23	5,21	- ,15	+ ,23	5,22	- ,15	+ ,23	5,57	- ,15	+ ,23	
Argus	ν 154,17	9 42	54,01	- ,15	+ ,23	54,25	- ,15	+ ,23	54,21	- ,15	+ ,23	
Hydri	ξ 158,59	2 36	2,03	- ,18	+ ,30	2,49	- ,18	+ ,30	2,24	- ,18	+ ,30	
Argus	β 159, 1	9 11	19,45	- ,18	+ ,30	19,73	- ,18	+ ,30	19,57	- ,18	+ ,30	

Taking the mean of each set we obtain the following Equations:

$$30^{\circ},57 + ,11 a + ,22 \beta = 30^{\circ},63 + ,11 a' + ,22 \beta' = 30^{\circ},22 + ,11 a'' + ,22 \beta''$$

$$24,90 - ,12 a + ,19 \beta = 24,97 + ,12 a' + ,19 \beta' = 25,05 - ,12 a'' + ,19 \beta''$$

by subtraction we have

$$5^{\circ},67 + ,23 a + ,03 \beta = 5^{\circ},66 + ,23 a' + ,03 \beta' = 5^{\circ},17 + 23 a'' + ,03 \beta''$$

showing that the Azimuth correction for 1832 differs insensibly from that of 1831 and may consequently be assumed = 0, and that the observations for 1833 stand in need of the correction 2',22 of space on this account. Now the observations constituting the above result for 1833, were made between the 20th January and the end of October, during which period we employed the formulæ $\frac{96^{\circ},20 - N - S}{2}$ for computing the Azimuthal error, and traced *as well as circumstances would then permit* (see pages 41 and 42.) that an alteration took place in the situation of the meridian marks in the same direction and to nearly the same amount with the correction now found on or about the 12th November; our present result however fixes the date of the alteration in question at a much earlier period; probably at the commencement of the year.

By adding the above Equations together we determine :

$$55^{\circ},47 - ,01 a + ,41 \beta = 55^{\circ},60 - ,01 a' + ,41 \beta' = 55^{\circ},27 - ,01 a'' + ,41 \beta''$$

from whence (assuming as above that 1831 is devoid of error i. e. that $a = \beta = 0$) we determine that the observations for 1832 require a correction for Collimation to the amount $- 0^{\circ},32$ and that those for 1833 require a correction $+ 0^{\circ},50$ or it appears on the whole, *that the observations of 1832 require no correction of consequence to reduce them to the tenor of those of 1831, and that those of 1833 up to November 12, require correction as follows.*

North Polar Distance.	Correction.	North Polar Distance.	Correction.
.	"	.	"
20	+ ,414	95	+ ,022
25	+ ,328	100	+ ,013
30	+ ,265	105	+ ,004
35	+ ,224	110	- ,007
40	+ ,187	115	- ,017
45	+ ,161	120	- ,028
50	+ ,139	125	- ,042
55	+ ,120	130	- ,055
60	+ ,104	135	- ,069
65	+ ,089	140	- ,087
70	+ ,077	145	- ,108
75	+ ,064	150	- ,133
80	+ ,054	155	- ,168
85	+ ,043	160	- ,220
90	+ ,033		

REMARKS UPON THE CATALOGUE OF FIXED STARS. cxxxiii

The above corrections not having been taken into account in obtaining the column "mean," our ' difference from Greenwich' and " difference from A. S." exhibits the *true* difference for those cases where the Star has been observed in 1831 and 1832 only; as the Catalogue now stands out of the 687 comparisons between the Madras and Greenwich Catalogues there are

280 cases which do not exceed ,05s. of time.
 or 470 ——— ——— ——— ——— ,10s. ———
 or 615 ——— ——— ——— ——— ,20s. ———
 and 72 ——— ——— ——— ——— exceed ,20s. ———

Admitting the accuracy of the above corrections, (for it must I think be readily conceded that some such sort of correction is necessary) and neglecting those which are below ,05s. except in the case of the principal Stars; the following corrections of the column " Mean A. R. January 1, 1832" become necessary.

No.	Correc- tion.	No.	Correc- tion.	No.	Correc- tion.	No.	Correc- tion.	No.	Correc- tion.
46	— 0,15	459	+ 0,09	637	+ 0,10	776	+ 0,10	976	+ 0,09
89	+ 0,09	462	+ 0,06	638	+ 0,07	779	+ 0,09	979	+ 0,09
113	+ 0,08	479	+ 0,06	640	+ 0,07	780	+ 0,05	992	+ 0,09
153	+ 0,07	481	+ 0,06	649	+ 0,07	783	+ 0,05	998	+ 0,07
157	+ 0,07	484	+ 0,09	651	+ 0,06	785	+ 0,05	999	+ 0,11
180	+ 0,06	497	+ 0,11	654	+ 0,06	790	+ 0,03	1023	+ 0,06
211	+ 0,08	505	+ 0,03	655	+ 0,07	796	+ 0,09	1025	+ 0,09
220	+ 0,05	509	+ 0,06	656	+ 0,06	799	+ 0,08	1028	+ 0,06
242	+ 0,08	511	+ 0,05	657	+ 0,09	800	+ 0,10	1032	— 0,05
247	+ 0,18	512	+ 0,06	670	+ 0,06	801	+ 0,08	1054	+ 0,08
253	+ 0,09	516	+ 0,06	671	+ 0,08	803	+ 0,08	1055	+ 0,05
259	+ 0,07	519	+ 0,06	674	+ 0,09	809	+ 0,07	1058	+ 0,08
262	+ 0,06	522	+ 0,10	678	+ 0,08	811	+ 0,07	1082	+ 0,05
270	+ 0,06	531	+ 0,06	683	+ 0,07	817	+ 0,10	1099	+ 0,06
271	+ 0,05	536	+ 0,06	687	+ 0,10	823	+ 0,07	1102	— 0,05
277	+ 0,09	537	+ 0,06	688	+ 0,09	843	+ 0,06	1103	+ 0,05
283	+ 0,08	538	— 0,05	701	+ 0,06	851	+ 0,06	1120	— 0,09
301	+ 0,07	552	+ 0,24	704	+ 0,06	859	+ 0,07	1124	+ 0,07
317	+ 0,08	572	+ 0,09	706	+ 0,07	861	+ 0,09	1131	+ 0,06
342	+ 0,07	573	+ 0,06	707	+ 0,07	864	+ 0,09	1133	— 0,12
344	+ 0,08	577	+ 0,13	708	+ 0,06	870	+ 0,09	1135	+ 1,24
346	+ 0,08	580	+ 0,05	716	+ 0,06	871	+ 0,09	1137	— 0,06
358	+ 0,09	581	+ 0,05	717	+ 0,10	889	+ 0,09	1144	— 0,06
362	+ 0,08	582	+ 0,05	720	+ 0,06	891	+ 0,10	1146	+ 0,09
369	+ 0,09	594	+ 0,09	721	+ 0,08	900	+ 0,06	1148	+ 0,22
374	+ 0,05	595	+ 0,08	723	+ 0,06	912	+ 0,05	1156	— 0,08
376	+ 0,09	596	+ 0,10	727	+ 0,11	919	+ 0,06	1158	+ 0,09
401	+ 0,08	601	+ 0,05	731	+ 0,08	922	+ 0,10	1175	+ 0,04
413	+ 0,09	602	+ 0,08	734	+ 0,04	926	+ 0,06	1179	+ 0,08
419	+ 0,09	609	+ 0,06	743	+ 0,10	927	+ 0,06	1181	+ 0,10
423	+ 0,09	611	+ 0,09	745	+ 0,08	931	+ 0,11	1182	— 0,07
430	+ 0,06	620	+ 0,06	754	+ 0,09	933	+ 0,05	1196	— 0,05
435	+ 0,08	623	+ 0,08	755	+ 0,07	939	+ 0,06	1209	+ 0,03
439	— 0,06	624	+ 0,07	766	+ 0,09	940	+ 0,07	1221	+ 0,06
451	+ 0,09	629	+ 0,07	768	+ 0,08	946	+ 0,09	1225	— 0,05
453	+ 0,10	634	+ 0,10	769	+ 0,08	948	+ 0,06	1229	— 0,06
457	+ 0,08	636	+ 0,10	771	+ 0,05	951	+ 0,07	1230	+ 0,07

CXXXIV REMARKS UPON THE CATALOGUE OF FIXED STARS.

No.	Correc- tion.	No.	Correc- tion.	No.	Correc- tion.	No.	Correc- tion.	No.	Correc- tion.
1236	+ 0 05	1575	+ 0 12	1870	+ 0 10	2091	+ 0,10	2342	+ 0,07
1240	+ 0 07	1599	+ 0,07	1876	+ 0,17	2094	+ 0 08	2347	+ 0,07
1255	+ 0,13	1607	+ 0 17	1879	+ 0,11	2111	+ 0,18	2353	+ 0 07
1266	+ 0,22	1616	+ 0 06	1880	+ 0,12	2113	+ 0 06	2358	+ 0,05
1273	- 0 06	1621	+ 0 07	1887	+ 0,13	2116	+ 0,10	2359	+ 0,09
1281	- 0,06	1625	+ 0 04	1895	+ 0,62	2117	+ 0,08	2362	+ 0,06
1285	+ 0,08	1628	+ 0,07	1899	+ 0,05	2121	+ 0 08	2370	+ 0 08
1291	+ 0,05	1639	+ 0,11	1903	+ 0,40	2134	+ 0 08	2377	+ 0 09
1294	- 0,05	1645	+ 0,08	1906	+ 0 21	2143	+ 0,13	2379	+ 0,09
1301	+ 0,11	1655	- 0 15	1918	+ 0,33	2146	+ 0,08	2380	+ 0,09
1305	+ 0,14	1658	+ 0,06	1926	+ 0,17	2156	+ 0,06	2381	+ 0,05
1329	+ 0,06	1664	+ 0,05	1927	+ 0,07	2158	- 0,12	2385	+ 0,08
1330	+ 0,06	1685	+ 0,10	1932	+ 0 05	2171	+ 0,13	2392	+ 0,09
1331	+ 0,06	1700	+ 0,34	1950	+ 0,33	2172	+ 0,23	2394	+ 0,17
1335	- 0 05	1708	+ 0,05	1955	+ 0,05	2173	+ 0,07	2398	- 0,11
1350	+ 0,23	1714	+ 0 06	1963	+ 0 13	2177	+ 0,05	2410	+ 0,05
1360	- 0 06	1722	- 0 16	1964	+ 1,33	2182	+ 0,08	2419	+ 0 05
1370	+ 0 09	1734	- 0 05	1976	+ 0 09	2192	+ 0,26	2420	+ 0,18
1379	+ 0,12	1735	- 0,07	1980	+ 0 05	2194	+ 0,06	2421	- 0,17
1404	+ 0,85	1745	+ 0 06	1982	+ 0 11	2200	+ 0,05	2428	+ 0,05
1406	+ 0 06	1748	+ 0,08	1987	+ 0,09	2209	+ 0,45	2435	+ 0,05
1407	+ 0,08	1765	+ 0 07	1996	+ 0 09	2219	+ 0,05	2436	+ 0,08
1421	+ 0 06	1766	- 0 06	1999	+ 0 06	2231	+ 0,08	2438	+ 0 08
1426	- 0 11	1767	+ 0 06	2001	- 0,07	2242	+ 0,13	2442	+ 0,06
1435	+ 0 06	1770	+ 0 05	2012	- 0 06	2259	+ 0,09	2458	+ 0 06
1436	+ 0 09	1775	+ 0 05	2013	+ 0 10	2263	+ 0 06	2480	+ 0 06
1440	+ 0 05	1783	+ 0 06	2014	+ 0 06	2267	+ 0 10	2485	+ 0 05
1447	+ 0,09	1784	+ 0 06	2015	+ 0,05	2279	+ 0,09	2488	+ 0 08
1451	+ 0 05	1786	+ 0,08	2016	+ 0 14	2283	+ 0,08	2499	+ 0 06
1455	+ 0 06	1791	+ 0 03	2026	+ 0 09	2292	+ 0 08	2526	- 0 17
1461	+ 0,08	1793	+ 0 07	2030	+ 0,39	2296	+ 0,10	2536	+ 0 16
1464	+ 0,05	1801	+ 0 07	2034	+ 0,07	2298	+ 0,07	2541	+ 0 09
1501	- 0,09	1809	+ 0,08	2038	+ 0,09	2303	+ 0 08	2559	+ 0,14
1504	+ 0,08	1819	+ 0,06	2041	+ 0,39	2307	+ 0 10	2571	+ 0 15
1505	+ 0,09	1822	+ 0 06	2053	+ 0 09	2314	+ 0,09	2588	+ 0,08
1507	+ 0,10	1830	+ 0 07	2059	+ 0,23	2315	+ 0 09	2612	+ 0,06
1508	+ 0,10	1832	+ 0 08	2060	+ 0 10	2324	+ 0,05	2667	+ 0 10
1512	- 0 06	1842	+ 0 26	2068	+ 0,07	2325	- 0 25	2751	+ 0 05
1520	+ 0,05	1843	+ 0 05	2071	+ 0 06	2330	+ 0 05	2755	+ 0 02
1534	+ 0,07	1844	+ 0 05	2081	+ 0 08	2332	+ 0 08	2854	+ 0,08
1541	+ 0,05	1867	+ 0,08	2087	+ 0 08	2341	+ 0 05		

When the corrections in the foregoing table are applied there appears to be

475 cases which do not exceed ,10s. of time.
 or 627 ——— ——— ——— ——— ,20s. ———
 and 60 ——— ——— ——— ——— exceed ,20s. ———

Among the latter class the following are those most deserving of notice.

No.			s.		h. m. s.
No. 115	α Urs. Min.	—	0,95	Madras	1 0 2,19
				Greenwich	— — 3,14
				Cambridge	— — 1,80
No. 340	* Persei	+ 0,65		Madras Results of 1832 and 1833 agree.	
No. 371	* Camelopard.	+ 0,47		Madras Result differs only ,05s. from the Greenwich Observations of 1831.	
No. 400	* Persei	— 0,37		Madras Result differs only ,02s. from the Greenwich Observations of 1831.	

			<i>s.</i>										
No. 552:	*	Camelopard.	+ 0,54	This must be examined.									
No. 603	β	Eridani	- 0,29	The Greenwich Result is no doubt about $\frac{1}{4}$ second too large.									
No. 832	28	Geminorum	+ 0,26	Must be examined.									
No. 874		Camelopardi	- 0,28	The situation of this Star (being only $7^{\circ} 18'$ from the Pole) fully accounts for the difference.									
No. 1106.	κ	Cancri	+ 0,36	Must be examined.									
No. 1386	π	Virginis	+ 0,66	The Greenwich place must be wrong									
No. 1417	ϵ	Virginis	+ 0,95	<table border="0" style="display: inline-table; vertical-align: middle;"> <tr> <td rowspan="4" style="font-size: 3em; vertical-align: middle;">}</td> <td>Madras Result</td> <td>49^o.42s.</td> </tr> <tr> <td>Greenwich</td> <td>— 48.47s.</td> </tr> <tr> <td>Cambridge</td> <td>— 49,22s.</td> </tr> <tr> <td colspan="2">Greenwich is about 1s. wrong.</td> </tr> </table>	}	Madras Result	49 ^o .42s.	Greenwich	— 48.47s.	Cambridge	— 49,22s.	Greenwich is about 1s. wrong.	
}	Madras Result	49 ^o .42s.											
	Greenwich	— 48.47s.											
	Cambridge	— 49,22s.											
	Greenwich is about 1s. wrong.												
No. 1451	k	Comæ Ber.	- 0,23	The Greenwich Observations for 1831 differ + ,08s. from the Madras Result.									
No. 1545.	l^b	Virginis	- 0,27	The Greenwich Result is probably too large.									
No. 1639.	θ	Bootis.	+ 0,36	The Greenwich Result is probably too small.									
No. 1754	γ^a	Urs. Min.	- 0,78	Only one observation: the Greenwich Observations for 1831 differ 0,29s. from the Madras Result.									
No. 1832	π	Serpentis	- 0,39	The Greenwich Observations of 1831 differ 0,18s. from the Madras Result.									
No. 1915	ϵ	Scorpii	- 0,46	The Greenwich Observations of 1831 differ 0,19s. from the Madras Result.									
No. 1964.	ϵ	Urs. Min.	+ 0,56	One observation.									
No. 2032	D	Ophiuchi	+ 0,43	The Greenwich Observations of 1831 differ 0,11s. and the Cambridge ,08s. from the Madras Result.									
No. 2043	γ	Telescopii	+ 0,29	The Greenwich place is probably too small.									
No. 2090	S ^a	Ophiuchi	- 0,24	The Greenwich place is probably too large.									
No. 2148	δ	Urs. Min.	- 0,67	<table border="0" style="display: inline-table; vertical-align: middle;"> <tr> <td rowspan="3" style="font-size: 3em; vertical-align: middle;">}</td> <td>Madras Result</td> <td>29 54s.</td> </tr> <tr> <td>Greenwich</td> <td>— 30.21s.</td> </tr> <tr> <td>Cambridge</td> <td>— 29,31s.</td> </tr> </table>	}	Madras Result	29 54s.	Greenwich	— 30.21s.	Cambridge	— 29,31s.		
}	Madras Result	29 54s.											
	Greenwich	— 30.21s.											
	Cambridge	— 29,31s.											
No. 2274	π .	Draconis	- 0,47	Requires examination.									
No. 2321	δ	Cygni	- 0,33	Do. Do.									
No. 2371	ρ	Draconis	- 0,45	The Greenwich Observations for 1831 differ only 0,10s. from the Madras Result.									
No. 2454	a	Microscopii	+ 0,29	Requires examination.									
No. 2465	η	Cephei	- 0,30	The Greenwich Observations for 1831 differ only 0,10s. from the Madras Result.									
No. 2489	ν	Cygni.	- 0,25	The Greenwich place is probably too large.									
No. 2593	π^a	Cygni	- 0,35.	Requires examination.									
No. 2676	c	Lacertæ	- 0,49	The Greenwich place must be too large.									
No. 2724.	τ^1	Aquarii	- 0,81	The Greenwich place is 1s. wrong.									
No. 2756	h^3	Aquarii	+ 0,41	The Greenwich place is probably too small.									
No. 2776	χ	Aquarii	+ 0,27	The Greenwich Observations of 1831 differ 0,15s. from the Madras Result.									
No. 2826	γ	Cephei	- 0,44	Requires to be examined.									

In the above comparison between the Madras and Greenwich Catalogues there now remain only 18 cases in which the difference exceeds a quarter of a second of time, the greater part of which will I apprehend be found to arise from *error* of result and *not* from the *uncertainty* attendant upon observation.

On looking over the column "Difference from A. S. C." a mere glance is sufficient to shew that the two Catalogues are not reckoned from the same Equinoctial point, a correction of about 0,30s. being necessary to reduce either Catalogue to the other; independant of this, it must be recollected that in the construction of the Society's Catalogue, proper motion was allowed only in 57 cases where it had been determined from "accurate observations" "to amount to 0,50s. in space (or 0,33s. in *time*)" since however accurate observations adapted to the purpose have not in many cases been available, it necessarily follows that the column of difference is further encumbered with the accumulated effect of proper motion, and consequently the discordances large or small cannot be looked upon as throwing any sort of doubt upon the accuracy of the Madras Results.

The discordancies most deserving attention are as follows.

No.			s.	
No. 21	ζ	Tucanæ	+ 2,97	About one years precession.
No. 55		Ceti	+ 3,57	Do. Do.
No. 79		Piscium	+ 2,03	Do. Do.
No. 91		Cephei	+ 5,62	N. P. D. 4° 39'.
No. 147	δ	Cassiopeiæ	— 1,42	Greenwich differs 0,19s. from the Madras Result.
No. 256	κ	Eridani	+ 1,96	Observed in two separate years at Madras.
No. 268		Ceti	+ 4,13	Do. Do.
No. 296	ξ	Hydri	+ 1,64	Observed in three separate years at Madras.
No. 340		Persei	+ 4,78	Madras confirmed by Greenwich.
No. 346		Arietis	+ 4,36	Observed in two separate years.
No. 439	γ	Hydri	+ 1,38	Do. Do.
No. 442	34	Tauri	+ 1,55	Do. Do.
No. 455	γ	Reticuli	+ 2,11	Do. Do.
No. 500	θ	Reticuli	+ 1,83	Madras Result re-examined;
No. 570	ι	Aurigæ	+ 3,25	Do. Do.
No. 610	ζ	Doradus	— 1,26	Madras confirmed by Greenwich.
No. 658		Leporis	+ 1,35	Observed in two separate years.
No. 661	χ	Aurigæ	+ 8,21	Greenwich differs 0,14s. from the Madras Result.
No. 721		Tauri	+ 1,50	} Madras Results re-examined.
No. 741	ε	Doradus	+ 1,54	
No. 805		This place has apparently been put in by mistake, the observations no doubt pertain to No. 799.		
No. 835	q	Camelopardi	— 1,96	Greenwich differs 0,08s. from the Madras Result.
No. 996	16	Argus	+ 1,38	} Madras Results re-examined.
No. 1132		Leonis	— 1,68	
No. 1135		Draconis	— 1,54	Observed in two separate years.
No. 1137	ι	Argus	+ 1,87*	Observed in three separate years.
No. 1156	n	Arg. in Car.	— 3,11	Do. Do.
1160	N	Arg. in Vel.	+ 1,50	Observed in two separate years.
1182	l	Arg. in Car.	+ 1,53	Do. Do.
1183	e	Sextantis	+ 1,91	Madras Result re-examined.
08	f	Sextantis	+ 1,49	Observed in three separate years.
29	q	Arg. in Car.	+ 1,54	Madras Result re-examined.

REMARKS UPON THE CATALOGUE OF FIXED STARS. CXXXVII

		<i>s.</i>		
No. 1234	T Arg. in Vel.	+	1,80	Observed in two separate years.
No. 1240	G Leonis Min.	+	1,48	Madras confirmed by Greenwich.
No. 1247	I Arg. in Car.	+	2,15	Madras Result re-examined.
No. 1270	34 Sextantis	+	1,70	Observed in two separate years.
No. 1273	θ^a Argus	+	1,97	Do. Do.
No. 1276	θ^b Argus	+	1,87	Do. Do.
No. 1360	λ Centauri	+	1,82	Do. Do.
No. 1406	δ Crucis	+	2,29	Do. Do.
No. 1421	ϵ Crucis	+	1,73	Do. Do.
No. 1426	α^d Crucis	+	1,68	Do. Do.
No. 1427	α^c Crucis	+	1,37	Observed in three separate years.
No. 1493	r Comæ Ber.	+	1,61	Observed in two separate years.
No. 1651	Libræ	—	1,55	
No. 1653	α^d Centauri	—	6,22	} Madras Results re-examined.
No. 1654	α^b Centauri †	—	6,35	
No. 1655	α Circini	—	2,12	
No. 1714	i Bootis	+	1,44	Greenwich differs 0,47s.
No. 1964	α Urræ Min.	+	3,96	Do, Do. 0,66s.
No. 1992	δ Aræ	—	1,59	Observed in two separate years.
No. 2067	α Sagittarii †	+	1,50	Re-examined.
No. 2148	δ Urræ Min.	+	7,33	Cambridge confirms Madras.
No. 2158	θ Pavonis	—	3,97	Observed in two separate years.
No. 2166	λ Pavonis	—	1,56	
No. 2278	ϵ Aquilæ	+	1,38	} Madras Results re-examined.
No. 2325	ϵ Pavonis	+	1,16	
No. 2378	R Sagittarii	+	2,99	
No. 2399	κ Cephei	+	3,29	Greenwich differs 0,17s.
No. 2421	ν Pavonis	+	1,56	Madras Result re-examined.
No. 2496	76 Draconis	—	1,97	Greenwich Observations for 1831 differ + 0,98s.
No. 2499	h Cephei	+	9,48	Do. Do. Do. Do. — 0,26s.
No. 2590	τ Cephei	+	1,70	Do. Do. Do. Do. — 0,19s.
No. 2621	Aquarii	+	3,54**	Madras Result re-examined.

	Madras.	Nautical Almanac 1834.	Society's Catalogue.
	<i>h. m. s.</i>	<i>s.</i>	<i>s.</i>
* Mean A. R.	9 12 35,75	38,19	33,88
† January 1, 1832.	14 28 16,25	18,58	22,60

† On the 27th of August 1834, I observed the interval between the transit of this Star and of No. 2078 with the Mural Circle Telescope,

	<i>m. s.</i>
To be.....	2 18,46
Interval in the Madras Catalogue.....	2 18,65
— — — Ast. Soc. Catalogue.....	2 19,65

** On the 16 h of October 1834, I observed the interval between the transit of this Star and of No. 2630 with the Mural Circle Telescope,

	<i>m. s.</i>
To be.....	4 17,60
Interval in the Madras Catalogue.....	4 17,30
— — — Ast. Soc. Catalogue.....	4 20,64

CXXXVIII REMARKS UPON THE CATALOGUE OF FIXED STARS.

No. 2625	ξ Cephei	+	1,49	Greenwich Observations for 1831. differ + 0,20s.
No. 2629	36 Aquarii ***	+	3,24	Piazzi's Catalogue confirms the Madras Result.
No. 2639	Aquarii	+	1,69	} Madras Results re-examined.
No. 2765	ι Gruis	+	3,61	
No. 2787	τ Pegasi	+	1,54	

We now come to the examination of the North Polar Distance column and the "difference from Greenwich." From the results of Pages 131, &c. we might naturally expect that (the computations being correctly performed) the result of one years observation if derived from 3 or 4 observations would never disagree to the amount of two seconds; whereas from the anomalies of Pages 122 and 123, a much larger amount of difference must be expected: selecting those cases where the largest difference of result occurs we have as follows.

No.	Obsers. vations.	Seconds of N. P. D.	Differ- ence.
		<i>s.</i>	<i>s.</i>
352	5	give 48,04	} 3,70
	1	44,34	
898	5	44,69	} 3,63
	5	48,32	
928	5	53,50	} 4,26
	5	49,24	
1015	5	35,39	} 3,40
	5	31,99	
1110	5	38,07	} 4,32
	5	33,75	
1223	5	32,76	} 3,74
	5	29,02	
1333	2	13,31	} 4,17
	4	9,14	
1360	5	28,74	} 3,49
	2	25,25	
1526	4	58,91	} 4,04
	1	54,87	
1700	3	31,71	} 5,16
	11	36,87	
1904	1	34,74	} 6,44
	3	41,18	
1981	2	26,66	} 4,02
	3	30,68	
2301	2	45,55	} 3,73
	6	49,28	
2433	3	58,08	} 3,46
	2	61,54	
2561	5	60,11	} 3,70
	2	56,41	

*** On the 16th of October 1834, I observed the interval between the transit of this Star and of No. 2630 with the Mural Circle Telescope.

	<i>m.</i>	<i>s.</i>
To be.....	1	0,20
Interval in the Madras Catalogue.....	1	0,08
— — — Ast. Soc. Catalogue.....	1	3,04

Neglecting the result No. 1904 from one observation, (for the Instrument might possibly have moved in the interval between the bisection and reading off) the largest discordance now met with is 5,16s. of which, more presently—from the remaining discordances (which it must be recollected are *extreme* ones,) it may be fairly inferred, *that the extreme error to which any result is liable does not often amount to two seconds of space and never exceeds two seconds and a half*—in comparing then two Catalogues such as the Madras and Greenwich, in which (in extreme cases) the errors may be expected to enter with opposite signs, it is proper to charge only *half* of the discordance as an error to either Catalogue; so that the extreme error of two seconds and a half just mentioned will occasionally give rise to a discordance of five seconds: in the column “difference from Greenwich” in which we can better learn the amount of these discordances

there are 177 cases in which the difference is less than 1s.

—	356	—	—	—	—	2s.
—	492	—	—	—	—	3s.
—	592	—	—	—	—	4s.
—	631	—	—	—	—	5s.
and	36	—	—	—	is greater	5s.

Which are as follows:

		s.				
No. 59	+	5,37	Greenwich Observations for 1831 differ	+	1,89s.	
No. 162	+	8,33	Do.	Do.	Do.	+ 3,14s.
No. 178	—	5,65	Do.	Do.	Do.	— 3,09s.
No. 217	—	11,09	Do.	Do.	Do.	— 2,63s.
No. 269	+	5,06	Must be examined.			
No. 280	+	7,81	Greenwich Observations for 1831 differ	—	4,03s.	
No. 595	—	5,16	Must be examined.			
No. 735	+	5,10*	Do.	Do.		
No. 757	+	5,54	Do.	Do.		
No. 791	+	5,57	Do.	Do.		
No. 877	—	6,57	Greenwich Observations for 1831 differ	—	4,30s.	
No. 1179	+	5,04	Must be examined.			
No. 1254	+	5,01	Do.	Do.		
No. 1289	+	5,66	Do.	Do.		
No. 1607	+	5,39	Do.	Do.		
No. 1619	+	5,81	Do.	Do.		
No. 1665	+	5,63	Do.	Do.		
No. 1700	+	7,86	Do.	Do.		
No. 1803	+	5,30	Do.	Do.		
No. 1806	+	5,63	Do.	Do.		
No. 1816	+	7,73	Do.	Do.		

* The Cambridge place of this Star is about 50s. in error.

REMARKS UPON THE CATALOGUE OF FIXED STARS.

		s.			
No. 1837	+	5,28	Must be examined.		
No. 1986	+	5,69	Do. Do.		
No. 2028	—	5,03	Do. Do.		
No. 2079	+	10,18	Greenwich Observations for 1831 differ	+	4,63s.
No. 2105	+	5,11	Must be examined.		
No. 2187	+	9,48	Greenwich Observations for 1831 differ	+	1,03s.
No. 2196	+	5,87	Do. Do. Do. Do.	+	3,32s.
No. 2198	+	5,57	Must be examined.		
No. 2371	—	5,44	Do. Do.		
No. 2561	+	6,89	Greenwich Observations for 1817 differ	—	8,17s.
No. 2562	—	5,13	Must be examined.		
No. 2661	+	16,33	Greenwich Observations for 1831 differ	—	2,63s.
No. 2696	+	10,15	Do. Do. for 1817 differ	+	1,35s.
No. 2710	+	6,39	Must be examined.		
No. 2754	+	5,21	Do. Do.		

In the above list there are six Stars which have been observed at the Cambridge Observatory see (Vol. VI for 1833) of which No. 162 differs from the Madras Catalogue 0,94s. and No. 2754 by 1,68s. ; of the other four cases, Nos. 1254, 1607 and 1619, agree to a fraction of a second with the Greenwich Catalogue; with regard to the remaining case (that of No. 1700 which likewise agrees with the Greenwich place to 1s.) it will be as well here to give the result of each observation made at Madras.

Mean N. P. D. of β Ursa Minoris reduced to January 1, 1832.

1832			1833		
	°	' "		°	' "
January 24	15	9 36,11	January 11	15	9 36,69
May 11	15	9 30,84	13	15	9 37,56
12	15	9 33,13	14	15	9 36,94
20	15	9 31,15	15	15	9 37,07
			16	15	9 36,73
			17	15	9 36,78
			18	15	6 37,17
			20	15	9 36,78
			21	15	9 37,37
			23	15	9 37,05
			25	15	9 36,40
			May 12	15	9 32,31
Mean.....		32,81		15	9 36,49
Or taking the general Mean we have.....				15	9 35,57*
Differing + 7,67s. from the Greenwich Observations.					
And + 6,69s. — Cambridge —					

* Differing a little from the result given at Page LXXVII in which I had rejected the observation of January 24, 1832 and of May 12, 1833.

On recomputing the observations of this Star my attention was arrested by noticing that the observations or rather the results of January 1832 agreed with those of January 1833, and that those of May 1832 agreed with those of the same month in 1833—could the change from the damp morning air of the N. E. Monsoon in January to the hot and dry winds from the S. W. and West in the month of May have any thing to do with it? were the corrections for Aberration, &c. correctly computed? these with several other possible sources of error have been very frequently and carefully examined without eliciting any cause to explain this very extraordinary disagreement—the observations of Polaris above and below the pole in January 1833 when applied to the determination of the Index Error agreed perfectly* with the Stars otherwise situated but, β Ursæ Minoris disagreed to the above amount:

With regard to the column “difference from A. S. C.” the same objections applies to the N. P. D. as to the A. R. it will consequently only be necessary here to note the larger discordances, which are as follows.

	<i>m.</i>	<i>s.</i>	
No. 21	+	2	1,51 Or an Error of 2 <i>m.</i>
No. 40	+	0	55,60 Do. Do. 1 <i>m.</i>
No. 79	+	0	40,79 Had the proper motion mentioned by Piazzi been allowed in constructing the A. S. C. (— 1,25 <i>s.</i>), the difference would have been + 0,79 <i>s.</i>
No. 124	—	4	49,84 A discordance of 5 <i>m.</i> which must be re-examined.
No. 268	—	0	45,96 Presumed amount of Proper Motion.
No. 337	—	0	25,00 Do. Do.
No. 368	—	0	28,32 Do. Do.
No. 439	+	5	28,12 Only one observation—possibly the wrong Star.
No. 483	—	0	20,20 Presumed amount of Proper Motion.
No. 610	+	0	41,63 Do. Do.
No. 630	+	0	20,73 Do. Do.
No. 671	+	0	25,29 Do. Do.
No. 1141	+	0	16,32 One observation gave 29 <i>m.</i> 8,60 <i>s.</i> there are probably two Stars.
No. 1247	—	0	27,53
No. 1414	+	0	8,35 Another Star has been observed twice which gives 66° 1 <i>m.</i> 31,75 <i>s.</i>
No. 1436	—	0	51,77 Presumed amount of Proper Motion.
No. 1655	+	0	22,21 Do. Do.
No. 1690	+	0	53,93 With reference to the difference in A. R. as well as in N. P. D. see Piazzi † our observations give about 1½ <i>s.</i> for P. M. in A. R. and about 2 <i>s.</i> in N. P. D.

* In selecting a Catalogue of Stars to be observed for the purpose of determining the Index Error I have rejected those which from the observations of 1831 differed to the amount of 2*s.* from the Madras Result.

† Piazzi says—Exnostris observationibus annorum 1800 — 2 — 8 — 9, A. R. et declinatio crescere videntur, & magis declinatio, cujus annua variatio foret 1,0*s.* circiter: idem proxime habetur ex Monierio, qui unus stellam hanc ante nos observavit. Eadem duplex, & ipse comes 0,7*s.* temporis præcedit parumper ad Anstrum.


	<i>m.</i>	<i>s.</i>	
No. 1752	+	0 58,36	This has been repeatedly re-observed in 1835.
No. 2120	+	0 34,53	Presumed amount of Proper Motion.
No. 2155	—	0 23,07	Do. Do.
No. 2158	—	0 25,78	Do. Do.
No. 2351	+	0 26,22	Do. Do.
No. 2663	+	2 2,64	Greenwich confirms the Madras Result—with reference to Piazzi remark upon this Star ‡ it would appear that we have each observed the N. P. D. of the small Star, but it will I think be found that Piazzi is in Error.

‡ Præcedit 26s. temporis, alia 9æ magnitud. 2m. circiter ad Austrum.



*Curve traced by a Dot situated .001 Inch from the Centre of the Ax.
of the Western Part of the Madras Transit Instrument.*

(see Page 6)



Collimation Telescope Supports. (see Page 125)

